

Penetration of the human nail plate: the effects of vehicle pH on the permeation of miconazole

KENNETH A. WALTERS*, GORDON L. FLYNN, JOHN R. MARVEL†, *University of Michigan, College of Pharmacy, Ann Arbor, Michigan 48109, †Dermatological Division, Ortho Pharmaceutical Corporation, Raritan, New Jersey, USA*

In order to assess the relative permeability of the nail plate to ionized and unionized drugs the permeation of miconazole at varying pH has been followed as a function of time. The pH was adjusted from 3.1 to 8.2 to obtain between 5 and 100% dissociation of the drug. No significant difference in the rate of permeation was demonstrated. The data suggests that the ionic form of miconazole dissolves as easily in the nail plate as the free base and, therefore, topical bioavailability can be enhanced by decreasing the formulation pH thereby increasing drug solubility.

Previous studies on the physicochemical properties of the human nail plate have indicated a marked difference between the permeability characteristics of this tissue and that of the epidermis (Walters et al 1983). Whereas the stratum corneum behaves as a lipid barrier to the permeation of low molecular weight chemicals (Scheuplein 1965; Durrheim et al 1980) the nail plate exhibits behaviour similar to that of a hydrogel of high ionic strength. Furthermore chemicals, such as dimethylsulphoxide, which have been shown to be remarkably effective as enhancers of skin penetration (Scheuplein & Ross 1970; Astley & Levine 1976) have shown little promise as accelerators of nail plate permeability (Kligman 1965; Walters & Flynn 1981). These differences between two tissues of such intimately related origins cannot be attributed to any obvious differences in morphogenesis; rather the inconsistencies appear to be due to differences in the relative amounts of the lipid and protein regimes (Walters et al 1983) and perhaps the physicochemical qualities of each of these phases. Stratum corneum contains at least 10% lipid, most of which is believed to be intercellular, whereas the nail plate contains no more than 1% (Baden et al 1973). The low lipid fraction of the nail is consistent in makeup and amount with lipids derived mostly from the residual cell membranes. In its normal state the nail also contains less water than the stratum corneum, about 10% moisture by actual estimate (Baden et al 1973). Permeation data for the homologous nonelectrolytes, the n-alkanols, indicate the relative importance of the various regimes of the nail plate in determining its barrier properties and show the nail to be a far different membrane than the stratum corneum. In this communi-

cation further evidence of unique behaviour of the nail plate membrane is presented. Specifically the permeation of a weak electrolyte, miconazole, does not appear to be influenced by its degree of ionization.

Method

Tritiated miconazole (1-[2,4-dichloro- β -(2,4-dichlorobenzyloxy)phenethyl]imidazole) was supplied by Ortho Pharmaceutical Corporation. The radiolabelled chemical was diluted with 0.9% NaCl before use. Details of the diffusion cell and permeation procedures have been given elsewhere (Walters et al 1981). The permeation of [^3H]miconazole through hydrated nail plate was followed as a function of pH of the bathing medium at 37 °C. The pH was adjusted from 3.1 to 8.2, with citrate/phosphate buffer, to obtain between 5 and 100% dissociation. During these experiments [^{14}C]ethanol (New England Nuclear) was the internal reference.




FIG. 1. pH-permeation profile for miconazole and ethanol data are mean \pm s.d. ($n = 5$). Figures beneath the miconazole curve are % dissociations.

Results and discussion

The effect of pH on the permeability coefficients of miconazole and ethanol are shown in Fig. 1. It is expected that the permeability of a weak electrolyte should vary as a function of pH, providing the membrane behaves as a lipoidal structure. It has been demonstrated, for example, that organic ions can permeate the stratum corneum, but the rate of permeation is only a fraction of that of the undissociated form

* Correspondence and present address: Fisons plc, Pharmaceutical Division, Bakewell Road, Loughborough, UK.

(Arita et al 1970; Wallace & Barnett 1978; Dyer et al 1981). The low permeation rates of dissociated compounds through skin are assumed to be due to a relative inability of ions to partition into the lipid phases of the horny layer. The lack of importance of lipid phases as a medium of transport across the nail, as suggested previously (Walters et al 1983), is reflected in the ability of both the dissociated and the undissociated species of miconazole to permeate at near equivalent rates (Fig. 1). Miconazole is a weak base with a pK_a of 6.65 and, therefore, the more acidic the medium the greater is the degree of ionization. Yet the flux of miconazole, through different nail plates, is invariant at low pH where ionization is near complete. Moreover, the permeability coefficients of the reference compound, ethanol, follow the same pattern as a function of pH. The ratio, $P_{\text{miconazole}}/P_{\text{ethanol}}$ is essentially invariant. Thus the ionic form of miconazole dissolves as easily in the nail plate as the free base. Since there is little or no dependency of permeability on pH, these data suggest that the overriding aspect in increasing topical bio-availability of miconazole, for the treatment of onychomycoses, is increasing the solubility of the drug in a formulation, which can be done by decreasing the pH.

This work was supported through the generosity of the Ortho Pharmaceutical Corporation, Raritan, New Jersey, USA.

REFERENCES

- Arita, T., Hori, R., Anmo, T., Washitake, M., Akatsu, M., Yajima, T. (1970) *Chem. Pharm. Bull.* 18: 1045-1049
- Astley, J. P., Levine, H. (1976) *J. Pharm. Sci.* 65: 210-215
- Baden, H. P., Goldsmith, L. A., Fleming, B. (1973) *Biochim. Biophys. Acta* 322: 269-278
- Durrheim, H., Flynn, G. L., Higuchi, W. I. Behl, C. R. (1980) *J. Pharm. Sci.* 69: 781-786
- Dyer, A., Hayes, G. G., Wilson, J. G., Catterall, R. (1981) *Int. J. Cosmet. Sci.* 3: 271-278
- Kligman, A. M. (1965) *J. Am. Med. Ass.* 193: 796-804
- Scheuplein, R. J. (1965) *J. Invest. Dermatol.* 45: 334-345
- Scheuplein, R. J., Ross, L. (1970) *J. Soc. Cosmet. Chem.* 21: 853-873
- Wallace, S. M., Barnett, G. (1978) *J. Pharmacokinet. Biopharm.* 6: 315-325
- Walters, K. A., Flynn, G. L. (1981) *J. Pharm. Pharmacol.* 33: 6P
- Walters, K. A., Flynn, G. L., Marvel, J. R. (1981) *J. Invest. Dermatol.* 76: 76-79
- Walters, K. A., Flynn, G. L., Marvel, J. R. (1983) *J. Pharm. Pharmacol.* 35: 28-33

J. Pharm. Pharmacol. 1985, 37: 499-501
Communicated January 7, 1985

© 1985 J. Pharm. Pharmacol.

The affinity and efficacy of naturally occurring catecholamines at β -adrenoceptor subtypes

GRANT A. MCPHERSON*, PETER MOLENAAR, ERROL MALTA, *School of Pharmacology, Victorian College of Pharmacy, 381 Royal Parade, Parkville 3052, Victoria, Australia*

The contribution of affinity and efficacy to the agonistic actions of the naturally occurring catecholamines, (-)-noradrenaline and (-)-adrenaline at β -adrenoceptor sites were assessed in guinea-pig driven left atrial (β_1) and K^+ -depolarized uterine (β_2) preparations. The dissociation constants of each agonist, required in these calculations, were calculated using radioligand binding techniques. [^{125}I]iodocyanopindolol bound to sites in membrane preparations of each tissue which have been shown to represent β_1 - (atria) and β_2 - (uterus) adrenoceptors. It was found that (-)-noradrenaline was approximately 10-fold more selective for the β_1 - as opposed to the β_2 -adrenoceptor in the pharmacological studies. Affinity/efficacy calculations indicated that this selectivity was entirely due to a selective affinity for the β_1 -adrenoceptor subtype. (-)-Noradrenaline, (-)-adrenaline and the reference compound (-)-isoprenaline all had approximately the same efficacy at either β -adrenoceptor subtype.

As noted by Lands et al (1967) in their work on the subclassification of β -adrenoceptors, the naturally occurring catecholamines, (-)-noradrenaline and (-)-adrenaline display a marked difference in their potency at so called β_1 - and β_2 -adrenoceptors. Thus at β_1 -

adrenoceptors (e.g. heart) (-)-noradrenaline has a similar potency to (-)-adrenaline whereas at β_2 -adrenoceptors (e.g. blood vessels) (-)-adrenaline is considerably more potent than (-)-noradrenaline as an agonist.

The purpose of this study was to ascertain whether the differing relative potencies of (-)-noradrenaline and (-)-adrenaline result from differences in affinity (i.e. different abilities of the agonists to form drug receptor complexes) or differing efficacies (i.e. different abilities of the drug receptor complex to initiate pharmacological activity). These studies were performed using the non-selective β -adrenoceptor agonist (-)-isoprenaline as the reference agonist.

Methods

Details of the methods used in the present study have been described previously (McPherson et al 1984). Guinea-pig left atrial and uterine tissues were used since previous pharmacological (Broadley 1982; Krstew et al 1982; McPherson et al 1984) and radioligand binding studies (McPherson et al 1984) have indicated the presence of homogeneous populations of β_1 - and

* Correspondence.