

## Antimalarial compounds isolated from plants used in traditional medicine

Joanne Bero<sup>a</sup>, Michel Frédérick<sup>b</sup> and Joëlle Quetin-Leclercq<sup>a</sup>

<sup>a</sup>Université catholique de Louvain, Louvain Drug Research Institute, Analytical Chemistry, Drug Analysis and Pharmacognosy Unit, Brussels and <sup>b</sup>University of Liège, Natural and Synthetic Drugs Research Center, Laboratory of Pharmacognosy, Liège, Belgium

### Abstract

**Objectives** This review covers the compounds with antiplasmodial activity isolated from plants published from 2005 to the end of 2008, organized according to their phytochemical classes. Details are given for substances with IC<sub>50</sub> values  $\leq 11 \mu\text{M}$ .

**Key findings** Malaria is a major parasitic disease in many tropical and subtropical regions and is responsible for more than 1 million deaths each year in Africa. The rapid spread of resistance encourages the search for new active compounds. Nature and particularly plants used in traditional medicine are a potential source of new antimalarial drugs as they contain molecules with a great variety of structures and pharmacological activities.

**Summary** A large number of antimalarial compounds with a wide variety of structures have been isolated from plants and can play a role in the development of new antimalarial drugs. Ethnopharmacological approaches appear to be a promising way to find plant metabolites that could be used as templates for designing new derivatives with improved properties.

**Keywords** antiplasmodial; malaria; plant compounds; *Plasmodium falciparum*; traditional medicine

### Introduction

Malaria is a parasitic disease caused by *Plasmodium* species transmitted from the blood of an infected person and passed to a healthy human by a female *Anopheles* mosquito. There are four types of human malaria and *Plasmodium falciparum* is responsible for the most severe cases, and so most studies have evaluated the activity of compounds on this species. Malaria affects 350–500 million people per year worldwide and is responsible for 1.1 million deaths per year. In many parts of the world the parasites have developed resistance to a number of antimalarials such as chloroquine and derivatives, the most widely used treatment for malaria, and so there is an urgent need to discover new compounds with an original mode of action. Plants commonly used in traditional medicine are a source of active new compounds. For example, artemisinin isolated from *Artemisia annua* and used in China to treat malaria is a sesquiterpene lactone prescribed in combination therapies to fight chloroquine-resistant *P. falciparum*. In this review, all new active metabolites isolated from plants used in traditional medicine to treat malaria are described and organised according to their phytochemical classes. All the activities described were determined *in vitro* on *P. falciparum* strains, unless otherwise specified, and bio-guided fractionation was also based on this antimalarial test. Activities were assessed on different strains, among which are chloroquine sensitive (NF54, NF54/64, 3D7, D6, F32, D10, HB3, FCC1-HN, Ghana), chloroquine resistant (FcB1, W2, FCM29, BH26/86, Dd2, EN36, ENT30, FCR3, FCR-3/A2) and/or multidrug resistant (K1, TM91C235) strains, to find effective compounds against resistant malaria. We considered that those having an IC<sub>50</sub>  $\leq 11 \mu\text{M}$  may have some interest for further development, while those with lower activity were less interesting. We only give structures for promising compounds, the others are cited in tables. As reviews already exist for compounds published before 2005,<sup>[1–7]</sup> we focused on those published from 2005 to the end of 2008. Some examples of recent natural antiplasmodial compounds are also cited in Mambu and Grellier<sup>[8]</sup> and, more recently, in Kaur *et al.*<sup>[9]</sup>

**Correspondence:** J. Bero,  
Catholic University of Louvain,  
Analytical Chemistry, Drug  
Analysis and Pharmacognosy  
Unit, Avenue E. Mounier,  
72, B-1200 Brussels, Belgium.  
E-mail: joanne.bero@uclouvain.be

## Phenolic derivatives

### Flavonoid derivatives

Bioassay-guided fractionation of the ethyl acetate extract of the leaves of *Piptadenia pervillei* Vatke (Leguminosae) led to the isolation of two phenolic compounds: (+)-catechin 5-gallate (**1**) and (+)-catechin 3-gallate (**2**). Compounds **1** and **2** displayed high activity, with IC<sub>50</sub> values of 1.2  $\mu\text{M}$  and 1.0  $\mu\text{M}$  (FcB1), respectively, and no significant cytotoxicity.<sup>[10]</sup> New secondary metabolites were isolated from the root extract of *Bauhinia purpurea* L. (Leguminosae). Among the isolated metabolites, a flavanone exhibited moderate antimalarial activity, demethoxymatteucinol (**3**) (IC<sub>50</sub> = 9.5  $\mu\text{M}$  against K1).<sup>[11]</sup> Phytochemical investigation of the stem bark and root bark of *Friesodielsia obovata* (Benth.) Verdc. (Annonaceae) also afforded demethoxymatteucinol (**3**), which possessed weak antiplasmodial activity, with IC<sub>50</sub> = 34.1 and 29.9  $\mu\text{M}$  against K1 and NF54, respectively.<sup>[12]</sup> Investigation of the chemical constituents of the root bark of *Artocarpus rigidus* Blume subsp. *rigidus* (Moraceae) led to the isolation of two known flavonoids, artonin F (**4**) and cycloartobiloxanthone (**5**), which exhibited antiplasmodial activity against K1 (4.8  $\mu\text{M}$  and 8.5  $\mu\text{M}$ , respectively).<sup>[13]</sup> Two new prenylated flavones, artocarpones A (**6**) and B (**7**) (IC<sub>50</sub> = 0.12 and 0.18  $\mu\text{M}$ , respectively), and seven known prenylated flavonoids, including artonin A (**8**) (IC<sub>50</sub> = 0.55  $\mu\text{M}$ ), cycloheterophyllin (**9**) (IC<sub>50</sub> = 0.02  $\mu\text{M}$ ), artoindonesianin R (**10**) (IC<sub>50</sub> = 0.66  $\mu\text{M}$ ), heterophyllin (**11**) (IC<sub>50</sub> = 1.04  $\mu\text{M}$ ), heteroflavanone C (**12**) and artoindonesianin A-2 (**13**) (IC<sub>50</sub> = 1.31  $\mu\text{M}$ ) were isolated from the stem bark of *Artocarpus champeden* Spreng. (Moraceae). The isolated compounds were tested for their inhibitory activity against 3D7. All possessed interesting activity with inhibitory concentrations from 0.001 to 1.31  $\mu\text{M}$ . Compound **12** was the most potent with an IC<sub>50</sub> of 1 nm. The inhibitory activity of these flavonoid derivatives supports the traditional use of the dried stem bark of *A. champeden* as an antimalarial drug.<sup>[14]</sup> Antitubercular and antimalarial activity-guided study of the dichloromethane extract of the roots of *Artocarpus altilis* (Parkinson) Fosberg (Moraceae) led to the isolation of nine prenylated flavones, including cycloartocarpin (**14**), artocarpin (**15**), chaplashin (**16**), morusin (**17**), cudraflavone B (**18**), artonin E (**19**) and artobiloxanthone (**20**). All compounds exhibited antiplasmodial activity against K1 with IC<sub>50</sub> values of 9.9, 6.9, 7.7, 4.5, 5.2, 6.4 and 6.9  $\mu\text{M}$ , respectively.<sup>[15]</sup> Ethyl acetate extract from the stem bark of *Erythrina fusca* Lour. (Leguminosae) showed antimalarial activity against K1, and lonchocarpol A (**21**) isolated from that extract showed notable antimalarial activity (IC<sub>50</sub> = 3.9  $\mu\text{M}$ ). However, two others flavonoids isolated from the same sample did not show any activity, even though these compounds possessed prenylated substitution.<sup>[16]</sup> Isoflavonoids and flavonoids were isolated from the root bark and the stem bark of *Erythrina saculeuxii* Hua (Leguminosae). The two most active against D6 and W2 were 5'-prenylpratensein (**22**, IC<sub>50</sub> on D6 = 6.3  $\mu\text{M}$  and IC<sub>50</sub> on W2 = 8.7  $\mu\text{M}$ ) and shinpterocarpin (**23**, IC<sub>50</sub> on D6 = 6.6  $\mu\text{M}$  and IC<sub>50</sub> on W2 = 8.3  $\mu\text{M}$ ).<sup>[17]</sup> Phytochemical investigation of the hexane and CH<sub>2</sub>Cl<sub>2</sub> extracts of *Erythrina*

*stricta* Roxb. (Leguminosae) roots and *Erythrina subumbrans* Merr. (Leguminosae) stems led to the isolation of two pterocarpan, erybraedin A (**24**) and erystagallin A (**25**), and one flavanone, 5-hydroxysophoranone (**26**). All of them exhibited reasonable antiplasmodial activity against K1 with IC<sub>50</sub> = 8.7, 9.0 and 5.3  $\mu\text{M}$ , respectively.<sup>[18]</sup> Vogelin C (**27**) and lespedezaflavanone B (**28**) were isolated from the bark of *Erythrina subumbrans* Merr. (Leguminosae) and possessed antiplasmodial activity against K1 with IC<sub>50</sub> values of 6.6 and 9.1  $\mu\text{M}$ , respectively.<sup>[19]</sup> A known compound, 6-prenylapigenin (**29**), was isolated from *Cannabis sativa* L. (Cannabaceae) and displayed notable antimalarial activity against D6 and W2 with IC<sub>50</sub> values of 6.7 and 4.8  $\mu\text{M}$ , respectively.<sup>[20]</sup> The 80% ethanol extract from the outer bark of *Ochna integerrima* Lour. (Merr.) (Ochnaceae) led to isolation of a biflavanone (**30**) that had not been found previously from a natural plant source and is a potent antimalarial ingredient against K1 (IC<sub>50</sub> = 157 nm). The stereoisomer of **30** (**31**) was also isolated from this plant but its activity was significantly lower than that of **30** (IC<sub>50</sub> = 10.2  $\mu\text{M}$ ).<sup>[21]</sup> The antiplasmodial activity of five natural biflavonoids was estimated on K1. Lanaroflavone (**32**) isolated from the aerial parts of *Camposperma panamensis* Standl. (Anacardiaceae) showed the highest antiplasmodial activity (IC<sub>50</sub> = 0.48  $\mu\text{M}$ ) and exhibited a high selectivity index value (SI = 159), indicating selective antiplasmodial activity. Ginkgetin (**33**), isoginkgetin (**34**), bilobetin (**35**) and sciadopitysin (**36**) isolated from the leaves of *Ginkgo biloba* L. (Ginkgoaceae) showed antiplasmodial activity (IC<sub>50</sub> = 2.0, 3.5, 6.7 and 1.4  $\mu\text{M}$  with SI = 4.1, 3.2, 4.0 and 49, respectively).<sup>[22,23]</sup> A new biflavanone, *ent*-naringeninyl-(I-3 $\alpha$ ,II-8)-4'-*O*-methylnaringenin (**37**) was isolated from the root bark of *Garcinia livingstonei* T. Anderson (Clusiaceae) collected in Tanzania. This compound showed reasonable activity against Ghana strain (IC<sub>50</sub> = 6.7  $\mu\text{M}$ ).<sup>[24]</sup> Phytochemical re-examination of the aerial exudates of *Polygonum senegalense* Meisn. (Polygonaceae) forma *senegalense* resulted in the isolation of two chalcones (**38** and **39**) active, respectively, with an IC<sub>50</sub> of 3.1  $\mu\text{M}$  on D6 and 2.4  $\mu\text{M}$  on W2, and 14.0  $\mu\text{M}$  on D6 and 9.5  $\mu\text{M}$  on W2.<sup>[25]</sup> The bioassay-guided purification of an *n*-hexane extract from the leaves of *Piper hostmannianum* C.DC. var. *berbicense* (Piperaceae) led to the isolation of four monoterpenes or prenyl-substituted dihydrochalcones as well as known compounds. (–)-Methylinderatin (**40**) and linderatone (**41**) exhibited moderate antiplasmodial activity with IC<sub>50</sub> values of 5.6 and 5.3  $\mu\text{M}$  (**40**) and 10.3 and 15.1  $\mu\text{M}$  (**41**), respectively, against F32 and FcB1. The activity of **40** was confirmed *in vivo* against *Plasmodium vinckei petteri* in mice (80% reduction of parasitemia) at a dose of 20 mg/kg per day intraperitoneally.<sup>[26]</sup> A prenylated chalcone, medicagenin (**42**), was isolated from *Crotalaria medicagenia* Lam. (Leguminosae). Antimalarial activity was evaluated against NF-54 and medicagenin exhibited 100% inhibition of schizont maturation at a concentration of 2  $\mu\text{g/ml}$ .<sup>[27]</sup> A prenylated chalcone, bartericin A (**43**), and three known natural products, stipulin (**44**), 4-hydroxy lonchocarpin (**45**) and kanzonol B (**46**) were isolated from the twigs of *Dorstenia barteri* var. *subtriangularis* (Engl.) Hijman & C.C. Berg (Moraceae). These compounds were evaluated against

W2 and found to be moderately active (IC<sub>50</sub> = 2.2, 5.1, 3.4 and 9.6  $\mu\text{M}$ , respectively).<sup>[28]</sup> Bioassay-directed fractionation of the EtOAc extract of the stem bark of *Hintonia latiflora* (Sessé & Moc. ex DC.) Bullock (Rubiaceae), using the in-vitro 16-h and the in-vivo 4-day suppression tests on *Plasmodium berghei* schizont numbers, led to the isolation of the new 5-*O*- $\beta$ -D-glucopyranosyl-7,4'-dimethoxy-3'-hydroxy-4-phenylcoumarin, along with the known 5-*O*- $\beta$ -D-glucopyranosyl-7-methoxy-3',4'-dihydroxy-4-phenylcoumarin. Both compounds suppressed the development of *P. berghei* schizonts with IC<sub>50</sub> values of 24.7 and 25.9  $\mu\text{M}$ , respectively, and the latter compound suppressed the development of schizonts by 70.8% at an oral dose of 40 mg/kg in the in-vivo assay.<sup>[29]</sup>

Figure 1 shows the flavonoid derivatives with moderate or promising activity *in vitro* against various strains of *P. falciparum*.

### Xanthones

The methanol extract of the stem bark of *Allanblackia monticola* Mildbr. (Clusiaceae) resulted in the isolation of a new prenylated xanthenedione, designated as allanxanthone C (47), together with the known compounds norcowanin (48), mangostin (49) and tovophyllin A (50). Compounds were assayed for their antiplasmodial activity and for their cytotoxicity. Three of these compounds (47–49) were found to be active against *Plasmodium*: 47, IC<sub>50</sub> on FcM29 = 1.3  $\mu\text{M}$  and IC<sub>50</sub> on F32 = 6.9  $\mu\text{M}$ ; 48, not tested on FcM29 and IC<sub>50</sub> on F32 = 6.3  $\mu\text{M}$ ; 49, IC<sub>50</sub> on FcM29 = 4.1  $\mu\text{M}$  and IC<sub>50</sub> on F32 = 7.8  $\mu\text{M}$ , and also showed weak cytotoxicity against human melanoma A375 cells.<sup>[30]</sup> Tovophyllin A (50) was the most interesting with promising antimalarial activity (IC<sub>50</sub> on FcM29 = 0.7  $\mu\text{M}$  and IC<sub>50</sub> on F32 = 20.3  $\mu\text{M}$ ) and relatively low cytotoxicity.<sup>[31]</sup> A new prenylated xanthone, 5-*O*-methylcelebixanthone (51), together with a known compound, cochinchinone C (52), were isolated from roots of *Cratoxylum cochinchinense* Blume (Clusiaceae). Compounds 51 and 52 exhibited antimalarial activity against K1 with IC<sub>50</sub> values of 8.9 and 6.3  $\mu\text{M}$ , respectively. IC<sub>50</sub> values for cytotoxicity were within the range of 5.6  $\mu\text{M}$  for 52. No cytotoxicity was observed with 51.<sup>[32]</sup> A xanthone derivative, gaboxanthone (53), was isolated from the seed shells of *Symphonia globulifera* L.f. (Clusiaceae), together with known compounds, symphonin (54) and globuliferin (55). The antiplasmodial activity of the phenolic compounds was evaluated against W2. Compounds 53–55 gave IC<sub>50</sub> values of 3.5, 1.3 and 3.9  $\mu\text{M}$ , respectively.<sup>[33]</sup> The whole plant of *Swertia alata* Royle ex D. Don (Gentianaceae) was investigated and three xanthonenes, swertiaperennine, swertianin and decussatin, were isolated and tested for antimalarial activity. The results indicated that all xanthonenes possessed superior IC<sub>50</sub> values at 50  $\mu\text{M}$ . However, swertiaperennine was tested *in vivo* in the *P. berghei* test model and reduced parasitemia by 17.60% at a dose of 10 mg/kg.<sup>[34]</sup> A new compound, garciniaxanthone (56), was isolated from the roots of *Garcinia polyantha* Oliv. (Clusiaceae), in addition to three known compounds, smeathxanthone A (57), smeathxanthone B (58) and chefouxanthone (59). They exhibited antimalarial activity against NF54 with IC<sub>50</sub> values ranging from 2.5 to 4.1  $\mu\text{M}$ .<sup>[35]</sup>

Figure 2 shows xanthonenes with moderate or promising activity *in vitro* against various strains of *P. falciparum*.

### Stilbenes

A new stilbene glycoside, piceid-(1 $\rightarrow$ 6)- $\beta$ -D-glucopyranoside (60), was isolated from the MeOH extract of the leaves of *Parthenocissus tricuspidata* Planch. (Vitaceae) together with three known compounds, piceid (IC<sub>50</sub> = 13.2  $\mu\text{M}$ ), longistylin A (IC<sub>50</sub> = 34.3  $\mu\text{M}$ ) and longistylin C (IC<sub>50</sub> = 19.2  $\mu\text{M}$ ). The antiplasmodial activity of isolated compounds was determined *in vitro* against D10. Among the compounds isolated, 60 was the best inhibitor with an IC<sub>50</sub> value of 5.3  $\mu\text{M}$ .<sup>[36]</sup> Compound 60 was tested *in vivo* against *P. berghei* in mice intraperitoneally and exhibited significant blood schizontocidal activity in 4-day early infection, in preventive and curative treatment, with chemosuppression of 59 and 44% at 5 mg/kg per day, respectively, and an LD<sub>50</sub> > 500 mg/kg.<sup>[37]</sup> A stilbene glycoside was isolated from an *n*-butanol-soluble fraction of the root of *Pleur-opterus ciliinervis* Nakai (Polygonaceae). The compound was identified as (*E*)-resveratrol-3-*O*- $\alpha$ -L-rhamnopyranosyl-(1-2)- $\beta$ -D-xylopyranoside (61). It showed only moderate cytotoxicity and antimalarial activity against D10 with an IC<sub>50</sub> of 3.9  $\mu\text{M}$ .<sup>[38]</sup> Compound 61 was also found to have moderate antimalarial activity *in vivo* when tested against *P. berghei* in mice intraperitoneally. It possessed useful blood schizontocidal effects when used at doses that cause no marked toxicity in mice.<sup>[39]</sup>

Figure 3 shows stilbenes with moderate or promising activity *in vitro* against various strains of *P. falciparum*.


### Coumarins

Biologically guided fractionation of the methanolic extract of the roots of *Zanthoxylum flavum* Vahl. (Rutaceae) led to the isolation of isoimperatorin (62) which displayed IC<sub>50</sub> values of 5.5 and 2.7  $\mu\text{M}$  against D6 and W2, respectively.<sup>[40]</sup> A new coumarinolignan was isolated from a sample of *Grewia bilamellata* Gagnep. (Tiliaceae), grewin (63), which displayed antimalarial activity against D6 and W2 (IC<sub>50</sub> 11.2  $\mu\text{M}$  and 5.5  $\mu\text{M}$ , respectively) without significant cytotoxicity.<sup>[41]</sup> The compound 1-*O*-galloyl-6-*O*-luteoyl- $\alpha$ -D-glucose (64) with an IC<sub>50</sub> value of 2.21  $\mu\text{M}$  (FCR3) was isolated from the boiled aqueous extract of the whole plant of *Phyllanthus niruri* L. (Euphorbiaceae).<sup>[42]</sup>

Figure 4 shows coumarins with moderate or promising activity *in vitro* against various strains of *P. falciparum*.

### Lignans

From the hexane extract of *Holostylis reniformis* Duch. (Aristolochiaceae), five lignans were isolated: (7'*R*,8*S*,8'*R*)-4,5-dimethoxy-3',4'-methylenedioxy-2,7'-cyclo lignan-7-one (65) (IC<sub>50</sub> = 0.26  $\mu\text{M}$ ), (7'*R*,8*S*,8'*R*)-3',4,4',5-tetramethoxy-2,7'-cyclo lignan-7-one (66) (IC<sub>50</sub> = 0.32  $\mu\text{M}$ ), (7'*R*,8*R*,8'*S*)-3',4,4',5-tetramethoxy-2,7'-cyclo lignan-7-one (67) (IC<sub>50</sub> = 0.20  $\mu\text{M}$ ), (7'*R*,8*S*,8'*S*)-3',4,4',5-tetramethoxy-2,7'-cyclo lignan-7-one (68) (IC<sub>50</sub> = 0.63  $\mu\text{M}$ ) and (7'*R*,8*S*,8'*S*)-3',4'-dimethoxy-4,5-methylenedioxy-2,7'-cyclo lignan-7-one (69) (IC<sub>50</sub> = 8.00  $\mu\text{M}$ ). Most compounds possessed high antiplasmodial activity against BH26/86 and low toxicity on hepatic cells. Therefore, these compounds are potential candidates for the development of antimalarial drugs.<sup>[43]</sup> Seven tetrahydrofuran lignans isolated from *Nectandra megapotamica* Mez (Lauraceae) were evaluated for their antimalarial activity.



**Figure 1** Flavonoid derivatives with moderate or promising activity *in vitro* against various strains of *P. falciparum*






Figure 1 (Continued)



**Figure 2** Xanthones with moderate or promising activity *in vitro* against various strains of *P. falciparum*




**Figure 3** Stilbenes with moderate or promising activity *in vitro* against various strains of *P. falciparum*

Among the evaluated compounds, calopectin (**70**) displayed moderate activity, with IC<sub>50</sub> values of 10.7  $\mu\text{M}$  (D6 clone) and 11.0  $\mu\text{M}$  (W2 clone), and no cytotoxicity.<sup>[44]</sup> Bioassay-directed fractionation of the antimalarial active CHCl<sub>3</sub> extract of the dried stems of *Rourea minor* (Gaertn.) Aubl. (Connaraceae) liana led to the isolation of rourinoside (**71**). This lignan showed activity with an IC<sub>50</sub> value of 3.7  $\mu\text{M}$  against D6 and 2.1  $\mu\text{M}$  against W2 strains.<sup>[45]</sup>


Figure 5 shows lignans with moderate or promising activity *in vitro* against various strains of *P. falciparum*.

#### Tannins

Partitioning of extracts of *Punica granatum* L. (Lythraceae) led to the isolation of ellagic acid, gallic acid (**72**), punicalin and punicalagin (**73**). Gallic acid and punicalagin exhibited moderate antiplasmodial activity against D6 (IC<sub>50</sub> of 10.9 and 10.6  $\mu\text{M}$ ) and W2 clones (IC<sub>50</sub> of 7.5 and 8.8  $\mu\text{M}$ ).<sup>[46]</sup>



**Figure 4** Coumarins with moderate or promising activity *in vitro* against various strains of *P. falciparum*



**Figure 5** Lignans with moderate or promising activity *in vitro* against various strains of *P. falciparum*

Figure 6 shows tannins with moderate or promising activity *in vitro* against various strains of *P. falciparum*.

### Other phenolic derivatives

The petroleum ether extract of *Viola websteri* Hemsl. (Violaceae) was investigated and the main antiplasmodial compound was 6-(8'*Z*-pentadecenyl)-salicylic acid (**74**) with an IC<sub>50</sub> of 10.1  $\mu\text{M}$  (D10).<sup>[47]</sup> Dictyochromenol (**75**) and a known compound, 2'*E*,6'*E* 2-farnesyl hydroquinone (**76**) obtained from the petroleum ether extract of the whole plant of *Piper tricuspe* C.DC. (Piperaceae) showed antimalarial activity against FcB1 with IC<sub>50</sub> values of 9.58 and 1.37  $\mu\text{M}$  while the selectivity index suggests their high toxicity.<sup>[48]</sup> 4-Nerolidylcatechol (**77**), isolated from the roots of *Pothomorphe peltata* (L.) Miq. (Piperaceae) presented significant inhibition (more active than quinine and chloroquine) against

K1 (IC<sub>50</sub> = 0.67  $\mu\text{M}$ ).<sup>[49]</sup> A new bischromone, chrobisiamone A (**78**), was isolated from the leaves of *Cassia siamea* Lam. (Leguminosae). Compound **78** displayed antiplasmodial activity against 3D7 with an IC<sub>50</sub> of 5.6  $\mu\text{M}$ .<sup>[50]</sup> A new cannabichromanone A derivative was isolated along with the known cannabichromanone C (**79**) from *Cannabis sativa* L. (Cannabaceae). Cannabichromanone A showed mild antimalarial activity against D6 and W2 clones with IC<sub>50</sub> values of 11.1 and 11.4  $\mu\text{M}$ , respectively, while cannabichromanone C had IC<sub>50</sub> values of 13.1 and 9.4  $\mu\text{M}$ , respectively.<sup>[51]</sup> Guttiferone A (**80**) was isolated from the seed shells of *Symphonia globulifera* L.f. (Clusiaceae). The antiplasmodial activity of compound **80** was evaluated against W2 and gave an IC<sub>50</sub> value of 3.2  $\mu\text{M}$ .<sup>[33]</sup> Isoxanthochymol (**81**) was isolated from the roots of *Garcinia polyantha* Oliv. (Clusiaceae) and exhibited antimalarial activity against NF54 with an IC<sub>50</sub> of 2.2  $\mu\text{M}$ .<sup>[35]</sup>

Figure 7 shows other phenolic derivatives with moderate or promising activity *in vitro* against various strains of *P. falciparum*.

Table 1 gives the tested phenolic derivatives presenting low or no activity *in vitro* against various strains of *P. falciparum*.<sup>[12,13,16,17,25,26,28,29,32,34,37,40–42,47,48,50–66]</sup>




Figure 6 Tannins with moderate or promising activity *in vitro* against various strains of *P. falciparum*




Figure 7 Other phenolic derivatives with moderate or promising activity *in vitro* against various strains of *P. falciparum*



**Table 1** Phenolic derivatives presenting low or no activity *in vitro* against various strains of *P. falciparum*

Compound	Plant	Family	IC50 ( $\mu\text{M}$ )	Reference no.
Lawinal	<i>Friesodielsia obovata</i> (Benth.) Verdc.	Annonaceae	27.5 (K1) 119.1 (NF54)	12
2',4'-Dihydroxy-3'-(2-hydroxybenzyl)-6'-methoxychalcone	<i>Ellipeiopsis cherrevensis</i> (Pierre ex Finet & Gagnep.) R.E.Fr.	Annonaceae	18.9 (K1)	52
Syringic acid	<i>Commiphora opobalsamum</i> Engl.	Burseraceae	17.7 (D6) 16.2 (W2)	53
Cannabichromanone A	<i>Cannabis sativa</i> L.	Cannabaceae	11.1 (D6) 11.4 (W2)	51
Celebixanthone	<i>Cratoxylum cochinchinense</i> Blume	Clusiaceae	14.3 (K1)	32
$\beta$ -Mangostin	<i>Cratoxylum cochinchinense</i> Blume	Clusiaceae	16.9 (K1)	32
1,5-Dihydroxy-3-methoxy-4-isoprenylxanthone	<i>Chrysochlamys tenuis</i> Hammel	Clusiaceae	31 (W2)	54
1,3,7-Trihydroxy-2,4-diisoprenylxanthone	<i>Chrysochlamys tenuis</i> Hammel	Clusiaceae	20 (W2)	54
Toxyloxanthone A	<i>Chrysochlamys tenuis</i> Hammel	Clusiaceae	16 (W2)	54
Combretastatin D-3	<i>Getonia floribunda</i> Roxb.	Combretaceae	–	55
Combretastatin D-4	<i>Getonia floribunda</i> Roxb.	Combretaceae	–	55
3,5-Di- <i>O</i> -galloylquinic acid	<i>Sloanea rhodantha</i> (Baker) Capuron var. <i>rhodantha</i>	Elaeocarpaceae	31.7 (HB3) 23.6 (FcM29)	56
1,6-Di- <i>O</i> -galloyl glucopyranoside	<i>Sloanea rhodantha</i> (Baker) Capuron var. <i>rhodantha</i>	Elaeocarpaceae	35.5 (HB3) 15.7 (FcM29)	56
3,4,5-Tri- <i>O</i> -galloylquinic acid	<i>Sloanea rhodantha</i> (Baker) Capuron var. <i>rhodantha</i>	Elaeocarpaceae	35.3 (HB3) 23.1 (FcM29)	56
1,2,3,6-Tetra- <i>O</i> -galloyl glucopyranoside	<i>Sloanea rhodantha</i> (Baker) Capuron var. <i>rhodantha</i>	Elaeocarpaceae	20.2 (HB3) 20.6 (FcM29)	56
3- <i>O</i> - $\beta$ -D-Glucopyranosyl-(2 $\rightarrow$ 1)- <i>O</i> - $\beta$ -D-xylopyranoside	<i>Phyllanthus niruri</i> L.	Euphorbiaceae	18.5 (FCR3)	42
$\beta$ -Glucogallin	<i>Phyllanthus niruri</i> L.	Euphorbiaceae	14.6 (FCR3)	42
( <i>E</i> )-3-(4-Methoxy-phenyl)-2-phenyl-acrylic acid	<i>Croton lobatus</i> L.	Euphorbiaceae	19.1 (K1)	57
Swertiaperennine	<i>Swertia alata</i> Royle ex <i>D.Don</i>	Gentianaceae	>50	34
Swertianin	<i>Swertia alata</i> Royle ex <i>D.Don</i>	Gentianaceae	>50	34
Decussatin	<i>Swertia alata</i> Royle ex <i>D.Don</i>	Gentianaceae	>50	34
Luteolin	<i>Satureja parvifolia</i> (Phil.) Epling	Lamiaceae	22.3 (K1)	58
Lupinifolin	<i>Erythrina fusca</i> Lour.	Leguminosae	30.8 (K1)	16
Citflavanone	<i>Erythrina fusca</i> Lour.	Leguminosae	14.8 (K1)	16
8-Prenylaidzein	<i>Erythrina fusca</i> Lour.	Leguminosae	12.1 (K1)	16
5-Deoxy-3'-prenylbiochanin	<i>Erythrina sacleuxii</i> Hua	Leguminosae	17.6 (D6) 22.5 (W2)	17
Corylin	<i>Erythrina sacleuxii</i> Hua	Leguminosae	16.6 (D6) 19.7 (W2)	17
Erysubin F	<i>Erythrina sacleuxii</i> Hua	Leguminosae	12.0 (D6) 12.8 (W2)	17
3'-Prenylbiochanin A	<i>Erythrina sacleuxii</i> Hua	Leguminosae	23.7 (D6) 28.4 (W2)	17
7-Demethylrobustigenin	<i>Erythrina sacleuxii</i> Hua	Leguminosae	27.2 (D6) 31.7 (W2)	17
5'-Formylpratensein	<i>Erythrina sacleuxii</i> Hua	Leguminosae	21.7 (D6) 27.9 (W2)	17
2,3-Dehydrokeivetone	<i>Erythrina sacleuxii</i> Hua	Leguminosae	15.1 (D6) 12.7 (W2)	17
Prostratol C	<i>Erythrina sacleuxii</i> Hua	Leguminosae	17.6 (D6) 19.8 (W2)	17
Saclenone	<i>Erythrina sacleuxii</i> Hua	Leguminosae	24.2 (D6) 22.6 (W2)	17
2,3-Dihydro-7-demethylrobustigenin	<i>Erythrina sacleuxii</i> Hua	Leguminosae	28.0 (D6) 31.8 (W2)	17
(3 <i>R</i> )-7-Hydroxy-3',4'-dimethoxyisoflavan-2',5'-quinone	<i>Colutea istria</i> Mill.	Leguminosae	>50 (D6, W2)	59
6,3'-Dihydroxy-7,4'-dimethoxyisoflavone	<i>Colutea istria</i> Mill.	Leguminosae	>50 (D6, W2)	59

(Continued)

Table 1 (Continued)

Compound	Plant	Family	IC50 ( $\mu\text{M}$ )	Reference no.
2'-Methoxy-4',5'-methylenedioxy-7,8-[2-(1-methylethenyl)furo]isoflavone	<i>Millettia puguensis</i> J.B.Gillett	Leguminosae	>50	60
(-)-Maackiain	<i>Millettia puguensis</i> J.B.Gillett	Leguminosae	>50	60
6,7-Dimethoxy-3',4'-methylenedioxyisoflavone	<i>Millettia puguensis</i> J.B.Gillett	Leguminosae	>50	60
7,2'-Dimethoxy-4',5'-methylenedioxyisoflavone	<i>Millettia puguensis</i> J.B.Gillett	Leguminosae	>50	60
5-Acetyl-7-hydroxy-2-methylchromone	<i>Cassia siamea</i> Lam.	Leguminosae	19.4 (3D7)	50
Anhydrobarakol	<i>Cassia siamea</i> Lam.	Leguminosae	36.4 (3D7)	50
7-Demethylartanol E	<i>Artocarpus rigidus</i> Blume subsp. <i>rigidus</i>	Moraceae	18.0 (K1)	13
Bartericin B	<i>Dorstenia barteri</i> var. <i>subtriangularis</i> (Engl.) Hijman & C.C.Berg	Moraceae	19.3 (W2)	28
Isobavachalcone	<i>Dorstenia barteri</i> var. <i>subtriangularis</i> (Engl.) Hijman & C.C.Berg	Moraceae	19.0 (W2)	28
Talaumidin	<i>Pycnanthus angolensis</i> (Welw.) Warb.	Myristicaceae	60.5 (Dd2)	61
5-Galloylquercetin-3-O-R-L-arabinofuranoside	<i>Calycolpus warscewiczianus</i> O.Berg	Myrtaceae	14.5 (W2)	62
2',6'-Dihydroxy-4'-methoxydihydrochalcone	<i>Piper hostmannianum</i> C.DC. var. <i>berbicense</i>	Piperaceae	12.7 (F32) 16.9 (FcB1)	26
3-Farnesyl-p-hydroxybenzoic acid	<i>Piper tricuspe</i> C.DC.	Piperaceae	29.78 (FcB1)	48
Polygohomoisoflavanone	<i>Polygonum senegalense</i> Meisn. forma <i>senegalense</i>	Polygonaceae	19.2 (D6) 18.2 (W2)	25
2-Propen-1-one, 1-(2,4-dihydroxy-3,6-dimethoxyphenyl)-3-phenyl-	<i>Polygonum senegalense</i> Meisn. forma <i>senegalense</i>	Polygonaceae	16.6 (D6) 17.8 (W2)	25
2-Propen-1-one, 1-(2-hydroxy-4,6-dimethoxyphenyl)-3-phenyl-	<i>Polygonum senegalense</i> Meisn. forma <i>senegalense</i>	Polygonaceae	15.7 (D6) 11.3 (W2)	25
1-Propanone, 1-(2,6-dihydroxy-4-methoxyphenyl)-3-phenyl-	<i>Polygonum senegalense</i> Meisn. forma <i>senegalense</i>	Polygonaceae	23.5 (D6) 22.8 (W2)	25
1-Propanone, 1-(2,6-dihydroxy-3,4-dimethoxyphenyl)-3-phenyl-	<i>Polygonum senegalense</i> Meisn. forma <i>senegalense</i>	Polygonaceae	11.8 (D6) 12.9 (W2)	25
4H-1-Benzopyran-4-one, 2,3-dihydro-5-hydroxy-7-methoxy-2-phenyl-	<i>Polygonum senegalense</i> Meisn. forma <i>senegalense</i>	Polygonaceae	16.3 (D6) 21.8 (W2)	25
2-Propen-1-one, 1-(2,4-dihydroxy-6-methoxyphenyl)-3-phenyl-	<i>Polygonum senegalense</i> Meisn. forma <i>senegalense</i>	Polygonaceae	14.0 (D6) 9.5 (W2)	25
Quercetin	<i>Morinda morindoides</i> (Baker) Milne-Redh.	Rubiaceae	18.2 (K1)	63
5-O- $\beta$ -D-Glucopyranosyl-7,4'-dimethoxy-3'-hydroxy-4-phenylcoumarin	<i>Hintonia latiflora</i> (Sessé & Moc. ex DC.) Bullock	Rubiaceae	24.7 ( <i>P. berghei</i> )	29
5-O- $\beta$ -D-Glucopyranosyl-7-methoxy-3',4'-dihydroxy-4-phenylcoumarin	<i>Hintonia latiflora</i> (Sessé & Moc. ex DC.) Bullock	Rubiaceae	25.9 ( <i>P. berghei</i> )	29
Bipinnatones A	<i>Boronia bipinnata</i> Lindl.	Rutaceae	64 (Hbase II assay)	64
Bipinnatones B	<i>Boronia bipinnata</i> Lindl.	Rutaceae	51 (Hbase II assay)	64
Bergapten	<i>Zanthoxylum flavum</i> Vahl.	Rutaceae	21.8 (W2)	40
Syringaldehyde	<i>Vepris uguenensis</i> Engl.	Rutaceae	71.4 (3D7) 117.6 (FcM29)	65
Greveichromenol	<i>Harrisonia perforata</i> Merr.	Simaroubaceae	38.3 (K1)	66
Nitidanin	<i>Grewia bilamellata</i> Gagnep.	Tiliaceae	21.2 (D6) 18.4 (W2)	41
6-(8'Z,11'Z,14'Z-Heptadecatrienyl)-salicylic acid	<i>Viola websteri</i> Hemsl.	Violaceae	13.3 (D10)	47
Piceid	<i>Parthenocissus tricuspidata</i> Planch.	Vitaceae	13.2 (D10)	37
Longistylin A	<i>Parthenocissus tricuspidata</i> Planch.	Vitaceae	34.3 (D10)	37
Longistylin C	<i>Parthenocissus tricuspidata</i> Planch.	Vitaceae	19.2 (D10)	37

## Quinones

Primin (**82**), a natural benzoquinone occurring in *Primula obconica* Hance. (Primulaceae), was investigated for its antiprotozoal potential. Compound **82** showed moderate activity against K1 with an IC50 of 10.9  $\mu\text{M}$ .<sup>[67]</sup> A new

non-cannabinoid constituent was isolated from *Cannabis sativa* L. (Cannabaceae) namely 5-acetoxy-6-geranyl-3-n-pentyl-1,4-benzoquinone (**83**), which displayed notable antimalarial activity against D6 and W2 clones with IC50 values of 7.5 and 7.0  $\mu\text{M}$ , respectively.<sup>[20]</sup> New secondary

metabolites were isolated from the root extract of *Bauhinia purpurea* L. (Leguminosae). Among the isolated metabolites, two compounds exhibited antimalarial activity against K1, bauhinoxepin I (**84**) (IC<sub>50</sub> = 10.5  $\mu$ M) and bauhinoxepin J (**85**) (IC<sub>50</sub> = 5.8  $\mu$ M).<sup>[111]</sup> The ethanol extract of *Zhumeria majdae* Rech. f. & Wendelbo (Lamiaceae) showed potent antiplasmodial activity. Bioactivity-guided fractionation of the extract led to the isolation of 12,16-dideoxy aegyptinone B (**86**). This compound exhibited antiplasmodial activity with IC<sub>50</sub> values of 4.4 and 4.7  $\mu$ M against D6 and W2 strains, respectively. This compound was further found to have mild cytotoxicity towards cancer cell lines (IC<sub>50</sub> = 15.2–50.6  $\mu$ M).<sup>[68]</sup> From the roots of *Bulbine frutescens* Willd. (Asphodelaceae), the first sulfated phenylanthraquinones were isolated, together with their known sulfate-free analogues. Two of them, isoknipholone (**87**) and sodium 4'-*O*-demethylknipholone 6'-*O*-sulfate (**88**), presented promising activity against K1 with an IC<sub>50</sub> of 0.28  $\mu$ M for isoknipholone and an IC<sub>50</sub> of 7.9  $\mu$ M for the sulfated phenylanthraquinone.<sup>[69]</sup> From the roots of the African plant *Bulbine frutescens* Willd. (Asphodelaceae), two novel dimeric phenylanthraquinones, joziknipholones A (**89**) and B (**90**), were isolated. These two compounds exhibited strong activity against K1 with IC<sub>50</sub> values of 164 and 270 nM, respectively.<sup>[70]</sup> Two compounds, 10-(chrysophanol-7'-yl)-10-( $\xi$ )-hydroxychrysophanol-9-anthrone (**91**) and chryslandicin (**92**), were isolated from the dichloromethane extract of the roots of *Kniphofia foliosa* Hochst. (Asphodelaceae). They showed good activity against 3D7 with IC<sub>50</sub> values of 0.5 and 1.0  $\mu$ M, respectively.<sup>[71]</sup> Glaberianthrone (**93**), a new bianthrone, was isolated from the hexane extract of the stem bark of *Psorospermum glaberrimum* Hochr. (Clusiaceae) together with known compounds, 3-geranyloxyemodin anthrone (**95**), 3-prenyloxyemodin anthrone (**96**), 2-geranyloxyemodin (**97**) and bianthrone 1a (**94**). Their IC<sub>50</sub> values were 2.94, 1.68, 1.98, 5.34 and 2.53  $\mu$ M, respectively, against the W2 strain.<sup>[72]</sup> Bazouanthrone (**98**), a new anthrone derivative, was isolated from the root bark of *Harungana madagascariensis* Poir. (Clusiaceae), together with known compounds, feruginin A (**99**), harunganin (**100**), harunganol A (**101**) and harunganol B (**102**). All the compounds were found to be moderately active against W2: **98**, IC<sub>50</sub> = 1.8  $\mu$ M; **99**, IC<sub>50</sub> = 5.0  $\mu$ M; **100**, IC<sub>50</sub> = 2.7  $\mu$ M; **101**, IC<sub>50</sub> = 3.7  $\mu$ M; **102**, IC<sub>50</sub> = 3.7  $\mu$ M.<sup>[73]</sup> To discover antimalarial substances from plants cultivated in Thailand, 80% EtOH extracts from selected plants were screened against K1 strain. *Polyalthia viridis* Craib (Annonaceae) was found to show notable antimalarial activity. Marcanine A (**103**) (IC<sub>50</sub> = 10.5  $\mu$ M) was identified as its major active constituent.<sup>[74]</sup>

Figure 8 shows quinones and derivatives with moderate or promising activity *in vitro* against various strains of *P. falciparum*.


Table 2 gives the tested quinones presenting low or no activity *in vitro* against various strains of *P. falciparum*.<sup>[11,41,63,75,76]</sup>

## Terpenoid compounds

### Sesquiterpenes

Two sesquiterpenes, corymbolone (**104**) and mustakone (**105**), isolated from the chloroform extract of the rhizomes

of *Cyperus articulatus* L. (Cyperaceae), exhibited antiplasmodial properties (IC<sub>50</sub> = 4.53 and 0.64  $\mu$ M against NF54 and IC<sub>50</sub> = 8.14 and 1.15  $\mu$ M against EN36, respectively).<sup>[77]</sup> *Oncosiphon piluliferum* (L.f.) Källersjö (Asteraceae) is used traditionally to treat a variety of ailments, mainly fevers. Sesquiterpene lactones of the germacranolide and eudesmanolide types displaying antiplasmodial activity against D10 were isolated and identified: sivasinolide (**106**, IC<sub>50</sub> = 9.8  $\mu$ M), tatrudin A or tavulin (**107**, IC<sub>50</sub> = 1.5  $\mu$ M) and tanachin (**108**, IC<sub>50</sub> = 1.5  $\mu$ M). In addition, the cytotoxic effects of the active compounds against Chinese Hamster Ovarian cells were evaluated and the compounds were found to be toxic to mammalian cells at similar concentrations.<sup>[78]</sup> Two new helenanolide sesquiterpene lactones, helenalin-[2-(1-hydroxyethyl)acrylate] (**109**) and helenalin-[2-hydroxyethyl-3-methylacrylate] (**110**), as well as one known related structure, 11 $\alpha$ ,13-dihydrohelenalin-[2-(1-hydroxyethyl)acrylate] (**111**), were isolated from an ethyl acetate extract of leaves of *Vernoniopsis caudate* (Drake) Humbert (Asteraceae). The three lactones displayed strong antiplasmodial activity against FcB1, with IC<sub>50</sub> values of 1, 0.19 and 0.41  $\mu$ M, respectively. However, these compounds also exhibited considerable cytotoxicity on KB cells (IC<sub>50</sub> < 1  $\mu$ M in each case).<sup>[79]</sup> Leaves and flowers of *Artemisia gorgonum* Webb (Asteraceae) collected in Fogo, Cape Verde, were phytochemically investigated and resulted in the isolation of a known germacranolide, hanphyllin (**112**), which exhibited antiplasmodial activity with an IC<sub>50</sub> of 9.7  $\mu$ M against FcB1 and was weakly cytotoxic to the Vero cell line (IC<sub>50</sub> = 111.9  $\mu$ M).<sup>[80]</sup> Two new sesquiterpene lactones, wedelolides A (**113**) and B (**114**), were isolated from the leaves of *Wedelia trilobata* Hitchc. (Asteraceae). The two compounds displayed antimalarial activity with IC<sub>50</sub> values of 4.2 and 9.1  $\mu$ M, respectively.<sup>[81]</sup> A new sesquiterpene lactone as well as two known ones were isolated from the dichloromethane fraction of an aqueous extract from *Vernonia cinerea* Less. (Asteraceae). Three compounds, 8 $\alpha$ -tigloyloxy-hirsutinolide-13-*O*-acetate (**115**), 8 $\alpha$ -(4-hydroxyethacryloyloxy)-hirsutinolide-13-*O*-acetate (**116**) and vernolide D (**117**), were active against W2 with IC<sub>50</sub> values of 3.9, 3.7 and 3.5  $\mu$ M, respectively.<sup>[82]</sup> Fractionation of the dichloromethane extract of the leaves of *Vernonia staehelinoides* Mart. ex Baker (Asteraceae) allowed the isolation of two structurally related hirsutinolides. These compounds displayed strong antiplasmodial activity against D10 and were less effective against K1, compound **118**, 8 $\alpha$ -(2-methylacryloyloxy)-3-oxo-1-desoxy-1,2-dehydrohirsutinolide-13-*O*-acetate had an IC<sub>50</sub> = 0.6  $\mu$ M on D10 and IC<sub>50</sub> = 4.5  $\mu$ M on K1 and compound **119**, 8 $\alpha$ -(50-acetoxyseneciolyoxy)-3-oxo-1-desoxy-1,2-dehydrohirsutinolide-13-*O*-acetate had an IC<sub>50</sub>/D10 = 0.5  $\mu$ M and IC<sub>50</sub>/K1 = 5.5  $\mu$ M. These two compounds were found to be cytotoxic to mammalian Chinese Hamster Ovarian cells at similar concentrations.<sup>[83]</sup> Chemical exploration of *Camchaya calcarea* Kitam. (Asteraceae) led to the isolation of eight known sesquiterpene lactones, which exhibited moderate antiplasmodial activity against K1, including goyazensolide (**120**) (IC<sub>50</sub> = 3.3  $\mu$ M), lychnophorolide B (**122**) (IC<sub>50</sub> = 7.2  $\mu$ M), isogoyazensolide (**123**) (IC<sub>50</sub> = 4.4  $\mu$ M), isocentratherin (**124**) (IC<sub>50</sub> = 5.6  $\mu$ M), 5-epi-isogoyazensolide (**125**) (IC<sub>50</sub> = 4.4  $\mu$ M) and 5-epi-isocentratherin (**126**) (IC<sub>50</sub> = 8.0  $\mu$ M). The most promising was lychnophorolide A (**121**) with an IC<sub>50</sub> of



**Figure 8** Quinones and derivatives with moderate or promising activity *in vitro* against various strains of *P. falciparum*

**Table 2** Quinones presenting low or no activity *in vitro* against various strains of *P. falciparum*

Compound	Plant	Family	IC <sub>50</sub> ( $\mu\text{M}$ )	Reference no.
Newbouldiaquinone A	<i>Newbouldia laevis</i> Seem.	Bignoniaceae	78% at 20 $\mu\text{M}$ (NF54)	75
Bauhinoxepin H	<i>Bauhinia purpurea</i> L.	Leguminosae	11.2 (K1)	11
Rubiadin-1-methyl ether	<i>Prismatomeris malayana</i> Ridl.	Rubiaceae	–	76
Nordamnacanthal	<i>Prismatomeris malayana</i> Ridl.	Rubiaceae	–	76
Damnacanthal	<i>Prismatomeris malayana</i> Ridl.	Rubiaceae	–	76
Chrysarín	<i>Morinda morindoides</i> (Baker) Milne-Redh.	Rubiaceae	105.4 (NF54/64)	63
Alizarin	<i>Morinda morindoides</i> (Baker) Milne-Redh.	Rubiaceae	60.4 (NF54/64)	63
2,6-Dimethoxy-1-acetylquinol	<i>Grewia bilamellata</i> Gagnep.	Tiliaceae	42.2 (D6) 23.0 (W2)	41


0.8  $\mu\text{M}$ .<sup>[84]</sup> Bioactivity-guided isolation of the chloroform fractions of the whole plants of *Carpesium rosulatum* Miq. (Asteraceae) led to the isolation of a sesquiterpene lactone, ineupatorolide A (**127**), displaying high antiplasmodial activity against D10 with an IC<sub>50</sub> of 19 nM.<sup>[85]</sup> Ineupatorolide A was also found to have potential antimalarial activity *in vivo* when tested against *P. berghei* in mice. Compound **127** (2, 5, 10 mg/kg per day, intraperitoneally) exhibited a significant blood schizontocidal activity in 4-day early infection, preventive and curative treatment, with a significant mean survival time comparable with that of the standard drug, chloroquine (5 mg/kg per day). Ineupatorolide A possesses promising antiplasmodial activity that can be exploited in malaria therapy.<sup>[86]</sup> 4-Hydroxyanthecotulide (**128**), a linear sesquiterpene lactone

isolated from *Anthemis auriculata* Boiss. (Asteraceae), was evaluated against K1 and had an IC<sub>50</sub> of 7.6  $\mu\text{M}$ .<sup>[87]</sup>

Figure 9 shows sesquiterpenes with moderate or promising activity *in vitro* against various strains of *P. falciparum*.

### Diterpenes

Geranylgeraniol (**129**) was isolated from the stems and leaves of *Croton lobatus* L. (Euphorbiaceae), a medicinal plant used in western Africa in traditional folk medicine to cure malaria, pregnancy troubles and dysentery. The compound showed reasonable antiplasmodial activity against K1 with an IC<sub>50</sub> value of 3.7  $\mu\text{M}$  and good selectivity (SI value > 25).<sup>[57]</sup> A new diterpenoid, steenkrotin A (**130**), was isolated from an ethanol extract of the leaves of *Croton steenkampianus*



**Figure 9** Sesquiterpenes with moderate or promising activity *in vitro* against various strains of *P. falciparum*

Gerstner (Euphorbiaceae) and was tested against D10, D6, Dd2 and W2, showing moderate activity with IC50 values of 15.8, >30, 9.4 and 9.1  $\mu\text{M}$ , respectively.<sup>[88]</sup>

Labdane diterpenoid, 3-deoxyaulacocarpin A (**131**) was isolated from the seeds of *Aframomum zambesiacum* K. Schum (Zingiberaceae). It possessed antiplasmodial activity against FcB1 with an IC50 of 4.97  $\mu\text{M}$ .<sup>[89]</sup> Known compounds were isolated from the dried fruits of *Juniperus seravschanica* Komarov (Cupressaceae) and were tested. Three of them showed promising antimalarial activity: cedrol (**132**) with IC50 = 265, 1.4 and 10.2  $\mu\text{M}$ ; sugiol (**133**) with IC50 = 1.6, 3.4 and 1.4  $\mu\text{M}$ ; and 12,15-dihydroxyabda-8 (17),13-dien-19-oic acid (**134**) with IC50 = 2.2, 4.1 and 4.6  $\mu\text{M}$  against D6, TM91C235 and W2, respectively.<sup>[90]</sup> *Vitex rehmannii* Gürke (Lamiaceae) contained a labdane diterpene, 12*S*,16*S*/*R*-dihydroxy-*ent*-labda-7,13-dien-15,16-olide (**135**), which exists as an inseparable epimeric mixture. This mixture exhibited reasonable antimalarial activity against FCR-3 (IC50 = 7.2  $\mu\text{M}$ ). However, this was due to its cytotoxic properties.<sup>[91]</sup>

Bioassay-guided fractionation of a trunk bark extract of *Laetia procera* (Poepp.) Eichler (Flacourtiaceae) led to the isolation of six clerodane diterpenoids: casearluin A (**136**), casamembrol A (**137**), laetiaprocerine A–D (**138–141**). The diterpenoids exhibited antiplasmodial activity with IC50 values of 0.62, 0.57, 0.58, 4.44, 4.66 and 6.04  $\mu\text{M}$  on F32 strain, and 0.54, 0.59, 0.66, 6.08, 5.35, 3.79 and 27.5  $\mu\text{M}$  on FcB1 strain, but most of them were also cytotoxic. Compound **138** showed the best selectivity index of 6.8.<sup>[92]</sup> Bioactivity-guided fractionation of hexane and dichloromethane extracts of the bark of *Casearia grewiifolia* Vent. (Flacourtiaceae) afforded four new clerodane diterpenes, caseargrewiins A–D (**142–145**), and two known clerodane diterpenes, *rel*-(2*S*,5*R*,6*R*,8*S*,9*S*,10*R*,18*S*,19*R*)-18,19-diacetoxy-18,19-epoxy-6-methoxy-2-(2-methylbutanoyloxy)cleroda-3,13(16),14-triene (**146**) and *rel*-(2*S*,5*R*,6*R*,8*S*,9*S*,10*R*,18*S*,19*R*)-18,19-diacetoxy-18,19-epoxy-6-hydroxy-2-(2-methylbutanoyloxy)cleroda-3,13(16),14-triene (**147**). All compounds exhibited antimalarial activity against K1 with IC50 values of 5.5, 3.6, 5.2, 7.9, 6.0 and 6.0  $\mu\text{M}$ , but also cytotoxicity.<sup>[93]</sup>

A new diterpene, (1*S*,5*S*,9*S*,10*S*,11*R*,13*R*)-1,11-dihydroxypimara-8(14),15-diene (**148**) was isolated from the dichloromethane extract of whole plants of *Kaempferia marginata* Carey (Zingiberaceae) and had antimalarial activity against K1 (IC50 = 10.5  $\mu\text{M}$ ).<sup>[94]</sup>

Some 44 cassane- and norcassane-type diterpenes isolated from  $\text{CH}_2\text{Cl}_2$  extract of *Caesalpinia crista* L. (Caesalpiniaaceae) from Myanmar and Indonesia were evaluated for their antimalarial activity against FCR-3/A2 clone. Caesalpinins MA (**149**), ME–MJ (**150–155**), ML (**156**), norcaesalpinins MC (**157**) and MD (**158**), caesalpinins C–F (**159–162**), J–K (**163–164**), N (**165**) and P (**166**), norcaesalpinins A–F (**167–172**), caesalmins B (**173**) and C (**174**), caesaldehydricin E (**175**), 2-acetoxycasaldehydricin E (**176**), 2-acetoxy-3-deacetoxycaesaldehydricin E (**177**), 14(17)-dehydrocaesalmin F (**178**), bonducellpins B (**179**) and C (**180**), 7-acetoxybonducellpin C (**181**) and 1-deacetoxy-1-oxocaesalmin C (**182**) displayed antimalarial activity with IC50 values of: 3.5, 3.6, 4.1, 2.5, 7.0, 2.1, 1.9, 0.65, 3.1, 1.0, 0.76, 0.8, 6.5, 0.65, 1.0, 0.40, 0.12, 1.7, 0.8, 0.26, 5.0, 2.0, 0.09, 0.14, 0.80, 3.4, 4.0, 6.5, 0.098, 0.2,

0.24, 0.12, 0.6 and 2.9  $\mu\text{M}$ , respectively. Eighteen diterpenes possessed strong activity with IC50  $\leq$  1  $\mu\text{M}$ , with norcaesalpinin E (**171**) and 2-acetoxy-3-deacetoxycaesaldehydricin E (**177**) being the most potent.<sup>[95,96]</sup> In continuity with the previous work, a new furanocassane-type diterpene, caesalpinin H (**183**), was isolated from the  $\text{CH}_2\text{Cl}_2$  extract of the seed kernels of *Caesalpinia crista* L. (Caesalpiniaaceae) and showed an IC50 value of 5.2  $\mu\text{M}$  against FCR-3/A2.<sup>[97]</sup> Three new cassane furanoditerpenoids were isolated from the EtOAc extract of the seed kernels of *Caesalpinia bonduc* L. Roxb. (Caesalpiniaaceae). Bonducellpins E–G (**184–186**) exhibited antimalarial activity on K1 strain with IC50 values of 1.6, 5.8 and 3.8  $\mu\text{M}$ , respectively. None of the compounds were cytotoxic against any of the tumour cell lines tested.<sup>[98]</sup> Compound 6 $\alpha$ ,7 $\beta$ -diacetoxyvouacapane (**187**) was isolated from the seeds of *Bowdichia nitida* Spruce ex Benth. (Leguminosae), and showed promising antiplasmodial activity against 3D7 (IC50 = 968 nm) and a good selectivity index with regard to cytotoxicity (IC50 > 250  $\mu\text{M}$ ).<sup>[99]</sup> Bioactivity-guided fractionation of the petroleum ether extract of the leaves of *Hyptis suaveolens* (L.) Poit. (Lamiaceae), widely used in traditional medicine, led to the isolation of an abietane-type diterpenoid endoperoxide, 13 $\alpha$ -epi-dioxiabiet-8(14)-en-18-ol (**188**), displaying high antiplasmodial activity against D10 (IC50 = 344 nm).<sup>[100]</sup>

Five known abietane diterpenes (**189–193**) were isolated from five *Plectranthus* species (Lamiaceae), namely *Plectranthus hadiensis* C. Chr., *Plectranthus lucidus* Burch. ex Benth., *Plectranthus ecklonii* Benth., *Plectranthus purpuratus* Harv. subsp. *purpuratus* and *Plectranthus purpuratus* Harv. subsp. *tongaensis*. The compounds showed antiplasmodial activity against FCR-3 (IC50 = 4.6, 5.3, 3.1, 6.0 and 4.7  $\mu\text{M}$ , respectively). However, the cytotoxicity profile indicated a low degree of specificity towards the malaria parasite.<sup>[101]</sup> A bioassay-guided fractionation of *Juniperus procera* Hochst. ex Endl. (Cupressaceae) berries yielded pure compounds. Among these, abieta-7,13-diene (**194**) and ferruginol (**195**) demonstrated antimalarial activity against D6 and W2 strains with IC50 = 7.0 and 7.4  $\mu\text{M}$ , respectively, for **194** and IC50 = 12.3 and 4.9  $\mu\text{M}$ , respectively, for **195**.<sup>[102]</sup>

In 2006, the antimalarial activity of ferruginol (IC50 = 6.9  $\mu\text{M}$ ) isolated from *Fuerstia africana* T.C.E.Fr. (Lamiaceae) was correlated with cytotoxic activity and, therefore, it was not a promising antimalarial candidate.<sup>[103]</sup>

Five new poly-*O*-acylated jatropha diterpenes, including 1 $\alpha$ ,13 $\beta$ ,14 $\alpha$ -trihydroxy-3 $\beta$ ,7 $\beta$ -dibenzoyloxy-9 $\beta$ ,15 $\beta$ -diacetoxyjatropha-5,11 *E*-diene (**196**), 1 $\alpha$ ,8 $\beta$ ,9 $\beta$ ,14 $\alpha$ ,15 $\beta$ -pentaacetoxy-3 $\beta$ -benzoyloxy-7-oxojatropha-5,12-diene (**197**), 7,8 $\beta$ ,9 $\beta$ ,14 $\alpha$ ,15 $\beta$ -pentaacetoxy-3 $\beta$ -benzoyloxy-1 $\alpha$ ,5 $\beta$ -dihydroxyjatropha-6(7),12-diene (**198**) and 1 $\alpha$ ,7,8 $\beta$ ,9 $\beta$ ,14 $\alpha$ ,15 $\beta$ -hexaacetoxy-3 $\beta$ -benzoyloxy-5 $\beta$ -hydroxyjatropha-6(7),12-diene (**199**) were isolated from the white latex of *Pedilanthus tithymaloides* (L.) Poit. (Euphorbiaceae). These highly oxygenated diterpenes possess a rare *O*-acetyl enol moiety and showed antiplasmodial activity against K1 strain: **196** (IC50 = 5.9  $\mu\text{M}$ ), **197** (IC50 = 4.9  $\mu\text{M}$ ), **198** (IC50 = 6.0  $\mu\text{M}$ ) and **199** (IC50 = 5.8  $\mu\text{M}$ ).<sup>[104]</sup>

Figure 10 shows diterpenes with moderate or promising activity *in vitro* against various strains of *P. falciparum*.




Figure 10 Diterpenes with moderate or promising activity *in vitro* against various strains of *P. falciparum*




Figure 10 (Continued)

### Triterpenes

From the stem bark of *Ekebergia capensis* Sparrm. (Meliaceae), a triterpenoid derivative was isolated and screened against FCR-3 and K1 strains. 2,3,22,23-Tetrahydroxy-2,6,10,15,19,23-hexamethyl-6,10,14,18-tetracosatetraene

(3*R*,22*R*) (**200**) displayed an IC<sub>50</sub> of 18  $\mu$ M and 7  $\mu$ M on FCR-3 and K1, respectively, and showed *in vivo* parasitaemia suppression of 52.9% when given intraperitoneally.<sup>[105]</sup> Seven new and two known compounds were isolated from an ethyl acetate extract of the leaves of *Nuxia sphaerocephala*



Baker (Buddlejaceae) and have been considered moderately active, 3-oxolupenal (**201**) and 3 $\beta$ -hydroxy-lupenal (**202**) having the best activity with IC50 values of 3.6 and 7.2  $\mu\text{M}$ , respectively, against FcB1.<sup>[106]</sup> One new and eight known ceanothane- and lupane-type triterpenes were isolated from the root bark of *Ziziphus cambodianus* Pierre (Rhamnaceae). The new compound, 3-*O*-vanillylceanothic acid (**203**), and two known compounds, 2-*O*-*E*-*p*-coumaroyl aliphatic acid (**204**) and zizyberenic acid (**205**), exhibited notable antiplasmodial activity with IC50 values of 5.8, 1.5 and 6.6  $\mu\text{M}$ , respectively.<sup>[107]</sup>

Bioassay-directed fractionation led to the isolation of betulinic acid 3-caffeate (**206**) from a sample of the dried leaves, twigs, and branches of *Diospyros quaesita* Thwaites (Ebenaceae). This compound showed strong antimalarial activity against D6 and W2 with IC50 values of 1.40 and 0.98  $\mu\text{M}$ , respectively. Evaluation of **206** in the human oral epidermoid (KB) cancer cell line revealed cytotoxicity at an IC50 of 4.0  $\mu\text{M}$ .<sup>[108]</sup> Phytochemical investigation of the  $\text{CH}_2\text{Cl}_2$  extracts of *Erythrina stricta* Roxb. (Leguminosae) roots and *Erythrina subumbrans* Merr. (Leguminosae) stems led to the isolation of one triterpene, soyasapogenol B (**207**), which exhibited moderate antiplasmodial activity (10.0  $\mu\text{M}$ ) against K1.<sup>[18]</sup> Friedelan-3-one (**208**) was isolated from the root bark of *Harungana madagascariensis* Poir. (Clusiaceae). Its antiplasmodial activity was evaluated against W2 strain and gave an IC50 of 7.7  $\mu\text{M}$ .<sup>[73]</sup> Garcinane (**209**) was isolated from the roots of *Garcinia polyantha* Oliv. (Clusiaceae) and exhibited antimalarial activity against NF54 with IC50 ranging from 2.5 to 4.1  $\mu\text{M}$ .<sup>[35]</sup> A new bisnortriterpene quinone methide, 20-epi-isoiguesterinol (**211**) and a known compound, isoiguesterin (**210**), were isolated from the petroleum ether extract of the roots of *Salacia madagascariensis* DC. (Celastraceae). Compound **211** was active with IC50 of 0.16  $\mu\text{M}$  on D6 and W2, and **210** with IC50 of 0.50 and 0.42  $\mu\text{M}$  on D6 and W2, respectively.<sup>[109]</sup>

Bioassay-directed fractionation led to the isolation of 2 $\alpha$ ,3 $\beta$ -dihydroxyolean-12-en-28-oic acid (**212**) from a sample of *Grewia bilamellata* Gagnep. (Tiliaceae), which displayed antimalarial activity against D6 and W2 (21.1 and 8.6  $\mu\text{M}$ ) without significant cytotoxicity.<sup>[41]</sup> A bioassay-guided fractionation from *Morinda lucida* Benth. (Rubiaceae) leaves and from *Satureja parvifolia* (Phil.) Epling (Lamiaceae) resulted in the isolation of two known triterpenic acids, ursolic acid (**213**) and oleanolic acid. These two compounds had already been evaluated *in vitro* but in these two studies some activity was observed for the first time, with IC50 of 32.3 and 19.8  $\mu\text{M}$  for oleanolic acid, and 6.8 and 10.7  $\mu\text{M}$  for ursolic acid. *In vivo*, oleanolic acid at a daily dose of 200 mg/kg produced 37.4% chemosuppression.<sup>[58,110]</sup> Bioactivity-guided fractionation of the petroleum ether extracts of the whole plants of *Viola verecunda* A. Gray (Violaceae) led to the isolation of epi-oleanolic acid (**214**), a triterpenoid, displaying high antiplasmodial activity against FcB1 strain (IC50 = 39 nm).<sup>[111]</sup>

Bioassay-guided fractionation of the antimalarial-active  $\text{CHCl}_3$  extract of the dried stem of *Nauclea orientalis* (L.) L. (Rubiaceae) resulted in the isolation of a known compound, 3 $\alpha$ ,23-dihydroxyurs-12-en-28-oic acid (**215**), which showed

moderate activity with an IC50 of 9.7  $\mu\text{M}$  on D6 and 12.7  $\mu\text{M}$  on W2.<sup>[112]</sup> Three triterpenes were isolated from *Cogniauxia podolaena* Baill. (Cucurbitaceae), cucurbitacin B (**216**), cucurbitacin D (**217**) and 20-epibryonolic acid (**218**). All compounds obtained were assayed for antiplasmodial activity (on FcM29) and cytotoxicity. The IC50 values were 2.9, 7.8 and 3.7  $\mu\text{M}$ , respectively. Both **216** and **217** have high cytotoxicity, whereas **218** showed a better selectivity index.<sup>[113]</sup> A compound was isolated from the active fraction of *Salvia radula* Epling (Lamiaceae) and identified as betulafolientriol oxide (**219**). It displayed moderate antimalarial activity (IC50 = 10.4  $\mu\text{M}$ ).<sup>[114]</sup>

A quassinoid, neosergeolide (**220**), isolated from the roots and stems of *Picrolemma sprucei* Hook.f. (Simaroubaceae), possessed a significant inhibitory effect (more active than quinine and chloroquine) on K1 strain (IC50 = 2 nm).<sup>[49]</sup> Two new limonoids, domesticulide B (**221**) and C (**222**), and three more known ones, methyl 6-acetoxyangolensate (**223**), azadiradione (**224**) and dukunolide C (**225**), were isolated from seeds of *Lansium domesticum* Corrêa (Meliaceae) and showed antimalarial activity against K1 with IC50 values of 6.0, 4.1, 7.2, 6.4, 9.6  $\mu\text{M}$ , respectively.<sup>[115]</sup> Marked antimalarial activity was observed for anthothecol (**226**), a limonoid of *Khaya anthotheca* C.DC. (Meliaceae). IC50 values were 1.4 and 0.17  $\mu\text{M}$  against W2 strain using two different assays.<sup>[116]</sup> In the search for active principles from the stem bark of *Entandrophragma angolense* C.DC. (Meliaceae), 7 $\alpha$ -obacynyl acetate (**227**) was isolated and tested against W2 strain. It exhibited antimalarial activity with an IC50 of 4.0  $\mu\text{M}$ .<sup>[117]</sup> Two limonoids isolated from the seeds of *Chisocheton siamensis* Craib (Meliaceae) were tested for antimalarial activity. Dysobinin (**228**) and mahonin (**229**) showed an inhibitory effect against K1 with IC50 values of 4.2 and 5.7  $\mu\text{M}$ , respectively.<sup>[118]</sup> The dichloromethane extract of *Pseudocedrela kotschyi* Harms (Meliaceae) root commonly used in Malian traditional medicine led to the isolation of two known compounds, 7-deacetylgedunin (**230**) and 7-deacetyl-7-oxogedunin (**231**), which exhibited activity against K1 (IC50 = 3.1 and 4.1  $\mu\text{M}$ , respectively).<sup>[119]</sup> Compound **231** was also isolated from the stem bark of *Ekebergia capensis* Sparrm. (Meliaceae) and screened against FCR-3 strain with an IC50 of 6  $\mu\text{M}$ .<sup>[105]</sup>

Figure 11 shows triterpenes with moderate or promising activity *in vitro* against various strains of *P. falciparum*.

### Steroids

Four new pregnane glycosides, 12-*O*-benzoyl-20-*O*-acetyl 3 $\beta$ ,12 $\beta$ ,14 $\beta$ ,20 $\beta$ -tetrahydroxy-(20*S*)-pregn-5-ene-6-deoxy-3-*O*-methyl- $\beta$ -D-allopyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-cymaropyranosyl (1 $\rightarrow$ 4)- $\beta$ -D-cymaropyranoside (**232**), 12-*O*-benzoyl-20-*O*-acetyl 3 $\beta$ ,7 $\alpha$ ,12 $\beta$ ,14 $\beta$ ,20 $\beta$ -tetrahydroxy-(20*S*)-pregn-5-ene-6-deoxy-3-*O*-methyl- $\beta$ -D-allopyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-cymaropyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-cymaropyranoside (**233**), 12-*O*-benzoyl-20-*O*-acetyl 3 $\beta$ ,5 $\alpha$ ,12 $\beta$ ,14 $\beta$ ,20 $\beta$ -pentahydroxy-(20*S*)-pregn-6-ene- $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 4)-6-deoxy-3-*O*-methyl- $\beta$ -D-allopyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-cymaropyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-cymaropyranoside (**234**) and 12,20-dibenzoyl-3 $\beta$ ,12 $\beta$ ,14 $\beta$ ,20 $\beta$ -tetrahydroxy-(20*S*)-pregn-5-ene- $\beta$ -D-glucopyranosyl-(1 $\rightarrow$ 4)-6-deoxy-3-*O*-methyl- $\beta$ -D-allopyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-cymaropyranosyl-(1 $\rightarrow$ 4)- $\beta$ -D-cymaropyranoside

(235) were isolated from *Caralluma tuberculata* N.E.Br. (Asclepiadaceae), in addition to a known one, russelioside E (236). All the isolated compounds were tested for their antimalarial activity against K1 and had IC<sub>50</sub> = 7.4, 6.5, 9.6, 5.7 and 7.5 μM, respectively.<sup>[120]</sup>

Figure 12 shows steroids with moderate or promising activity *in vitro* against various strains of *P. falciparum*.

Table 3 gives the tested terpenoid compounds presenting low or no activity *in vitro* against various strains of *P. falciparum*.<sup>[41,58,65,74,78,80,87,89,92,94,105,110,115,117–119,121–140]</sup>




Figure 11 Triterpenes with moderate or promising activity *in vitro* against various strains of *P. falciparum*





Figure 11 (Continued)

Figure 12 Steroids with moderate or promising activity *in vitro* against various strains of *P. falciparum*

## Alkaloids

### Ornithine and lysine derivatives

The methanolic extract of *Albizia gummifera* C.A.Sm. (Leguminosae) was fractionated to isolate five known spermine alkaloids

from the alkaloidal fraction, budmunchiamine K (237), 6-hydroxybudmunchiamine K (238), 5-normethylbudmunchiamine K (239), 6-hydroxy-5-normethylbudmunchiamine K (240) and 9-normethylbudmunchiamine K (241). These alkaloids exhibited good activity with IC<sub>50</sub> values of 0.18, 0.29, 0.20, 0.33

**Table 3** Terpenoid compounds presenting low or no activity *in vitro* against various strains of *P. falciparum*

Compound	Plant	Family	IC <sub>50</sub> (μM)	Reference no.
16-Hydroxycyclo-3,13(14)Z-dien-15,16-olide	<i>Goniothalamus marcanii</i> Craib	Annonaceae	11.3 (K1)	74
Lupeol	<i>Holarrhena floribunda</i> T.Durand & Schinz	Apocynaceae	>50 (FCR-3/3D7)	121
3-O-(3'-Hydroxyeicosanoyl)lupeol	<i>Holarrhena floribunda</i> T.Durand & Schinz	Apocynaceae	>50 (FCR-3/3D7)	121
3-O-[(2'-(Tetracosyloxy)acetyl]lupeol	<i>Holarrhena floribunda</i> T.Durand & Schinz	Apocynaceae	>50 (FCR-3/3D7)	121
3-O-[(1''-Hydroxyoctadecyloxy)-2'-hydroxypropanoyl]lupeol	<i>Holarrhena floribunda</i> T.Durand & Schinz	Apocynaceae	>50 (FCR-3/3D7)	121
Uzariogenin	<i>Calotropis gigantea</i> (L.) W.T.Aiton	Asclepiadaceae	>50 (K1)	122
Calactin	<i>Calotropis gigantea</i> (L.) W.T.Aiton	Asclepiadaceae	>50 (K1)	122
Calotropin	<i>Calotropis gigantea</i> (L.) W.T.Aiton	Asclepiadaceae	>50 (K1)	122
Taraxasteryl acetate	<i>Calotropis gigantea</i> (L.) W.T.Aiton	Asclepiadaceae	>50 (K1)	122
Esacetyl-α-cyclopyrethrosin	<i>Oncosiphon piluliferum</i> (L.f.) Källersjö	Asteraceae	16.7 (D10)	78
1,10-Dioxo-1,10-deoxy-1,10-secogorgonolide	<i>Artemisia gorgonum</i> Webb	Asteraceae	50.0 (FcB1)	80
3β,4β-Epoxy-1β,10β-epiarborescin	<i>Artemisia gorgonum</i> Webb	Asteraceae	50.4 (FcB1)	80
Arborescin	<i>Artemisia gorgonum</i> Webb	Asteraceae	15.3 (FcB1)	80
1β,10β-Epoxy-2α-hydroxykauniolide	<i>Artemisia gorgonum</i> Webb	Asteraceae	22.1 (FcB1)	80
1α,4α,10α-Trihydroxy-5α,11βH-guaia-2-en-12,6α-olide	<i>Artemisia gorgonum</i> Webb	Asteraceae	32.1 (FcB1)	80
Ridentin	<i>Artemisia gorgonum</i> Webb	Asteraceae	21.4 (FcB1)	80
Anthecularin	<i>Anthemis auriculata</i> Boiss.	Asteraceae	>50 (K1)	123
Anthecotulide	<i>Anthemis auriculata</i> Boiss.	Asteraceae	16.1 (K1)	87
4-Acetoxyanthecotulide	<i>Anthemis auriculata</i> Boiss.	Asteraceae	16.7 (K1)	87
Methyl populnoate	<i>Austroplenckia populnea</i> (Reissek) Lundell	Celastraceae	>50 (D6)	124
Populnoic acid	<i>Austroplenckia populnea</i> (Reissek) Lundell	Celastraceae	>50 (D6)	124
Stigmast-5-en-3-O-β-(D-glucopyranoside)	<i>Austroplenckia populnea</i> (Reissek) Lundell	Celastraceae	>50 (D6)	124
Endodesmiadiol	<i>Endodesmia calophylloides</i> Benth.	Clusiaceae	13.0 (W2)	125
Canophyllal	<i>Endodesmia calophylloides</i> Benth.	Clusiaceae	18.2 (W2)	125
Cerin	<i>Endodesmia calophylloides</i> Benth.	Clusiaceae	14.1 (W2)	125
Morelloflavone	<i>Endodesmia calophylloides</i> Benth.	Clusiaceae	23.6 (W2)	125
3β-Acetoxyoleanolic acid	<i>Endodesmia calophylloides</i> Benth.	Clusiaceae	13.1 (W2)	125
Sutherlandioside A	<i>Sutherlandia frutescens</i> (L.) R. Br.	Fabaceae	>50 (D6/W2)	126
Sutherlandioside B	<i>Sutherlandia frutescens</i> (L.) R. Br.	Fabaceae	>50 (D6/W2)	126
Sutherlandioside C	<i>Sutherlandia frutescens</i> (L.) R. Br.	Fabaceae	>50 (D6/W2)	126
Sutherlandioside D	<i>Sutherlandia frutescens</i> (L.) R. Br.	Fabaceae	>50 (D6/W2)	126
Laetianolide A	<i>Laetia procera</i> (Poepp.) Eichler	Flacourtiaceae	57.6 (F32) 27.5 (FCb1)	92
15-O-Ethylleopersin C	<i>Leonurus cardiaca</i> L.	Lamiaceae	>50 (D6/W2)	127
15-O-Methylleopersin C	<i>Leonurus cardiaca</i> L.	Lamiaceae	>50 (D6/W2)	127
15-Epi-O-methylleopersin C	<i>Leonurus cardiaca</i> L.	Lamiaceae	>50 (D6/W2)	127
12,17-Diacetoxy,15-hydroxy,2-oxo,3,13 E(14)-diene clerodane	<i>Gomphostemma crinitum</i> Wall.	Lamiaceae	22.14 (MRC-02)	128
Fagraldehyde	<i>Fagraea fragrans</i> Roxb.	Loganiaceae	>50 (W2)	129
Geranylarnesol	<i>Thalia geniculata</i> L.	Marantaceae	12.7	130
Kurubasch aldehyde	<i>Trichilia emetica</i> Vahl.	Meliaceae	76 (3D7)	131
16-Oxolabda-8(17),12(E)-dien-15-oic acid	<i>Turraeanthus africana</i> Pellegr.	Meliaceae	83.1 (F32)	132
Methyl 14,15-epoxyabda-8(17),12E-diene-16-oate	<i>Turraeanthus africana</i> Pellegr.	Meliaceae	>50 (F32)	133
Ekeberin D4	<i>Ekebergia capensis</i> Sparrm.	Meliaceae	40 (FCR-3)	105
Domesticulide D	<i>Lansium domesticum</i> Corrêa	Meliaceae	11.8 (K1)	115
6-Acetoxy-mexicanolide	<i>Lansium domesticum</i> Corrêa	Meliaceae	18.4 (K1)	115
Walsuronoid	<i>Walsura robusta</i> Roxb.	Meliaceae	40% at 40 μM (3D7)	134
Walsuronoid B	<i>Walsura robusta</i> Roxb.	Meliaceae	40% at 40 μM (3D7)	134
24-Methylene cycloartenol	<i>Entandrophragma angolense</i> C.DC.	Meliaceae	12.3 (W2)	117
7α-Acetoxydihydranomilin	<i>Entandrophragma angolense</i> C.DC.	Meliaceae	34.9 (W2)	117
Methylangolensate	<i>Entandrophragma angolense</i> C.DC.	Meliaceae	48.7 (W2)	117
6α-Acetoxyepoxyazadiradione	<i>Chisocheton siamensis</i> Craib	Meliaceae	12.0 (K1)	118
Kotschyin A	<i>Pseudocedrela kotschyi</i> Harms	Meliaceae	> 5 (K1)	119
Methyl 3,4-dihydroxy-5-(3-methyl-2-butenyl)benzoate	<i>Piper glabratum</i> Kunth	Piperaceae	17.4 (F32)	135
Methyl 4-hydroxy-3-(3-methyl-2-butenyl)benzoate	<i>Piper glabratum</i> Kunth	Piperaceae	12.7 (F32)	135

(Continued)

Table 3 (Continued)

Compound	Plant	Family	IC50 ( $\mu\text{M}$ )	Reference no.
Benzoic acid,3-[2-(acetyloxy)-3-methyl-3-buten-1-yl]-4,5-dihydroxy-,methyl ester	<i>Piper glabratum</i> Kunth	Piperaceae	34.0 (F32)	135
3-(Z)-Caffeoyllupeol	<i>Bruguiera parviflora</i> Wight	Rhizophoraceae	14.6 (K1)	136
Oleanolic acid	<i>Morinda lucida</i> Benth.	Rubiaceae	32.3 (CQS)	110
	<i>Satureja parvifolia</i> (Phil.) Epling	Lamiaceae	19.8 (3D7)	58
$\beta$ -Acetylolean-12-en-28-olic acid	<i>Prismatomeris fragrans</i> Geddes	Rubiaceae	11.9 (K1)	137
Methyl uguenenoate	<i>Vepris uguenensis</i> Engl.	Rutaceae	20.7 (3D7)	65
			27.5 (FCM29)	
3,4-Dihydro-methylcatalpol	<i>Scrophularia lepidota</i> Boiss.	Scrophulariaceae	>50 (K1)	138
Scrolepidoside	<i>Scrophularia lepidota</i> Boiss.	Scrophulariaceae	>50 (K1)	138
Catalpol	<i>Scrophularia lepidota</i> Boiss.	Scrophulariaceae	>50 (K1)	138
6-O-Methylcatalpol	<i>Scrophularia lepidota</i> Boiss.	Scrophulariaceae	>50 (K1)	138
Sinuato	<i>Scrophularia lepidota</i> Boiss.	Scrophulariaceae	>50 (K1)	138
aucubin	<i>Scrophularia lepidota</i> Boiss.	Scrophulariaceae	>50 (K1)	138
6-O- $\beta$ -D-Xylopyranosylaucubin	<i>Scrophularia lepidota</i> Boiss.	Scrophulariaceae	>50 (K1)	138
Ajugol	<i>Scrophularia lepidota</i> Boiss.	Scrophulariaceae	>50 (K1)	138
Ajugoside	<i>Scrophularia lepidota</i> Boiss.	Scrophulariaceae	>50 (K1)	138
Iridoid-related aglycone	<i>Scrophularia lepidota</i> Boiss.	Scrophulariaceae	>50 (K1)	138
3-O- $\beta$ -D-glucopyranosylpseudojubenin	<i>Bacopa monniera</i> Wettst	Scrophulariaceae	>50	139
Bacoside A3	<i>Bacopa monniera</i> Wettst	Scrophulariaceae	>50	139
Bacoside A6	<i>Bacopa monniera</i> Wettst	Scrophulariaceae	>50	139
Bacopaside II	<i>Bacopa monniera</i> Wettst	Scrophulariaceae	>50	139
Bacopaside IV	<i>Bacopa monniera</i> Wettst	Scrophulariaceae	>50	139
Bacopaside V	<i>Bacopa monniera</i> Wettst	Scrophulariaceae	>50	139
Bacopaside X	<i>Bacopa monniera</i> Wettst	Scrophulariaceae	>50	139
Bacopaside N2	<i>Bacopa monniera</i> Wettst	Scrophulariaceae	>50	139
Bacopasaponin C	<i>Bacopa monniera</i> Wettst	Scrophulariaceae	>50	139
Bacopasaponin G	<i>Bacopa monniera</i> Wettst	Scrophulariaceae	>50	139
Buddlejasaponin	<i>Scrophularia cryptophilla</i> Boiss. & Heldr.	Scrophulariaceae	24.5 (K1)	140
3 $\alpha$ ,20-Lupandiol	<i>Grewia bilamellata</i> Gagnep.	Tiliaceae	19.8 (D6)	41
			19.1 (W2)	
Aulacocarpin A	<i>Aframomum zambesiacum</i> K. Schum	Zingiberaceae	13.7 (FcB1)	89
Zambesiacolactone A	<i>Aframomum zambesiacum</i> K. Schum	Zingiberaceae	17.2 (FcB1)	89
Zambesiacolactone B	<i>Aframomum zambesiacum</i> K. Schum	Zingiberaceae	15.5 (FcB1)	89
(1R,2S,5S,9S,10S,11R,13R)-1,2,11-trihydroxypimara-8(14),15-diene	<i>Kaempferia marginata</i> Carey	Zingiberaceae	27.5 (K1)	94


and 0.24  $\mu\text{M}$ , respectively, on NF54, and 1.43, 1.73, 1.88, 1.72 and 1.79, respectively, on ENT30. Four of these alkaloids were further evaluated intraperitoneally for activity against *P. berghei* *in vivo*. The alkaloids showed chemosuppression percentages of parasitaemia in mice ranging from 43 to 72% at 20 mg/kg per day. The use of the extracts of *A. gummifera* for the treatment of malaria in traditional medicine seems to have a scientific basis.<sup>[141]</sup> Vertine (**242**) and epi-lyfoline (**243**) were isolated from *Heimia salicifolia* Link & Otto (Lythraceae) and showed antimalarial activity with IC50 values of 10.9 and 6.7  $\mu\text{M}$ , respectively.<sup>[142]</sup> A new lycorine derivative LT1 (**244**), 1-O-(30S)-hydroxybutanoyllycorine was isolated from the aerial part and bulbs of *Lycoris traubii* Hayward (Amaryllidaceae). It showed significant activity against FCR-3 and K1 strains (IC50 = 1.2 and 1.6  $\mu\text{M}$ , respectively).<sup>[143]</sup>

Figure 13 shows alkaloids derived from ornithine and lysine with moderate or promising activity *in vitro* against various strains of *P. falciparum*.

### Phenylalanine and tyrosine derivatives

A novel alkaloid with an unprecedented tricyclic skeleton, cassiarin A (**245**), was isolated from the leaves of *Cassia*

*siamea* Lam. (Leguminosae). It showed promising antiplasmodial activity (IC50 23.5 nM).<sup>[144]</sup> *Zanthoxylum rhoifolium* Lam. (Rutaceae) bark is a medicinal plant traditionally used in French Guiana to treat and prevent malaria. Bioassay-guided fractionation of the alkaloid extract yielded three benzophenanthridine alkaloids. Dihydroneitidine (**246**) was evaluated against FcB1 (IC50 = 4.9  $\mu\text{M}$ ).<sup>[145]</sup> Biologically guided fractionation of the methanolic extract of the roots of *Zanthoxylum flavum* Vahl (Rutaceae) led to the isolation of two alkaloids, dihydrochelerythrin (**247**) and chelerythrine acetate (**248**). Compound **247** was only moderately active with an IC50 of 10.6  $\mu\text{M}$  on D6. Compound **248** displayed greater activity with IC50 values of 5.7 and 3.4  $\mu\text{M}$  on D6 and W2 strains, respectively.<sup>[40]</sup> Decoction of *Strychnopsis thouarsii* Baill. (Menispermaceae) is used in Malagasy traditional medicine to fight malaria. It has been shown that this traditional remedy prevents malaria infection by targeting *Plasmodium* at its early liver stage. Bioassay-guided fractionation of *S. thouarsii* stem barks extracts, using a rodent *Plasmodium yoelii* liver stage parasites inhibition assay, led to the isolation of the new morphinan alkaloid tazopsine (**249**) together with sinococuline (**250**).



**Figure 13** Alkaloids derived from ornithine and lysine with moderate or promising activity *in vitro* against various strains of *P. falciparum*


Compounds **249** and **250** exhibited selective inhibitory activity (SI > 13.8) against *P. yoelii* liver stage *in vitro*. Tazopsine showed the most potent inhibitory activity with an IC<sub>50</sub> value of 3.1 μM. Sinococuline was both slightly less active (IC<sub>50</sub> = 4.5 μM) and less toxic.<sup>[146,147]</sup> Dehydroroemerine (**251**) was isolated from *Stephania rotunda* Lour. (Menispermaceae) and was found to be the most active against W2 with an IC<sub>50</sub> value of 0.36 μM.<sup>[148]</sup> Two new alkaloids, desmorostratine (**252**) and discretine *N*-oxide (**253**), were isolated from the stem bark of *Desmos rostrata* (Merr. & Chun) P.T.Li (Annonaceae), together with five known alkaloids including discretine (**254**) and dehydrodiscretine (**255**). Compounds **253**, **254** and **255** inhibited FcB1 with IC<sub>50</sub> values of 4.2, 1.6 and 0.9 μM, respectively, and showed weak cytotoxic activity. On the other hand, **252** had an IC<sub>50</sub> of 3.6 μM but was also moderately toxic (IC<sub>50</sub> = 2.4 μM).<sup>[149]</sup> A new dimeric aporphine alkaloid, bidebiline E (**256**), was isolated from the roots of *Polyalthia cerasoides* (Roxb.) Bedd. (Annonaceae). It exhibited antimalarial activity against K1 (IC<sub>50</sub> = 7.7 μM).<sup>[150]</sup> Three alkaloids, melosmine (**257**), atherospermidine (**258**) and isomoschatoline (**259**), isolated from the ethyl acetate extract of the stem of *Rollinia pittieri* Saff. and *Pseudomalmea boyacana* (J.F. Macbr.) Chatrou (Annonaceae) exhibited moderate activity with IC<sub>50</sub> values of 12.2, 10.6 and 10.9 μM, respectively, against F32, and 10.4, 12.8 and 27.8 μM against W2.<sup>[151]</sup> Two bioactive bisbenzylisoquinolines, magnoline (**260**) and magnolamine (**261**), were isolated from the leaves of *Michelia figo* (Lour.) Spreng (Magnoliaceae). Magnolamine showed a significant IC<sub>50</sub> of 1.28 μM on K1 and less than 0.16 μM on FCR3. Magnoline also inhibited both strains with an IC<sub>50</sub> of 1.51 μM on FCR3 and less than 0.16 μM on K1.<sup>[152]</sup> A new diastereoisomer of the bis-benzylisoquinoline alkaloid rodiasine, 1*S*,1'*R*-rodiasine (**262**), was isolated from *Pseudoxandra cuspidata* Maas (Annonaceae) bark, used in French Guiana as an antimalarial. The antimalarial activity of this bark was mostly due to **262** (IC<sub>50</sub> = 1.14 μM), which also displayed low cytotoxicity.<sup>[153]</sup> The antiplasmodial activity of *Triclisia saclexii* Diels (Menispermaceae) was investigated on 3D7 and W2 strains. Phytochemical analysis of the root tertiary alkaloids fraction yielded four major compounds, phaeanthine (**263**), *N*-methylapateline (**264**), 1,2-dehydroapateline (**265**)

and 1,2-dehydrotelobine (**266**). They demonstrated antiplasmodial activity with IC<sub>50</sub> values of 1.72, 0.93, 1.39 and 12.4 μM, respectively, on 3D7, and 0.35, 1.10, 1.63 and 1.52 μM, respectively, on W2.<sup>[154]</sup> Three new fully dehydrogenated naphthylisoquinoline alkaloids, the 7,1'-coupled *ent*-dioncophylleine A (**267**), the 7,1'-coupled 5'-*O*-demethyl-*ent*-dioncophylleine A (**268**), and the 7,8'-linked dioncophylleine D (**269**), were isolated from the leaves of the liana *Ancistrocladus benomensis* Rischer & G.Bringmann (Ancistrocladaceae). Compounds **267** and **268** exhibited moderate antiplasmodial activity against K1 with IC<sub>50</sub> values of 10.5 and 8.6 μM, respectively. Compound **269** showed better activity with a low IC<sub>50</sub> (1.3 μM).<sup>[155]</sup> From the roots of a recently discovered *Ancistrocladus* taxon with close affinity to *Ancistrocladus congolensis* J. Léonard (Ancistrocladaceae), six new naphthylisoquinoline alkaloids, 5'-*O*-demethylhamatine (**270**), 5'-*O*-demethylhamatinine (**271**), 6-*O*-demethylancistroealaine A (**272**), 6,5'-*O*,*O*-didemethylancistroealaine A (**273**), 5-epi-6-*O*-methylancistrobertsonine A (**274**), and 5-epi-4'-*O*-demethylancistrobertsonine C (**275**), and also the known 6-*O*-demethylancistrobrevine A (**276**), were isolated. All of the naphthylisoquinoline alkaloids tested were found to exhibit antiplasmodial activity against K1 strain: IC<sub>50</sub> = 2.5, 7.2, 4.4, 5.4, 4.4, 6.2 and 5.0 μM, respectively.<sup>[156]</sup>

Figure 14 shows alkaloids derived from phenylalanine and tyrosine with moderate or promising activity *in vitro* against various strains of *P. falciparum*.

### Tryptophane derivatives

Bioassay-guided fractionation of the antimalarial CHCl<sub>3</sub> extract of the dried stem of *Nauclea orientalis* (L.) L. (Rubiaceae) resulted in the isolation of a novel tetrahydro-β-carboline monoterpene alkaloid glucoside, naucleaorine (**277**), which showed activity with an IC<sub>50</sub> of 6.9 μM on D6 and 8.0 on W2.<sup>[112]</sup> An indole alkaloid, naucleofficine A (**278**), was isolated from the stems (with bark) of *Nauclea officinalis* Pierre ex Pitard (Rubiaceae). Compound **278** exhibited moderate antimalarial activity against FCC1-HN with an IC<sub>50</sub> value of 9.7 μM and no cytotoxic effect was observed.<sup>[157]</sup> A dimeric indoloquinoline alkaloid, biscryptolepine (**279**), originally obtained from the plant *Cryptolepis sanguinolenta* (Lindl.) Schltr. (Asclepiadaceae), showed



**Figure 14** Alkaloids derived from phenylalanine and tyrosine with moderate or promising activity *in vitro* against various strains of *P. falciparum*



good antiplasmodial activity against K1 (IC<sub>50</sub> of 0.27  $\mu\text{M}$ ), while the cytotoxicity (L6 cells) was 13.62  $\mu\text{M}$ .<sup>[158]</sup> Biologically guided fractionation of the methanolic extract of the roots of *Zanthoxylum flavum* Vahl (Rutaceae) led to the isolation of dihydrorutaecarpine (**280**) which displayed higher activity with IC<sub>50</sub> values of 5.7 and 3.4  $\mu\text{M}$  on D6 and W2 strains, respectively.<sup>[40]</sup> Isosungucine is a quasi-symmetric bisindolomonoterpenoid alkaloid isolated from the roots of *Strychnos icaja* Baill. (Loganiaceae). The antimalarial activity against the *P. vinckei petteri* murine strain was determined *in vivo*. In the Peters 4-day suppressive test, the compound suppressed parasitaemia by almost 50% on day 4 at a dose of 30 mg/kg given intraperitoneally.<sup>[159]</sup>

Figure 15 shows alkaloids derived from tryptophane with moderate or promising activity *in vitro* against various strains of *P. falciparum*.

### Steroidal alkaloids

Bioguided phytochemical investigation of *Sarcococca hookeriana* Baill. (Buxaceae) yielded two new pregnane-type steroidal alkaloids hookerianamide H (**281**) and hookerianamide I (**282**), along with three known alkaloids, Namethylepipachysamine D (**283**), sarcovagine C (**284**) and dictyophlebine (**285**). These compounds showed reasonable antiplasmodial activity with IC<sub>50</sub> values of 3.5, 6.6, 10.3, 3.4

and 2.4, respectively, against W2 strain.<sup>[160]</sup> Bioassay-guided fractionation of the EtOH extract of the stem bark of *Funtumia elastica* Stapf (Apocynaceae) resulted in the isolation of four steroidal alkaloids, holarrhetine (**286**), conessine (**287**), holarrhesine (**288**) and isoconessimine (**289**). They exhibited antiplasmodial activity against FcB1 strain with IC<sub>50</sub> values of 1.13, 1.04, 0.97 and 3.39  $\mu\text{M}$  and weak cytotoxicity (L-6 cell line) with IC<sub>50</sub> values of 5.13, 14.6, 7.49 and 36.55  $\mu\text{M}$ .<sup>[161]</sup>

Figure 16 shows steroidal alkaloids with moderate or promising activity *in vitro* against various strains of *P. falciparum*.

### Other N-containing compounds

Bioassay-guided fractionation of the EtOAc extract of the roots of Thai *Ziziphus oenoplia* L. Mill. var. *brunoniana* (Rhamnaceae) resulted in the isolation of two new 13-membered cyclopeptide alkaloids of the 5(13) type, ziziphine N (**290**) and Q (**291**), which exhibited notable antiplasmodial activity with IC<sub>50</sub> values of 6.4 and 5.9  $\mu\text{M}$ , respectively.<sup>[162]</sup> Three alkamides were isolated from the leaves of *Zanthoxylum syncarpum* Tul. (Rutaceae). Compound **292**, the racemic form of the known compound syncarpamide, showed moderate antiplasmodial activity, with IC<sub>50</sub> values of 4.2 and 6.1  $\mu\text{M}$  against D6 and W2 clones, respectively. Cytotoxicity was evaluated at an IC<sub>50</sub> of 10.3  $\mu\text{M}$ .<sup>[163]</sup>




Figure 15 Alkaloids derived from tryptophane with moderate or promising activity *in vitro* against various strains of *P. falciparum*





Figure 16 Steroidal alkaloids with moderate or promising activity *in vitro* against various strains of *P. falciparum*





290.  $R_1 = R_2 = \text{CH}_3$ ,  $R_3 = \text{CH}_2\text{CH}(\text{CH}_3)_2$

291.  $R_1 = R_2 = \text{CH}_3$ ,  $R_3 = \text{CH}(\text{CH}_3)_2$

**Figure 17** Other *N*-containing compounds with moderate or promising activity *in vitro* against various strains of *P. falciparum*

**Table 4** Alkaloids presenting low or no activity *in vitro* against various strains of *P. falciparum*

Compound	Plant	Family	IC <sub>50</sub> ( $\mu\text{M}$ )	Reference no.
Predicentrine	<i>Desmos rostrata</i> (Merr. & Chun) P.T.Li	Annonaceae	27.8 (FcB1)	149
Aristolactam	<i>Desmos rostrata</i> (Merr. & Chun) P.T.Li	Annonaceae	>37 (FcB1)	149
Octadeca-9,11,13-triynoic acid	<i>Polyalthia cerasoides</i> (Roxb.) Bedd.	Annonaceae	18.4 (K1)	150
N-methylouregidione	<i>Pseuduvaria setosa</i> (King) J. Sinclair.	Annonaceae	>50 (K1)	164
Ouregidione	<i>Pseuduvaria setosa</i> (King) J. Sinclair.	Annonaceae	>50 (K1)	164
Oxostephanine	<i>Pseuduvaria setosa</i> (King) J. Sinclair.	Annonaceae	>50 (K1)	164
Ellipticine	<i>Aspidosperma vargasii</i> A.DC.	Apocynaceae	>50 (K1)	49
	<i>Aspidosperma desmanthum</i> Benth.			
Aspidocarpine	<i>Aspidosperma vargasii</i> A.DC.	Apocynaceae	19 (K1)	49
	<i>Aspidosperma desmanthum</i> Benth.			
Integerrimide A	<i>Jatropha integerrima</i> Jacq.	Euphorbiaceae	>50 (K1)	165
Integerrimide A	<i>Jatropha integerrima</i> Jacq.	Euphorbiaceae	>50 (K1)	165
(-)-Phoebescoretechiniine	<i>Phoebe scortechinii</i> (Gamb.) Kochummen	Lauraceae	Nt	166
Phoebegrandine A	<i>Phoebe scortechinii</i> (Gamb.) Kochummen	Lauraceae	Alkaloidal extract =	166
Phoebegrandine B	<i>Phoebe scortechinii</i> (Gamb.) Kochummen	Lauraceae	6.11 g/ml (Gombak A)	166
Tetrahydropronuciferine	<i>Phoebe scortechinii</i> (Gamb.) Kochummen	Lauraceae	0.69 g/ml (D10)	166
Cassiarin B	<i>Cassia siamea</i> Lam.	Leguminosae	22.0	144
10-Epi-tazopsine	<i>Strychnopsis thouarsii</i> Baill.	Menispermaceae	16.1 ( <i>P. yoelii</i> )	147
Tetrahydropalmatine	<i>Stephania rotunda</i> Lour.	Menispermaceae	32.6 (W2)	148
Xylopinine	<i>Stephania rotunda</i> Lour.	Menispermaceae	>50 (W2)	148
Cycleanine <i>N</i> -oxide	<i>Epinetrum villosum</i> (Exell) Troupin	Menispermaceae	13.5 (FcB1)	167
Epimethoxy-naucleaorine	<i>Nauclea orientalis</i> (L.) L.	Rubiaceae	12.4 (D6)	112
			13.2 (W2)	
Naucleofficine B	<i>Nauclea officinalis</i> Pierre ex Pitard	Rubiaceae	42.1 (FCC1-HN)	157
Naucleofficine C	<i>Nauclea officinalis</i> Pierre ex Pitard	Rubiaceae	40.5 (FCC1-HN)	157
Naucleofficine D	<i>Nauclea officinalis</i> Pierre ex Pitard	Rubiaceae	39.8 (FCC1-HN)	157
Naucleofficine E	<i>Nauclea officinalis</i> Pierre ex Pitard	Rubiaceae	38.3 (FCC1-HN)	157
Uncarine D	<i>Mitragyna inermis</i> (Willd.) Kuntze	Rubiaceae	46.2 (W2)	168
Maculosidine	<i>Vepris uguenensis</i> Engl.	Rutaceae	>50 (3D7/FCM29)	65
Flindersiamine	<i>Esenbeckia febrifuga</i> A.Juss.	Rutaceae	>50 (3D7/W2)	169
r-Fagarine	<i>Esenbeckia febrifuga</i> A.Juss.	Rutaceae	>50 (3D7/W2)	169
Rutaevine	<i>Esenbeckia febrifuga</i> A.Juss.	Rutaceae	>50 (3D7/W2)	169
Tegerrardin A	<i>Teclea gerrardii</i> Verdoorn	Rutaceae	12.3 (D10)	170
Dihydroavicine	<i>Zanthoxylum rhoifolium</i> Lam.	Rutaceae	33.0 (FcB1)	145
Fagaridine	<i>Zanthoxylum rhoifolium</i> Lam.	Rutaceae	38.0 (FcB1)	145
7,9-Dimethoxy-2,3-methylenedioxybenzophenanthridine	<i>Zanthoxylum rubescens</i> Planch. ex Hook.	Rutaceae	>50 (3D7/FCM29)	171
Bis[6-(5,6-dihydrochelerythrinyl)] ether	<i>Zanthoxylum rubescens</i> Planch. ex Hook.	Rutaceae	15.3 (3D7)	171
			14.9 (FCM29)	
Zanthomamide	<i>Zanthoxylum rubescens</i> Planch. ex Hook.	Rutaceae	>50 (3D7/FCM29)	171
Lemairamide	<i>Zanthoxylum rubescens</i> Planch. ex Hook.	Rutaceae	>50 (3D7/FCM29)	171
Harmine	<i>Peganum harmala</i> L.	Zygophyllaceae	37.7	172
Harmaline	<i>Peganum harmala</i> L.	Zygophyllaceae	>50	172


Figure 17 shows other *N*-containing compounds with moderate or promising activity *in vitro* against various strains of *P. falciparum*.

Table 4 gives the tested alkaloids presenting low or no activity *in vitro* against various strains of *P. falciparum*.<sup>[49,65,112,144,145,147–150,157,164–172]</sup>

### Other metabolites

Two compounds were isolated from the stems and leaves of *Croton lobatus* L. (Euphorbiaceae), a medicinal plant used in western Africa in traditional folk medicine to cure malaria, pregnancy troubles and dysentery. (*Z,Z,Z*)-9,12,15-Octadecatrienoic acid methyl ester (**293**) and 8,11,17,21-tetramethyl-(*E,E,E,E*)-8,10,17,21-tetraentetracosanoic acid (**294**) showed some antiplasmodial activity on K1 strain, with IC<sub>50</sub> values of 10.9 and 9.1  $\mu\text{M}$ , respectively, and SI values of 18.4 and over 20, respectively.<sup>[57]</sup> Bioassay-directed fractionation of the  $\text{CHCl}_3$  extract of the dried stems of *Rourea minor* (Gaertn.) Aubl. (Connoraceae) liana led to the isolation of rouremin (**295**), as well as a known

compound, 1-(26-hydroxyhexacosanoyl)-glycerol (**296**). They showed activity with IC<sub>50</sub> values of 5.1 and 9.5  $\mu\text{M}$  against D6 and 4.5 and 12.7  $\mu\text{M}$  against W2 strains.<sup>[45]</sup> Cryptophilic acid C (**297**) isolated from the resin of *Scrophularia cryptophilla* Boiss. & Heldr. (Scrophulariaceae) showed antimalarial activity with an IC<sub>50</sub> of 4.1  $\mu\text{M}$ .<sup>[140]</sup> Three new ether diglycosides (**298–300**), namely matayosides A–B and D, were isolated from the root bark of *Matayba guianensis* Aubl. (Sapindaceae), a plant exhibiting antiplasmodial activity. These compounds were evaluated for their antiplasmodial activity against FcB1 with IC<sub>50</sub> values of 8.1, 4.7 and 3.5  $\mu\text{M}$ .<sup>[173]</sup> Guieranone A (**301**), a naphthyl butenone, was purified from the leaves and roots of *Guiera senegalensis* J. F. Gmel. (Combretaceae). Guieranone A showed notable antiplasmodial activity (IC<sub>50</sub> = 4.1  $\mu\text{M}$ ) associated with high cytotoxicity towards human monocytes.<sup>[174]</sup> From the petroleum ether extract of the root bark of *Cussonia zimmermannii* Harms (Araliaceae), two polyacetylenes were isolated, 8-hydroxyheptadeca-1-ene-4,6-diyn-3-yl ethanoate (**302**) and 16-acetoxy-11-hydroxyoctadeca-17-ene-12,14-diynyleth-



**Figure 18** Other metabolites with moderate or promising activity *in vitro* against various strains of *P. falciparum*

**Table 5** Other metabolites presenting low or no activity *in vitro* against various strains of *P. falciparum*

Compound	Plant	Family	IC <sub>50</sub> ( $\mu\text{M}$ )	Reference no.
19-(2-Furyl)nonadeca-5,7-dienoic acid	<i>Polyalthia evecta</i> Finet & Gagnep.	Annonaceae	>50 (K1)	176
8-Hydroxyheptadeca-4,6-dien-3-yl ethanoate	<i>Cussonia zimmermannii</i> Harms	Arialaceae	19	175
Scleropyric acid	<i>Scleropyrum wallichianum</i> Arn.	Santalaceae	27.3 (K1)	177
Matayoside C	<i>Matayba guianensis</i> Aubl.	Sapindaceae	11.7 (FcB1)	173

anoate (**303**). They were active with IC<sub>50</sub> values of 1.4 and 2.2  $\mu\text{M}$ , respectively.<sup>[175]</sup>


Figure 18 shows other metabolites with moderate or promising activity *in vitro* against various strains of *P. falciparum*.

Table 5 gives the tested other metabolites presenting low or no activity *in vitro* against various strains of *P. falciparum*.<sup>[173,175–177]</sup>

## Discussion

Several plants are used in traditional medicine in many countries for the treatment of malaria or symptoms of the

disease. This review focused on publications from 2005 to the end of 2008, and shows that the ethnopharmacological and bio-guided fractionation approaches have led to the isolation of some promising new antimalarial compounds with a large variety of structures. About 480 compounds were isolated and evaluated for antimalarial activity *in vitro*. These compounds possess low ( $11 < \text{IC}_{50} \leq 50 \mu\text{M}$ ), moderate ( $2 < \text{IC}_{50} \leq 11 \mu\text{M}$ ) or promising ( $\text{IC}_{50} \leq 2 \mu\text{M}$ ) activity *in vitro* against various strains of *P. falciparum*, which is responsible for the most severe cases of malaria. Some of the active compounds were also tested for their cytotoxicity but only a few of them were tested for their antimalarial activity *in vivo*. Nevertheless, in these cases, the



**Figure 19** Number of compounds with high ( $\text{IC}_{50} \leq 2 \mu\text{M}$ ) or moderate ( $2 < \text{IC}_{50} \leq 11 \mu\text{M}$ ) activity *in vitro* against various strains of *P. falciparum*, classified according to their chemical classes


promising in-vitro activity of several compounds from different chemical classes could be confirmed by in-vivo tests. Among them are several phenolic compounds: a chalcone, two stilbenes and a coumarine; terpenoids: a sesqui- and two tri-terpenes; and alkaloids: four ornithine/lysine and one tryptophane derivative. Moreover, several compounds already isolated and tested *in vitro* before 2005 were evaluated for in-vivo antimalarial activity, for example, simalikalactone D with an ED<sub>50</sub> of 3.7 mg/kg per day orally.<sup>[178]</sup>

In traditional medicine, it is often the aqueous decoction of the plant that is used as a treatment to fight malaria. In the present study, we observed that most of the promising compounds are aglycones or other lipophilic compounds. Nevertheless, it is known that other compounds present in crude extracts, for example flavonoid heterosides or saponosides, could promote the solubilisation of more lipophilic compounds in water, supporting the implication of lipophilic compounds in the activity of water extracts.<sup>[179]</sup> We also observed that several extracts that are used in traditional medicine do not possess interesting activity *in vitro*. This may be explained by the fact that some compounds can only become active after metabolisation so they do not display

good activity *in vitro*. This absence of activity could also be explained by an action on another stage of the *Plasmodium* (sporozoites, gametocytes), the in-vitro test focusing on the erythrocytic stage of the parasite (trophozoites, schizonts and merozoites). Moreover, the plants can also be used because they are effective on other symptoms of the illness such as fever, vomiting, abdominal pains and cephalgias. It can also be noted that compounds that have weak activity could promote their potential with other metabolites through a synergy of action as is often the case with crude extracts possessing several active molecules or adjuvants.

Among the compounds we reviewed, only a few of them exhibited high activity and should be considered as lead compounds for further investigation (Figures 19 and 20). When considering chemical classes and families from which active compounds ( $IC_{50} \leq 2 \mu M$ ) were isolated, we observed that they belong mainly to the diterpenes and alkaloids (between 20 and 30 compounds), flavonoids and sesquiterpenes (between 10 and 15 compounds), and they were isolated from several plant families. Nevertheless, some chemotaxonomical correlations could be observed.

Figures 19 and 20 show the number of compounds with high ( $IC_{50} \leq 2 \mu M$ ) or moderate ( $2 < IC_{50} \leq 11 \mu M$ ) activity



**Figure 20** Number of compounds with high ( $IC_{50} \leq 2 \mu M$ ) or moderate ( $2 < IC_{50} \leq 11 \mu M$ ) activity *in vitro* against various strains of *P. falciparum*, classified according to the plant family from which they were isolated

*in vitro* against various strains of *P. falciparum* classified according to their chemical classes and according to the plant family from which they were isolated, respectively.

The majority of active diterpenes were isolated from the Caesalpinaceae family and particularly from *Caesalpinia crista* L. This plant should be further investigated and its several active cassane- and norcassane-compounds tested clinically.

There are different sources of alkaloids that have strong activity, Leguminosae, Apocynaceae, Menispermaceae and Annonaceae. Leguminosae is one of the greatest families with active alkaloids and flavonoids. The main active alkaloids were derivatives from phenylalanine and tyrosine.

Sesquiterpene lactones, like the famous artemisinin from *Artemisia annua*, have been isolated from a few plants of the Asteraceae family but most of them are also highly cytotoxic. Clusiaceae is an interesting plant family which has allowed the isolation of active quinones and xanthenes. A review that covers antimalarial plant metabolites from 1990 to 2000 highlighted the importance of this family with five new active xanthenes.<sup>[3]</sup>

Moraceae is a family with several interesting compounds and particularly flavonoids from *A. champeden* Spreng. In earlier years, active stilbenes were isolated from *Artocarpus integer*.<sup>[3]</sup> When examining the structures of the most active flavonoids from the last four years, we observed that most of them are prenylated compounds, thus being generally more lipophilic than non-prenylated ones. This higher lipophilicity probably increases the resorption through cell membrane, which may explain the higher activity often observed. Moreover, common dietary flavonoids possess antimalarial activity with IC50 values of between 11 and 66  $\mu\text{M}$ .<sup>[180]</sup>

From the 1990s, the major compounds that showed promising activity were alkaloids from Apocynaceae and Loganiaceae, flavonoids from Fabaceae, quinones from Asphodelaceae, tannins and triterpenes from Combretaceae and butenolides from Monimiaceae.<sup>[2]</sup> In 2003, a review showed that three classes of secondary plant metabolites were mostly responsible for antimalarial activity: alkaloids, quassinoids and sesquiterpenes lactones.<sup>[5]</sup> Indeed, non-nitrogenous natural products were also shown to be promising.<sup>[7]</sup> In the present study, very active alkaloids and quinones were also isolated from Apocynaceae and Asphodelaceae families, respectively, showing the potential of these families, but other families must not be underestimated and are also worthy of evaluation.

In traditional medicine, essential oils containing mono- and sesquiterpenes are also used as antimalarials. Several studies showed the antimalarial activity of these oils from different plant species. For example, the essential oil of *Salvia repens* exhibited an IC50 of 1.7  $\mu\text{g}/\text{ml}$  with  $\beta$ -phellandrene,  $\beta$ -caryophyllene, limonene and camphor as major components.<sup>[181]</sup> Nevertheless, the effect of isolated compounds was not evaluated and so they are not listed here.

## Conclusions

A large number of antimalarial compounds with a wide variety of structures have been isolated from plants and can play a role in the development of new antimalarial drugs. Ethnopharmacological approach appears to be a promising

way to find plant metabolites that could be used as templates for designing new derivatives with improved properties.

## Declarations

### Conflict of interest

Michel Frédéricich is a senior research associate from the FNRS.

### Funding

This review received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

## References

1. Phillipson JD. Natural products as drugs. *Trans R Soc Trop Med Hyg* 1994; 88: 17–19.
2. Fournet A, Munoz V. Natural products as trypanocidal, antileishmanial and antimalarial drugs. *Curr Top Med Chem* 2002; 2: 1215–1238.
3. Schwikkard S, van Heerden FR. Antimalarial activity of plant metabolites. *Nat Prod Rep* 2002; 19: 675–692.
4. Caniato R, Puricelli L. Review: natural antimalarial agents (1995–2001). *Crit Rev Plant Sci* 2003; 22: 79–105.
5. Saxena S et al. Antimalarial agents from plant sources. *Curr Sci* 2003; 85: 1314–1329.
6. Wright CW. Plant derived antimalarial agents: new leads and challenges. *Phytochem Rev* 2005; 4: 55–61.
7. Bilia AR. Non-nitrogenous plant-derived constituents with antiplasmodial activity. *Nat Prod Commun* 2006; 1: 1181–1204.
8. Mambu L, Grellier P. Antimalarial compounds from traditionally used medicinal plants. In: Colegate SM, Molyneux RJ, eds. *Bioactive Natural Products. Detection, Isolation and Structural Determination*, 2nd edn. Boca Raton, USA: CRC press, 2008: 491–529.
9. Kaur K et al. Antimalarials from nature. *Bioorg Med Chem* 2009; 17: 3229–3256.
10. Ramanandralbe V et al. Antiplasmodial phenolic compounds from *Piptadenia pervillei*. *Planta Med* 2008; 74: 417–421.
11. Boonphong S et al. Bioactive compounds from *Bauhinia purpurea* possessing antimalarial, antimycobacterial, antifungal, anti-inflammatory, and cytotoxic activities. *J Nat Prod* 2007; 70: 795–801.
12. Joseph CC et al. A novel antiplasmodial 3',5'-diformylchalcone and other constituents of *Friesodielsia obovata*. *Nat Prod Res* 2007; 21: 1009–1015.
13. Namdaung U et al. Bioactive constituents of the root bark of *Artocarpus rigidus* subsp. *rigidus*. *Chem Pharm Bull* 2006; 54: 1433–1436.
14. Widyawaruyanti A et al. New prenylated flavones from *Artocarpus champeden*, and their antimalarial activity *in vitro*. *J Nat Med* 2007; 61: 410–413.
15. Boonphong S et al. Antitubercular and antiplasmodial prenylated flavones from the roots of *Artocarpus altilis*. *Chiang Mai J Sci* 2007; 34: 339–344.
16. Khaomek P et al. *In vitro* antimalarial activity of prenylated flavonoids from *Erythrina fusca*. *J Nat Med* 2008; 62: 217–220.
17. Andayi AW et al. Antiplasmodial flavonoids from *Erythrina saculeuxii*. *Planta Med* 2006; 72: 187–189.
18. Rukachaisirikul T et al. Biological activities of the chemical constituents of *Erythrina stricta* and *Erythrina subumbrans*. *Arch Pharm Res* 2007; 30: 1398–1403.

19. Rukachaisirikul T *et al.* Erythrina alkaloids and a pterocarpan from the bark of *Erythrina subumbrans*. *J Nat Prod* 2008; 71: 156–158.
20. Radwan MM *et al.* Non-cannabinoid constituents from a high potency *Cannabis sativa* variety. *Phytochemistry* 2008; 69: 2627–2633.
21. Ichino C *et al.* Antimalarial activity of biflavonoids from *Ochna integerrima*. *Planta Med* 2006; 72: 611–614.
22. Weniger B *et al.* A bioactive biflavonoid from *Camposperma panamense*. *Fitoterapia* 2004; 75: 764–767.
23. Weniger B *et al.* Comparative antiplasmodial, leishmanicidal and antitrypanosomal activities of several biflavonoids. *Phytomedicine* 2006; 13: 176–180.
24. Mbwambo ZH *et al.* Antiparasitic activity of some xanthenes and biflavonoids from the root bark of *Garcinia livingstonei*. *J Nat Prod* 2006; 69: 369–372.
25. Midiwo JO *et al.* The first 9-hydroxyhomoisoflavanone, and antiplasmodial chalcones, from the aerial exudates of *Polygonum senegalense*. *Arkivoc* 2007; ix: 21–27.
26. Portet B *et al.* Activity-guided isolation of antiplasmodial dihydrochalcones and flavanones from *Piper hostmannianum* var. *berbicense*. *Phytochemistry* 2007; 68: 1312–1320.
27. Narender T *et al.* Prenylated chalcones isolated from *Crotalaria* genus inhibits *in vitro* growth of the human malaria parasite *Plasmodium falciparum*. *Bioorg Med Chem Lett* 2005; 15: 2453–2455.
28. Ngameni B *et al.* Antimalarial prenylated chalcones from the twigs of *Dorstenia barteri* var. *subtriangularis*. *Arkivoc* 2007; xiii: 116–123.
29. Argotte-Ramos R *et al.* Antimalarial 4-phenylcoumarins from the stem bark of *Hintonia latiflora*. *J Nat Prod* 2006; 69: 1442–1444.
30. Azebaze AGB *et al.* Prenylated xanthone derivatives with antiplasmodial activity from *Allanblackia monticola* Staner LC. *Chem Pharm Bull* 2006; 54: 111–113.
31. Azebaze AGB *et al.* Antimalarial and vasorelaxant constituents of the leaves of *Allanblackia monticola* (Guttiferae). *Ann Trop Med Parasitol* 2007; 101: 23–30.
32. Laphookhieo S *et al.* Cytotoxic and antimalarial prenylated xanthenes from *Cratogeomys cochinchinense*. *Chem Pharm Bull* 2006; 54: 745–747.
33. Ngouela S *et al.* Anti-plasmodial and antioxidant activities of constituents of the seed shells of *Symphonia globulifera* Linn f. *Phytochemistry* 2006; 67: 302–306.
34. Karan M *et al.* Phytochemical and antimalarial studies on *Swertia alata* Royale. *Acta Hort* 2005; 675: 139–145.
35. Lannang AM *et al.* Antimalarial compounds from the root bark of *Garcinia polyantha* Oliv. *J Antibiot* 2008; 61: 518–523.
36. Son IH *et al.* Antiplasmodial activity of novel stilbene derivatives isolated from *Parthenocissus tricuspidata* from South Korea. *Parasitol Res* 2007; 101: 237–241.
37. Park WH *et al.* Antimalarial activity of a new stilbene glycoside from *Parthenocissus tricuspidata* in mice. *Antimicrob Agents Chemother* 2008; 52: 3451–3453.
38. Lee SI *et al.* Antimalarial activity of a stilbene glycoside from *Pleuropterus ciliinervis*. *Ann Trop Med Parasitol* 2008; 102: 181–184.
39. Moon HI, Sim J. Antimalarial activity in mice of resveratrol derivative from *Pleuropterus ciliinervis*. *Ann Trop Med Parasitol* 2008; 102: 447–450.
40. Ross SA *et al.* Constituents of *Zanthoxylum flavum* and their antioxidant and antimalarial activities. *Nat Prod Commun* 2008; 3: 791–794.
41. Ma CY *et al.* Antimalarial compounds from *Grewia bilamellata*. *J Nat Prod* 2006; 69: 346–350.
42. Subeki S *et al.* Anti-babesial and anti-plasmodial compounds from *Phyllanthus niruri*. *J Nat Prod* 2005; 68: 537–539.
43. Andrade-Neto VF *et al.* Antiplasmodial activity of aryltetralone lignans from *Holostylis reniformis*. *Antimicrob Agents Chemother* 2007; 51: 2346–2350.
44. da Silva AA *et al.* *In vitro* antileishmanial and antimalarial activities of tetrahydrofuran lignans isolated from *Nectandra megapotamica* (Lauraceae). *Phytother Res* 2008; 22: 1307–1310.
45. He ZD *et al.* Rourinoside and rouremin, antimalarial constituents from *Rourea minor*. *Phytochemistry* 2006; 67: 1378–1384.
46. Reddy MK *et al.* Antioxidant, antimalarial and antimicrobial activities of tannin-rich fractions, ellagitannins and phenolic acids from *Punica granatum* L. *Planta Med* 2007; 73: 461–467.
47. Lee SJ *et al.* Bioassay-guided isolation of antiplasmodial anacardic acids derivatives from the whole plants of *Viola websteri* Hemsl. *Parasitol Res* 2009; 104: 463–466.
48. Vega AS *et al.* Antimalarials and antioxidants compounds from *Piper tricuspe* (Piperaceae). *Pharmacologyonline* 2008; 1: 1–8.
49. Andrade-Neto VF *et al.* *In vitro* inhibition of *Plasmodium falciparum* by substances isolated from Amazonian antimalarial plants. *Mem Inst Oswaldo Cruz* 2007; 102: 359–365.
50. Oshimi S *et al.* Chrobisiamone A, a new bischromone from *Cassia siamea* and a biomimetic transformation of 5-acetyl-7-hydroxy-2-methylchromone into cassiarin A. *Bioorg Med Chem Lett* 2008; 18: 3761–3763.
51. Ahmed SA *et al.* Structure determination and absolute configuration of cannabichromanone derivatives from high potency *Cannabis sativa*. *Tetrahedron Lett* 2008; 49: 6050–6053.
52. Wirasathien L *et al.* Cytotoxic C-benzylated chalcone and other constituents of *Ellipeiopsis cherrevensis*. *Arch Pharm Res* 2006; 29: 497–502.
53. Abbas FA *et al.* Phytochemical and biological studies on Saudi *Commiphora opobalsamum* L. *Nat Prod Res* 2007; 21: 383–391.
54. Molinar-Toribio E *et al.* Antiprotozoal activity against *Plasmodium falciparum* and *Trypanosoma cruzi* of xanthenes isolated from *Chrysochlamys tenuis*. *Pharm Biol* 2006; 44: 550–553.
55. Vongvanich N *et al.* Combretastatins D-3 and D-4 new macrocyclic lactones from *Getonia floribunda*. *Planta Med* 2005; 71: 191–193.
56. Cao SG *et al.* Antiplasmodial activity of compounds from *Sloanea rhodantha* (Baker) Capuron var. *rhodantha* from the Madagascar rain forest. *Planta Med* 2006; 72: 1438–1440.
57. Attioua B *et al.* Antiplasmodial activity of constituents isolated from *Croton lobatus*. *Pharm Biol* 2007; 45: 263–266.
58. van Baren C *et al.* Triterpenic acids and flavonoids from *Satureja parvifolia*. Evaluation of their antiprotozoal activity. *Z Naturforsch C* 2006; 61: 189–192.
59. Radwan MM. Isoflavonoids from an Egyptian collection of *Colutea istria*. *Nat Prod Commun* 2008; 3: 1491–1494.
60. Kapingu MC *et al.* A novel isoflavonoid from *Milletia puguensis*. *Planta Med* 2006; 72: 1341–1343.
61. Abrantes M *et al.* Antiplasmodial activity of lignans and extracts from *Pycnanthus angolensis*. *Planta Med* 2008; 74: 1408–1412.
62. Torres-Mendoza D *et al.* Weakly antimalarial flavonol arabinofuranosides from *Calycolpus warszewiczianus*. *J Nat Prod* 2006; 69: 826–828.
63. Cimanga RK *et al.* Antimalarial activity of some extracts and isolated constituents from *Morinda morindoides* leaves. *J Nat Remedies* 2008; 8: 191–202.

64. Carroll AR *et al.* Prenylated dihydrochalcones from *Boronia bipinnata* that inhibit the malarial parasite enzyme target hemoglobinase II. *J Nat Prod* 2008; 71: 1479–1480.
65. Cheplogoi PK *et al.* An azole, an amide and a limonoid from *Vepris uguenensis* (Rutaceae). *Phytochemistry* 2008; 69: 1384–1388.
66. Tuntiwachwuttikul P *et al.* Chromones from the branches of *Harrisonia perforata*. *Chem Pharm Bull* 2006; 54: 44–47.
67. Tasdemir D *et al.* Antituberculous and antiprotozoal activities of primin, a natural benzoquinone: *in vitro* and *in vivo* studies. *Chem Biodivers* 2006; 3: 1230–1237.
68. Moein MR *et al.* Antileishmanial, antiplasmodial and cytotoxic activities of 12,16-dideoxy aegyptinone B from *Zhumeria majdae* Rech.f. & Wendelbo. *Phytother Res* 2008; 22: 283–285.
69. Mutanyatta J *et al.* The first 6'-*O*-sulfated phenylanthraquinones: isolation from *Bulbine frutescens*, structural elucidation, enantiomeric purity, and partial synthesis. *Tetrahedron* 2005; 61: 8475–8484.
70. Bringmann G *et al.* Joziknipholones A and B: the first dimeric phenylanthraquinones, from the roots of *Bulbine frutescens*. *Chemistry* 2008; 14: 1420–1429.
71. Wube AA *et al.* Antimalarial compounds from *Kniphofia foliosa* roots. *Phytother Res* 2005; 19: 472–476.
72. Lenta BN *et al.* Anti-plasmodial and cholinesterase inhibiting activities of some constituents of *Psorospermum glaberrimum*. *Chem Pharm Bull* 2008; 56: 222–226.
73. Lenta BN *et al.* Anti-plasmodial activity of some constituents of the root bark of *Harungana madagascariensis* LAM. (Hypericaceae). *Chem Pharm Bull* 2007; 55: 464–467.
74. Ichino C *et al.* Screening of Thai medicinal plant extracts and their active constituents for *in vitro* antimalarial activity. *Phytother Res* 2006; 20: 307–309.
75. Eyong KO *et al.* Newbouldiaquinone A: a naphthoquinone-anthraquinone ether coupled pigment, as a potential antimicrobial and antimalarial agent from *Newbouldia laevis*. *Phytochemistry* 2006; 67: 605–609.
76. Tuntiwachwuttikul P *et al.* Anthraquinones from the roots of *Prismatomeris malayana*. *Nat Prod Res* 2008; 22: 962–968.
77. Rukunga GM *et al.* Anti-plasmodial activity of the extracts and two sesquiterpenes from *Cyperus articulatus*. *Fitoterapia* 2008; 79: 188–190.
78. Pillay P *et al.* Isolation and identification of antiplasmodial sesquiterpene lactones from *Oncosiphon piluliferum*. *J Ethnopharmacol* 2007; 112: 71–76.
79. Ramanandraibe V *et al.* Pseudoguanolide sesquiterpene lactones from *Vernoniopsis caudata* and their *in vitro* antiplasmodial activities. *J Nat Prod* 2005; 68: 800–803.
80. Ortet R *et al.* Sesquiterpene lactones from the endemic Cape Verdean *Artemisia gorgonum*. *Phytochemistry* 2008; 69: 2961–2965.
81. That QT *et al.* Wedelolides A and B: novel sesquiterpene delta-lactones, (9R)-eudesman-9,12-olides, from *Wedelia trilobata*. *J Org Chem* 2007; 72: 7102–7105.
82. Chea A *et al.* Antimalarial activity of sesquiterpene lactones from *Vernonia cinerea*. *Chem Pharm Bull* 2006; 54: 1437–1439.
83. Pillay P *et al.* Antiplasmodial hirsutinolides from *Vernonia staehelinoides* and their utilization towards a simplified pharmacophore. *Phytochemistry* 2007; 68: 1200–1205.
84. Vongvanich N *et al.* Antiplasmodial, antimycobacterial, and cytotoxic principles from *Camchaya calcarea*. *Planta Med* 2006; 72: 1427–1430.
85. Moon HI. Antiplasmodial activity of ineupatorolides A from *Carpesium rosulatum*. *Parasitol Res* 2007; 100: 1147–1149.
86. Chung IM *et al.* Antiplasmodial activity of sesquiterpene lactone from *Carpesium rosulatum* in mice. *Parasitol Res* 2008; 103: 341–344.
87. Karioti A *et al.* Inhibiting enoyl-ACP reductase (FabI) across pathogenic microorganisms by linear sesquiterpene lactones from *Anthemis auriculata*. *Phytomedicine* 2008; 15: 1125–1129.
88. Adelekan AM *et al.* Bioactive diterpenes and other constituents of *Croton steenkampianus*. *J Nat Prod* 2008; 71: 1919–1922.
89. Kenmogne M *et al.* Five labdane diterpenoids from the seeds of *Aframomum zambesiacum*. *Phytochemistry* 2006; 67: 433–438.
90. Okasaka M *et al.* Terpenoids from *Juniperus polycarpus* var. *seravschanica*. *Phytochemistry* 2006; 67: 2635–2640.
91. Nyiligira E *et al.* Phytochemistry and *in vitro* pharmacological activities of South African Vitex (Verbenaceae) species. *J Ethnopharmacol* 2008; 119: 680–685.
92. Jullian V *et al.* New clerodane diterpenoids from *Laetia procera* (Poepp.) Eichler (Flacourtiaceae), with antiplasmodial and antileishmanial activities. *Bioorg Med Chem Lett* 2005; 15: 5065–5070.
93. Kanokmedhakul S *et al.* New bioactive clerodane diterpenoids from the bark of *Casearia grewiifolia*. *J Nat Prod* 2005; 68: 183–188.
94. Thongnest S *et al.* Oxygenated pimarane diterpenes from *Kaempferia marginata*. *J Nat Prod* 2005; 68: 1632–1636.
95. Linn TZ *et al.* Cassane- and norcassane-type diterpenes from *Caesalpinia crista* of Indonesia and their antimalarial activity against the growth of *Plasmodium falciparum*. *J Nat Prod* 2005; 68: 706–710.
96. Kalauni SK *et al.* Antimalarial activity of cassane- and norcassane-type diterpenes from *Caesalpinia crista* and their structure–activity relationship. *Biol Pharm Bull* 2006; 29: 1050–1052.
97. Awale S *et al.* Constituents of *Caesalpinia crista* from Indonesia. *Chem Pharm Bull* 2006; 54: 213–218.
98. Pudhom K *et al.* Cassane furanoditerpenoids from the seed kernels of *Caesalpinia bonduc* from Thailand. *J Nat Prod* 2007; 70: 1542–1544.
99. Matsuno Y *et al.* Sucutiniranes A and B, new cassane-type diterpenes from *Bowdichia nitida*. *Bioorg Med Chem Lett* 2008; 18: 3774–3777.
100. Chukwujekwu JC *et al.* Antiplasmodial diterpenoid from the leaves of *Hyptis suaveolens*. *J Ethnopharmacol* 2005; 102: 295–297.
101. Van Zyl RL *et al.* Antiplasmodial activities of some abietane diterpenes from the leaves of five *Plectranthus* species. *S Afr J Sci* 2008; 104: 62–64.
102. Samoylenko V *et al.* Antiparasitic, nematicidal and antifouling constituents from *Juniperus* berries. *Phytother Res* 2008; 22: 1570–1576.
103. Koch A *et al.* An antimalarial abietane diterpene from *Fuerstia africana* TCE fries. *Biochem Syst Ecol* 2006; 34: 270–272.
104. Mongkolvisut W, Sutthivaiyakit S. Antimalarial and antituberculous poly-*O*-acylated jatrophane diterpenoids from *Pedilanthus thymaloides*. *J Nat Prod* 2007; 70: 1434–1438.
105. Murata T *et al.* Antiplasmodial triterpenoids from *Ekebergia capensis*. *J Nat Prod* 2008; 71: 167–174.
106. Mambu L *et al.* Clerodane and labdane diterpenoids from *Nuxia sphaerocephala*. *Phytochemistry* 2006; 67: 444–451.
107. Suksamrarn S *et al.* Ceanothane- and lupane-type triterpenes with antiplasmodial and antimycobacterial activities from *Ziziphus cambodiana*. *Chem Pharm Bull* 2006; 54: 535–537.
108. Ma CY *et al.* Study of antimalarial activity of chemical constituents from *Diospyros quaesita*. *Chem Biodivers* 2008; 5: 2442–2448.

109. Thiem DA *et al.* Bisnortriterpenes from *Salacia madagascariensis*. *J Nat Prod* 2005; 68: 251–254.
110. Cimanga RK *et al.* Bioassay-guided isolation of antimalarial triterpenoid acids from the leaves of *Morinda lucida*. *Pharm Biol* 2006; 44: 677–681.
111. Moon HI *et al.* Antiplasmodial activity of triterpenoid isolated from whole plants of *Viola* genus from South Korea. *Parasitol Res* 2007; 100: 641–644.
112. He ZD *et al.* Antimalarial constituents from *Nauclea orientalis* (L.) L. *Chem Biodivers* 2005; 2: 1378–1386.
113. Banzouzi JT *et al.* *Cogniauxia podolaena*: bioassay-guided fractionation of defoliated stems, isolation of active compounds, antiplasmodial activity and cytotoxicity. *Planta Med* 2008; 74: 1453–1456.
114. Kamatou GPP *et al.* Antimalarial and anticancer activities of selected South African *Salvia* species and isolated compounds from *S. radula*. *S Afr J Bot* 2008; 74: 238–243.
115. Saewan N *et al.* Antimalarial tetranortriterpenoids from the seeds of *Lansium domesticum* Corr. *Phytochemistry* 2006; 67: 2288–2293.
116. Lee SE *et al.* Antimalarial activity of anthothecol derived from *Khaya anthotheca* (Meliaceae). *Phytomedicine* 2008; 15: 533–535.
117. Bickii J *et al.* The antiplasmodial agents of the stem bark of *Entandrophragma angolense* (Meliaceae). *Afr J Tradit Complement Altern Med* 2007; 4: 135–139.
118. Maneerata W *et al.* Antimalarial, antimycobacterial and cytotoxic limonoids from *Chisocheton siamensis*. *Phytomedicine* 2008; 15: 1130–1134.
119. Hay AE *et al.* Limonoid orthoacetates and antiprotozoal compounds from the roots of *Pseudocedrela kotschyi*. *J Nat Prod* 2007; 70: 9–13.
120. Abdel-Sattar E *et al.* Acylated pregnane glycosides from *Caralluma tuberculata* and their antiparasitic activity. *Phytochemistry* 2008; 69: 2180–2186.
121. Fotie J *et al.* Lupeol long-chain fatty acid esters with antimalarial activity from *Holarrhena floribunda*. *J Nat Prod* 2006; 69: 62–67.
122. Lhinhatrakool T, Sutthivaiyakit S. 19-Nor- and 18,20-epoxycardenolides from the leaves of *Calotropis gigantea*. *J Nat Prod* 2006; 69: 1249–1251.
123. Karioti A *et al.* Anthecularin: a novel sesquiterpene lactone from *Anthemis auriculata* with antiprotozoal activity. *J Org Chem* 2007; 72: 8103–8106.
124. Andrade SF *et al.* Antileishmanial, antimalarial and antimicrobial activities of the extract and isolated compounds from *Austroplenckia populnea* (Celastraceae). *Z Naturforsch C, J Biosci* 2008; 63: 497–502.
125. Ngouamegne ET *et al.* Endodesmiadiol, a friedelane triterpenoid, and other antiplasmodial compounds from *Endodesmia calophylloides*. *Chem Pharm Bull* 2008; 56: 374–377.
126. Fu X *et al.* Cycloartane glycosides from *Sutherlandia frutescens*. *J Nat Prod* 2008; 71: 1749–1753.
127. Agnihotri VK *et al.* New labdane diterpenes from *Leonurus cardiaca*. *Planta Med* 2008; 74: 1288–1290.
128. Acharya BN *et al.* Antimalarial activity of *Gomphostemma crinitum* leaf extracts. *Med Chem Res* 2008; 17: 530–540.
129. Jonville MC *et al.* Fagraldehyde, a secoiridoid isolated from *Fagraea fragrans*. *J Nat Prod* 2008; 71: 2038–2040.
130. Lagnika L *et al.* Phytochemical study and antiprotozoal activity of compounds isolated from *Thalia geniculata*. *Pharm Biol* 2008; 46: 162–165.
131. Traore M *et al.* Cytotoxic kurubasch aldehyde from *Trichilia emetica*. *Nat Prod Res* 2007; 21: 13–17.
132. Ngemenya MN *et al.* Antiplasmodial activities of some products from *Turreanthus africanus* (Meliaceae). *Afr J Health Sci* 2006; 13: 33–39.
133. Akam TM *et al.* A pregnane derivative and an anti-plasmodial labdane diterpenoid from the stem bark of *Turraenthus africanus*. *Nat Prod Commun* 2006; 1: 449–452.
134. Yin S *et al.* The first limonoid peroxide in the Meliaceae family: walsuronoid a from *Walsura robusta*. *Org Lett* 2007; 9: 2353–2356.
135. Flores N *et al.* Benzoic acid derivatives from *Piper* species and their antiparasitic activity. *J Nat Prod* 2008; 71: 1538–1543.
136. Chumkaew P *et al.* A new triterpenoid ester from the fruits of *Bruguiera parviflora*. *Chem Pharm Bull* 2005; 53: 95–96.
137. Kanokmedhakul K *et al.* Biological activity of anthraquinones and triterpenoids from *Prismatomeris fragrans*. *J Ethnopharmacol* 2005; 100: 284–288.
138. Tasdemir D *et al.* Anti-protozoal and plasmodial FabI enzyme inhibiting metabolites of *Scrophularia lepidota* roots. *Phytochemistry* 2005; 66: 355–362.
139. Pawar RS *et al.* Glycosides of 20-deoxy derivatives of jujubogenin and pseudojujubogenin from *Bacopa monniera*. *Planta Med* 2007; 73: 380–383.
140. Tasdemir D *et al.* Evaluation of antiprotozoal and antimycobacterial activities of the resin glycosides and the other metabolites of *Scrophularia cryptophila*. *Phytomedicine* 2008; 15: 209–215.
141. Rukunga GM *et al.* The antiplasmodial activity of spermine alkaloids isolated from *Albizia gummifera*. *Fitoterapia* 2007; 78: 455–459.
142. Rumalla CS *et al.* Alkaloids from *Heimia salicifolia*. *Phytochemistry* 2008; 69: 1756–1762.
143. Toriizuka Y *et al.* New lycorine-type alkaloid from *Lycoris traubii* and evaluation of antitrypanosomal and antimalarial activities of lycorine derivatives. *Bioorg Med Chem* 2008; 16: 10 182–10 189.
144. Morita H *et al.* Cassiarins A and B, novel antiplasmodial alkaloids from *Cassia siamea*. *Org Lett* 2007; 9: 3691–3693.
145. Jullian V *et al.* Validation of use of a traditional antimalarial remedy from French Guiana, *Zanthoxylum rhoifolium* Lam. *J Ethnopharmacol* 2006; 106: 348–352.
146. Mazier D *et al.* Alkaloid compounds and their use as antimalarial drugs. International Patent Application No. PCT/EP2005/00523, 21 April 2005.
147. Carraz M *et al.* Isolation and antimalarial activity of new morphinan alkaloids on *Plasmodium yoelii* liver stage. *Bioorg Med Chem* 2008; 16: 6186–6192.
148. Chea A *et al.* Antimalarial activity of alkaloids isolated from *Stephania rotunda*. *J Ethnopharmacol* 2007; 112: 132–137.
149. Nguyen NT *et al.* Antiplasmodial alkaloids from *Desmos rostrata*. *J Nat Prod* 2008; 71: 2057–2059.
150. Kanokmedhakul S *et al.* Bioactive constituents of the roots of *Polyalthia cerasoides*. *J Nat Prod* 2007; 70: 1536–1538.
151. Osorio E *et al.* Actividad antiplasmodial de alcaloides aporfínicos de *Rollinia pittieri* y *Pseudomalmea boyacana* (Annonaceae). *Vitae. Rev Fac Química Farm* 2006; 13: 49–54.
152. Phrutivorapongkul A *et al.* Anti-plasmodial activity of bisbenzylisoquinoline alkaloids from *Michelia figo* leaves. *Thai J Health Res* 2006; 20: 121–128.
153. Roumy V *et al.* Isolation and antimalarial activity of alkaloids from *Pseudoxandra cuspidata*. *Planta Med* 2006; 72: 894–898.
154. Murebwayire S *et al.* Antiplasmodial and antitrypanosomal activity of *Triclisia saculeuxii* (Pierre) Diels. *Phytomedicine* 2008; 15: 728–733.
155. Bringmann G *et al.* ent-Dioncophylleine A and related dehydrogenated naphthylisoquinoline alkaloids, the first Asian dioncophyllaceae-type alkaloids, from the ‘new’ plant species *Ancistrocladus benomensis*. *J Nat Prod* 2005; 68: 686–690.



156. Bringmann G *et al.* Six naphthylisoquinoline alkaloids and a related benzopyranone from a Congolese *Ancistrocladus* species related to *Ancistrocladus congolensis*. *Phytochemistry* 2008; 69: 1065–1075.
157. Sun JY *et al.* Indole alkaloids from *Nauclea officinalis* with weak antimalarial activity. *Phytochemistry* 2008; 69: 1405–1410.
158. Van Miert S *et al.* Isoneocryptolepine, a synthetic indoloquinoline alkaloid, as an antiplasmodial lead compound. *J Nat Prod* 2005; 68: 674–677.
159. Philippe G *et al.* *In vivo* antimalarial activity of isosungucine, an indolomonoterpenic alkaloid from *Strychnos icaja*. *Planta Med* 2007; 73: 478–479.
160. Devkota KP *et al.* Cholinesterase inhibiting and antiplasmodial steroidal alkaloids from *Sarcococca hookeriana*. *Chem Pharm Bull* 2007; 55: 1397–1401.
161. Zirihi GN *et al.* Isolation, characterization and antiplasmodial activity of steroidal alkaloids from *Funtumia elastica* (Preuss) Stapf. *Bioorg Med Chem Lett* 2005; 15: 2637–2640.
162. Suksamrarn S *et al.* Ziziphine N, O, P and Q, new antiplasmodial cyclopeptide alkaloids from *Ziziphus oenoplia* var. *brunoniana*. *Tetrahedron* 2005; 61: 1175–1180.
163. Ross SA *et al.* Alkamides from the leaves of *Zanthoxylum syncarpum*. *J Nat Prod* 2005; 68: 1297–1299.
164. Wirasathien L *et al.* Biological activities of alkaloids from *Pseuduvaria setosa*. *Pharm Biol* 2006; 44: 274–278.
165. Mongkolvisut W *et al.* Integerrimides A and B, cyclic heptapeptides from the latex of *Jatropha integerrima*. *J Nat Prod* 2006; 69: 1435–1441.
166. Awang K *et al.* New alkaloids from *Phoebe scortechinii*. *Nat Prod Res* 2007; 21: 704–709.
167. Otshudi AL *et al.* Biologically active bisbenzylisoquinoline alkaloids from the root bark of *Epinetrum villosum*. *J Ethnopharmacol* 2005; 102: 89–94.
168. Fiot J *et al.* HPLC quantification of uncarine D and the anti-plasmodial activity of alkaloids from leaves of *Mitragyna inermis* (Willd.) O. Kuntze. *Phytochem Anal* 2005; 16: 30–33.
169. Dolabela MF *et al.* *In vitro* antiplasmodial activity of extract and constituents from *Esenbeckia febrifuga*, a plant traditionally used to treat malaria in the Brazilian Amazon. *Phytomedicine* 2008; 15: 367–372.
170. Waffo AFK *et al.* Acridone and furoquinoline alkaloids from *Teclea gerrardii* (Rutaceae: Toddalioideae) of southern Africa. *Phytochemistry* 2007; 68: 663–667.
171. Penali L *et al.* Low antiplasmodial activity of alkaloids and amides from the stem bark of *Zanthoxylum rubescens* (Rutaceae). *Parasite* 2007; 14: 161–164.
172. Astulla A *et al.* Alkaloids from the seeds of *Peganum harmala* showing antiplasmodial and vasorelaxant activities. *J Nat Med* 2008; 62: 470–472.
173. de Mesquita ML *et al.* New ether diglycosides from *Matayba guianensis* with antiplasmodial activity. *Bioorg Med Chem* 2005; 13: 4499–4506.
174. Fiot J *et al.* Phytochemical and pharmacological study of roots and leaves of *Guiera senegalensis* JF Gmel (Combretaceae). *J Ethnopharmacol* 2006; 106: 173–178.
175. Senn M *et al.* Antiprotozoal polyacetylenes from the Tanzanian medicinal plant *Cussonia zimmermannii*. *J Nat Prod* 2007; 70: 1565–1569.
176. Kanokmedhakul S *et al.* 2-Substituted furans from the roots of *Polyalthia evecta*. *J Nat Prod* 2006; 69: 68–72.
177. Suksamrarn A *et al.* Antimycobacterial and antiplasmodial unsaturated carboxylic acid from the twigs of *Scleropyrum wallichianum*. *Chem Pharm Bull* 2005; 53: 1327–1329.
178. Bertani S *et al.* Simalikalactone D is responsible for the antimalarial properties of an amazonian traditional remedy made with *Quassia amara* L. (Simaroubaceae). *J Ethnopharmacol* 2006; 108: 155–157.
179. Elford BC *et al.* Potentiation of the antimalarial activity of ginghamosu by methoxylated flavones. *Trans R Soc Trop Med Hyg* 1987; 81: 434–436.
180. Lehane AM, Saliba KJ. Common dietary flavonoids inhibit the growth of the intraerythrocytic malaria parasite. *BMC Res Notes* 2008; 1: 26
181. Kamatou GPP *et al.* The *in vitro* pharmacological activities and a chemical investigation of three South African *Salvia* species. *J Ethnopharmacol* 2005; 102: 382–390.