

Subject Index of Volume 172

- Absorption cross-section
High-resolution absorption cross-section of glyoxal in the UV–vis and IR spectral ranges, 35
- Absorption spectra
The synthesis and photochemical characterization of *meso*-tetra-thienyl porphyrins, 283
- Absorption spectrum
Spectral characteristics of 2-(3',5'-diaminophenyl)benzothiazole: effects of solvents and acid–base concentrations, 185
Excited state intramolecular proton transfer in anionic and cationic species of 2-(2'-amino-3-pyridyl)benzimidazole, 196
- Acid Orange 7
Role of primary active species and TiO₂ surface characteristic in UV-illuminated photodegradation of Acid Orange 7, 47
- Acridine
Synthesis and fluorescent properties of novel biotinylated labels. Prospects for application in bioanalytical detections, 215
- Advanced oxidation processes
Dechlorination of pulp bleaching plant E-1 effluent by ArF* and KrF* excimer laser photolysis. Part 1. Dechlorination of chlorophenols identified in E-1 effluent, 222
Dechlorination of pulp bleaching plant E-1 effluent by ArF* and KrF* excimer laser photolysis. 2. Dechlorination of polychlorinated oxylicnins, 231
- Aggregation
Dye sensitization of nanocrystalline TiO₂: enhanced efficiency of unsymmetrical versus symmetrical squaraine dyes, 63
- Alkali nitrate crystals
The peroxyinitrite formation under photolysis of alkali nitrates, 289
- 2-(2'-Amino-3'-pyridyl)benzimidazole
Excited state intramolecular proton transfer in anionic and cationic species of 2-(2'-amino-3-pyridyl)benzimidazole, 196
- 193 and 248 nm photons
Dechlorination of pulp bleaching plant E-1 effluent by ArF* and KrF* excimer laser photolysis. Part 1. Dechlorination of chlorophenols identified in E-1 effluent, 222
- Aniline
TiO₂—photocatalyzed oxidation of aniline, 207
- Anthracene
Singlet and triplet energy transfer in a bichromophoric system with anthracene covalently linked through sulfonamide to a *meso*-tetraphenylporphyrin, 151
- ArF* and KrF* excimer laser photolyses
Dechlorination of pulp bleaching plant E-1 effluent by ArF* and KrF* excimer laser photolysis. Part 1. Dechlorination of chlorophenols identified in E-1 effluent, 222
- ArF* (193 nm) excimer laser photolysis
Dechlorination of pulp bleaching plant E-1 effluent by ArF* and KrF* excimer laser photolysis. 2. Dechlorination of polychlorinated oxylicnins, 231
- Avidin
Synthesis and fluorescent properties of novel biotinylated labels. Prospects for application in bioanalytical detections, 215
- Azo dye
Effect of supporting electrolytes in electrochemically-assisted photodegradation of an azo dye, 316
- Azobenzene groups
Thermo-, chiro- and photo-optical properties of cholesteric azobenzene-containing copolymer in thin films, 140
- Bacteriochlorophyll *c*
Properties of bacteriochlorophyll *c* and bacteriopheophytin *c* in resting and stimulated lymphocytes, 161
- Bacteriopheophytin *c*
Properties of bacteriochlorophyll *c* and bacteriopheophytin *c* in resting and stimulated lymphocytes, 161
- Bandgap energies
Photocatalytic activity of sol–gel derived titania converted into nanocrystalline powders by supercritical drying, 19
- Benzonitrile
Dissociative excitation of benzonitrile by ultraviolet multiphoton absorption, 241
- Binary liquids
Slow solvation dynamics of 4-AP and DCM in binary mixtures, 180
- Biotin
Synthesis and fluorescent properties of novel biotinylated labels. Prospects for application in bioanalytical detections, 215
- Bisphenol A
Rapid degradation of bisphenol A using air as the oxidant catalyzed by polynuclear phthalocyanine complexes under visible light irradiation, 275
- β-Methyl-β-nitrostyrene
Different relative rates for photo-rearrangements of (*E*)- and (*Z*)-β-nitrostyrene derivatives to oximinoketones, 331
- β-Nitrostyrene
Different relative rates for photo-rearrangements of (*E*)- and (*Z*)-β-nitrostyrene derivatives to oximinoketones, 331
- Bond cleavage
Photochemistry of 4-(methylthio)phenylacetic acid. Steady-state and laser flash photolysis studies, 322
- Charge transfer
Mechanism of the excited singlet and triplet states quenching by molecular oxygen in acetonitrile, 170
- Charge-carrier lifetimes
Photocatalytic activity of sol–gel derived titania converted into nanocrystalline powders by supercritical drying, 19
- Cholesteric phase
Thermo-, chiro- and photo-optical properties of cholesteric azobenzene-containing copolymer in thin films, 140

- Cis-trans* isomerization
 Different relative rates for photo-rearrangements of (*E*)- and (*Z*)- β -nitrostyrene derivatives to oximinoketones, 331
- CN($A^2\Pi_i - X^2\Sigma^+$)
 Dissociative excitation of benzonitrile by ultraviolet multiphoton absorption, 241
- CN($B^2\Sigma^+ - X^2\Sigma^+$)
 Dissociative excitation of benzonitrile by ultraviolet multiphoton absorption, 241
- Co-polymerization
 Novel functionalized 2-(2-hydroxyphenyl)-benzotriazole – benzo[*de*]isoquinoline-1,3-dione fluorescent UV absorbers. Synthesis and photostabilizing efficiency, 308
- Daylight irradiation
 Self-cleaning of wool-polyamide and polyester textiles by TiO₂-rutile modification under daylight irradiation at ambient temperature, 27
- Dechlorination
 Dechlorination of pulp bleaching plant E-1 effluent by ArF* and KrF* excimer laser photolysis. Part 1. Dechlorination of chlorophenols identified in E-1 effluent, 222
 Dechlorination of pulp bleaching plant E-1 effluent by ArF* and KrF* excimer laser photolysis. 2. Dechlorination of polychlorinated oxylignins, 231
- Decolorization
 Dechlorination of pulp bleaching plant E-1 effluent by ArF* and KrF* excimer laser photolysis. 2. Dechlorination of polychlorinated oxylignins, 231
- Degradation
 Rapid degradation of bisphenol A using air as the oxidant catalyzed by polynuclear phthalocyanine complexes under visible light irradiation, 275
- 2-(3',5'-Diaminophenyl)benzothiazole
 Spectral characteristics of 2-(3',5'-diaminophenyl)benzothiazole: effects of solvents and acid–base concentrations, 185
- Diazo dyes
 The effect of operational parameters on the photocatalytic degradation of three textile azo dyes in aqueous TiO₂ suspensions, 89
- Dimerization
 Selective photochemical synthesis of 3,3'-dimethoxy-4,2'-dihydroxybiphenyl, 115
- 3,3'-Dimethoxy-4,2'-dihydroxybiphenyl
 Selective photochemical synthesis of 3,3'-dimethoxy-4,2'-dihydroxybiphenyl, 115
- 1,6-Dioxapyrene
 Synthesis, characterization, and optical properties of a cyano-functionalized 2,3,7,8-tetraaryl-1,6-dioxapyrene, 97
- Dissociation dynamics
 Dissociative excitation of benzonitrile by ultraviolet multiphoton absorption, 241
- Dissociative excitation
 Dissociative excitation of benzonitrile by ultraviolet multiphoton absorption, 241
- DMABN
 Field diagram for a molecule appearing in charge transfer state in polar solution, 1
- Dye sensitization
 Dye sensitization of nanocrystalline TiO₂: enhanced efficiency of unsymmetrical versus symmetrical squaraine dyes, 63
- Dye-sensitized solar cell
 Improvement of photovoltaic performance of solid-state dye-sensitized solar cells by iodine doping in conjugated polymer, 135
- E-Z* isomerization
 Thermo-, chiro- and photo-optical properties of cholesteric azobenzene-containing copolymer in thin films, 140
- Electrochemically-assisted
 Effect of supporting electrolytes in electrochemically-assisted photodegradation of an azo dye, 316
- Electrochemistry
 Synthesis, characterization, and optical properties of a cyano-functionalized 2,3,7,8-tetraaryl-1,6-dioxapyrene, 97
- Electrodeless discharge lamp
 The electrodeless discharge lamp: a prospective tool for photochemistry. Part 6. Photochemistry of valerophenone and 4-nitroanisole in high-temperature water, 146
- Electron transfer
 The photodegradation of pyromethene 567 and pyromethene 597 by pyromethene 546, 109
- Electronic spectra
 Synthesis, characterization, and optical properties of a cyano-functionalized 2,3,7,8-tetraaryl-1,6-dioxapyrene, 97
- Energy transfer
 Mechanism of the excited singlet and triplet states quenching by molecular oxygen in acetonitrile, 170
- ESIPT
 Excited state intramolecular proton transfer in anionic and cationic species of 2-(2'-amino-3-pyridyl)benzimidazole, 196
- Excitation energies
 Theoretical study of hypericin, 293
- Field diagram
 Field diagram for a molecule appearing in charge transfer state in polar solution, 1
- Flavin
 Single-molecule dynamics of conformational changes in flavin adenine dinucleotide, 73
- Fluorescence spectra
 The synthesis and photochemical characterization of *meso*-tetra-thienyl porphyrins, 283
- Fluorescence spectrum
 Spectral characteristics of 2-(3',5'-diaminophenyl)benzothiazole: effects of solvents and acid–base concentrations, 185
 Excited state intramolecular proton transfer in anionic and cationic species of 2-(2'-amino-3-pyridyl)benzimidazole, 196
- Fluorescence
 Photophysics of mesoporphyrin IX in solution and confined in sol-gel-derived matrices, 7
 Singlet and triplet energy transfer in a bichromophoric system with anthracene covalently linked through sulfonamide to a *meso*-tetraphenylporphyrin, 151
 Synthesis and fluorescent properties of novel biotinylated labels. Prospects for application in bioanalytical detections, 215
- Gas-phase
 High-resolution absorption cross-section of glyoxal in the UV–vis and IR spectral ranges, 35
- Glyoxal
 High-resolution absorption cross-section of glyoxal in the UV–vis and IR spectral ranges, 35
- Guaiaacol
 Selective photochemical synthesis of 3,3'-dimethoxy-4,2'-dihydroxybiphenyl, 115
- 2'-Halobenzanilides
 Photochemistry of 2'-halobenzanilides: reaction mechanism, 258
- Heterocycles
 Photochemical and thermal behaviour of new photochromic indeno-fused naphthopyrans, 300
- High-temperature water
 The electrodeless discharge lamp: a prospective tool for photochemistry. Part 6. Photochemistry of valerophenone and 4-nitroanisole in high-temperature water, 146

- Hydrogen bonding
Proton transfer reaction in 2-hydroxy-3-formyl benzoic acid at room temperature and 77 K and some study on AM1 potential surfaces, 250
- Hydrogen peroxide
Dechlorination of pulp bleaching plant E-1 effluent by ArF* and KrF* excimer laser photolysis. Part 1. Dechlorination of chlorophenols identified in E-1 effluent, 222
Dechlorination of pulp bleaching plant E-1 effluent by ArF* and KrF* excimer laser photolysis. 2. Dechlorination of polychlorinated oxylinins, 231
- Hydrothermal synthesis
Synthesis and photocatalytic activity of ZnO/ZnO₂ composite, 269
- 2-(2-Hydroxyphenyl)-benzotriazole
Novel functionalized 2-(2-hydroxyphenyl)-benzotriazole – benzo[de]-isoquinoline-1,3-dione fluorescent UV absorbers. Synthesis and photostabilizing efficiency, 308
- Hypericin
Theoretical study of hypericin, 293
- Infrared
High-resolution absorption cross-section of glyoxal in the UV–vis and IR spectral ranges, 35
- Inhomogeneous broadening of spectra
Field diagram for a molecule appearing in charge transfer state in polar solution, 1
- Initial step
Role of primary active species and TiO₂ surface characteristic in UV-illuminated photodegradation of Acid Orange 7, 47
- Internal energy distribution
Dissociative excitation of benzonitrile by ultraviolet multiphoton absorption, 241
- Iodine doping
Improvement of photovoltaic performance of solid-state dye-sensitized solar cells by iodine doping in conjugated polymer, 135
- Kinetics
Dechlorination of pulp bleaching plant E-1 effluent by ArF* and KrF* excimer laser photolysis. Part 1. Dechlorination of chlorophenols identified in E-1 effluent, 222
Dechlorination of pulp bleaching plant E-1 effluent by ArF* and KrF* excimer laser photolysis. 2. Dechlorination of polychlorinated oxylinins, 231
- LE (local-excited) and CT (charge transfer) states
Field diagram for a molecule appearing in charge transfer state in polar solution, 1
- Liquid crystalline copolymer
Thermo-, chiro- and photo-optical properties of cholesteric azobenzene-containing copolymer in thin films, 140
- Luminescence
Field diagram for a molecule appearing in charge transfer state in polar solution, 1
Synthesis, characterization, and optical properties of a cyano-functionalized 2,3,7,8-tetraaryl-1,6-dioxapyrene, 97
- MEHPPV
Improvement of photovoltaic performance of solid-state dye-sensitized solar cells by iodine doping in conjugated polymer, 135
- Mesoporphyrin IX
Photophysics of mesoporphyrin IX in solution and confined in sol–gel-derived matrices, 7
- 2-Methoxyphenol
Selective photochemical synthesis of 3,3'-dimethoxy-4,2'-dihydroxy-biphenyl, 115
- Methyl methacrylate
Photopolymerization of methyl methacrylate using dye-sensitized semiconductor based photocatalyst, 121
- 4-(Methylthio)phenylacetic acid
Photochemistry of 4-(methylthio)phenylacetic acid. Steady-state and laser flash photolysis studies, 322
- Microwave
The electrodeless discharge lamp: a prospective tool for photochemistry. Part 6. Photochemistry of valerophenone and 4-nitroanisole in high-temperature water, 146
- Nanocrystalline TiO₂
Dye sensitization of nanocrystalline TiO₂: enhanced efficiency of unsymmetrical versus symmetrical squaraine dyes, 63
- Nanoparticles
Physicochemical and photocatalytic characterizations of TiO₂/Pt nanocomposites, 81
- Nanowires
Physicochemical and photocatalytic characterizations of TiO₂/Pt nanocomposites, 81
- 1,8-Naphthalimide yellow–green emitting fluorophores
Novel functionalized 2-(2-hydroxyphenyl)-benzotriazole – benzo[de]-isoquinoline-1,3-dione fluorescent UV absorbers. Synthesis and photostabilizing efficiency, 308
- Naphthopyrans
Photochemical and thermal behaviour of new photochromic indeno-fused naphthopyrans, 300
- Nitro-nitrite rearrangement
Different relative rates for photo-rearrangements of (*E*)- and (*Z*)-β-nitrostyrene derivatives to oximinoketones, 331
- Nitroolefins
Different relative rates for photo-rearrangements of (*E*)- and (*Z*)-β-nitrostyrene derivatives to oximinoketones, 331
- NMR
Different relative rates for photo-rearrangements of (*E*)- and (*Z*)-β-nitrostyrene derivatives to oximinoketones, 331
- Out-of-plane order
Thermo-, chiro- and photo-optical properties of cholesteric azobenzene-containing copolymer in thin films, 140
- Oxygen
Dechlorination of pulp bleaching plant E-1 effluent by ArF* and KrF* excimer laser photolysis. Part 1. Dechlorination of chlorophenols identified in E-1 effluent, 222
Dechlorination of pulp bleaching plant E-1 effluent by ArF* and KrF* excimer laser photolysis. 2. Dechlorination of polychlorinated oxylinins, 231
- Peroxynitrite
The peroxynitrite formation under photolysis of alkali nitrates, 289
- Phenol photodegradation
Photocatalytic activity of sol–gel derived titania converted into nanocrystalline powders by supercritical drying, 19
- Phosphorescence
The photodegradation of pyromethene 567 and pyromethene 597 by pyromethene 546, 109
Singlet and triplet energy transfer in a bichromophoric system with anthracene covalently linked through sulfonamide to a *meso*-tetraphenylporphyrin, 151
- Photocatalysis
Self-cleaning of wool-polyamide and polyester textiles by TiO₂-rutile modification under daylight irradiation at ambient temperature, 27
Role of primary active species and TiO₂ surface characteristic in UV-illuminated photodegradation of Acid Orange 7, 47
Synthesis and photocatalytic activity of ZnO/ZnO₂ composite, 269
Rapid degradation of bisphenol A using air as the oxidant catalyzed by polynuclear phthalocyanine complexes under visible light irradiation, 275
- Photocatalyst
Photopolymerization of methyl methacrylate using dye-sensitized semiconductor based photocatalyst, 121

- Photochemical synthesis
 Selective photochemical synthesis of 3,3'-dimethoxy-4,2'-dihydroxy-biphenyl, 115
- Photochemistry
 Single-molecule dynamics of conformational changes in flavin adenine dinucleotide, 73
 The electrodeless discharge lamp: a prospective tool for photochemistry. Part 6. Photochemistry of valerophenone and 4-nitroanisole in high-temperature water, 146
- Photochromism
 Photochemical and thermal behaviour of new photochromic indeno-fused naphthopyrans, 300
- Photodegradation
 Physicochemical and photocatalytic characterizations of TiO₂/Pt nanocomposites, 81
 Effect of supporting electrolytes in electrochemically-assisted photodegradation of an azo dye, 316
- Photodynamic diagnosis
 Properties of bacteriochlorophyll *c* and bacteriopheophytin *c* in resting and stimulated lymphocytes, 161
- Photodynamic therapy
 The convenient screening method using albumin for the tumor localizing property of Ga-porphyrin complexes, 55
 Properties of bacteriochlorophyll *c* and bacteriopheophytin *c* in resting and stimulated lymphocytes, 161
- Photokinetics
 Photochemical and thermal behaviour of new photochromic indeno-fused naphthopyrans, 300
- Photolysis
 The peroxy-nitrite formation under photolysis of alkali nitrates, 289
 Photochemistry of 4-(methylthio)phenylacetic acid. Steady-state and laser flash photolysis studies, 322
- Photooxidation
 TiO₂—photocatalyzed oxidation of aniline, 207
- Photophysics
 Singlet and triplet energy transfer in a bichromophoric system with anthracene covalently linked through sulfonamide to a *meso*-tetraphenylporphyrin, 151
- Photopolymerization
 Photopolymerization of methyl methacrylate using dye-sensitized semiconductor based photocatalyst, 121
- Photo-rearrangement
 Different relative rates for photo-rearrangements of (*E*)- and (*Z*)- β -nitrostyrene derivatives to oximinoketones, 331
- Photoreduction
 Photochemistry of 2'-halobenzanilides: reaction mechanism, 258
- Photosensitizer
 The convenient screening method using albumin for the tumor localizing property of Ga-porphyrin complexes, 55
- Photostability
 The photodegradation of pyromethene 567 and pyromethene 597 by pyromethene 546, 109
 Novel functionalized 2-(2-hydroxyphenyl)-benzotriazole – benzo[*de*]isoquinoline-1,3-dione fluorescent UV absorbers. Synthesis and photostabilizing efficiency, 308
- Photosubstitution
 Photochemistry of 2'-halobenzanilides: reaction mechanism, 258
- Phototoxicity
 Theoretical study of hypericin, 293
- pK_a values
 Spectral characteristics of 2-(3',5'-diaminophenyl)benzothiazole: effects of solvents and acid–base concentrations, 185.
- Platinisation
 Physicochemical and photocatalytic characterizations of TiO₂/Pt nanocomposites, 81
- Polychlorinated oxylignins
 Dechlorination of pulp bleaching plant E-1 effluent by ArF* and KrF* excimer laser photolysis. 2. Dechlorination of polychlorinated oxylignins, 231
- Polymerizable adducts
 Novel functionalized 2-(2-hydroxyphenyl)-benzotriazole – benzo[*de*]isoquinoline-1,3-dione fluorescent UV absorbers. Synthesis and photostabilizing efficiency, 308
- Porphyrazine
 Theoretical investigation on the one- and two-photon absorption properties of a series of porphyrazines with annulated 1,2,5-thiadiazole and 1,4-dimethoxybenzene moieties, 126
- Porphyrin
 The synthesis and photochemical characterization of *meso*-tetra-thienyl porphyrins, 283
- Porphyrins
 Singlet and triplet energy transfer in a bichromophoric system with anthracene covalently linked through sulfonamide to a *meso*-tetraphenylporphyrin, 151
- Potential energy surface
 Proton transfer reaction in 2-hydroxy-3-formyl benzoic acid at room temperature and 77 K and some study on AM1 potential surfaces, 250
- Protein conformation
 Single-molecule dynamics of conformational changes in flavin adenine dinucleotide, 73
- Proton transfer
 Proton transfer reaction in 2-hydroxy-3-formyl benzoic acid at room temperature and 77 K and some study on AM1 potential surfaces, 250
- Prototropic equilibrium
 Excited state intramolecular proton transfer in anionic and cationic species of 2-(2'-amino-3-pyridyl)benzimidazole, 196
- Pulp bleaching plant E-1 effluent
 Dechlorination of pulp bleaching plant E-1 effluent by ArF* and KrF* excimer laser photolysis. Part 1. Dechlorination of chlorophenols identified in E-1 effluent, 222
 Dechlorination of pulp bleaching plant E-1 effluent by ArF* and KrF* excimer laser photolysis. 2. Dechlorination of polychlorinated oxylignins, 231
- Pyromethene dyes
 The photodegradation of pyromethene 567 and pyromethene 597 by pyromethene 546, 109
- Quantum yield
 Photochemistry of 4-(methylthio)phenylacetic acid. Steady-state and laser flash photolysis studies, 322
- Quenching
 The photodegradation of pyromethene 567 and pyromethene 597 by pyromethene 546, 109
 Mechanism of the excited singlet and triplet states quenching by molecular oxygen in acetonitrile, 170
- Radical
 Photochemistry of 4-(methylthio)phenylacetic acid. Steady-state and laser flash photolysis studies, 322
- Rates of reaction
 Different relative rates for photo-rearrangements of (*E*)- and (*Z*)- β -nitrostyrene derivatives to oximinoketones, 331
- Resting cell
 Properties of bacteriochlorophyll *c* and bacteriopheophytin *c* in resting and stimulated lymphocytes, 161
- Rutile loaded synthetic fibers
 Self-cleaning of wool-polyamide and polyester textiles by TiO₂-rutile modification under daylight irradiation at ambient temperature, 27

- Scavenger
Role of primary active species and TiO₂ surface characteristic in UV-illuminated photodegradation of Acid Orange 7, 47
- Screening method
The convenient screening method using albumin for the tumor localizing property of Ga-porphyrin complexes, 55
- Self-cleaning
Self-cleaning of wool-polyamide and polyester textiles by TiO₂-rutile modification under daylight irradiation at ambient temperature, 27
- Semiconductor
Photopolymerization of methyl methacrylate using dye-sensitized semiconductor based photocatalyst, 121
- Semi-empirical calculations
Proton transfer reaction in 2-hydroxy-3-formyl benzoic acid at room temperature and 77 K and some study on AM1 potential surfaces, 250
- Single molecule
Single-molecule dynamics of conformational changes in flavin adenine dinucleotide, 73
- Singlet oxygen
The photodegradation of pyrromethene 567 and pyrromethene 597 by pyrromethene 546, 109
Mechanism of the excited singlet and triplet states quenching by molecular oxygen in acetonitrile, 170
Rapid degradation of bisphenol A using air as the oxidant catalyzed by polynuclear phthalocyanine complexes under visible light irradiation, 275
- Singlet state
Singlet and triplet energy transfer in a bichromophoric system with anthracene covalently linked through sulfonamide to a *meso*-tetraphenylporphyrin, 151
Mechanism of the excited singlet and triplet states quenching by molecular oxygen in acetonitrile, 170
- Smectic phase
Thermo-, chiro- and photo-optical properties of cholesteric azobenzene-containing copolymer in thin films, 140
- Solar cells
Dye sensitization of nanocrystalline TiO₂: enhanced efficiency of unsymmetrical versus symmetrical squaraine dyes, 63
- Sol-gel
Photocatalytic activity of sol-gel derived titania converted into nanocrystalline powders by supercritical drying, 19
- Solvation dynamics
Slow solvation dynamics of 4-AP and DCM in binary mixtures, 180
- Solvatochromism
Synthesis, characterization, and optical properties of a cyano-functionalized 2,3,7,8-tetraaryl-1,6-dioxapyrene, 97
- Solvent effect
Photophysics of mesoporphyrin IX in solution and confined in sol-gel-derived matrices, 7
The synthesis and photochemical characterization of *meso*-tetra-thienyl porphyrins, 283
- Squaraine dyes
Dye sensitization of nanocrystalline TiO₂: enhanced efficiency of unsymmetrical versus symmetrical squaraine dyes, 63
- Stimulated cell
Properties of bacteriochlorophyll *c* and bacteriopheophytin *c* in resting and stimulated lymphocytes, 161
- Sum-over-states
Theoretical investigation on the one- and two-photon absorption properties of a series of porphyrazines with annulated 1,2,5-thiadiazole and 1,4-dimethyloxybenzene moieties, 126
- Sunlight
TiO₂-photocatalyzed oxidation of aniline, 207
Rapid degradation of bisphenol A using air as the oxidant catalyzed by polynuclear phthalocyanine complexes under visible light irradiation, 275
- Supercritical drying
Photocatalytic activity of sol-gel derived titania converted into nanocrystalline powders by supercritical drying, 19
- Supporting electrolytes
Effect of supporting electrolytes in electrochemically-assisted photodegradation of an azo dye, 316
- Surface modification
Role of primary active species and TiO₂ surface characteristic in UV-illuminated photodegradation of Acid Orange 7, 47
- Surface photovoltage
Improvement of photovoltaic performance of solid-state dye-sensitized solar cells by iodine doping in conjugated polymer, 135
- Synthesis
Synthesis, characterization, and optical properties of a cyano-functionalized 2,3,7,8-tetraaryl-1,6-dioxapyrene, 97
- Synthetic fibers
Self-cleaning of wool-polyamide and polyester textiles by TiO₂-rutile modification under daylight irradiation at ambient temperature, 27
- TD-DFT
Theoretical study of hypericin, 293
- Theoretical calculations
Spectral characteristics of 2-(3',5'-diaminophenyl)benzothiazole: effects of solvents and acid-base concentrations, 185
Excited state intramolecular proton transfer in anionic and cationic species of 2-(2'-amino-3-pyridyl)benzimidazole, 196
- Thermal bleaching
Photochemical and thermal behaviour of new photochromic indeno-fused naphthopyrans, 300
- Thienyl
The synthesis and photochemical characterization of *meso*-tetra-thienyl porphyrins, 283
- TiO₂ clusters
Self-cleaning of wool-polyamide and polyester textiles by TiO₂-rutile modification under daylight irradiation at ambient temperature, 27
- TiO₂ colloids
Self-cleaning of wool-polyamide and polyester textiles by TiO₂-rutile modification under daylight irradiation at ambient temperature, 27
- TiO₂ photocatalysis
Photocatalytic activity of sol-gel derived titania converted into nanocrystalline powders by supercritical drying, 19
- TiO₂
Physicochemical and photocatalytic characterizations of TiO₂/Pt nanocomposites, 81
TiO₂-photocatalyzed oxidation of aniline, 207
Effect of supporting electrolytes in electrochemically-assisted photodegradation of an azo dye, 316
- Titanium dioxide
The effect of operational parameters on the photocatalytic degradation of three textile azo dyes in aqueous TiO₂ suspensions, 89
- Triplet state
Singlet and triplet energy transfer in a bichromophoric system with anthracene covalently linked through sulfonamide to a *meso*-tetraphenylporphyrin, 151
Mechanism of the excited singlet and triplet states quenching by molecular oxygen in acetonitrile, 170
- Tumor localizing property
The convenient screening method using albumin for the tumor localizing property of Ga-porphyrin complexes, 55
- Two-photon absorption
Theoretical investigation on the one- and two-photon absorption properties of a series of porphyrazines with annulated 1,2,5-thiadiazole and 1,4-dimethyloxybenzene moieties, 126
- Ultrafiltration
Dechlorination of pulp bleaching plant E-1 effluent by ArF* and KrF* excimer laser photolysis. 2. Dechlorination of polychlorinated oxylinins, 231

Ultraviolet

High-resolution absorption cross-section of glyoxal in the UV-vis and IR spectral ranges, 35

UV light

TiO₂—photocatalyzed oxidation of aniline, 207

UV-photodegradation

The effect of operational parameters on the photocatalytic degradation of three textile azo dyes in aqueous TiO₂ suspensions, 89

Valerophenone

The electrodeless discharge lamp: a prospective tool for photochemistry. Part 6. Photochemistry of valerophenone and 4-nitroanisole in high-temperature water, 146

Visible light

Rapid degradation of bisphenol A using air as the oxidant catalyzed by polynuclear phthalocyanine complexes under visible light irradiation, 275

ZINDO

Theoretical investigation on the one- and two-photon absorption properties of a series of porphyrazines with annulated 1,2,5-thiadiazole and 1,4-dimethoxybenzene moieties, 126

ZnO

Synthesis and photocatalytic activity of ZnO/ZnO₂ composite, 269

ZnO₂

Synthesis and photocatalytic activity of ZnO/ZnO₂ composite, 269