

EFFECT OF O₂ ON MOLECULAR IODINE PRODUCTION FOLLOWING FLASH PHOTODISSOCIATION OF CF₃I

JULIE E. STRONG, HARRY F. PREST and ROBERT L. STRONG

Department of Chemistry, Rensselaer Polytechnic Institute, Troy, N.Y. 12181 (U.S.A.)

(Received August 18, 1978)


Summary

The temporal production of molecular iodine following the gas phase flash photodissociation of CF₃I (CO₂ inert diluent gas) has been studied. In the absence of O₂ there is an induction period of about 200 μs determined by quenching of I(²P_{1/2}) atoms, followed by three-body recombination of ground state I(²P_{3/2}) atoms. Addition of small amounts of O₂ greatly enhances the rate of I₂ production, such that at P(O₂) ≥ 0.2 Torr I₂ formation occurs during the photolyzing flash. These results are discussed in terms of formation of a strong intermediate exciplex ($K \geq 10^4 \text{ dm}^3 \text{ mol}^{-1}$) between I(²P_{1/2}) and O₂ as precursor to I₂ formation.

1. Introduction

The optically pumped atomic iodine laser, operating on the magnetic dipole allowed transition ²P_{1/2} → ²P_{3/2} and oscillating at 1315 nm, was the first laser to utilize photodissociation as a mode of excitation [1]. The long natural lifetime of I(²P_{1/2}) resulting from the electric dipole forbidden character of the radiative decay, the superior beam distortion and power-dissipating characteristics of gas relative to solid state lasers, the potential for nearly full extraction of the stored energy and the exceedingly high gain and potential for high power density operation make this laser a good prospect for repetitively pulsed laser plasma and fusion research. Pulse powers of 1 TW have recently been reported [2].

The photodissociation and stimulated emission steps



where RI is an alkyl or perfluoroalkyl iodide molecule, I* = I(²P_{1/2}) and I = I(²P_{3/2}), are well established in the literature [1]. In many cases the fraction

x of excited atoms produced in reaction (1) is very high, *e.g.* 0.92 for CH_3I , 0.91 for CF_3I and greater than 0.99 for $n\text{-C}_3\text{F}_7\text{I}$ [3], leading to a very high degree of population inversion and high gain stimulated emission. Premature termination of laser oscillation is now generally accepted [4] to occur from a high temperature rise and pyrolytic build-up of I_2 which is an extremely efficient quencher of I^* atoms.

It has also been suggested [5] that population inversion and hence potential stimulated emission in the iodine atom system may be possible through energy transfer from $\text{O}_2(^1\Delta_g)$:




Continuous (c.w.) laser action has now been accomplished by generating $\text{O}_2(^1\Delta_g)$ chemically from the reaction of chlorine with H_2O_2 [6], and population inversion has been observed when $\text{O}_2(^1\Delta_g)$ is produced by reaction between chlorine fluorosulfate and H_2O_2 [7]; ground state iodine atoms are produced by the "energy-pooling" process [8]



Energy transfer between $\text{O}_2(^1\Sigma_g^+)$ and I_2 leading to excited I_2 (but not necessarily dissociation) is extremely efficient, occurring at every collision [9].

Reaction (3) is exothermic by only 3.3 kJ, and therefore nearly resonant with the reverse I^* quenching reaction. Possibly because of this molecular oxygen is also a very efficient quencher of $\text{I}(^2\text{P}_{1/2})$ atoms, although the O_2 quenching mechanism is not clear [10]. At sufficiently high concentrations of $\text{O}_2(^1\Delta_g)$, and hence $\text{O}_2(^1\Sigma_g^+)$ by energy pooling, I_2 is kept almost completely dissociated by reaction (5). We report here temporal studies on the formation of I_2 following production of I^* by the flash photolysis of CF_3I in the presence of O_2 with no (initially) $\text{O}_2(^1\Delta_g)$ or $\text{O}_2(^1\Sigma_g^+)$, but under conditions of flash energy and "inert" third-body pressures such that pyrolytic production of I_2 was insignificant.

The reaction of CF_3 radicals with O_2 in low intensity (continuous) photolyses has been studied in several laboratories. Francis and Hazeldine [11] obtained CF_2O , CO_2 and SiF_4 as the only detected products in the photolysis of $\text{CF}_3\text{I-O}_2$ mixtures. They proposed a reaction scheme involving $\text{CF}_3 + \text{O}_2$ additives:



Similarly, CF_3O_2 formation by the addition reaction (6) has been suggested in the photochemical fluorination of fluoroform in the presence of O_2 [12]. Other products identified were $\text{CF}_3\text{O}_2\text{CF}_3$ and $\text{CF}_3\text{O}_3\text{CF}_3$, with the latter undergoing partial conversion to CF_2O in the IR spectrophotometer detec-

tion cell. However, Heicklen [13] proposed direct formation of CF_2O by an exchange reaction



Presumably F_2 and O_2 are then formed by



although F_2 was not found as a reaction product.

2. Experimental

The flash spectrophotometric apparatus has been described elsewhere [14]. A xenon-oxygen (approximately 80:1 mixture) flash lamp was discharged at 900 J ($12.5 \mu\text{F}$, 12 kV) and its light filtered through 10 mm of a UV filter solution ($200 \text{ g dm}^{-3} \text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ and $100 \text{ g dm}^{-3} \text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ in 1 M HCl, upper wavelength cut-off approximately 380 nm). The continuous analyzing beam from a 450 W Osram xenon arc was split after passing through the cell: time-resolved generation of I_2 following flash photodissociation of CF_3I was monitored at 498 nm ($\epsilon(\text{I}_2, 498) = 585 \text{ dm}^3 \text{ mol}^{-1} \text{ cm}^{-1}$) with the Zeiss MM12 monochromator bandwidth maintained at about 1 nm; simultaneously, the flash profile was monitored at 360 nm with a Bausch and Lomb grating monochromator.

All gases were introduced into a vacuum transfer system through a phosphorous pentoxide drying column. Oxygen (Matheson Research Grade minimum purity 99.99%) was used without further purification. CO_2 (from dry ice), CF_3I (PCR Incorporated), SF_6 (Matheson, Instrument Purity minimum purity 99.99%), xenon (Airco Research Grade, minimum purity 99.99%) and propane (Matheson, Instrument Purity, minimum purity 99.5%) were outgassed by several freeze-pump-thaw cycles prior to being condensed into the cylindrical quartz photolysis cell (100 mm light path, 25 mm internal diameter). The total amount of I_2 produced after each flash or series of flashes was measured spectrophotometrically at 498 nm on a Beckman DU Spectrophotometer.

The initial flashing of each I_2 -free cell was carried out at room temperature. Molecular iodine accumulates by irreversible processes as detailed in Section 3, and following flashes on the same cell maintained at room temperature led to some transitory I_2 dissociation that masked I_2 photogeneration. (This dissociation probably occurs by energy transfer since direct photodissociation was prevented by the 380 nm cut-off filter; that no direct I_2 photodissociation occurred was confirmed by flashing a cell containing only CO_2 and I_2 .) Accordingly, subsequent flashes were carried out after the condensable contents of the cell had first been frozen into a side-arm at -196°C and then maintained at -39°C with a benzyl chloride slush bath (vapor pressure of I_2 at -39°C , 4.5×10^{-4} Torr). Under these conditions, results on cells in which I_2 had been produced previously were consistent with initial-flash temporal I_2 production results.

3. Results and discussion

3.1. Photolysis of CF_3I in the absence of O_2

Figure 1 is a typical oscillogram monitored at 498 nm showing temporal behavior of I_2 production following UV flash photodissociation of CF_3I in the presence of about 400 Torr CO_2 ; comparable oscillograms were obtained with xenon or SF_6 as the inert diluent gas. It is seen that generation of I_2 is preceded by an induction period of approximately 200 μs . This induction period decreases with increasing pressure of CF_3I , ranging from 190 μs at 24 Torr CF_3I to 425 μs at 4 Torr CF_3I .

A similar delay in I_2 production occurs in the iodine photodissociation laser [15] where accumulation of I_2 is attributed primarily to thermal decomposition of the active medium in the laser; the induction period is the time needed to reach a pyrolysis threshold temperature (about 1000 K). Pyrolysis of CF_3I clearly cannot be the source of I_2 in the non-laser flash results reported here, however. Under the flash conditions typically used (total flash energy of the order of 1 kJ with non-collimated light output distributed roughly uniformly over the UV and visible spectral regions), the temperature rise within the photolysis cell was limited to less than 1 $^\circ C$ per flash by the gas pressure (greater than 0.5 atm).

In the absence of CF_3I pyrolysis, three-body atom recombination is the major source of I_2 . Photodissociation of CF_3I leads to over 90% I^* atoms [3]; these excited atoms do not combine themselves and they combine with




Fig. 1. I_2 formation following flash photodissociation of CF_3I in CO_2 : upper trace, scattered light from cell (monitored at 360 nm) showing time of flash initiation; lower trace, absorption at 498 nm. Sweep time, 50 μs per major division; vertical sensitivity, 4.9×10^{14} I_2 molecules cm^{-3} per major division; $P(CF_3I) = 16$ Torr; $P(CO_2) = 400$ Torr.

ground state I atoms two orders of magnitude slower than recombination of two ground state atoms [16]. The induction period in I₂ production in the absence of O₂ thus results from time-delayed deactivation of I* :



Trifluoromethyl radicals are very efficient quenchers of I* : $k_{Q,CF_3} = 2.9 \times 10^{-12} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ [17]. However, the high pressure second order recombination $2CF_3 \rightarrow C_2F_6$ is at least four times as fast as the deactivation $CF_3 + I^* \rightarrow CF_3 + I$ [17, 18] so that under the conditions of the work reported here it is assumed that no quenching by CF₃ occurs. (It should be noted that if appreciable deactivation by CF₃ did occur during the short lifetime of the CF₃ radicals in the system, then I₂ production would have occurred by recombination at a much shorter time than actually observed, as for example in Fig. 1). Conversely, CO₂ and CF₃I are weak quenchers of I* : $k_{Q,M} = 1.3 \times 10^{-16}$ and $3.5 \times 10^{-16} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ for CO₂ [19] and CF₃I [20] respectively. The results are qualitatively consistent, however, with the following mechanism utilizing the extremely efficient quenching by I₂ ($k_{Q,I_2} = 3.6 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ [21]):



The third-order recombination rate constants $k_{R,M}$ for CO₂, CF₃I and I₂ are 5.0×10^{-32} (this work, in agreement with earlier results [22]), 3.1×10^{-31} [23] and 3.0×10^{-30} [24] $\text{cm}^6 \text{ molecule}^{-2} \text{ s}^{-1}$. In all of the work reported, $P(\text{CO}_2)/P(\text{I}_2) > 6000$ and $P(\text{CO}_2)/P(\text{CF}_3\text{I}) > 15$, so that CO₂ is the predominant third body in reaction (14). We have simulated this mechanism on an EAI TR-10 analog computer using k_{Q,I_2} as an adjustable parameter, assuming M = CO₂ only and that the concentration of I* at the time of maximum flash intensity is $[I^*]_0 = 0.91 (2 \times [I_2]_\infty)$, where $[I_2]_\infty$ is the total amount of I₂ produced for one flash. Results for the flash oscillogram in Fig. 1 are shown in Fig. 2, giving a best fit value for k_{Q,I_2} of $7.4 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. Computer simulations for CF₃I pressures of 8 Torr and 24 Torr give best fit values of 4.0×10^{-11} and $2.5 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ for k_{Q,I_2} . In view of the major assumptions involved in the simulations, these best fit values are in satisfactory agreement with the literature value given above and support mechanism (12) - (14) as the major source of the induction period preceding I₂ production.

Propane is an efficient quencher of I* : $k_{Q,C_3H_8} = 1.6 \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ [25]. Addition of about 3 Torr propane to the reaction mixture of Fig. 1 greatly reduces the induction period but has negligible effect on the subsequent rate of I₂ production, as seen in Fig. 3. Inclusion of




Fig. 2. Computer simulation (solid curve) of experimental (○) formation of I₂ following flash photodissociation of CF₃I in CO₂: $P(\text{CF}_3\text{I}) = 16$ Torr; $P(\text{CO}_2) = 400$ Torr.




Fig. 3. I₂ formation following flash photodissociation of CF₃I in CO₂ with C₃H₈ quencher: upper trace, scattered light at 360 nm; lower trace, absorption at 498 nm. Sweep time, 100 μs per major division; vertical sensitivity, 4.9×10^{14} I₂ molecules cm⁻³ per major division; $P(\text{CF}_3\text{I}) = 16$ Torr; $P(\text{CO}_2) = 400$ Torr; $P(\text{C}_3\text{H}_8) = 3$ Torr.



in the computer simulation gives a best fit value for k_{Q, C_3H_8} of $5.6 \times 10^{-13} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. Substitution of propane (400 Torr) for CO_2 completely eliminates the induction period and leads immediately following the flash to the expected third order recombination (Fig. 4).




Fig. 4. I_2 formation following flash photodissociation of CF_3I in C_3H_8 : upper trace, scattered light at 360 nm; lower trace, absorption at 498 nm. Sweep time, 500 μs per major division; vertical sensitivity, $4.9 \times 10^{14} \text{ I}_2 \text{ molecule cm}^{-3}$ per major division; $P(\text{CF}_3\text{I}) = 16 \text{ Torr}$; $P(\text{C}_3\text{H}_8) = 400 \text{ Torr}$.

3.2. Photolysis of CF_3I in the presence of O_2

Molecular oxygen is an extremely efficient quencher of I^* ($k_{Q, \text{O}_2} = 2.5 \times 10^{-11} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ [26]) but is a poor third body in ground state I atom recombination, being only about half as effective as CO_2 [27]. Addition of about 2 Torr O_2 to the system represented in Fig. 1 should be as effective as 400 Torr propane in quenching I^* ; on this basis I_2 production following flash photodissociation of CF_3I in the presence of O_2 might be expected to follow that of Fig. 4. The results, however, are quite different, as shown by the oscillogram in Fig. 5. Although the induction period has been eliminated (less than 10 μs) by addition of 2 Torr O_2 to CF_3I (16 Torr) and CO_2 (400 Torr), the rate of I_2 production is rapid and essentially that of the activating flash intensity, *i.e.* about $7 \times 10^{19} \text{ I}_2 \text{ molecules cm}^{-3} \text{ s}^{-1}$. This behavior is followed at all higher pressures of O_2 and to O_2 pressures less than 0.2 Torr. At $P(\text{O}_2) \approx 0.03 \text{ Torr}$, the rate of I_2 production is less than that of the light intensity: initial rate = $R_0(\text{I}_2) \approx 4 \times 10^{18} \text{ molecules cm}^{-3} \text{ s}^{-1}$. Similarly, at $P(\text{O}_2) = 1 \text{ Torr}$, $P(\text{CF}_3\text{I}) = 4 \text{ Torr}$, the rate of I_2 production is slightly less than the flash intensity profile.




Fig. 5. I_2 formation following flash photodissociation of CF_3I in CO_2 with O_2 quencher: upper trace, scattered light at 360 nm; lower trace, absorption at 498 nm. Sweep time, 50 μs per major division; vertical sensitivity, 4.9×10^{14} I_2 molecules cm^{-3} per major division; $P(CF_3I) = 16$ Torr; $P(CO_2) = 400$ Torr; $P(O_2) = 2$ Torr.

At $P(O_2) \geq 1$ Torr, the molecular iodine concentration reaches a maximum value at about 60 μs after flash initiation followed by a first order disappearance ($k_1 \approx 10^4$ s^{-1}) that is independent of CF_3I or O_2 pressure. The results are summarized in Table 1. It is seen that both the maximum I_2 concentration and the equilibrium concentration reached several hundred microseconds after the flash are roughly independent of O_2 pressure, but do increase with increasing CF_3I pressure (and hence increased light absorption). The fall-off in a linear dependence at the highest (23 Torr) CF_3I pressure is due in part to high, and hence non-linear, absorption by CF_3I (decadic $\epsilon = 155$ dm^3 $mole^{-1}$ cm^{-1} at 270 nm).

This rapid I_2 production cannot be due solely to three-body atom recombination of ground state atoms. Assuming that all of the I^* atoms are quenched in a time short compared with the flash duration and that the initial iodine atom concentration is twice that of $[I_2]_{max}$, then the initial rate of I_2 production by ground state recombination at $P(CO_2) \approx 400$ Torr would be about 2.5×10^{18} molecules cm^{-3} s^{-1} , or approximately 30 times slower than actually observed.

Enhanced I atom recombination in a large excess of O_2 has been observed when the flash radiation extends as low as 200 nm [28]. This results from generation of more effective third bodies such as O_3 and singlet $O_2(^1\Delta_g)$:



TABLE 1

Molecular iodine production following room temperature flash photolysis of CF₃I in the presence of O₂

Diluent gas (~400 Torr)	P(O ₂) (Torr)	[I ₂] _{max} × 10 ⁻¹⁴ (molecules cm ⁻³)	[I ₂] _∞ × 10 ⁻¹⁴ (molecules cm ⁻³)
<i>P</i> (CF ₃ I) ≈ 10 Torr			
CO ₂	0.03		6
CO ₂	0.2		8
CO ₂	1	9	7
Xe	1	10	6
SF ₆	1	8	6
CO ₂	2	7	4
CO ₂	2	9	6
CO ₂	5	10	7
CO ₂	8	12	9
<i>P</i> (CF ₃ I) ≈ 16 Torr			
CO ₂	1	13	10
C ₃ H ₈	1	16	9
CO ₂	2	14	10
CO ₂	4	14	10
CO ₂	6	17	13
CO ₂	9	17	13
Diluent gas (~400 Torr)	<i>P</i> (CF ₃ I) (Torr)	[I ₂] _{max} × 10 ⁻¹⁴ (molecules cm ⁻³)	[I ₂] _∞ × 10 ⁻¹⁴ (molecules cm ⁻³)
<i>P</i> (O ₂) ≈ 1 Torr			
CO ₂	4	4	2
Xe	4	4	2
CO ₂	8	9	7
Xe	8	10	6
SF ₆	8	8	6
CO ₂	16	13	10
SF ₆	15	16	9
CO ₂	23	16	11



or by inclusion of the rapid bimolecular step



The O₃, O₂(¹Δ_g) and IO species result from photodissociation of O₂ in the Herzberg region near 200 nm; Antrim *et al.* [28] obtained no kinetic evidence for IO formation from I or I* atoms reacting with ground state (³Σ_g⁻) O₂ even though energetically favorable for I* atoms (and possibly I + I* atoms, depending on IO bond energy). (It should also be noted that in fact IO disappears by first order kinetics with *t*_{1/2} ≈ 85 μs [29], rather than

by the second order combination required by reaction (18); this IO lifetime is too long to account for the rapid I₂ generation reported here.) Since the UV filter solution used in the work reported here limited radiation to $\lambda > 235$ nm and since inclusion of reactions (16) and (17) would require unreasonably large ground state I atom recombination rate constants (more than 5000 times that for O₂(³Σ_g⁻) as a third body), we rule out formation of O₃, O₂(¹Δ_g) and IO as significant intermediate contributors to I₂ formation.

Although I* atoms in general recombine much slower than I atoms [16], a possible mechanism for the rapid build-up of I₂ in the presence of O₂ is the formation of a strong intermediate complex (exciplex) between I* and ground state O₂:



followed by



or



A similar exciplex between I₂ and O₂ has been proposed in the laser-induced photodecomposition of CHI₃ in the presence of trace amounts of I₂ and O₂ [30].

Stability of (O₂ · I)* may be enhanced by the near-resonant energies of I* and O₂(¹Δ_g). However, although near resonance enhances quenching of I* [31] and Br(²P_{1/2}) [32] atoms, it does not appear to be a major factor in O₂ quenching of triplet states [33]. More reasonable perhaps is a high level of charge transfer complexation resulting from the large electron affinity of the I* atom, which is greater than that of ground state I atoms by the excitation energy (0.94 eV).

We may estimate a lower limit for the strength of (O₂ · I)* if $R_0(\text{I}_2) \geq 7 \times 10^{19}$ molecules cm⁻³ s⁻¹ (i.e. that of the photolysis flash). Assuming that reaction (19) is in equilibrium (K_{19}), that reaction (20) occurs at every encounter and that $[\text{I}^*]_0 = 0.91 (2 \times [\text{I}_2]_{\text{max}})$, then it follows from simple collision frequency for $P(\text{O}_2) = 1$ Torr, $P(\text{CF}_3\text{I}) = 8$ Torr ($[\text{I}_2]_{\text{max}} \approx 10^{16}$ molecules cm⁻³, Table 1) that $K_{19} \geq 10^4$ dm³ mol⁻¹. This is very large in general compared with that for ground state I atom charge transfer complexes which in many cases may be solely contact interactions [34] although stability constants in excess of 10⁶ dm³ mol⁻¹ at 400 K have been obtained for complexes between I and NO or NO₂ [35].

3.3. Reactions following flash-initiated I₂ production

Undoubtedly other undetected transient species such as CF₃ and CF₃O₂ may contribute to the overall mechanism, as evidenced (Fig. 5) by the relatively slow first order disappearance of approximately 30% of the I₂ following the photolyzing flash. The rate constant for the reaction of trifluoromethyl radicals with iodine



is at least nine times smaller than that for two-body CF_3 recombination [36], so that an unreasonably large CF_3 concentration is required in order for reaction (22) to contribute substantially to I_2 disappearance. This is also shown directly in other work in this laboratory, to be reported later, in which CF_3I or CF_3COCF_3 is flash photolyzed with UV light in the presence of I_2 ; although I_2 is dissociated within the lifetime of the flash with a quantum yield suggesting very efficient collision energy transfer leading to I_2 dissociation in addition to reaction (22), there is no induction period in the reappearance of I_2 as would be required if reaction (22) is important several hundred microseconds after the flash.

Trifluoromethyl radicals react with O_2 to form CF_3O_2 radicals. Very little is known of the reactions of CF_3O_2 , but undoubtedly they will react to some extent with I_2 in addition to leading to fluorinated and oxygenated products such as CF_2O . If F_2 is an intermediate, such as by reaction (10), then I_2 may be decomposed by the bimolecular reaction [36, 37]



Unidentified solid products were produced in all O_2 -containing reaction mixtures, as shown by light scattering of the continuous analyzing beam immediately following the photolyzing flash. This cloudiness disappeared when the reaction mixture was frozen out with liquid nitrogen and then maintained at -39°C , or after 24 h at room temperature for cells with $P(\text{O}_2)$ less than 2 Torr. Mass spectral analyses suggested products considerably larger than CF_2O but yields were too small and variable for characterization.

Acknowledgments

Preliminary results were obtained by Mr. Raymond F. Brown (senior thesis, submitted to the Department of Chemistry, Rensselaer Polytechnic Institute). One of us (J.E.S.) was a National Science Foundation Undergraduate Research Participant, Summer, 1977.

References

- 1 J. V. V. Kasper and G. C. Pimentel, *Appl. Phys. Lett.*, 5 (1964) 231.
- 2 G. Brederlow, K. J. Witte, E. Fill, K. Hohla and R. Volk, *Digest of Technical Papers of the 1977 IEEE/OSA Conf. on Laser Engineering and Applications*, Washington, D.C., June 1 - 3, 1977, *IEEE J. Quantum Electron.*, QE-13 (9) (1977) 61D.
- 3 T. Donahue and J. R. Wiesenfeld, *J. Chem. Phys.*, 63 (1975) 3130.
- 4 C. C. Davis, R. J. Pirkle, R. A. McFarlane and G. J. Wolga, *IEEE J. Quantum Electron.*, QE-12 (1976) 334.
- 5 R. G. Derwent and B. A. Thrush, *Chem. Phys. Lett.*, 9 (1971) 591.
- 6 W. E. McDermott, N. R. Pchelkin, D. J. Benard and R. R. Bousek, *Appl. Phys. Lett.*, 32 (1978) 469.

- 7 A. T. Pritt, Jr., R. D. Coombe, D. Pilipovich, R. I. Wagner, D. Benard and C. Dymek, *Appl. Phys. Lett.*, 31 (1977) 745.
- 8 S. J. Arnold, N. Finlayson and E. A. Ogryzlo, *J. Chem. Phys.*, 44 (1966) 2529.
- 9 R. G. Derwent and B. A. Thrush, *J. Chem. Soc. Faraday Trans. 2*, 68 (1972) 720.
- 10 J. J. Deakin and D. Husain, *J. Chem. Soc. Faraday Trans. 2*, 68 (1972) 1603.
- 11 W. C. Francis and R. N. Hazeldine, *J. Chem. Soc.*, (1955) 2151.
- 12 V. I. Vedenev, M. A. Teitel'boim and A. A. Shoikhet, *Bull. Acad. Sci. USSR*, 25 (1976) 1850; 26 (1977) 480.
- 13 J. Heicklen, *J. Phys. Chem.*, 70 (1966) 112.
- 14 V. A. Brosseau, J. R. Basila, J. F. Smalley and R. L. Strong, *J. Am. Chem. Soc.*, 94 (1972) 716.
- 15 I. M. Belousova, N. G. Gorshkov, O. B. Danilov, V. Yu. Zaleskii and I. L. Yachnev, *Sov. Phys.-JETP*, 38 (1974) 254.
- 16 A. A. Filyukov, V. B. Mitrofanov and T. V. Mishchenko, *Khim. Vys. Energ.*, 10 (1976) 35.
- 17 G. A. Skorobogatov, V. G. Seleznev and O. N. Slesar, *Dokl. Akad. Nauk SSSR*, 231 (1976) 1407.
- 18 G. A. Skorobogatov, V. S. Komarov and V. G. Seleznev, *Dokl. Akad. Nauk SSSR*, 218 (1974) 886.
- 19 R. J. Donovan and D. Husain, *Trans. Faraday Soc.*, 62 (1966) 2023.
- 20 D. Husain and J. R. Wiesenfeld, *Trans. Faraday Soc.*, 63 (1967) 1349.
- 21 D. H. Burde, R. A. McFarlane and J. R. Wiesenfeld, *Chem. Phys. Lett.*, 32 (1975) 296.
- 22 H. W. Chang and G. Burns, *Can. J. Chem.*, 51 (1973) 3394.
- 23 B. P. Dymov, G. A. Skorobogatov, V. E. Khomenko and S. A. Shchukarev, *Zh. Obshch. Khim.*, 47 (1977) 848.
- 24 H. Hippler, K. Luther and J. Troe, *Ber. Bunsenges. Phys. Chem.*, 77 (1973) 1104.
- 25 J. R. Wiesenfeld and G. L. Volk, *J. Chem. Phys.*, 67 (1977) 509.
- 26 D. H. Burde and R. A. McFarlane, *J. Chem. Phys.*, 64 (1976) 1850.
- 27 K. E. Russell and J. Simons, *Proc. R. Soc. London, Ser. A*, 217 (1953) 271.
- 28 R. E. Antrim, G. Burns and J. K. K. Ip, *Can. J. Chem.*, 55 (1977) 749.
- 29 D. Yu. Stupin, V. N. Tezиков and O. M. Kaplii, *Khim. Vys. Energ.*, 7 (1973) 327.
- 30 H. Kornweitz, M. Brith and H. Friedmann, *J. Photochem.*, 6 (1977) 221.
- 31 R. J. Butcher, R. J. Donovan and R. H. Strain, *J. Chem. Soc. Faraday Trans. 2*, 70 (1974) 1839.
- 32 F. J. Wodarczyk and P. B. Sackett, *Chem. Phys.*, 12 (1976) 65.
- 33 O. L. J. Gijzeman, F. Kaufman and G. Porter, *J. Chem. Soc. Faraday Trans. 2*, 69 (1973) 708.
- 34 M. Tamres and R. L. Strong, Contact charge-transfer spectra, in R. Foster (ed.), *Molecular Association*, Vol. 2, Academic Press, London, 1979, p. 331.
- 35 H. van den Bergh and J. Troe, *J. Chem. Phys.*, 64 (1976) 736.
- 36 J. W. Birks, S. D. Gabelnick and H. S. Johnston, *J. Mol. Spectrosc.*, 57 (1975) 23.
- 37 M. J. Coggiola, J. J. Valentini and Y. T. Lee, *Int. J. Chem. Kinet.*, 8 (1976) 605.
J. J. Valentini, M. J. Coggiola and Y. T. Lee, *Discuss. Faraday Soc.*, 62 (1977) 232.