

UNIMOLECULAR FRAGMENTATION KINETICS BY MULTIPHOTON IONIZATION*

D. PROCH⁺, D.M. RIDER and R.N. ZARE

Department of Chemistry, Stanford University, Stanford, CA 94305 (U.S.A.)

Summary

The multiphoton ionization spectra of aniline ($C_6H_5NH_2$) and its perdeutero analog were investigated as a function of laser wavelength (266 - 300 nm) using a time-of-flight mass spectrometer. The appearance of a broad asymmetrically distorted peak at $m/e = 66$ is interpreted to be the slow fragmentation of the parent ion, $C_6H_7N^+ \rightarrow C_5H_6^+ + HCN$, from which a unimolecular decomposition rate is estimated to be about $2 \times 10^6 \text{ s}^{-1}$.

1. Introduction

Ever since Hipple *et al.* [1] identified that the diffuse peaks in a mass spectrum are caused by the decomposition of ions in the course of mass analysis, the importance of understanding unimolecular reactions of ions has been recognized. While electron impact mass spectrometry was the first method for observing these "metastable" ions, many new techniques have been developed to study this phenomenon. The most informative are those in which the reactant ions are created with a well-characterized distribution of internal energies and in which this distribution can be readily controlled. This usually calls for the use of vacuum UV photoionization [2 - 4]. Unfortunately, the number of such studies is very limited because of the difficulty of producing and manipulating sufficiently intense tunable vacuum UV radiation.

An alternative to one-photon vacuum UV photoionization is multiphoton ionization (MPI) in which the absorption of more than one visible or near-UV photon causes ionization [5]. The MPI technique has many advantages [6]. Because of the resonant enhancement provided by intermediate states, MPI can produce large numbers of ions with moderately intense laser sources [7]. Moreover, MPI studies only require conventional optics.

In this paper we present the first use of the MPI technique to study the unimolecular decomposition rates of ions. Aniline ($C_6H_5NH_2$) is ionized by the multiphoton absorption of pulsed UV laser light (266 - 300 nm) and the resulting ions are mass analyzed by time of flight (TOF). We investigated the elimination of HCN from the aniline parent ion $C_6H_7N^+$ to yield $C_5H_6^+$. Because of

* Paper presented at the Xth International Conference on Photochemistry, Iraklion, Crete, Greece, September 6 - 12, 1981.

⁺Permanent address: Max-Planck-Institut für Quantenoptik, 8046 Garching, F.R.G.

fragmentation during ion acceleration the daughter ion peak is asymmetrically distorted. Moreover, the extent of distortion changes with laser wavelength in a manner suggesting that the rate of fragmentation increases with photon energy.

2. Experimental details

Figure 1 shows a schematic diagram of the experimental set-up. The unfocused output of a tunable frequency-doubled dye laser (Quanta-Ray PDL-1) pumped by an Nd-YAG laser (Quanta-Ray DCR-1A) is stopped down to a diameter of 1.75 mm with two irises before passing through a TOF mass spectrometer. The wavelength range (266 - 300 nm) is obtained by frequency doubling the output of the dyes C500, R590, R610 and Kiton Red (Exciton Chemical Co.). The pulse width of the laser is 5 - 7 ns. Power densities vary depending on the wavelength. Most spectra are recorded at the power level of 14 MW cm^{-2} .

The TOF mass spectrometer was designed and built in this laboratory. The spectrometer utilizes a double-electric-field ion source [8] to accelerate the ions into a 1.5 m field-free drift region. A relatively small electric field of 126 V cm^{-1} is applied in region I (Fig. 2). This low field region serves to minimize the TOF spread of ions of a single m/e ratio formed over the diameter of the laser beam. In region II there is a much higher electric field (4710 V cm^{-1}). The ions acquire most of their kinetic energy in this region. The resolution of the instrument is better than 200, *i.e.* the value of $M/\Delta M$ at 10% of the peak height is greater than 200.

The ion bunches are detected and amplified by a microchannel plate electron multiplier (Varian model 8992-2). The signal from the multiplier is amplified and averaged with a boxcar integrator (Princeton Applied Research model 162 with model 165 plug-in). A photodiode triggers an adjustable delay gener-




Fig. 1. Schematic diagram of the MPI TOF mass spectrometer.




Fig. 2. Detailed drawing of the MPI TOF mass spectrometer ion source.

ator which in turn triggers the boxcar. The output of the boxcar drives a strip-chart recorder.

The TOF mass spectrometer is evacuated by a diffusion pump through a liquid-nitrogen-cooled cryotrap. The base pressure of the instrument is 3×10^{-8} Torr. Samples are admitted through a small tube with its orifice near the ionization region. The unfocused laser beam enters and exits through quartz windows in the vacuum chamber.

Reagent grade aniline (J. T. Baker Chemical Co.) and aniline- d_7 (98% D, Merck) were used without further purification. Both compounds were thoroughly degassed by several freeze-pump-thaw cycles before introduction into the TOF mass spectrometer. During a run, the partial pressure of aniline was 1×10^{-7} Torr, as measured with an ionization gauge.

3. Results and discussion

3.1. The time-of-flight mass spectrum of aniline

Aniline is efficiently ionized by the absorption of two 266 - 300 nm photons because the process is resonantly enhanced by one-photon absorption to the 1B_2 state [9]. The origin of the 1B_2 state is at 293.9 nm [8] and the ionization potential of aniline is 8.32 eV [10]. As expected, the parent ion signal varies with the square of the laser power.

Figure 3 shows the TOF MPI mass spectrum of aniline taken at a laser wavelength of 266 nm. The most intense feature is the parent ion ($m/e = 93$ for $C_6H_7N^+$) which is accompanied by the ${}^{13}C^{12}C_5H_7N^+$ ion at $m/e = 94$. In addition, there is extensive fragmentation. The most intriguing aspect of this fragmentation pattern is the appearance of a distorted peak in the vicinity of $m/e = 66$.

It is well known that the extent of fragmentation in MPI depends strongly on laser power [7, 9, 12, 13]. In the present study, only the parent ion of aniline appears at low power densities while, as the power density is increased, fragment ions appear. Figure 3 was recorded under increased power density at 20 MW cm^{-2} .

Figure 4 illustrates the TOF MPI mass spectrum in the region of $m/e = 64 - 68$ at several laser wavelengths. As the wavelength decreases the $m/e =$

66 peak broadens and becomes asymmetric, trailing off toward higher masses (longer arrival times). This distortion begins to appear near 285 nm. Under the same condition that the $m/e = 66$ peak broadens and distorts, the other peaks in the mass spectrum remain well resolved. Thus the appearance of the $m/e = 66$ peak is not caused by limited resolution of the TOF mass spectrometer. There are two possible isobaric components of the $m/e = 66$ peak (neglecting ^{13}C -containing ions): C_5H_6^+ and $\text{C}_4\text{H}_4\text{N}^+$. However, the equivalent ions of aniline- d_7 , C_5D_6^+ and $\text{C}_4\text{D}_4\text{N}^+$, are at $m/e = 72$ and $m/e = 70$ respectively. In the 266 nm




Fig. 3. MPI TOF mass spectrum of aniline: laser wavelength, 266 nm; laser power, 20 MW cm^{-2} .




Fig. 4. Portion of the MPI TOF mass spectrum of aniline in the region of the $m/e = 66$ peak as a function of laser wavelength. Each trace was recorded at the same laser power level (14 MW cm^{-2}).

MPI TOF mass spectrum of aniline- d_7 $m/e = 70$ is a well-resolved peak while $m/e = 72$ is quite distorted. Hence we conclude that the formation of $C_5H_6^+$ is the slow fragmentation step which causes the observed distortion. This is supported by the 70 eV electron impact mass spectrum [14] where $C_5H_6^+$ contributes 93.4% and $C_4H_4N^+$ 6.6% at a nominal value of $m/e = 66$.

In the protonated compound the $m/e = 66$ distorted peak shape has a contribution from $m/e = 67$ ($C_4H_5N^+$), which causes the appearance of a small peak or shoulder on the trailing portion of the $m/e = 66$ peak. Often the presence of the $m/e = 67$ peak is obscured by overlap with the $m/e = 66$ peak, but TOF mass spectra taken at longer wavelengths clearly show its presence. Although this complicates the $m/e = 66$ profile, the $m/e = 67$ peak does not prevent us from determining qualitatively the distortion of the $m/e = 66$ peak as a function of laser wavelength.

When $C_5H_6^+$ is produced by electron impact ionization of aniline it is found to be formed from the metastable parent ion [14, 15]. This implies that the rate of fragmentation to form $C_5H_6^+$ is relatively slow compared with non-metastable fragments. It is the relatively slow rate of fragmentation that causes the $C_5H_6^+$ to be a distorted peak in the MPI TOF mass spectrum. This can be ascertained by an analysis of ion trajectories in our TOF mass spectrometer.

Consider some parent and daughter ions that are formed at the center of region I (Fig. 2) during the laser pulse. These two types of ions will be accelerated through regions I and II and enter the field-free region with the same kinetic energy but different velocities since they have unequal masses. Traversing the field-free region they will separate in time and space and be detected at a time difference that depends only on the ratio of the square roots of their masses. If, however, some of the parent ions fragment while they are being accelerated, the resulting daughter ions enter the field-free region with a velocity which is less than that of a daughter ion formed promptly in the center of region I but greater than that of the parent ion. The resulting daughter ion arrives at the detector at some intermediate time that depends on the precise position of the fragmentation as well as the masses of the daughter and parent ions.

An ion gains most of its kinetic energy in region II and only a small fraction in region I. Decomposition of parent ions as they traverse region I produces fragments that are detected in an interval near the arrival time of a promptly formed fragment. Fragment ions formed in region II are detected in a time interval between those formed at grid 1 and the arrival time of the parent ion. Fragments formed in the field-free region arrive at the detector simultaneously with the parent ion and thus are indistinguishable from them. A fragment peak shape can thus have two detectable components. It is estimated for prompt ions with $m/e = 66 - 93$ that residence times in regions I, II and the field-free region are 400 - 500 ns, 115 - 130 ns and 18 - 20 μs respectively. The major component is caused by fragmentation in region I. The component formed by fragmentation in region II contributes less because the parent ions spend less time in this region and the flight time of the fragments is distributed over a longer time interval. The transition from the first component to the second according to calculations is 90 ns after the arrival time of a promptly formed $m/e = 66$ ion. It is observed as a sharp drop in intensity in Fig. 4. The shape of component I is a measure of

the relative number of fragments formed at each position in region I and is directly related to the fragmentation rate.

The unimolecular fragmentation rate of an ion depends on the amount of internal energy with which it is prepared. In the statistical case the rate increases rapidly with increasing internal energy [16]. Thus, with decreasing laser wavelength, more energy is deposited in the $m/e = 93$ precursor and hence we would expect that the fragmentation rate increases. Inspection of Fig. 4 shows this trend, as can readily be seen by the change in slope between the two vertical dotted lines. At the shortest wavelength, the slope is the steepest, corresponding to the most rapid rate of fragmentation.

The structure of the $C_5H_6^+$ ion is uncertain. Ocolowitz and White [17] argue that the $C_5H_6^+$ ion from the electron impact ionization of aniline is linear and suggest that it is the 3-penten-1-yne cation. In any case they determine an appearance potential from aniline for this ion to be 12.56 eV. This value is energetically equivalent to three photons at 298 nm. The distortion of the $m/e = 66$ peak first appears near 285 nm, three photons of which are equivalent to 13.1 eV. In the wavelength range studied we conclude that $C_5H_6^+$ must be formed by the absorption of three photons. To the red of 285 nm the fragmentation rate may be too slow for the production of an appreciable number of $C_5H_6^+$ ions in region I of the TOF mass spectrometer.

The calculation of a fragmentation rate from the data in Fig. 4 is complicated by the mapping of the arrival times t of the daughter ion at the detector into formation times τ of the daughter ion in the ion source. We assume that this unimolecular decomposition process is described by a single exponential. Then the number $D(\tau)$ of daughter ions formed per unit time at time τ is related to the initial number $P(0)$ of parent ions by

$$D(\tau) = kP(0) \exp(-k\tau) \quad (1)$$

where k is the rate constant. The ratio of the daughter ion formation rates at two times τ_1 and τ_2 is given by

$$\frac{D(\tau_1)}{D(\tau_2)} = \exp\{-k(\tau_1 - \tau_2)\} \quad (2)$$

We approximate the mapping of $\Delta\tau$ in the ion source to Δt at the detector by a linear transformation. Then the corresponding signal ratio at the detector is

$$\frac{S(t_1)}{S(t_2)} = \exp\{-k\alpha(t_1 - t_2)\} \quad (3)$$

where under our operating conditions the scaling constant α is approximately 5. We find from applying eqn. (3) to the 266 nm signal shape of the $C_5H_6^+$ ion shown in Fig. 4 the fragmentation rate

$$k \approx 2 \times 10^6 \text{ s}^{-1} \quad (4)$$

This value of k represents the phenomenological rate constant for the appearance of $C_5H_6^+$ ions from aniline at this wavelength and power level. Because of the various approximations introduced in this simple analysis, eqn. (4) may represent no better than an order of magnitude estimate. Computer simulations are in

progress to refine this value of k . Preliminary results suggest that the uncertainty in k may be less than a factor of 2.

The present study concerns the production of a metastable precursor to $C_5H_6^+$ in the multiphoton ionization of aniline and the observation of its fragmentation rate as a function of laser wavelength. Multiphoton ionization can prepare parent ions with known internal energies and this can be accomplished with extreme spatial definition and temporal resolution. Thus, it would appear that MPI has a bright future for the study of ionic rearrangement and/or fragmentation kinetics.

Acknowledgments

This research was supported in part by the U.S. Air Force Office of Scientific Research under Grant AFOSR-81-0053 and by the U.S. Army Research Office under Grant DAAG-29-80-K-0097.

References

- 1 J. A. Hipple, R. E. Fox and E. U. Condon, *Phys. Rev.*, **69** (1946) 347.
- 2 W. A. Chupka, Photoionization and fragmentation of polyatomic molecules, in C. Sandorfy, P. J. Ausloos and M. B. Robin (eds.), *Chemical Spectroscopy and Photochemistry in the Vacuum Ultraviolet*, Reidel, Boston, MA, 1974, pp. 431 - 463.
- 3 A. S. Werner and T. Baer, *J. Chem. Phys.*, **62** (1975) 2900.
- 4 H. M. Rosenstock, *Int. J. Mass. Spectrom. Ion Phys.*, **20** (1976) 139.
- 5 P. M. Johnson, *Acc. Chem. Res.*, **13** (1980) 20, and references therein.
- 6 V. S. Antonov and V. S. Letokhov, *Appl. Phys.*, **24** (1981) 89.
- 7 D. M. Lubman, R. Naaman and R. N. Zare, *J. Chem. Phys.*, **72** (1980) 3034.
- 8 W. C. Wiley and I. H. McLaren, *Rev. Sci. Instrum.*, **26** (1955) 1150.
- 9 C. T. Rettner and J. H. Brophy, *Chem. Phys.*, **56** (1981) 53.
- 10 J. H. Brophy and C. T. Rettner, *Opt. Lett.*, **4** (1979) 337; *Chem. Phys. Lett.*, **67** (1979) 351.
- 11 T. Cuitas, J. M. Hollas and G. H. Kirby, *Mol. Phys.*, **19** (1970) 305.
- 12 H. M. Rosenstock, K. Draxl, B. W. Steiner and J. T. Herron, *J. Phys. Chem. Ref. Data*, **6**, Suppl. 1 (1977) I-235.
- 13 L. Zandee and R. B. Bernstein, *J. Chem. Phys.*, **70** (1979) 2574; **71** (1979) 1359.
- 14 S. Rockwood, J. P. Reilly, K. Hohla and K. L. Kompa, *Opt. Commun.*, **28** (1979) 175.
- 15 A. V. Robertson and C. Djerassi, *J. Am. Chem. Soc.*, **90** (1968) 6992.
- 16 P. N. Rylander, S. Meyerson, E. L. Eliel and J. D. McCollum, *J. Am. Chem. Soc.*, **8** (1963) 2723.
- 17 F. W. McLafferty, *Interpretation of Mass Spectra*, Benjamin, Reading, MA, 1973.
- 18 J. L. Occolowitz and G. L. White, *Aust. J. Chem.*, **21** (1968) 997.