

## Theoretical Thermochemistry of the 1-Buten-3-yn-1-yl Radical and Its Chloro Derivatives

Jerzy Cioslowski,\* Guanghua Liu, and David Moncrieff

Department of Chemistry and Supercomputer Computations Research Institute, Florida State University, Tallahassee, Florida 32306-3006

Received: August 24, 1999; In Final Form: October 19, 1999

High-level electronic structure calculations combined with empirical adjustments predict the standard enthalpy of the C–H bond dissociation in  $\text{HC}\equiv\text{C}-\text{CH}=\text{CH}_2$  to be equal to  $115.1 \pm 1.4$  kcal/mol, i.e., ca. 4.0 kcal/mol higher than that of the analogous bond cleavage in ethene. This difference in bond strengths stems from resonance stabilization of the parent molecule. The standard enthalpy of formation of the 1-buten-3-yn-1-yl radical is estimated at  $133.8 \pm 1.5$  kcal/mol, which is significantly higher than all of the previously published values. As in the case of polychlorinated alkanes, the BLYP approximation is found to seriously underestimate the strengths of the C–H and C–Cl bonds in chloro derivatives of  $\text{HC}\equiv\text{C}-\text{CH}=\text{CH}_2$ . On the other hand, the BLYP/6-311G\*\*, MP2/6-311G\*\*, QCISD/6-311G\*\*, and CCSD(T)/6-311G\*\* predictions for the standard enthalpy of ethyne dimerization all closely match their experimental counterpart.

## Introduction

The 1-buten-3-yn-1-yl radical ( $\text{HC}\equiv\text{C}-\text{CH}=\text{CH}\cdot$ ,  $n\text{-C}_4\text{H}_3\cdot$ ) is believed to play a pivotal role in combustion of small hydrocarbons such as  $\text{CH}_4$  and  $\text{C}_2\text{H}_2$ . For many years, the reaction



has been regarded as the initiation step of the thermally induced radical-chain polymerization of ethyne.<sup>1</sup> However, the rate of formation of  $n\text{-C}_4\text{H}_3\cdot$  offered by this bimolecular process has been shown to be much too low to account for the observed polymerization kinetics.<sup>2–4</sup> Invoking reaction mechanisms that involve autocatalytic processes<sup>5</sup> does not eliminate this discrepancy.<sup>6</sup>

In fuel-rich  $\text{C}_2\text{H}_2/\text{O}_2$  and  $\text{CH}_4/\text{O}_2$  flames, benzene is produced primarily via recombination of propargyl radicals.<sup>7,8</sup> However, in fuel-lean  $\text{C}_2\text{H}_2/\text{O}_2$  flames the addition of  $\text{C}_2\text{H}_2$  to  $n\text{-C}_4\text{H}_3\cdot$  constitutes an important reaction pathway to benzene.<sup>7,9</sup> Analogous reactions take place during pyrolysis and combustion of polychlorinated hydrocarbons, such as  $\text{C}_2\text{HCl}_3$  and  $\text{C}_2\text{Cl}_4$ .<sup>10–12</sup> Thus, the formation of hexachlorobenzene during pyrolysis of those compounds most probably involves the  $n\text{-C}_4\text{Cl}_3\cdot$  radical pathway,<sup>11</sup> which begins with dichloroethyne.<sup>10</sup>

Despite its importance to modeling of combustion reaction kinetics, the standard enthalpy of formation  $\Delta H_f^\circ(\text{HC}\equiv\text{C}-\text{CH}=\text{CH}\cdot)$  of the 1-buten-3-yn-1-yl radical has not been accurately established. The few estimates of  $\Delta H_f^\circ(\text{HC}\equiv\text{C}-\text{CH}=\text{CH}\cdot)$  that have been published so far vary widely: 124,<sup>1</sup> 125,<sup>8</sup> 127,<sup>13</sup> and 130 kcal/mol.<sup>6</sup> Even worse, the thermochemistry of chloro derivatives of  $n\text{-C}_4\text{H}_3\cdot$  has not been investigated either experimentally or theoretically. This lack of data has prompted the research described in this paper.

## Details of Calculations

All the calculations were carried out with the Gaussian 94 suite of programs.<sup>14</sup> Full geometry optimizations were performed

\* To whom all correspondence should be addressed. E-mail: jerzy@kyoko.chem.fsu.edu. WWW: <http://www.scri.fsu.edu/~jerzy>.

TABLE 1: Computed Standard Enthalpies of the C–H Bond Dissociation in the  $\text{CX}_1\text{X}_2=\text{CX}_3\text{H}$  Molecules<sup>a</sup>

$\text{X}_1$	$\text{X}_2$	$\text{X}_3$	BLYP	MP2	QCISD	CCSD(T)
H	H	H	106.72	112.21	107.13	107.91
H	H	Cl	104.70	112.04	106.03	106.64
H	Cl	Cl	106.53	115.08	108.10	108.76
Cl	H	Cl	104.35	113.70	106.79	107.36
H	$\text{HC}\equiv\text{C}$	H	109.50	123.60	112.12	112.90
$\text{HC}\equiv\text{C}$	H	H	109.52	123.33	112.12	112.87
Cl	$\text{ClC}\equiv\text{C}$	H	107.75	124.79	112.54	113.22
$\text{ClC}\equiv\text{C}$	Cl	H	109.40	124.96	113.32	114.06
H	$\text{ClC}\equiv\text{C}$	Cl	106.59	122.83	110.10	110.76
$\text{ClC}\equiv\text{C}$	H	Cl	106.35	122.73	109.82	110.49
Cl	$\text{HC}\equiv\text{C}$	Cl	104.91	124.16	110.57	111.19
$\text{HC}\equiv\text{C}$	Cl	Cl	106.54	125.10	111.34	112.06

<sup>a</sup> All values in kcal/mol. The substituents  $\text{X}_1$  and  $\text{X}_3$  are in the *cis* position, the latter being attached to the carbon atom of the C–H bond undergoing dissociation.

at the BLYP/6-311G\*\*, MP2/6-311G\*\*, and QCISD/6-311G\*\* levels of theory. All structures were confirmed to be local minima on potential energy hypersurfaces by the computed BLYP/6-311G\*\* and MP2/6-311G\*\* vibrational frequencies. CCSD(T)/6-311G\*\* energies at the optimized QCISD/6-311G\*\* geometries were also obtained. Spin-unrestricted wave functions were used as reference states in all calculations on radicals.

The standard enthalpies of all the species under study were calculated from the respective geometries and unscaled harmonic vibrational frequencies. The MP2/6-311G\*\* zero-point energies and thermal corrections were used to convert the QCISD/6-311G\*\* and CCSD(T)/6-311G\*\* energies to the corresponding standard enthalpies.

## Results and Discussion

The standard enthalpy  $\Delta H_{\text{C-H}}^\circ(\text{H}_2\text{C}=\text{CH}\cdot\text{H})$  of the C–H bond dissociation in ethene equals  $111.2 \pm 0.8$  kcal/mol.<sup>15</sup> Among the four levels of theory employed in the present study, the MP2 approximation used in conjunction with the 6-311G\*\* basis set appears to match this experimental figure most closely (Table 1). However, this agreement undoubtedly arises from a

**TABLE 2: Computed Standard Enthalpies of the C–Cl Bond Dissociation in the  $CX_1X_2=CX_3Cl$  Molecules<sup>a</sup>**

X <sub>1</sub>	X <sub>2</sub>	X <sub>3</sub>	BLYP	MP2	QCISD	CCSD(T)
H	H	H	88.14	98.63	86.40	89.13
H	Cl	H	86.83	99.93	86.52	89.38
Cl	H	H	85.06	98.65	85.49	88.23
H	Cl	Cl	81.51	97.61	82.35	85.34
Cl	H	Cl	79.31	95.82	80.96	83.82
H	C≡C	H	89.87	109.65	90.24	93.18
C≡C	H	H	89.99	109.41	90.56	93.43
Cl	HC≡C	H	85.63	109.67	89.31	92.19
HC≡C	Cl	H	87.30	110.23	90.48	93.42
H	HC≡C	Cl	82.78	107.17	85.78	88.80
HC≡C	H	Cl	82.65	107.02	85.76	88.75
Cl	C≡C	Cl	79.13	106.20	83.72	86.87
C≡C	Cl	Cl	80.79	107.58	84.90	88.14

<sup>a</sup> All values in kcal/mol. The substituents X<sub>1</sub> and X<sub>3</sub> are in the *cis* position, the latter being attached to the carbon atom of the C–Cl bond undergoing dissociation.

**TABLE 3: Empirically Adjusted Predictions for the Standard Enthalpies of the C–H Bond Dissociation in the  $CX_1X_2=CX_3H$  Molecules<sup>a</sup>**

X <sub>1</sub>	X <sub>2</sub>	X <sub>3</sub>	BLYP	MP2	QCISD	CCSD(T)
H	H	H	111.2 <sup>b</sup>	111.2 <sup>b</sup>	111.2 <sup>b</sup>	111.2 <sup>b</sup>
H	H	Cl	109.2	111.0	110.1	109.9
H	Cl	Cl	111.0	114.1	112.2	112.1
Cl	H	Cl	108.8	112.7	110.9	110.7
H	HCC	H	114.0	122.6	116.2	116.2
HCC	H	H	114.0	122.3	116.2	116.2
Cl	CICC	H	112.2	123.8	116.6	116.5
CICC	Cl	H	113.9	124.0	117.4	117.4
H	CICC	Cl	111.1	121.8	114.2	114.1
CICC	H	Cl	110.8	121.7	113.9	113.8
Cl	HCC	Cl	109.4	123.2	114.6	114.5
HCC	Cl	Cl	111.0	124.1	115.4	115.4

<sup>a</sup> All values in kcal/mol. The substituents X<sub>1</sub> and X<sub>3</sub> are in the *cis* position, the latter being attached to the carbon atom of the C–H bond undergoing dissociation. <sup>b</sup> Assumed (see the text for explanation).

fortuitous cancellation of two errors, namely, the underestimation of stability of the radical due to spin contamination and the relative underestimation of stability of the parent molecule due to the low level of electron correlation treatment and the moderate size of the basis set. Indeed, a closer inspection of the data computed for the radical reveals the Hartree–Fock ( $\langle\hat{S}^2\rangle$ ) value of 0.921 and the difference between the MP2 and the spin-projected MP2 (PMP2) energies amounting to 6.0 kcal/mol.

The standard enthalpy of the C–Cl bond dissociation in chloroethene (vinyl chloride) can be readily computed from the published values of  $\Delta H_f^\circ$  for  $H_2C=CH^\bullet$ , Cl<sup>•</sup>, and  $H_2C=CHCl$ , which equal  $71.6 \pm 0.8$ , 29.0,<sup>15,20</sup> and  $5.5 \pm 0.5$  kcal/mol,<sup>16</sup> respectively. The resulting  $\Delta H_{C-Cl}^\circ(H_2C=CH\cdot Cl)$  of  $95.1 \pm 0.9$  kcal/mol is again best reproduced at the MP2/6-311G\*\* level of theory (Table 2). Both the C–H and C–Cl bonds are predicted to weaken upon chlorine substitution in the parent molecules. This trend is much more prominent in the BLYP/6-311G\*\* predictions than in those obtained with the conventional treatments of electron correlation. This steep decline in the BLYP/6-311G\*\* standard enthalpies of the C–Cl bond dissociation is reminiscent of a similar artifact present in the analogous data for the  $C_2H_{6-n}Cl_n$  series, which has been explained by the inability of DFT-based methods to account for the stabilizing dispersion interactions between proximate chlorine atoms.<sup>17</sup>

The known values of  $\Delta H_{C-H}^\circ(H_2C=CH\cdot H)$  and  $\Delta H_{C-Cl}^\circ(H_2C=CH\cdot Cl)$  can be used to correct the theoretical

**TABLE 4: Empirically Adjusted Predictions for the Standard Enthalpies of the C–Cl Bond Dissociation in the  $CX_1X_2=CX_3Cl$  Molecules<sup>a</sup>**

X <sub>1</sub>	X <sub>2</sub>	X <sub>3</sub>	BLYP	MP2	QCISD	CCSD(T)
H	H	H	95.1 <sup>b</sup>	95.1 <sup>b</sup>	95.1 <sup>b</sup>	95.1 <sup>b</sup>
H	Cl	H	93.8	96.4	95.2	95.3
Cl	H	H	92.0	95.1	94.2	94.2
H	Cl	Cl	88.5	94.1	91.0	91.3
Cl	H	Cl	86.3	92.3	89.7	89.8
H	C≡C	H	96.8	106.1	98.9	99.2
C≡C	H	H	96.9	105.9	99.3	99.4
Cl	HC≡C	H	92.6	106.1	98.0	98.2
HC≡C	Cl	H	94.3	106.7	99.2	99.4
H	HC≡C	Cl	89.7	103.6	94.5	94.8
HC≡C	H	Cl	89.6	103.5	94.5	94.7
Cl	C≡C	Cl	86.1	102.7	92.4	92.8
C≡C	Cl	Cl	87.8	104.0	93.6	94.1

<sup>a</sup> All values in kcal/mol. The substituents X<sub>1</sub> and X<sub>3</sub> are in the *cis* position, the latter being attached to the carbon atom of the C–Cl bond undergoing dissociation. <sup>b</sup> Assumed (see the text for explanation).

**TABLE 5: QCISD/6-311G\*\* Optimized Geometries of the  $X_1C_a=C_b-C_cX_2=C_dX_3X_4$  Molecules<sup>a</sup>**

X <sub>1</sub>	X <sub>2</sub>	X <sub>3</sub>	X <sub>4</sub>	C <sub>a</sub> –C <sub>b</sub>	C <sub>b</sub> –C <sub>c</sub>	C <sub>c</sub> –C <sub>d</sub>	C <sub>a</sub> –C <sub>b</sub> –C <sub>c</sub>	C <sub>b</sub> –C <sub>c</sub> –C <sub>d</sub>
H	H	H	H	1.213	1.441	1.344	178.2	123.4
H	H	H	•	1.209	1.449	1.325	178.7	122.5
H	H	•	H	1.208	1.445	1.325	177.6	124.1
H	H	Cl	Cl	1.212	1.431	1.344	177.3	124.0
H	H	Cl	•	1.208	1.442	1.330	178.6	123.6
H	H	•	Cl	1.208	1.441	1.328	177.6	122.6
H	Cl	H	Cl	1.212	1.434	1.344	179.5	120.7
H	Cl	H	•	1.208	1.441	1.320	178.9	122.8
H	Cl	•	Cl	1.207	1.436	1.326	179.8	122.0
H	Cl	Cl	H	1.211	1.430	1.343	179.5	124.9
H	Cl	Cl	•	1.207	1.436	1.328	179.0	123.7
H	Cl	•	H	1.207	1.439	1.318	180.0	124.4
Cl	H	H	Cl	1.211	1.435	1.341	177.8	121.9
Cl	H	H	•	1.207	1.447	1.325	178.3	122.6
Cl	H	•	Cl	1.206	1.438	1.329	177.1	122.7
Cl	H	Cl	H	1.211	1.432	1.343	176.8	125.0
Cl	H	Cl	•	1.207	1.440	1.330	178.1	123.7
Cl	H	•	H	1.207	1.443	1.325	177.1	124.2
Cl	Cl	H	H	1.210	1.434	1.341	179.8	123.9
Cl	Cl	H	•	1.206	1.438	1.320	178.9	123.0
Cl	Cl	•	H	1.205	1.436	1.318	179.8	124.5
Cl	Cl	Cl	Cl	1.210	1.426	1.350	178.9	122.7
Cl	Cl	Cl	•	1.206	1.433	1.328	179.1	123.8
Cl	Cl	•	Cl	1.206	1.433	1.326	179.6	122.1

<sup>a</sup> Bond lengths in angstroms and bond angles in degrees. The substituents X<sub>2</sub> and X<sub>4</sub> are in the *cis* position. The dot denotes the unpaired electron.

predictions by uniformly shifting them to match the experimental data (note that applying such a correction is equivalent to employing the isodesmic reaction  $H_2C=CH\cdot X + R^\bullet \rightarrow H_2C=CH^\bullet + R\cdot X$ , X = H or Cl, in the calculation of  $\Delta H_{C-X}^\circ$ ). The resulting adjusted standard enthalpies of the C–H and C–Cl bond dissociations are listed in Tables 3 and 4. There is a reasonable agreement among the adjusted data computed for the chloro derivatives of ethene, except for the BLYP/6-311G\*\* predictions pertaining to the polychlorinated compounds. On the other hand, the adjusted MP2/6-311G\*\* standard enthalpies of bond dissociations in *n*-C<sub>4</sub>H<sub>3</sub>• and its chloro analogs deviate strongly from their BLYP, QCISD, and CCSD(T) counterparts. These discrepancies are most probably attributable to the difference in the degree of spin contamination in the unrestricted Hartree–Fock electronic wave functions of the  $H_2C=CH^\bullet$  and *n*-C<sub>4</sub>H<sub>3</sub>• radicals. Indeed, the MP2 energy of *n*-C<sub>4</sub>H<sub>3</sub>• exceeds its spin-projected MP2 (PMP2) counterpart by 10.0 kcal/mol, which is 4.0 kcal/mol greater than the corresponding value for

TABLE 6: Computed Standard Enthalpies of Dimerization<sup>a</sup>

monomer	dimer	BLYP	MP2	QCISD	CCSD(T)
HC≡CH	HC≡C-CH=CH <sub>2</sub>	-39.54	-38.36	-38.11	-38.69
HC≡CCl	HC≡C-CH=CCl <sub>2</sub>	-50.50	-52.04	-50.50	-51.60
	(Z)-HC≡C-CCl=CHCl	-50.25	-52.01	-50.77	-51.83
	(E)-HC≡C-CCl=CHCl	-50.20	-51.55	-50.33	-51.39
	(Z)-ClC≡C-CH=CClH	-47.18	-47.31	-46.23	-47.08
	(E)-ClC≡C-CH=CHCl	-47.26	-47.25	-46.48	-47.31
	ClC≡C-CCl=CH <sub>2</sub>	-45.62	-46.42	-45.37	-46.35
ClC≡CCl	ClC≡C-CCl=CCl <sub>2</sub>	-54.57	-58.74	-55.71	-57.37

<sup>a</sup> All values in kcal/mol.

H<sub>2</sub>C=CH• (see above). This difference in spin contamination, which is very small at the BLYP/6-311G\*\* level of theory, is much less important than that in the case of the QCISD and CCSD(T) approximations.

In light of the above discussion,  $\Delta H_{C-H}^{\circ}(\text{HC}\equiv\text{C}-\text{CH}=\text{CH}\cdot\text{H}) = 115.1 \pm 1.4$  kcal/mol (which is an average of the corrected CCSD(T)/6-311G\*\* and BLYP/6-311G\*\* predictions with the error bars arrived at by combining the experimental uncertainty in  $\Delta H_{C-H}^{\circ}(\text{H}_2\text{C}=\text{CH}\cdot\text{H})$  employed in the aforedescribed adjustment procedure with the difference between these predictions) constitutes an accurate estimate of the standard enthalpy of the C-H bond dissociation that produces *n*-C<sub>4</sub>H<sub>3</sub>•. In order to compute  $\Delta H_f^{\circ}(\text{HC}\equiv\text{C}-\text{CH}=\text{CH}\cdot)$ , this estimate has to be combined with  $\Delta H_f^{\circ}(\text{H}) = 52.1$  kcal/mol (refs 15 and 20) and  $\Delta H_f^{\circ}(\text{HC}\equiv\text{C}-\text{CH}=\text{CH}_2) = 70.8 \pm 0.5$  kcal/mol, the latter value being readily inferred from the standard enthalpy of hydrogenation of H<sub>2</sub>C=CH-C≡CH (100.8 ± 0.5 kcal/mol (ref 18) and  $\Delta H_f^{\circ}(n\text{-C}_4\text{H}_{10}) = -30.0 \pm 0.2$  kcal/mol (ref 19)). The resulting  $\Delta H_f^{\circ}(\text{HC}\equiv\text{C}-\text{CH}=\text{CH}\cdot) = 133.8 \pm 1.5$  kcal/mol is significantly higher than all of the known estimates.<sup>1,6,8,13</sup>

The key difference between the present value of  $\Delta H_f^{\circ}(\text{HC}\equiv\text{C}-\text{CH}=\text{CH}\cdot)$  and the previously published data lies in the standard enthalpy of the C-H bond dissociation in the parent molecule, which has been usually assumed to be identical with that in ethene. However, the data compiled in Tables 3 and 4 unequivocally show that the terminal C-H and C-Cl bonds of the ethene moieties in HC≡C-CH=CH<sub>2</sub> and its chloro derivatives are stronger than those in the corresponding chloroethenes. This bond strengthening is caused by a resonance stabilization of HC≡C-CH=CH<sub>2</sub> and its congeners, which is conspicuously reflected in their QCISD/6-311G\*\* optimized geometries (Table 5). Thus, the C≡C bond length in HC≡C-CH=CH<sub>2</sub> equals 1.213 Å, as opposed to 1.209, 1.208, and 1.210 Å in the two geometrical isomers of HC≡C-CH=CH• and in HC≡CH. Similarly, the C=C bond in HC≡C-CH=CH<sub>2</sub> is longer (1.344 Å) than those in the HC≡C-CH=CH• radicals (1.325 Å), the H<sub>2</sub>C=CH• radical (1.320 Å), and H<sub>2</sub>C=CH<sub>2</sub> (1.339 Å). As expected, the C-C bond in HC≡C-CH=CH<sub>2</sub> lengthens from 1.441 Å to 1.445/1.449 Å upon the C-H bond dissociation due to the concomitant diminution of conjugation.

It is of interest to compare the predictions for the standard enthalpy of the ethyne dimerization with the corresponding experimental value. By combining  $\Delta H_f^{\circ}(\text{HC}\equiv\text{CH}) = 54.2 \pm 0.2$  kcal/mol<sup>20</sup> with the previously mentioned  $\Delta H_f^{\circ}(\text{HC}\equiv\text{C}-\text{CH}=\text{CH}_2) = 70.8 \pm 0.5$  kcal/mol, one arrives at  $\Delta H_{\text{dim}}^{\circ}(\text{HC}\equiv\text{CH}) = 37.6 \pm 0.6$  kcal/mol, which is in an excellent agreement with the data computed at all four levels of theory (Table 6). The computed reaction enthalpies also imply the dimerization of ClC≡CCl to be much more exothermic than that of HC≡CH. Among the products of the HC≡CCl,

HC≡C-CH=CCl<sub>2</sub> and the two geometrical isomers of HC≡C-CCl=CHCl are found to be equally stable and preferred by 4–5 kcal/mol over the other three species.

## Conclusions

High-level electronic structure calculations combined with empirical adjustments predict the standard enthalpy of the C-H bond dissociation in HC≡C-CH=CH<sub>2</sub> to be equal to 115.1 ± 1.4 kcal/mol, i.e., ca. 4.0 kcal/mol higher than that of the analogous bond cleavage in ethene. This difference in bond strengths stems from resonance stabilization of the parent molecule. The standard enthalpy of formation of the 1-buten-3-yn-1-yl radical is estimated at 133.8 ± 1.5 kcal/mol, which is significantly higher than all of the previously published values.

As in the case of polychlorinated alkanes, the BLYP approximation is found to seriously underestimate the strengths of the C-H and C-Cl bonds in chloro derivatives of HC≡C-CH=CH<sub>2</sub>. On the other hand, the BLYP/6-311G\*\*, MP2/6-311G\*\*, QCISD/6-311G\*\*, and CCSD(T)/6-311G\*\* predictions for the standard enthalpy of ethyne dimerization all closely match their experimental counterpart.

**Acknowledgment.** The research described in this publication has been supported by the Office of Energy Research, Office of Basic Energy Sciences, Division of Chemical Sciences, U.S. Department of Energy under Grant DE-FG02-97ER14758. Support by Florida State University through the allocation of supercomputer resources on the SGI Power Challenge computer is acknowledged.

## References and Notes

- Colket, M. B., III; Seery, D. J. *Combust. Flame* **1989**, *75*, 343 and references therein.
- Duran, R. P.; Amorebieta, V. T.; Colussi, A. J. *Int. J. Chem. Kinet.* **1989**, *21*, 947.
- Ghibaudi, E.; Colussi, A. J. *J. Phys. Chem.* **1988**, *92*, 5839.
- Duran, R. P.; Amorebieta, V. T.; Colussi, A. J. *J. Phys. Chem.* **1988**, *92*, 636.
- Benson, S. W. *Int. J. Chem. Kinet.* **1992**, *24*, 217.
- Kiefer, J. H. *Int. J. Chem. Kinet.* **1993**, *25*, 215 and references therein.
- Lindstedt, R. P.; Skevis, G. *Combust. Sci. Technol.* **1997**, *125*, 73.
- Leung, K. M.; Lindstedt, R. P. *Combust. Flame* **1995**, *102*, 129.
- Westmoreland, P. R.; Dean, A. M.; Howard, J. B.; Longwell, J. P. *J. Phys. Chem.* **1989**, *93*, 8171 and references therein.
- Chang, W. D.; Senkan, S. M. *Environ. Sci. Technol.* **1989**, *23*, 442.
- Taylor, P. H.; Tirey, D. A.; Rubey, W. A.; Dellinger, B. *Combust. Sci. Technol.* **1994**, *101*, 75.
- Tirey, D. A.; Taylor, P. H.; Kasner, J.; Dellinger, B. *Combust. Sci. Technol.* **1990**, *74*, 137.
- Wang, H.; Frenklach, M. *J. Phys. Chem.* **1994**, *98*, 11465.
- Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Gill, P. M. W.; Johnson, B. G.; Robb, M. A.; Cheeseman, J. R.; Keith, T. A.; Petersson, G. A.; Montgomery, J. A.; Raghavachari, K.; Al-Laham, M. A.; Zakrzewski, V. G.; Ortiz, J. V.; Foresman, J. B.; Cioslowski, J.; Stefanov, B. B.; Nanayakkara, A.; Challacombe, M.; Peng, C. Y.; Ayala, P. Y.; Chen, W.; Wong, M. W.; Andres, J. L.; Replogle, E. S.; Gomperts, R.; Martin, R. L.;

Fox, D. J.; Binkley, J. S.; Defrees, D. J.; Baker, J.; Stewart, J. J. P.; Head-Gordon, M.; Gonzalez, C.; Pople, J. A. *Gaussian 94*; Gaussian, Inc.: Pittsburgh, PA, 1995.

(15) Berkowitz, J.; Ellison, G. B.; Gutman, D. *J. Phys. Chem.* **1994**, *98*, 2744.

(16) Kolesov, V. P.; Papina, T. S. *Russ. Chem. Rev.* **1983**, *52*, 425; see Ochterski, J. W.; Petersson, G. A.; Wiberg, K. B. *J. Am. Chem. Soc.* **1995**, *117*, 11299 and references therein for the comments on the standard enthalpy of formation of  $\text{H}_2\text{C}=\text{CHCl}$ .

(17) Cioslowski, J.; Liu, G.; Moncrieff, D. *J. Am. Chem. Soc.* **1997**, *119*, 11452.

(18) Roth, W. R.; Adamczak, O.; Breuckmann, R.; Lennartz, H.-W.; Boese, R. *Chem. Ber.* **1991**, *124*, 2499.

(19) Pedley, J. B.; Naylor, R. D.; Kirby, S. P. *Thermochemical Data of Organic Compounds*, 2nd ed.; Chapman and Hall: New York, 1986.

(20) Chase, M. W.; Davies, C. A.; Downey, J. R.; Frurip, D. R.; McDonald, R. A.; Syverud, A. N. JANAF Thermochemical Tables, 3rd ed. *J. Phys. Chem. Ref. Data* **1985**, *1*, 14 (Suppl. 1).