COMMENTS

Comment on "Theoretical Investigation of the Formation Mechanism of Metallofullerene $Y@C_{82}$ "

Douglas L. Strout*

Department of Physical Sciences, Alabama State University, Montgomery, Alabama 36101

Received: June 30, 2005

A recent study¹ by Gan and Wang characterized a formation mechanism for the metallofullerene Y@C₈₂, and the energetically favorable path was determined to be the reaction C_{76} + YC₆ \rightarrow Y@C₈₂. The YC₆ reactant was portrayed as a sixmembered ring of carbon coordinated to a yttrium atom in η^6 fashion, with $C_{6\nu}$ or near- $C_{6\nu}$ point group symmetry. However, what is the stability of this proposed reactant relative to other YC₆ isomers? Will the YC₆ exist in the proposed form with a sufficient lifetime to perform its proposed role in the mechanism? By the use of the B3LYP/LANL2DZ method^{2,3} of the previous study, this isomer of YC₆ is compared to two other YC₆ isomers from a previous study⁴ by Strout and Hall. These two other isomers are shown in Figures 1 and 2. Figure 1 shows




Figure 1. Fan isomer of YC₆ (C_{2v} point group symmetry).




Figure 2. Planar ring isomer of YC_6 ($C_{2\nu}$ point group symmetry).

the so-called "fan isomer" in which the yttrium atom is coordinated to a linear chain of six carbons in such a way as to have Y-C bonding distances with all six carbon atoms. Figure 2 shows a planar isomer in which the yttrium atom is coordinated to a six-membered carbon ring in η^2 fashion.

The first major result is that the geometry optimization of a C_{6v} isomer was unsuccessful due to gradients that suggest that the six Y–C distances should be nonidentical. The stationary

point most similar to the previous authors' $C_{6\nu}$ isomer was found in $C_{2\nu}$ symmetry and is shown in Figure 3. Even this structure




Figure 3. Nonplanar ring isomer of YC₆ ($C_{2\nu}$ point group symmetry). This structure resulted from unsuccessful attempts to optimize the structure with $C_{6\nu}$ symmetry.

is not quite a local minimum, having a single imaginary frequency of 135*i*. The molecules in Figures 1 and 2 are local minima at the B3LYP/LANL2DZ level of theory. Table 1 shows

TABLE 1: Relative Energies of Three Isomers of YC6(calculated with the B3LYP/LANL2DZ method, energies in kcal/mol)

| isomer | energy |
|----------------------------------|--------|
| fan isomer (Figure 1) | 0.0 |
| planar ring isomer (Figure 2) | +19.3 |
| nonplanar ring isomer (Figure 3) | +63.1 |

the relative energies of these three stationary points. The fan isomer is the lowest in energy, followed by the planar η^2 ring isomer, with the nonplanar isomer lying much higher in energy. Given this energy ordering of the isomers, it is plausible to envision a reaction path whereby the yttrium atom of the nonplanar isomer slides down to its position on the planar ring, followed by insertion of the yttrium into the ring to form a fan isomer. If the barrier between nonplanar ring isomer and planar ring isomer is a high one, then the nonplanar ring isomer may be stable enough to perform its proposed role in the formation of Y@C₈₂. However, that would have to be demonstrated to be the case in order for the nonplanar YC₆ to be a plausible reactant in a Y@C₈₂ reaction mechanism.

Acknowledgment. The Alabama Supercomputer Authority is gratefully acknowledged for a grant of computer time on the SGI Altix operated in Huntsville, AL. This work is also supported by the National Institutes of Health (NIH/NCMHD grant 1P20MD000547-01). The taxpayers of the state of Alabama in particular and the U.S. in general are also gratefully acknowledged.

References and Notes

- (1) Gan, L.-H.; Wang, C.-R. J. Phys. Chem. A 2005, 109, 3980.
- (2) Becke, A. D. J. Chem. Phys. **1993**, 98, 5648. Lee, C.; Yang, W.; Parr, R. G. Phys. Rev. B **1988**, 37, 785.
 - (3) Hay, P. J.; Wadt, W. R. J. Chem. Phys. **1985**, 82, 299.
 - (4) Strout, D. L.; Hall, M. B. J. Phys. Chem. 1996, 100, 18007.

^{*} E-mail: dstrout@alasu.edu.