

Kinetics and mechanism for oxime formation from methyl pyruvate

T. Córdova, A. J. Peraza, M. Calzadilla and A. Malpica*

Escuela de Química, Facultad de Ciencias, Universidad Central de Venezuela, Caracas, 47102, Venezuela


Received 26 February 2001; revised 2 July 2001; accepted 20 July 2001

ABSTRACT: Rate and equilibrium constants for methyl pyruvate oxime formation were determined as a function of pH over the range 0–7 in aqueous solution at 30 °C and ionic strength 0.5 by spectrophotometric methods. The reaction occurs with rate-determining carbinolamine dehydration over the entire range of pH investigated. Carbinolamine dehydration is not susceptible to detectable general acid–base catalysis by a carboxylic acid buffer or hydroxylamine/hydroxylammonium ion buffer. Specific acid catalysis for carbinolamine formation is dominant at pH values below 5. Above that value, a pH-independent, water-catalyzed reaction becomes apparent. The pH-independent carbinolamine dehydration is unusually important with this substrate. Copyright © 2001 John Wiley & Sons, Ltd.

KEYWORDS: methyl pyruvate; oxime; kinetics; mechanism

INTRODUCTION

Imine formation from moderately basic amines and moderately reactive carbonyl compounds occurs with a single break in the pH–rate profile over the pH range 0–7.^{1–3} This break reflects a transition from rate-determining amine addition under acidic conditions to rate-determining carbinolamine dehydration as the pH is increased.



When the electrophilicity of the carbonyl group and the nucleophilicity of the amine are enhanced, the rate of carbinolamine formation is increased without a compensating increase in the rate of carbinolamine dehydration. Thus, carbinolamine dehydration becomes rate determining over this pH range and the break in the pH–rate profile disappears.^{4–8}

In the work reported herein, we examine the kinetics and mechanism for addition of hydroxylamine to methyl pyruvate. This work is motivated by the desire to provide an additional example of mechanism and reactivity for imine formation to test and extend the detailed mechanistic proposal of Jencks and Sayer.^{1,2} In addition, the

choice of methyl pyruvate as substrate is motivated by the need to simulate certain aspects of pyruvic acid oxime formation. In a previous study of pyruvic acid oxime formation, it proved impossible to determine directly the equilibrium constant for carbinolamine formation from pyruvic acid (as opposed to the pyruvate anion).⁷ The low values of pH required to form pyruvic acid convert substantially all the hydroxylamine to its conjugate acid. Consequently, accumulation of the carbinolamine from pyruvic acid and hydroxylamine free base could not be observed. This equilibrium constant was estimated from: (i) the relative extents of hydration of pyruvic acid and its anion; (ii) the equilibrium constant for addition of hydroxylamine to pyruvate; (iii) the assumption that the ratio of equilibrium constants for addition of water and hydroxylamine to pyruvic acid and pyruvate is the same.⁷ Methyl pyruvate is expected to be a good model for pyruvic acid.

EXPERIMENTAL

Materials

Methyl pyruvate and hydroxylamine were obtained commercially. Solutions of these reagents were prepared just prior to use to minimize the possibility of decomposition. Buffer solutions of dilute hydrochloric acid, chloroacetic acid, acetic acid and hydroxylamine were employed in the appropriate ranges of pH. Glass-distilled water was used throughout. Values of pH were measured with an Orion pH-meter model 420-A.

*Correspondence to: A. Malpica, Mr Anais Ron Calzadilla, 8824 NW Street Suite CCS 00123, Miami, FL 33166, USA.
Contract/grant sponsor: Consejo de Desarrollo Científico y Humanístico de la Universidad Central de Venezuela.




Figure 2. Logarithms of second-order rate constants for oxime formation from methyl pyruvate plotted as a function of pH. Measurements were made at 30 °C and ionic strength 0.5. The solid line is a theoretical line based on Eqn. (1) and values in Table 1

This behavior is interpreted in terms of the mechanism outlined in Scheme 1, in which it is proposed that carbinolamine dehydration is the sole rate-determining step over the entire range of pH investigated.

The rate law for the mechanism outlined in Scheme 1 is:

$$k_{\text{obs}}/(\text{amine})_{\text{fb}} = K_{\text{add}}^{\text{exp}}[k_{\text{H}}(\text{H}^+) + k_{\text{o}}] \quad (1)$$

At low pH Eqn. (1) becomes:

$$k_{\text{obs}}/(\text{amine})_{\text{fb}} = K_{\text{add}}^{\text{exp}}k_{\text{H}}(\text{H}^+) \quad (2)$$

The limiting value of Eqn. (1) at high pH is:

$$k_{\text{obs}}/(\text{amine})_{\text{fb}} = K_{\text{add}}^{\text{exp}}k_{\text{o}} \quad (3)$$

Scheme 1 includes the conversion of the hydrate of methyl pyruvate to the reactive unhydrated form. Therefore, the equilibrium constant for addition of hydroxylamine to the keto ester is corrected for the hydration of the substrate:

$$K_{\text{add}} = K_{\text{add}}^{\text{exp}}(1 + K_{\text{H}}) \quad K_{\text{H}} = 2.8 \quad (\text{see Ref. 11})$$

The equilibrium constant for addition of hydroxylamine to methyl pyruvate was monitored by the disappearance of the chromophore of the unhydrated keto ester at pH 6.75. Injecting a small amount of a solution of the substrate in dioxane into aqueous solutions of hydroxylamine led to a rapid drop in absorbance followed by a slower increase in absorbance attributed to oxime formation (see Experimental section). According to Pocker and coworkers,^{11,12} the hydration of methyl pyruvate is catalyzed by water, hydronium ions and

[†] Pocker *et al.* report several values of k_{f} over a temperature range of 0–25.6 °C, pH 4.6 and $\mu = 0.1$. A simple Arrhenius plot from their data allowed us to obtain $k_{\text{f}} = 0.15 \text{ s}^{-1}$ at 30 °C.

Table 1. Summary of equilibrium and rate constants of methyl pyruvate oxime formation measured in aqueous solution at 30 °C, and ionic strength 0.5

$K_{\text{add}}^{\text{exp}} (\text{M}^{-1})$	713.00 ^a
$k_{\text{H}} (\text{M}^{-1} \text{ s}^{-1})$	200.00 ^b
$k_{\text{o}} (\text{s}^{-1})$	2.3×10^{-4b}
$K_{\text{add}} (\text{M}^{-1})$	2700.00 ^a

^a Standard deviation of $K_{\text{add}}^{\text{exp}}$ was about 5%.

^b In most cases the data agree within 3%.

general bases [$k_{\text{f}} = k_{\text{o}} + k_{\text{H}}(\text{H}^+) + k_{\text{B}}(\text{B}^+)$].[†] Both catalytic constants are significant and their contribution to k_{f} depends on the pH and buffer concentration. Therefore, hydration is important over the entire pH range investigated. Thus the hydrate is included in the formulation of the reaction mechanism and the addition constant $K_{\text{add}}^{\text{exp}}$ is corrected for hydration.

In Fig. 2, logarithms of second-order rate constants [$k_{\text{obs}}/(\text{amine})_{\text{fb}}$] for methyl pyruvate oxime formation are plotted as a function of pH. The second-order rate constants decrease linearly with the concentration of the hydrated proton down to a value of pH ~4.0. Above this limiting pH value the logarithm of the second-order rate constants deviates from linearity with pH. These deviations are more pronounced as the pH increases.

Table 1 is a summary of the rate and equilibrium constants for methyl pyruvate oxime formation. The solid line in Fig. 2 is a theoretical line based in Eqn. (1) and values in Table 1. The agreement of theory with experimental data is satisfying.

Values of K_{add} and k_{H} reported for pyruvic acid oxime formation are respectively $3.5 \times 10^3 \text{ M}^{-1}$ and $2.43 \times 10^2 \text{ M}^{-1} \text{ s}^{-1}$.⁷ A comparison with the same constants reported here for methyl pyruvate oxime formation, $K_{\text{add}} = 2.7 \times 10^3 \text{ M}^{-1}$ and $k_{\text{H}} = 2 \times 10^2 \text{ M}^{-1} \text{ s}^{-1}$, demonstrate that methyl pyruvate is a good model for pyruvic acid.

To our knowledge, there are no studies that relate the values of pH in which catalysis by water is important to the carbinolamine structure. Generally, the pH-independent route of carbinolamine dehydration is usually unimportant, compared with the specific acid-catalyzed pathway, and it is ordinarily observed at values of pH close to 8. Carbinolamine formed from methyl pyruvate and hydroxylamine has a carbomethoxy group bonded directly to the carbonyl moiety. The strong electron-withdrawing inductive effect of this group decreases the carbinolamine oxygen basicity, making hydronium ion catalysis difficult. This is reflected in the low value of k_{H} reported here ($k_{\text{H}} = 2 \times 10^2 \text{ M}^{-1} \text{ s}^{-1}$) compared with the same process using pyruvate anion as substrate ($k_{\text{H}} = 5.75 \times 10^4 \text{ M}^{-1} \text{ s}^{-1}$).⁷ Therefore, the presence of the carbomethoxy group increases the relative importance of spontaneous dehydration; the percent of contribution of k_{o} from pH 5.0 to 6.75 in this study varies from 10 to 87%. Further investigations of imine

formation from activated substrates at pH >5 are desirable in order to test the importance of the carbinolamine structure for spontaneous catalysis to become apparent at early pH values.

Acknowledgements

We are indebted to Professor Eugene H. Cordes for helpful comments concerning aspects of this work. This work was supported in part by the Consejo de Desarrollo Científico y Humanístico de la Universidad Central de Venezuela.

REFERENCES

1. Rosenberg S, Silver SM, Sayer SJ, Washtien W. *J. Am. Chem. Soc.* 1974; **96**: 7988.
2. Sayer JM, Pinsky B, Schonbrum A, Jencks WP. *J. Am. Chem. Soc.* 1974; **96**: 7998.
3. Calzadilla M, Malpica A, Cordova T. *J. Phys. Org. Chem.* 1999; **12**: 708.
4. Sojo P, Viloria F, Malave L, Possamai R, Calzadilla M, Baumrucker J, Malpica A, Moscovici R, do Amaral L. *J. Am. Chem. Soc.* 1976; **98**: 4519.
5. Malpica A, Calzadilla M, Baumrucker J, Jimenez J, Lopez L, Escobar G, Montes C. *J. Org. Chem.* 1994; **59**: 3398.
6. Calzadilla M, Malpica A, Diaz P. *Int. J. Chem. Kinet.* 1996; **28**: 687.
7. Malpica A, Calzadilla M, Cordova T. *J. Phys. Org. Chem.* 2000; **13**: 162.
8. Malpica A, Calzadilla M, Cordova T, Torres S, Saulny CH. *Int. J. Chem. Kinet.* 1999; **31**: 387.
9. Jencks WP. *Proc. Phys. Org. Chem.* 1994; **2**: 63.
10. Jencks WP. *Catalysis in Chemistry and Enzymology*. McGraw-Hill: New York, 1969; Chapter 10, 463, and references cited therein.
11. Pocker Y, Meany JE, Zadoroj C. *J. Phys. Chem.* 1971; **75**: 792.
12. Pocker Y, Meany JE, Jones RC. *J. Am. Chem. Soc.* 1982; **104**: 4885.