

Synthesis and conformational analysis of chiral ureas incorporating *N*-1-phenylethyl groups. Manifestation of allylic 1,3-strain[†]

Marcos Hernández-Rodríguez, Roberto Melgar-Fernández and Eusebio Juaristi*

Departamento de Química, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Apdo. Postal 14-740, 07000 México, D. F., México

Received 1 February 2005; revised 20 February 2005; accepted 25 February 2005

ABSTRACT: The synthesis of novel chiral ureas (*R,R*)-**2**, (*S,S*)-**3** and (*R*)-**6** incorporating the α -phenylethyl group is described. Conformational analysis of these ureas, and of previously reported (*R,R*)-**1**, was carried out computationally, both at semiempirical (AM1 and PM3) and *ab initio* (HF and B3LYP) levels, and experimentally from x-ray crystallographic analysis of (*R,R*)-**2** and (*S,S*)-**3**, and in the case of (*R*)-**6** by means of NOE NMR spectroscopy. A substantial preference of 1.5–2.6 kcal mol^{−1} in favor of conformations with *syn*-periplanar arrangements between the C—H bond at the α -phenylethyl *N*-substituent and the N—C(O) segment was found, and this observation confirms the relevance of allylic A^{1,3} strain in this system. The possibility of hydrogen bonding in the *syn*-periplanar C—H...O=C—N arrangement was discarded in the light of topological analysis of (*R,R*)-**1**, within the frame of Bader's atoms in molecules theory. Copyright © 2005 John Wiley & Sons, Ltd.

KEYWORDS: chiral ureas; conformational analysis; allylic 1,3-strain; theoretical chemistry; density functional theory; *ab initio* calculations

INTRODUCTION

Urea and its derivatives constitute an important class of organic compounds with a great variety of applications in fundamental and applied science. They find extensive application as antioxidants in gasoline,¹ dyes for cellulose fibers,¹ non-linear optical devices,² resin precursors³ and synthetic intermediates.⁴ Moreover, the presence of the urea moiety in many biologically important natural compounds, such as nucleotides, vitamin B₁₃ and enzymes,⁵ renders it a subject of great interest in biochemistry and related areas. Furthermore, urea-based drugs have been developed that are effective antitumor agents⁶ and HIV protease inhibitors.⁷ Finally, the anion binding properties of urea, owing to its ability to form two hydrogen bonds,⁸ have been exploited in several relevant areas such as anion recognition and sensing,⁹ enantioselective asymmetric synthesis¹⁰ and kinetic resolution of chiral compounds.¹¹

In this context, the synthesis of new chiral ureas (*R,R*)-**1** and (*S,S*)-**1**, incorporating the 1-phenylethyl group at the *N*-atoms was recently reported¹² [Scheme 1(a)]. The

potential of these chiral Lewis bases as promoters of stereoselective reactions will depend on the possibility that the *N*-1-phenylethyl chiral adjuvant (for recent reviews on applications of 1-phenylethylamine in the preparation of enantiomerically pure compounds, see Ref. 13) adopts a single or predominant conformation in the molecule. In particular, conformations presenting coplanar orientation between the C—H bond and the N—C(O) segment should be preferred in order to prevent allylic A^{1,3} strain,¹⁴ which would be present in the other possible conformations around the N—CHMePh bond. Indeed, N→C=O conjugation in the urea segment effectively places the phenethyl group in an allylic-like position, so that the C—H bond, being the smallest substituent at the stereogenic carbon, should adopt a *syn*-periplanar arrangement in relation to the N—C(O) segment [Scheme 1(b)].

This paper concerns the preparation and conformational analysis of several chiral ureas structurally related to **1**.

RESULTS AND DISCUSSION


Preparation of chiral ureas 2–5

Acyclic chiral urea (*R,R*)-**2** was obtained by reaction of (*R*)-1-phenylethylamine with triphosgene in the presence

*Correspondence to: E. Juaristi, Departamento de Química, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Apdo. Postal 14-740, 07000 México, D. F., México.
E-mail: ejuaristi@cinvestav.mx

Contract/grant sponsor: Conacyt; Contract grant number: 45157.

[†]Selected paper presented for a special issue dedicated to Professor Otto Exner on the occasion of his 80th birthday.




Scheme 1

of triethylamine and methylene chloride as solvent, according to the procedure described in the literature¹⁵ [Scheme 2(a)]. Similarly, *N,N'*-bis[(*S*)-1-phenylethyl]ethane-1,2-diamine¹⁶ was the starting material for the preparation of five-membered urea (*S,S*)-**3** [Scheme 2(b)]. Non- C_2 -symmetric chiral urea (*R*)-**6** was synthesized from acrylonitrile via conjugate addition of (*R*)-1-phenylethylamine followed by catalytic hydrogenation and condensation with triphosgene [Scheme 2(c)].

syn-periplanar to the N—C(O) segment.¹² This solid-state conformation is also predicted to be the most stable in the gas phase, at various levels of theory (Table 1).

All *ab initio* methods predict conformation **A** to be significantly more stable than conformation **B**, by 2.2–2.6 kcal mol^{−1} (1 kcal = 4.184 kJ). For comparison, Broeker *et al.*¹⁷ calculated (4*S*)-phenyl-(2*Z*)-pentene to prefer the a *syn*-periplanar (C=C/C—H) arrangement **C** by 1.5 kcal mol^{−1} over arrangement **D** [Eqn (1)]:




whereas Tietze and Schulz¹⁸ found from *ab initio* calculations on 3-methyl-1-butene rotamer **E** (*syn*-periplanar


Conformational analysis

N,N'-Bis[(*R*)-1-phenylethyl]propyleneurea, (*R,R*)-**1**

As expected from allylic A^{1,3} strain, x-ray analysis of single crystals of (*R,R*)-**1** showed an orientation of the 1-phenylethyl groups in which the C—H bonds are nearly




Scheme 2


Table 1. Calculated (gas-phase) conformational preference of the *N*-(1-phenylethyl) *N*-substituents in (*R,R*)-**1**


Method	ΔE (kcal mol ⁻¹)	Dihedral [H—C*—N—C(O)] angles (°)	
		A	B
AM1	1.69	28.9 and 26.5	28.6 and -163.7
PM3	1.27	27.1 and 32.4	34.0 and -176.7
HF/3-21G	2.19	14.7 and 14.9	5.9 and -165.1
HF/6-31G**	2.54	2.9 and -14.5	-3.6 and 170.4
B3LYP/6-31G*	2.35	3.0 and 7.9	2.8 and -169.4
B3LYP/6-31G**	2.33	3.0 and 7.6	2.8 and -170.1
B3LYP/6-311+G**//B3LYP/6-31G*	2.59	3.0 and 7.9	2.8 and -169.4

orientation between the C=C double bond and the methine C—H bond) to be 2.48 kcal mol⁻¹ more stable than rotamer **F** (*anti*-periplanar orientation) [Eqn (2)]:



The experimentally obtained [x-ray diffraction analysis of crystalline (*R,R*)-**1**; see Ref. 12] interatomic distance between the carbonyl oxygen and the methine C—H hydrogen on the 1-phenylethyl group is 2.27 Å, which is less than the sum of the atomic van der Waals radii for oxygen and hydrogen, 1.50 and 1.20 Å, respectively.¹⁹ Hence the possibility of the existence of a hydrogen bonding interaction that could contribute to the stabilization of conformer **A** in the equilibrium depicted in Table 1 was considered. Nevertheless, it must be pointed out that two recent, high-level theoretical studies²⁰ of the *N*-methyl rotational barriers in amide and urea derivatives concluded that staggered conformers **G** are 0.3–0.8 kcal mol⁻¹ more stable than conformers **H** [Eqn (3)], in which one could expect that hydrogen bonding stabilization would be more efficient.



To obtain information regarding the nature of the C—H···O=C interaction in **A** (Table 1), theoretical analysis within the frame of the topological theory of atoms in molecules²¹ was carried out. The AIM2000²² set of programs was used, obtaining the properties of atoms and critical points (cps) in the charge density: electron density (ρ), Laplacians ($\nabla^2\rho$) and ellipticities (ε).

Two of the most important contributions of the AIM theory are the precise definition of an atom in a molecule and the definition of the chemical bond. These concepts correspond to the topological properties of the electron density.²¹ The chemical structure of a molecule is unambiguously described determining the critical points through the electron density (ρ) and corresponds to the gradient zero density points. These points, and also the first and second derivatives, can be determined with the use of the AIM2000²² program. From the second derivative, it is possible to determine the three principal curvatures associated with a critical point due to its index (the algebraic addition of the curvature sign). In addition to density, ellipticity is an important property of a critical point. This is defined as the coefficient of the negative curves along the perpendicular axis to the bonding path, $\varepsilon = \lambda(1)/\lambda(2) - 1$, and should be considered as an index of a bond's anisotropy.

Figure 1 shows the calculated C—H···O=C bond trajectories and critical points between atoms in the electron density analysis of (*R,R*)-**1**. Most relevant, the C—H···O=C bond trajectories present low electronic


**Figure 1.** Critical points and bonding trajectories in the electron density of (*R,R*)-**1** calculated at the HF/6-311+G**/HF/6-31G** level of theory (for clarity, Ph groups have been deleted)

Table 2. Properties of critical points associated with weak interactions in (*R,R*)-**1** (see Fig. 1)

Point	ρ ($\text{e } \text{\AA}^{-3}$)	ϵ	x^a	y^a	z^a
a^b	0.022	1.865	−0.0230	−0.0080	0.1326
b^b	0.0213	3.752	−0.0221	−0.0046	0.1301
c^c	0.0215		−0.022	0.0097	0.1272
d^c	0.0213		−0.0215	0.0052	0.1275
e^c	0.0197		−0.0148	0.0769	0.0825


^a Eigenvalues of the Hessian at critical point.^b Critical bond point (3, −1).^c Critical ring point (3, +1).

density at the critical points *a* and *b* ($\rho = 0.022$ and $0.021 \text{ e } \text{\AA}^{-3}$, respectively) which is indicative of a very weak bonding interaction. In contrast, bonding interactions are usually associated with much larger electronic densities; for example, a typical $\text{A} \cdots \text{H} \cdots \text{B}$ hydrogen bond presents $\rho \geq 0.36 \text{ e } \text{\AA}^{-3}$ (Ref. 21) (Fig. 1 and Table 2). Furthermore, the large ellipticity parameters for critical points *a* and *b* in (*R,R*)-**1** (1.865 and 3.752,

respectively) (Table 2) are indicative of a minor interaction.²³

***N,N'*-Bis[(*R*)-1-phenylethyl]urea, (*R,R*)-2.** Relative to cyclic urea (*R,R*)-**1** (see above), the conformational behavior of the acyclic urea (*R*)-**2** is much more complex since rotation around the amidic segments leads to various possible arrangements of the 1-phenylethyl group in combination with *E* and *Z* conformations (see, for example, Ref. 24). An *ab initio* DFT (B3LYP/6–31G* level) evaluation of all rotamers originating from rotation around the $\text{N}—\text{C}(\text{O})$ and $\text{N}—\text{C}(\text{H})$ bonds afforded the seven energy minima ($E < 5.0 \text{ kcal mol}^{-1}$), which are given in Table 3. This table does not include conformer *EEa*, whose energy was estimated to be $> 12.0 \text{ kcal mol}^{-1}$.

It can be appreciated from Table 3 that the two conformers of lowest energy, i.e. *EZa* and *ZZa* ($E_{\text{rel}} = 0.0$ and $0.2 \text{ kcal mol}^{-1}$, respectively), fulfil expectations in terms of the concept of allylic $\text{A}^{1,3}$ strain. Indeed, the $\text{C}—\text{H}$ bonds in these conformers are oriented in a

Table 3. Relative energies for the conformers of lowest energy of (*R,R*)-**2**, according to *ab initio* DFT B3LYP/6–31G* calculations

Conformer	Total energy (hartree)	$E_{\text{rel.}}$ (kcal mol^{-1})	Dihedral $\text{H}—\text{C}^*—\text{N}—\text{C}(\text{O})$ angles ($^\circ$)
<i>ZZa</i>	−844.272822	0.22	−24.6 and −24.6
<i>ZZb</i>	−844.268623	2.86	−13.4 and −167.5
<i>ZZc</i>	−844.266308	4.31	166.7 and −166.7
<i>EZa</i>	−844.273178	0.00	−26.4 and −38.4
<i>EZb</i>	−844.268699	2.81	39.8 and 173.4
<i>EZc</i>	−844.26946	2.33	164.0 and −39.5
<i>EZd</i>	−844.265698	4.69	167.7 and 171.6





Figure 2. X-ray crystallographic structure and solid-state conformation of *N,N'*-bis[(*R*)-1-phenylethyl]urea, (*R,R*)-**2**

syn-periplanar manner to the N—C(O) segment. In the rest of the isomers, allylic $A^{1,3}$ strain induced by the methyl and phenyl substituents results in higher energy content (lower stability).

Figure 2 presents the x-ray crystallographic structure and the solid-state conformation of (*R,R*)-**2**. The structure was solved and refined using SHELX-97,^{25a} within WinGX program version 1.64.05.28.^{25b} Crystal data: $C_{17}H_{20}N_2O_1$, molecular weight = 268.35, orthorhombic $P2_12_12_1$, $a = 4.6697(2)$ Å, $b = 9.8125(4)$ Å, $c = 16.3979(7)$ Å, $\alpha = 90.0^\circ$, $\beta = 90.0^\circ$, $\gamma = 90.0^\circ$, $V = 751.38(5)$ Å³, crystal size $0.17 \times 0.30 \times 0.48$ mm³, $R1 = 0.0401$ ($wR2 = 0.0982$). [Atomic coordinates for the structures reported in this paper have been deposited at the Cambridge Crystallographic Data Centre. The coordinates can be obtained, on request from the Director, Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK (Fax +44 1223 336036; E-mail: deposit@ccdc.cam.ac.uk; deposition number CCDC 263119.) The C—H bonds at the *N*-1-phenylethyl groups orient themselves in a *syn*-periplanar manner relative to the N=C=O segment. Therefore, as a consequence of allylic $A^{1,3}$ strain, the molecule adopts a quasi- C_2 arrangement in the crystal.


N,N'-Bis[(*S*)-1-phenylethyl]propyleneurea, (*S,S*)-**3**.

This chiral urea was modeled at the B3LYP/6-31G* level. The optimized structure (Table 4) presents a half-chair ring conformation with the carbonyl group oriented along the C_2 symmetry axis. The calculated energy difference between conformers **I** and **J** [Eqn (4)] is $1.27 \text{ kcal mol}^{-1}$, the former being more stable.



Again, in conformer **I** [*syn*-periplanar arrangement between the C—H bond and the N—C(O) segment] allylic $A^{1,3}$ strain is minimized. Nevertheless, it can be appreciated that $\Delta E(\mathbf{I} \rightleftharpoons \mathbf{J}) = 1.27 \text{ kcal mol}^{-1}$ in (*S,S*)-**3** is significantly lower than $\Delta E(\mathbf{A} \rightleftharpoons \mathbf{B}) = 2.35 \text{ kcal mol}^{-1}$ calculated for six-membered (*R,R*)-**1** at the same level

Table 4. Optimized geometry of lowest energy in (*R,R*)-**1**, (*R,R*)-**2**, (*S,S*)-**3** and (*R*)-**6** at the B3LYP/6-31G* level (distances in Å, angles in degrees; the phenyl ring is not included).



(*R,R*)-**1**, $R = \text{Ph(Me)CH}$, $n = 3$
 (*R,R*)-**2**, $R = \text{Ph(Me)CH}$, $n = 0$
 (*S,S*)-**3**, $R = \text{Ph(Me)CH}$, $n = 2$
 (*R*)-**6**, $R = \text{H}$, $n = 3$

	(<i>R,R</i>)- 1	(<i>R,R</i>)- 2	(<i>S,S</i>)- 3	(<i>R</i>)- 6
N ₁ —C ₂	1.385	1.378	1.387	1.387
C ₂ —N ₃	1.385	1.391	1.387	1.381
N ₁ —C ₄	1.478	1.472	1.470	1.478
C ₄ —H ₅	1.091	1.092	1.094	1.092
C ₂ —O ₆	1.238	1.230	1.228	1.232
N ₁ —C ₂ —N ₃	117.3	115.3	108.0	116.5
N ₁ —C ₄ —H ₅	104.0	104.4	103.5	103.6
N ₁ —C ₂ —O ₆	121.3	123.8	126.0	123.2
N ₃ —C ₂ —O ₆	121.4	120.9	126.0	120.3
H ₅ —C ₄ —N ₁ —C ₂	7.6	−26.4	−26.6	−12.7
C ₄ —N ₁ —C ₂ —O ₆	1.3	8.7	1.3	5.6
C ₄ —N ₁ —C ₂ —N ₃	−179.6	−168.9	162.2	−177.3

of theory. This difference in conformational bias can easily be accounted for in terms of the structural differences in the five- and six-membered rings. In particular, the interatomic distance between the carbonyl oxygen and the C—H hydrogen of interest are estimated as 2.15 Å in (*R,R*)-**1** and 2.39 Å in (*S,S*)-**3**. Hence, manifestation of the allylic $A^{1,3}$ strain effect is more evident in the former system.

Recrystallization of (*S,S*)-**3** afforded single crystals suitable for x-ray analysis (Fig. 3). The structure was solved and refined using SHELX-97,^{25a} within WinGX program version 1.64.05.28.^{25b} Crystal data: $C_{19}H_{22}N_2O_1$, molecular weight = 294.39, orthorhombic $P2_12_12_1$, $a = 8.6014(3)$ Å, $b = 12.1590(4)$ Å, $c = 15.8840(6)$ Å, $\alpha = 90.0^\circ$, $\beta = 90.0^\circ$, $\gamma = 90.0^\circ$, $V = 1661.22(10)$ Å³, crystal size $0.30 \times 0.37 \times 0.40$ mm³, $R1 = 0.0394$ ($wR2 = 0.0846$) (deposition number CCDC 263120). Most interesting is the propeller-like orientation




Figure 3. X-ray crystallographic structure and solid-state conformation of *N,N'*-bis[(*S*)-1-phenylethyl]ethyleneurea, (*S,S*)-**3**





Figure 4. Rotation around the N—C* bond in (*R*)-**6**, according to HF/3–21G* calculations (fully optimized geometries)

of the 1-phenylethyl groups, which are expected to lead to high enantioselectivities in reactions taking place with a suitable substrate coordinated to the C=O oxygen atom. Again, the solid-state structure of (*S,S*)-**3** fulfils expectation based on the concept of allylic A^{1,3} strain.

***N*–[(*R*)-1-Phenylethyl]propyleneurea, (*R*)-**6**.** The potential energy surface (PES) presented in Fig. 4 was obtained by constraining the H—C—N—C(O) dihedral angle for rotation about the C—N bond and fully optimizing (B3LYP/6–31G* level) the remaining internal coordinates. Intervals of 10° were used.

As seen in Fig. 2, the two lowest energy rotamers present H—C—N—C(O) dihedral angles of 0° and 195°, the former being 2.04 kcal mol^{−1} more stable. Once again, this observation is in agreement with expectation in terms of minimization of allylic A^{1,3} strain.

Support for this conclusion was acquired experimentally by means of ¹H NMR spectroscopy, and in particular from observation of a significant nuclear Overhauser effect (NOE) on one of the diastereotopic²⁶ hydrogens adjacent to the *N*-1-phenylethyl group. As illustrated in Eqn (5), a 2.4% enhancement at H_A upon irradiation of the methyl group is congruent with a predominance of conformer **K** and a dihedral angle $\tau = 0^\circ$.



Summary

Theoretical analysis by means of semiempirical (AM1 and PM3) and *ab initio* (HF and DFT) methods confirm the predominance of those conformations that minimize allylic A^{1,3} strain in *N*-(1-phenylethyl)-containing ureas **1–3** and **6**. This conformational bias suggests that chiral ureas such as **1–3** and **6** or their derivatives should be effective promoters of stereoselective reactions,²⁷ owing to the substantial steric and electronic differences presented by the hydrogen, methyl and phenyl groups in the 1-phenylethyl chiral auxiliary.

EXPERIMENTAL

General methods

TLC: silica gel F₂₅₄ plates; detection with UV radiation or iodine vapor. Flash column chromatography:²⁸ silica gel (230–400 mesh). Melting-points: not corrected. ¹H NMR spectra: 60, 270 and 400 MHz spectrometers. ¹³C NMR spectra: 22.5, 67.8, and 100 MHz spectrometers. Chemical shifts (δ) in ppm downfield from internal TMS reference; the coupling constants (*J*) are given in Hz. Optical rotations were measured in a polarimeter, using the sodium D-line (589 nm). Mass spectra: 20 eV.

Computational methods

Geometry optimizations (with no symmetry constraints) of all conformers were performed using semiempirical

(PM3 and AM1) and *ab initio* (HF and B3LYP) methods, using the Gaussian 98, Revision A.7,²⁹ series of programs. Thermodynamic corrections of energy in *ab initio* calculations were carried out by zero-point energy (ZPE) correction. The consideration of electronic correlation is very important in conformational studies;³⁰ thus, hybrid functionals B3LYP at the 6–31G(d,p) or 6–31G(d) level and also *ab initio* level were examined.

Topological analysis of (*R,R*)-**1** was possible by means of AIM2000²² (atoms in molecules package), a set of programs that defines the properties of atoms and bonding patterns [charge density (ρ), Laplacians ($\nabla^2\rho$) and ellipticities (ϵ) of critical points between atoms] in a molecule.

Syntheses

N,N'-Bis[(*R*)-1-phenylethyl]propyleneurea, (*R,R*)-**1**. This compound was prepared according to the procedure developed in our laboratory.¹²

N,N'-Bis[(*R*)-1-phenylethyl]urea, (*R,R*)-**2**. The procedure described in the literature¹⁵ was followed. The crude product was purified by silica gel flash chromatography²⁸ (CH_2Cl_2 –AcOEt, 8:2) to give a white solid (79% yield), m.p. 209–210 °C (lit.³¹ m.p. 207–209 °C). $[\alpha]_{\text{D}}^{25^\circ\text{C}} = +55.6$ ($c = 0.85$, EtOH) [lit.³¹ $[\alpha]_{\text{D}}^{25^\circ\text{C}} = +51.7$ (EtOH, concentration not given)]. ¹H NMR (400 MHz, DMSO- d_6): δ 1.26 (d, $J = 7.0$ Hz, 6H), 4.69 (dq, $J^1 = 8.0$ Hz, $J^2 = 7.0$ Hz, 2H), 6.25 (d, $J = 8.0$ Hz, 2H), 7.16–7.32 (m, 10H). ¹³C NMR (100 MHz, DMSO- d_6): δ 23.9, 40.6, 126.2, 126.9, 128.7, 146.2, 157.0.

N,N'-Bis[(*S*)-1-phenylethyl]ethyleneurea, (*S,S*)-**3**. In a 250 ml round-bottomed flask provided with a magnetic stirrer was placed 5.37 g (20.0 mmol) of *N,N'*-bis[(*S*)-1-phenylethyl]ethane-1,2-diamine¹⁶ dissolved in 100 ml of dichloromethane. The solution was cooled to 0 °C before the dropwise addition of 2.11 g (7.1 mmol) of triphosgene dissolved in 50 ml of dichloromethane. The ice–water bath was removed and stirring was continued for 24 h. Isolation of the product involved addition of 50 ml of 1.0 M HCl. The aqueous phase was separated and washed with 50 ml of dichloromethane and the combined organic layers were dried and concentrated in a rotary evaporator. The crude product was purified by flash chromatography²⁸ (hexane–EtOAc, 9:1) to give 2.36 g (40% yield) of (*S,S*)-**3** as white crystals, m.p. 73–74 °C. $[\alpha]_{\text{D}}^{25^\circ\text{C}} = -84.4$ ($c = 2.2$, CHCl_3). ¹H NMR (300 MHz, CDCl_3): δ 1.49 (d, $J = 7.1$ Hz, 6H), 2.86 (m, 2H), 3.17 (m, 2H), 5.32 (q, $J = 7.1$ Hz, 2H), 7.24–7.35 (m, 10H). ¹³C NMR (75 MHz, CDCl_3): δ 16.3, 37.7, 50.3, 127.2, 127.3, 128.4, 140.9, 160.1. MS: m/z 294 (M^+ , 53), 280 (14), 189 (42), 175 (45), 105 (100). Anal. calcd for $\text{C}_{19}\text{H}_{22}\text{N}_2\text{O}$: C, 77.51; H, 7.53. Found: C, 77.59; H, 7.75%.

3-[(*R*)-1-Phenylethylamino]propionitrile, (*R*)-**4**. In a 125 ml round-bottomed flask provided with a magnetic stirrer was placed 11.0 g (90.9 mmol) of (*R*)-1-phenylethylamine dissolved in 60 ml of methanol. To this solution was added 6.95 ml (100 mmol) of acrylonitrile and the resulting mixture was heated at reflux for 18 h. Concentration in a rotary evaporator afforded the crude product that was distilled in a Kugelrohr at 110–113 °C/0.1 mmHg to give 14.9 g (95% yield) of (*R*)-**4**, $[\alpha]_{\text{D}}^{25^\circ\text{C}} = +69.9$ ($c = 1.55$, CHCl_3). ¹H NMR (CDCl_3 , 400 MHz): δ 1.37 (d, $J = 6.6$ Hz, 3H), 1.55 (bs, 1H), 2.31–2.49 (m, 2H), 2.69–2.84 (m, 2H), 3.81 (q, $J = 6.6$, 1H), 7.24–7.37 (m, 5H). ¹³C NMR (CDCl_3 , 100 MHz): δ 18.9, 24.5, 42.9, 57.8, 118.8, 126.5, 127.3, 128.6, 144.8. MS: m/z 174 (M^+ , 1), 160 (12), 159 (100), 105 (41), 91 (10).

N[(*R*)-1-Phenylethyl]propylene-1,3-diamine, (*R*)-**5**. In a hydrogenation flask were placed 3.33 g (19.1 mmol) of (*R*)-**4** and 0.33 g of 5% rhodium on alumina before the addition of 60 ml of methanol saturated with ammonia. The flask was pressurized to 500 psi of hydrogen and shaken at ambient temperature for 18 h. The reaction mixture was filtered over Celite, concentrated and the crude product was purified by distillation in a Kugelrohr, b.p. 100–105 °C/1.0 mmHg, to afford 3.06 g (90% yield) of (*R*)-**5** as a colorless liquid, $[\alpha]_{\text{D}}^{25^\circ\text{C}} = +53.0^\circ$ ($c = 1.86$, CHCl_3). ¹H NMR (CDCl_3 , 300 MHz): δ 1.34 (d, $J = 6.6$ Hz, 3H), 1.59 (~q, $J = 6.8$ Hz, 2H), 2.06 (br, 3H), 2.40–2.59 (m, 2H), 2.72 (~t, $J = 6.8$, 2H); 3.73 (q, $J = 6.6$, 1H), 7.21–7.32 (m, 5H). ¹³C NMR (CDCl_3 , 75 MHz): δ 24.5, 34.0, 40.6, 45.7, 58.5, 126.9, 127.2, 128.8, 145.8. MS: m/z 179 ($\text{M}^+ + 1$, 5), 120 (100), 105 (94), 91 (34), 73 (60). HRMS (FAB): calcd for $\text{C}_{11}\text{H}_{19}\text{N}_2$ ($\text{M}^+ + \text{H}$): 179.1548. Found: 179.1541.

N[(*R*)-1-Phenylethyl]propyleneurea, (*R*)-**6**. In a 150 ml round-bottomed flask provided with a magnetic stirrer were placed 2.3 g (12.8 mmol) of (*R*)-**5**, 3.6 ml (26.1 mmol) of triethylamine and 8.0 ml of dichloromethane. The resulting solution was cooled to 0 °C before the dropwise addition of 1.3 g (4.5 mmol) of triphosgene dissolved in 30 ml of dichloromethane. Stirring was continued for 30 min at 0 °C and for 2 days at ambient temperature. The reaction mixture was treated with 50 ml of 1.0 M HCl and the organic phase was separated, washed with brine solution, dried and concentrated to give the crude product, which was purified by flash chromatography²⁸ with EtOAc as eluent. (*R*)-**6** was obtained (2.05 g, 78% yield) as white crystals, m.p. 94–96 °C $[\alpha]_{\text{D}}^{25^\circ\text{C}} = +99.2^\circ$ ($c = 1$, CHCl_3). ¹H NMR (CDCl_3 , 400 MHz): δ 1.50 (d, $J = 7.0$ Hz, 3H), 1.78 (m, 2H), 2.79 (m, 1H), 3.08 (m, 1H), 3.27 (m, 2H), 5.28 (bs, 1H), 5.89 (q, $J = 7.0$ Hz, 1H), 7.21–7.36 (m, 5H). ¹³C NMR (CDCl_3 , 100 MHz): δ 15.8, 22.3, 39.3, 40.5, 50.6, 127.0, 127.4, 128.4, 141.5, 156.5. MS: m/z 205 ($\text{M}^+ + 1$, 17), 204 (M^+ , 100), 189 (87), 161 (24), 127 (16), 105

(21). Anal. Calcd for $C_{12}H_{16}N_2O$: C, 70.56; H, 7.90; N, 13.70. Found: C, 70.45; H, 8.11; N, 13.78%

Acknowledgements

We are grateful to Professor Maria de Jesús Rosales for useful discussions, Q. C. Gabriela Avila-Ortiz for technical assistance and Conacyt, México, for generous financial support via grant 45157.

REFERENCES

- Vishnyakova TP, Golubeva IA, Glebova EV. *Russ. Chem. Rev. (Engl. Transl.)* 1985; **54**: 249–261.
- (a) Halbout JM, Tang CL. In *Nonlinear Optical Properties of Organic Molecules and Crystals*, Chemla DS, Zyss J (eds). Academic Press: New York, 1987; 385–404; (b) Dixon DA, Matsuzawa N. *J. Phys. Chem.* 1994; **98**: 3967–3977; (c) Manzor B, Daly JH, Islam MS, Pethrick RA, Sherwood JN. *J. Chem. Soc., Faraday Trans.* 1997; **93**: 3799–3805.
- Meier RJ, Maple JR, Hwang M-J, Hagler AT. *J. Phys. Chem.* 1995; **99**: 5445–5456.
- (a) Nicolaou KC, Baran PS, Zhong Y-L, Sugita K. *J. Am. Chem. Soc.* 2002; **124**: 2212–2220; (b) Yin J, Buchwald SL. *J. Am. Chem. Soc.* 2002; **124**: 6043–6048; (c) Humphrey JM, Liao Y, Ali A, Rein T, Wong Y-L, Chen H-J, Courtney AK, Martin SF. *J. Am. Chem. Soc.* 2002; **124**: 8584–8592; (d) Saluzzo C, Lamouille T, Le Guyader F, Lemaire M. *Tetrahedron: Asymmetry* 2002; **13**: 1141–1146; (e) Yoshida H, Shirakawa E, Honda Y, Hiyama T. *Angew. Chem. Int. Ed.* 2002; **41**: 3247–3249; (f) Lee S-H, Clapham B, Koch G, Zimmermann J, Janda KD. *Org. Lett.* 2003; **5**: 511–514; (g) Muccioli GG, Wouters J, Poupaert JH, Norberg B, Poppitz W, Scriba GKE, Lambert DM. *Org. Lett.* 2003; **5**: 3599–3602; (h) Sergeev AG, Artamkina GA, Beletskaya IP. *Tetrahedron Lett.* 2003; **44**: 4719–4723; (i) Dunetz JR, Danheiser RL. *Org. Lett.* 2003; **5**: 4011–4014.
- Flippin-Anderson JL. *Acta Crystallogr., Sect. C* 1987; **43**: 2228–2230.
- Theophanides T, Harvey PD. *Coord. Chem. Rev.* 1987; **76**: 237–264.
- (a) Ala P, Huston EE, Klabe RM, McCabe DD, Duke JL, Rizzo CJ, Korant BD, DeLoskey RJ, Lam PYS, Hodge CN, Chang C-H. *Biochemistry* 1997; **36**: 1573–1580; (b) Hodge CN, Lam PYS, Eyermann CJ, Jadhav PK, Ru Y, Fernandez CH, De Lucca GV, Chang C-H, Kaltenbach RF III, Holler ER, Woerner F, Daneker WF, Emmett G, Calabrese JC, Aldrich PE. *J. Am. Chem. Soc.* 1998; **120**: 4570–4581.
- (a) Smith PJ, Reddington MV, Wilcox CS. *Tetrahedron Lett.* 1992; **33**: 6085–6088; (b) Fan E, Van Arman SA, Kincaid S, Hamilton AD. *J. Am. Chem. Soc.* 1993; **115**: 369–370.
- (a) Fitzmaurice RJ, Kyne GM, Douheret D, Kilburn JD. *J. Chem. Soc., Perkin Trans. 1* 2002; 841–864; (b) Boicchi M, Del Boca L, Gómez DE, Fabbrizzi L, Licchelli M, Monzani E. *J. Am. Chem. Soc.* 2004; **126**: 16507–16514.
- (a) Sigman MS, Jacobsen EN. *J. Am. Chem. Soc.* 1998; **120**: 4901–4902; (b) Wenzel AG, Jacobsen EN. *J. Am. Chem. Soc.* 2002; **124**: 12964–12965; (c) Okino T, Nakamura S, Furukawa T, Takemoto Y. *Org. Lett.* 2004; **6**: 625–627.
- Berkessel A, Cleemann F, Mukherjee S, Mueller TN, Lex J. *Angew. Chem. Int. Ed.* 2005; **44**: 807–811.
- Juaristi E, Hernández-Rodríguez M, López-Ruiz H, Aviña J, Muñoz-Muñoz O, Hayakawa M, Seebach D. *Helv. Chim. Acta* 2002; **85**: 1999–2008.
- (a) Juaristi E, Escalante J, León-Romo JL, Reyes A. *Tetrahedron: Asymmetry* 1998; **9**: 715–740; (b) Juaristi E, León-Romo JL, Reyes A, Escalante J. *Tetrahedron: Asymmetry* 1999; **10**: 2441–2495.
- (a) Johnson F. *Chem. Rev.* 1968; **68**: 375–413; (b) Hoffmann RW. *Chem. Rev.* 1989; **89**: 1841–1860; (c) Seebach D, Lamatsch B, Amstutz R, Beck AK, Dobler M, Egli M, Fitzi R, Gautschi M, Herradón B, Hidber PC, Irwin JJ, Locher R, Maestro M, Maetzke T, Mouriño A, Pfammatter E, Plattner DA, Schickli C, Schweizer WB, Seiler P, Stucky G, Petter W, Escalante J, Juaristi E, Quintana D, Miravittles C, Molins E. *Helv. Chim. Acta* 1992; **75**: 913–934.
- Grotjahn DB, Joubran C. *Tetrahedron: Asymmetry* 1995; **6**: 745–752.
- Hulst R, de Vries K, Feringa BL. *Tetrahedron: Asymmetry* 1994; **5**: 699–708.
- Broecker JL, Hoffmann RW, Houk KN. *J. Am. Chem. Soc.* 1991; **113**: 5006–5017.
- Tietze LF, Schulz G. *Liebigs Ann.* 1996; 1575–1579.
- Bondi A. *J. Phys. Chem.* 1964; **68**: 441–451.
- (a) Wiberg KB, Rush DJ. *J. Org. Chem.* 2002; **67**: 826–830; (b) Bryantsev VS, Firman TK, Hay BP. *J. Phys. Chem. A* 2005; **109**: 832–842.
- (a) Bader RFW. *Atoms in Molecules, a Quantum Theory*. Clarendon Press: Oxford, 1990; (b) Bader RFW. *Chem. Rev.* 1991; **91**: 893–928.
- Biegler-König F, Schönbohm J, Bayles D. *J. Comput. Chem.* 2001; **22**: 545–559.
- (a) Cioslowski J. *J. Am. Chem. Soc.* 1993; **115**: 5177–5179; (b) Madrid G, Rochín A, Juaristi E, Cuevas G. *J. Org. Chem.* 2001; **66**: 2925–2931.
- (a) Rang K, Sandström J, Svensson C. *Can. J. Chem.* 1998; **76**: 811–820; (b) Bharatam PV, Moudgil R, Kaur D. *J. Phys. Chem. A* 2003; **107**: 1627–1634.
- (a) Sheldrick GM. *Programs for Crystal Structure Analysis*, University of Göttingen: Göttingen, 1997; (b) Farrugia LJ. *J. Appl. Crystallogr.* 1999; **32**: 837–838.
- (a) Eliel EL, Wilen SH. *Stereochemistry of Organic Compounds*. Wiley: New York, 1994; chapt. 8, 465–538; (b) Juaristi E. *Introduction to Stereochemistry and Conformational Analysis*. Wiley: New York 1991; chapt. 6, 91–105.
- Bartlett PA. *Tetrahedron* 1980; **36**: 3–72.
- Still WC, Kahn M, Mitra A. *J. Org. Chem.* 1978; **43**: 2923–2925.
- Frisch MJ, Trucks GW, Schlegel HB, Scuseria GE, Robb MA, Cheeseman JR, Zakrzewski VG, Montgomery JA Jr, Stratmann RE, Burant JC, Dapprich S, Millam JM, Daniels AD, Kudin KN, Strain MC, Farkas O, Tomasi J, Barone V, Cossi M, Cammi R, Mennucci B, Pomelli C, Adamo C, Clifford S, Ochterski J, Petersson GA, Ayala PY, Cui Q, Morokuma K, Malick DK, Rabuck AD, Raghavachari K, Foresman JB, Cioslowski J, Ortiz JV, Baboul AG, Stefanov BB, Liu G, Liashenko A, Piskorz P, Komaromi I, Gomperts R, Martin RL, Fox DJ, Keith T, Al-Laham MA, Peng CY, Nanayakkara A, Gonzalez C, Challacombe M, Gill PMW, Johnson B, Chen W, Wong MW, Andres JL, Gonzalez C, Head-Gordon M, Replogle ES, Pople JA. *Gaussian 98, Revision A.7*. Gaussian: Pittsburgh, PA, 1998.
- Stephens PJ, Devlin FJ, Chabalowski CF, Frisch MJ. *J. Phys. Chem.* 1994; **98**: 11623–11627.
- Hiatt RR, Shaio M-J, Georges F. *J. Org. Chem.* 1979; **44**: 3265–3266.