IOPscience iopscience.iop.org

Home Search Collections Journals About Contact us My IOPscience

2D to 3D percolation crossover in the resistivity of co-evaporated Al-Ge mixture films

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1983 J. Phys. A: Math. Gen. 16 2889

(http://iopscience.iop.org/0305-4470/16/12/538)

View the table of contents for this issue, or go to the journal homepage for more

Download details:
IP Address: 129.252.86.83
The article was downloaded on 31/05/2010 at 06:28

Please note that terms and conditions apply.

J. Phys. A: Math. Gen. **16** (1983) 2889. Printed in Great Britain

Corrigenda

Langevin approach to the dynamics of interacting Brownian particles
Pusey P N and Tough R J A 1982 *J. Phys. A: Math. Gen.* **15** 1291–308

Equation (2.28) should read

$$\lim_{\tau\to 0^+}\frac{\mathrm{d}^3F(K,\tau)}{\mathrm{d}\tau^3}$$
$$= -D_0^3K^6 - \frac{3D_0^2K^4}{Nf}\sum_i\left\langle\frac{\partial^2U}{\partial r_{i1}^2}\right\rangle + \frac{2D_0^2K^3}{Nf}\sum_{i,j}\left\langle\sin(Kr_{ij1})\frac{\partial^3U}{\partial r_{i1}^2\,\partial r_{j1}}\right\rangle$$
$$-\frac{D_0K^2}{Nf^2}\sum_{i,j,k}\sum_\alpha\left\langle\exp(\mathrm{i}Kr_{ij1})\frac{\partial^2U}{\partial r_{i1}\,\partial r_{k\alpha}}\frac{\partial^2U}{\partial r_{k\alpha}\,\partial r_{j1}}\right\rangle.$$

Due to an oversight the third term was missing in the original paper. The correct version of this equation (as above) was obtained recently by J L Arauz-Lara and M Medina-Noyola (*Physica A* in press) and we are grateful to them for pointing out our error. Starting from the Smoluchowski equation, these authors have also obtained expressions for $\mathrm{d}F/\mathrm{d}\tau$, $\mathrm{d}^2F/\mathrm{d}\tau^2$ and $\mathrm{d}^3F/\mathrm{d}\tau^3$ for a suspension containing two types of particle.

2D to 3D percolation crossover in the resistivity of co-evaporated Al–Ge mixture films
Kapitulnik A and Deutscher G 1983 *J. Phys. A: Math. Gen.* **16** L243–8

In the figure caption to figure 1 the last sentence should be replaced by: 'The straight line is the best fit for the 2D regime.'

In figure 2 the vertical axis should be l(Å) rather than d(Å).

© 1983 The Institute of Physics