

Exact enumeration approach to fractal properties of the percolation backbone and $1/\sigma$ expansion

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1983 J. Phys. A: Math. Gen. 16 L475

(<http://iopscience.iop.org/0305-4470/16/13/007>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 129.252.86.83

The article was downloaded on 31/05/2010 at 06:28

Please note that [terms and conditions apply](#).

LETTER TO THE EDITOR

Exact enumeration approach to fractal properties of the percolation backbone and $1/\sigma$ expansion

Daniel C Hong and H Eugene Stanley

Center for Polymer Studies[†] and Department of Physics, Boston University, Boston, MA 02215, USA

Received 20 June 1983

Abstract. An exact enumeration approach is developed for the backbone fractal of the incipient infinite cluster at the percolation threshold. We use this approach to calculate exactly the first low-density expansion of $L_{\text{BB}}(p)$ for arbitrary system dimensionality d , where $L_{\text{BB}}(p)$ is the mean number of backbone bonds and p is the bond occupation probability. Standard series extrapolation methods provide estimates of the fractal dimension of the backbone for all d ; these disagree with the Sierpinski gasket model of the backbone. We also calculate the first low-density expansions of $L_{\text{min}}(p)$ and $L_{\text{red}}(p)$ which are, respectively, the mean number of bonds in the minimum path between i and j and the mean number of singly connected ('red') bonds.

How can one describe the flow of fluid in random porous media? This important question has long eluded explanation. Recently, considerable attention has been focused on the utility of fractals as models of random media. In particular, the topology of the network that exists just at the onset of fluid flow has been modelled by percolation theory. Bonds are considered intact if fluid can flow through them. When the fraction of bonds is small, the system consists of many small finite clusters. However, as the bond fraction approaches a critical value p_c the clusters grow large and ramified until at p_c fluid can flow. If we consider the network of intact bonds right at p_c , there will be a subset of bonds that carry fluid ('backbone' bonds) and a remainder that does not ('dangling ends'). The structure of the backbone remains an important open question. Two models of the backbone have been discussed in the literature. In one (Gefen *et al* 1981), the backbone is replaced by a d -dimensional Sierpinski gasket. In the other (Stanley 1977, Coniglio 1981, 1982, Pike and Stanley 1981, Stanley and Coniglio 1983), the backbone is considered to consist of an alternating sequence of singly connected ('red') bonds and multiply connected ('blue') bonds; these are shown in colour as figure 5 of Hamann (1983).

The advantage of the Sierpinski gasket model of the backbone is that one can calculate exactly its fractal dimension, $D_B = \ln(d+1)/\ln 2$. Hence it is important to obtain estimates of D_B for the actual backbone of percolation clusters. Thus far, the only efforts have been Monte Carlo simulations in $d = 2, 3$ for the backbone order parameter exponent β_B ; however, the order parameter is extremely difficult to calculate by Monte Carlo methods (Kirkpatrick 1978, Li and Streider 1982). Also, a limited attempt has been made to estimate the field-like scaling power y_h by large-cell

[†] Supported in part by grants from NSF, ARO and ONR.

position-space renormalisation group; however, this work was limited to $d = 2$. The Sierpinski gasket model gives reasonable quantitative values for $d = 2, 3$ but not for higher d . For this and other reasons, it is highly desirable to have calculations of d_f for the backbone for general d , in order that one can better assess the relative merits of various models of backbone topology. To this end, here we present an exact enumeration approach for the backbone fractal, and calculate the first ten terms in the low-density expansions for arbitrary d . We shall see that these ten terms behave sufficiently smoothly with increasing order that extrapolations to obtain the asymptotic behaviour can be made (table 1).

Table 1. Critical exponents characterising the backbone of the incipient infinite cluster in percolation. The basic quantities calculated are ζ_B and ζ_{\min} , since $\zeta_{\text{red}} = 1$ for all d (Coniglio 1982). In order to obtain the exponent for the derived quantities $L_{BB}(p)$ and $L_{\min}(p)$, we need γ_p , which is also tabulated. Finally, to obtain the backbone fractal dimension, D_B , we need ν .

	γ_p	ν	ζ_B	ζ_{\min}	$D_B = \frac{\rho_B}{\nu}$	$D_B = \frac{\ln(d+1)}{\ln 2}$
$d = 2$	$\frac{43^{(a)}}{18}$	$\frac{4^{(a)}}{3}$	$2.40 \pm 0.05^{(j)}$	$1.38 \pm 0.1^{(j)}$ $1.49 \pm 0.02^{(i)}$	$1.80 \pm 0.04^{(j)}$	$1.585^{(f)}$
$d = 3$	$1.73 \pm 0.03^{(b)}$ $1.66 \pm 0.07^{(c)}$ $1.66^{(d)}$	$0.88 \pm 0.02^{(b)}$	$1.61 \pm 0.07^{(j)}$	$1.18 \pm 0.07^{(j)}$	$1.83 \pm 0.08^{(j)}$	$2.000^{(f)}$
$d = 4$	$1.48 \pm 0.08^{(e)}$ $1.40^{(d)}$ $1.41 \pm 0.25^{(c)}$	$0.7^{(h)}$	$1.32^{+0.10}_{-0.15}$	$1.03 \pm 0.08^{(j)}$	$1.89^{+0.14}_{-0.21}$	$2.322^{(f)}$
$d = 5$	$1.16^{(d)}$ 1.18 ± 0.07 1.25 ± 0.15	$0.6^{(h)}$	$1.16 \pm 0.1^{(j)}$	$1.02 \pm 0.1^{(j)}$	$1.93 \pm 0.16^{(j)}$	$2.585^{(f)}$
$d = 6 - \epsilon$	$1 + \frac{1}{7}\epsilon + 0.046\epsilon^2$	$0.5 + \frac{5}{84}\epsilon$	$1 + \frac{1}{7}\epsilon^{(g)}$ $+ 0.025\epsilon^2$	—	$2 + 0.0476\epsilon$ $+ 0.0186\epsilon^2$	—
$d = 6$	$1.00^{(a)}$	$0.500^{(a)}$	$1.02 \pm 0.02^{(j)}$	$1.02 \pm 0.02^{(j)}$	$2.04 \pm 0.04^{(j)}$	$2.807^{(f)}$

^(a) Exact.

^(b) Gaunt and Sykes (1983).

^(c) Gaunt *et al* (1976).

^(d) Fisch and Harris (1978).

^(e) Gaunt *et al* (1976).

^(f) Gefen *et al* (1981).

^(g) Harris and Lubensky (1983).

^(h) Stauffer (1979).

⁽ⁱ⁾ Pike and Stanley (1981).

^(j) This work.

Since the 'thermal' scaling power $\gamma_T = 1/\nu$ is the same for the backbone as for the full cluster (Shlifer *et al* 1979), it is sufficient to calculate only one exponent in addition to ν . Since low-density expansions are generally more accurate than high-density expansions for the same effort, we focus our attention on the exponent ζ_B rather than on β_B .

In order to define clearly our approach, it is useful to review how one may calculate the exponent γ for the full cluster (Essam 1971, Dunn *et al* 1975). The pair connectedness P_{ij} is defined to be the probability that sites i and j are connected,

$$P_{ij} = [\nu_{ij}]_{\text{config}}, \quad (1a)$$

where $\nu_{ij} = 1$ if sites i and j are connected (belong to the same cluster) and $\nu_{ij} = 0$ otherwise. The square brackets denote a configurational average over all 2^N configurations of an N -bond system. For example, for $d = 1$, $P_{ij} = p^{|i-j|}$. The mean cluster size or 'susceptibility' is given by the fluctuation relation

$$S(p) = \chi_p(p) = \sum_{ij} P_{ij} \sim \varepsilon^{-\gamma}, \tag{1b}$$

where $\varepsilon = (p_c - p)/p_c \rightarrow 0^+$ and γ is the mean-size critical exponent. For $d = 1$, the known result (Reynolds *et al* 1977) $S(p) = (1+p)/(1-p) \sim \varepsilon^{-1}$ follows immediately from (1b).

These ideas can be extended to other quantities (Fisch and Harris 1978, Coniglio 1982). Thus for the backbone we may define

$$B_{ij} = [b_{ij}\nu_{ij}] \tag{2a}$$

where b_{ij} is the number of backbone bonds connecting sites i and j . For $d = 1$, $B_{ij} = |i-j|p^{|i-j|}$. In analogy with equation (1b), we may define the 'backbone susceptibility'

$$\chi_{BB}(p) = \sum_{ij} B_{ij}. \tag{2b}$$

For $d = 1$, we have $\chi_{BB}(p) = 2p/(1-p)^2$.

Below p_c , the mean number of backbone bonds L_{BB} connecting two sites i and j that are separated a distance of the order of the correlation length is (Coniglio 1982)

$$L_{BB}(p) = \sum_{ij} B_{ij} / \sum_{ij} P_{ij} \sim \varepsilon^{-\zeta_B}, \tag{3}$$

which defines the critical exponent ζ_B for the backbone.

Using the cumulant method, we have developed a low-density expansion for $\chi_{BB}(p)$ in the form

$$\chi_{BB}(p) = \sum_{n=1}^{\infty} B_n(d)p^n, \tag{4}$$

and have evaluated the coefficients $B_n(d)$ in closed form for all d for $1 \leq n \leq 10$. To this end, we have generalised the inclusion-exclusion principle for the pair connectedness function to the backbone. The generalisation arises from the fact that the number of backbone bonds between sites i and j , like the pair connectedness P_{ij} , is independent of any 'dangling ends'. Therefore we only need self-avoiding walks (SAWs) and loop diagrams that are constructed by the union of SAWs between i and j . Thus for any graph G the average B_{ij} is given by

$$B_{ij} = \sum_g C(g)E(g)p^{b(g)}, \tag{5a}$$

where the summation runs over all subgraphs of the graph G , $E(g)$ is the embedding constant, $b(g)$ is the number of bonds of g , and the cumulant $C(g)$ satisfies the recursive relation (Essam 1971, Fisch and Harris 1978)

$$C(g) = V(g) - \sum_{g'} C(g'). \tag{5b}$$

Here $V(g)$ is the value of g , while g' are the subgraphs of g .

For example, consider the simple four-bond graph $G(\square)$, where the crosses denote sites i and j . Applying (5), we find

$$\begin{aligned}
 B_{ij} &= C(\square) p^4 + 2C(\square) p^2 \\
 &= \left[V(\square) - 2V(\square) \right] p^4 + 2V(\square) p^2 \\
 &= [4 - 2 \times 2] p^4 + [2 \times 2] p^2.
 \end{aligned}
 \tag{6a}$$

Had we assigned the value $V(g) = 1$ for each graph, then we would have recovered the usual inclusion-exclusion principle,

$$P_{ij} = [1 - 2 \times 1] p^4 + [2 \times 1] p^2 = -p^4 + 2p^2.
 \tag{6b}$$

For the infinite lattice, with two fixed sites i and j , there are of course an *infinite* number of SAWS joining i and j (unless the lattice has directed bonds). Therefore we cannot perform an exact calculation, but instead must perform an expansion in increasing powers of p , by including in the calculation graphs with increasing numbers of bonds and multiplying the contribution of each graph by the appropriate lattice embedding constant (Fisch 1977). Thus the initial terms in this expansion are

$$\begin{aligned}
 \chi_{BB}(p) &= C_{BB}(\times - \times) [2d] p + C_{BB}(\times - \cdot - \times) [(2d)(2d - 1)] p^2 \\
 &\quad + C_{BB}(\times - \cdot - \cdot - \times) [(2d)(2d - 1)^2] p^3 \\
 &\quad + C_{BB}(\times - \cdot - \cdot - \cdot - \times) [2d(2d - 1)^3 - 4d(d - 1)] p^4 \\
 &\quad + \left[4C_{BB} \left(\begin{array}{c} \cdot - \times \\ \times - \cdot \end{array} \right) + 8C_{BB} \left(\begin{array}{c} \cdot - \cdot \\ \times - \times \end{array} \right) \right] d(d - 1) p^4.
 \end{aligned}
 \tag{7a}$$

Here the backbone cumulant $C_{BB}(g)$ for a diagram g is given by

$$C_{BB}(g) = V(g) - \sum_{g' \in g} C_{BB}(g').
 \tag{7b}$$

The expansion (7) increases rapidly in complexity, but the requisite diagrams and lattice embedding constants are known through order p^{10} (Baker *et al* 1967, Fisch 1977). Therefore it is necessary only to apply the recursive relation (6) to each graph g and then to calculate the number of backbone bonds in each subgraph g' . The general- d expressions in the form of (7) are given in table 2.

We find that the coefficients $B_n(d)$ in the series for the backbone $\chi_{BB}(p)$ are of the same form as the coefficients in the series for mean size $S(p)$,

$$B_n(d) = \sum_{k=1}^n W_{n,n-k+1} \binom{d}{k}.
 \tag{8}$$

The (d -independent) coefficients $W_{n,n-k+1}$ are given in table 2(a) up to $n = 10$. We find for the first two coefficients the simple results

$$W_{n1} = n 2^n n!
 \tag{9a}$$

$$W_{n2} = 2n(n - 1)^2 2^{n-2} (n - 1)!
 \tag{9b}$$

$$W_{n3} = \frac{1}{3} n 2^{n-4} (n - 2)! (6n^4 - 40n^3 + 96n^2 - 128n + 126) + 24(n - 3) 2^{n-4} (n - 2).
 \tag{9c}$$

In order to form the function $L_{BB}(p)$ of (3), we must also obtain general- d expressions for the mean-size function $S(p)$ for bond percolation with *site* counting (as opposed to *bond* counting, as in the conventional series expansions). Fortunately, Fisch and Harris (1978) tabulate the needed results.

The resulting series were analysed for $d = 2-6$ by the usual extrapolation procedures. We found that Padé approximants provided the most consistent estimates of the backbone exponent ζ_B defined in (3), and the results are shown in table 1. Table 1 also shows $D_B = \zeta_B/\nu$, the fractal dimension of the backbone. For comparison, we include the prediction $D_B = \ln(d+1)/\ln 2$ of the Sierpinski gasket model. We see that the discrepancy is most serious for large d .

The same procedure developed above for the number of backbone bonds B_{ij} can be readily extended to the *minimum* number of bonds M_{ij} (Middlemiss *et al* 1980, Pike and Stanley, 1981), where $M_{ij} = [m_{ij}\nu_{ij}]$ in analogy to (2a). The general- d results are given in table 2(b) while table 1 also gives the results of extrapolation procedures for the exponent ζ_{\min} defined through

$$L_{\min}(p) = \sum_j M_{ij} / \sum_j P_{ij} = \chi_{\min}(p) / \chi_p(p) \sim \varepsilon^{-\zeta_{\min}}, \quad (10a)$$

where

$$\chi_{\min}(p) = \sum_{n=1}^{\infty} M_n(d)p^n. \quad (10b)$$

Similarly we can define

$$L_{\text{red}}(p) = \sum_j R_{ij} / \sum_j P_{ij} = \chi_{\text{red}}(p) / \chi_p(p) \sim \varepsilon^{-\zeta_{\text{red}}}, \quad (11a)$$

$$\chi_{\text{red}}(p) = \sum_{n=1}^{\infty} R_n(d)p^n, \quad (11b)$$

where $R_{ij} = [r_{ij}\nu_{ij}]$ is the number of singly connected ('red') bonds or links between sites i and j . Pike and Stanley (1981) found that $L_{\text{red}}(p)$ diverges at p_c , contrary to some intuitive expectations, and Coniglio (1981, 1982) proved rigorously that $\zeta_{\text{red}} = 1$ for all spatial dimensionalities d . The proof was based on the 'Lemma' that $L_{\text{red}}(p) = p(d/dp) \log S(p)$. We find that Coniglio's lemma holds 'term-by-term' in the sense that for each graph in the low-density series expansion,

$$L_{ij} = p(d/dp) \log P_{ij}. \quad (12)$$

In table 2, we tabulate the general- d results for the function $\sum_j R_{ij}$.

Although less reliable than direct analysis, an approximation to the functions $\chi_{BB}(p)$, $\chi_{\min}(p)$ and $\chi_{\text{red}}(p)$ may be obtained by deriving expansions in the variable $1/\sigma$, where $\sigma = 2d-1$. Following the procedure used for $\chi_p(p)$ (Gaunt and Ruskin 1978), we find to $O(1/\sigma^2)$,

$$\chi_{BB} = \chi_{BB}^{\text{CT}} + (-5x^2/(1-x)^3 + x/(1-x)^2 - 18x/(1-x) + 7x + x^2)\sigma^{-2} + \dots, \quad (13a)$$

$$\chi_{\min} = \chi_{\min}^{\text{CT}} + (-5x^2/(1-x)^3 - 3x/(1-x)^2 - 6/(1-x) - x - 3x^2)\sigma^{-2} + \dots, \quad (13b)$$

$$\chi_{\text{red}} = \chi_{\text{red}}^{\text{CT}} + [-5x^2/(1-x)^3 - 5x/(1-x)^2 - 5x - 5x^2]\sigma^{-3} + \dots, \quad (13c)$$

where $x = \sigma p$. Following the procedure used in deriving

$$\chi_p^{\text{CT}}(p) = 1 + (\sigma^{-1} + 1) \sum_{r=1}^{\infty} (\sigma p)^r = (1+p)/(1-\sigma p), \quad (14a)$$

we obtain

$$\chi_{\text{BB}}^{\text{CT}} = \chi_{\text{min}}^{\text{CT}} = \chi_{\text{red}}^{\text{CT}} = (1 + \sigma^{-1}) \sum_{r=1}^{\infty} r(\sigma p)^r = (\sigma + 1)p/(1 - \sigma p)^2. \quad (14b)$$

In summary, then, we have calculated the first low-density expansions for three quantities characterising the backbone fractal of the incipient infinite cluster at the percolation threshold. These are L_{BB} , L_{min} and L_{red} which, respectively, are the mean number of backbone bonds, the mean length of minimum path, and the mean number of 'red' bonds between sites i and j separated by a correlation length. Extrapolations for the corresponding critical exponents, ζ_{BB} and ζ_{min} , are presented in table 1; ζ_{R} is known to diverge as $1/\varepsilon$ for all d (Coniglio 1982).

After this work was completed, we received two interesting preprints (Harris 1983, Harris and Lubensky 1983) which apply field-theoretic methods to calculate *one* of these quantities, L_{BB} , as a series in $(6-d)$; for $d > 3$, their results are not inconsistent with ours based on series expansions.

We wish to thank S Redner and especially A Coniglio, as well as A B Harris for providing us with a copy of Fisch (1977).

References

- Baker G A, Gilbert H E, Eve J and Rushbrooke G S 1967 *BNL 50053(T-460)* (Brookhaven National Laboratory)
- Coniglio A 1981 *Phys. Rev. Lett.* **46** 250
- 1982 *J. Phys. A: Math. Gen.* **15** 3829
- Dunn A G, Essam J W and Ritchie D S 1975 *J. Phys. C: Solid State Phys.* **8** 4219
- Essam J W 1971 *J. Math. Phys.* **12** 874
- Fisch R 1977 *PhD thesis* University of Pennsylvania
- Fisch R and Harris A B 1978 *Phys. Rev. B* **18** 416
- Gaunt D S and Ruskin H 1978 *J. Phys. A: Math. Gen.* **11** 1369
- Gaunt D S and Sykes M F 1983 *J. Phys. A: Math. Gen.* **16** 783
- Gaunt D S, Sykes M F and Ruskin H 1976, *J. Phys. A: Math. Gen.* **9** 1899
- Gefen Y, Aharony A, Mandelbrot B B and Kirkpatrick S 1981 *Phys. Rev. Lett.* **47** 1771
- Hamann D R 1983 *Phys. Today* **36** [5] 25
- Harris A B 1983 *Preprint*
- Harris A B and Lubensky T C 1983 *Preprint*
- Kirkpatrick S 1978 *AIP conference* **40** 99
- Li P S and Streider W 1982 *J. Phys. C: Solid State Phys.* **15** L1235
- Middlemiss K M, Whittington S G and Gaunt D S 1980 *J. Phys. A: Math. Gen.* **13** 1835
- Pike R and Stanley H E 1981 *J. Phys. A: Math. Gen.* **14** L169
- Reynolds P J, Stanley H E and Klein W 1977 *J. Phys. A: Math. Gen.* **11** L203
- Shlifer G, Klein W, Reynolds P J and Stanley H E 1979 *J. Phys. A: Math. Gen.* **12** L169
- Stanley H E 1977 *J. Phys. A: Math. Gen.* **10** L211
- Stanley H E and Coniglio A 1983 in *Percolation Structures and Processes* ed G Deutscher, R Zallen and J Adler (Bristol: Adam Hilger) p 101
- Stanley H E, Redner S and Yang Z R 1982 *J. Phys. A: Math. Gen.* **15** L569
- Stauffer D 1979 *Phys. Rep.* **54** 1