

The analogue of the Bäcklund transformation for integrable many-body systems

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1983 J. Phys. A: Math. Gen. 16 671

(<http://iopscience.iop.org/0305-4470/16/3/527>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 129.252.86.83

The article was downloaded on 30/05/2010 at 17:02

Please note that [terms and conditions apply](#).

Corrigendum

The analogue of the Bäcklund transformation for integrable many-body systems

Wojciechowski S 1982 *J. Phys. A: Math. Gen.* **15** L653–8

In equation (11) the summation \sum should be replaced by the product \prod .

Vacant animals and 2D to 1D crossover in the percolation problem

Gefen Y, Kanto Y and Deutscher G 1982 *J. Phys. A: Math. Gen.* **15** L553

Equation (4) should read

$$\sigma_{p \geq p_c}(L_y) \sim L_y^{-1/\nu} f[L_y^{1/\nu}(p - p_c)].$$

The reference Gefen Y 1981 *Phys. Rev. Lett.* **47** 1771 should read:

Gefen Y, Aharony A, Mandelbrot B B and Kirkpatrick S 1981 *Phys. Rev. Lett.* **47** 1771