

Asymptotic operator algebras in quantum mechanics

This article has been downloaded from IOPscience. Please scroll down to see the full text article. 1984 J. Phys. A: Math. Gen. 17 2363 (http://iopscience.iop.org/0305-4470/17/11/532)

View the table of contents for this issue, or go to the journal homepage for more

Download details: IP Address: 129.252.86.83 The article was downloaded on 31/05/2010 at 06:54

Please note that terms and conditions apply.

Corrigenda

Asymptotic operator algebras in quantum mechanics

Kay-Kong Wan and R G D McLean 1984 J. Phys. A: Math. Gen. 17 825-36

The last line on page 826 should read

$$\|(E_x(b_{r_k}) - E_x(b_{r_{k-1}}))\psi_{t_k}\|^2 > 1 - 1/k.$$

The expression in (1) of Definition 3 on page 829 should be

 $\mathscr{A}_{\Lambda} = \{ E_x(\Lambda) A E_x(\Lambda) \colon A \in B(\mathscr{H}) \}.$

The line immediately above Lemma 2 on page 831 should read:

by the lemma and theorem below.

On page 832 the expression at the end of the line labelled (4) in the Proof should be

 $A \in$, not $A \in \Delta$.

On page 834 the expression at the end of the first line in the last paragraph at the bottom should read

 $G \in$, not $G \in \Delta$.

Observables of asymptotically vanishing correlations, states at infinity and quantum separability

Kay-Kong Wan and R G D McLean 1984 J. Phys. A: Math. Gen. 17 837-46

The first line on page 839 should read

Postulate 1. A free quantum particle in \mathbb{R}^n has associated with it the C^* -algebra...