

Gauge invariance as the Lie-Backlund transformation group

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1988 J. Phys. A: Math. Gen. 21 L681

(<http://iopscience.iop.org/0305-4470/21/13/002>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 129.252.86.83

The article was downloaded on 31/05/2010 at 12:23

Please note that [terms and conditions apply](#).

LETTER TO THE EDITOR

Gauge invariance as the Lie-Bäcklund transformation group

K Kiiranen† and V Rosenhaus

Institute of Physics, Riia 142, Tartu 202400, Estonian SSR, USSR

Received 18 December 1987

Abstract. The gauge transformations in the extended (to potentials and their derivatives) space are shown to lead to the Lie-Bäcklund tangent transformation group.

Let us first consider the simplest gauge theory—electrodynamics. The Maxwell equation

$$\begin{aligned} \partial_\mu F_{\mu\nu} &= 0 \\ F_{\mu\nu} &= A_{\nu,\mu} - A_{\mu,\nu} \end{aligned} \tag{1}$$

is a well known invariant under the gauge (gradient) transformation

$$A_\mu \rightarrow A_\mu + \partial_\mu \varphi(x). \tag{2}$$

(We give all expressions for the Euclidean space, but the obtained results are valid also for the Minkowski space.) In the extended space $(x_\mu, A_\alpha, A_{\alpha,\beta}, A_{\alpha,\beta\gamma}, \dots)$ ($A_{\alpha,\beta} = \partial A_\alpha / \partial x_\beta$, one can write the transformation corresponding to (2):

$$A_\mu \rightarrow A_\mu + d_\mu F(x_\nu, A_\alpha, A_{\alpha,\beta}, A_{\alpha,\beta\gamma}, \dots) \tag{3}$$

where d_μ is the total derivative

$$d_\mu \equiv \frac{\partial}{\partial x_\mu} + A_{\alpha,\mu} \frac{\partial}{\partial A_\alpha} + A_{\alpha,\beta\mu} \frac{\partial}{\partial A_{\alpha,\beta}} + \dots$$

(transformations (3) and (2) and obviously invariant).

Now let us recall that the system of differential equations

$$\omega_p(x_\mu, A_\alpha, A_{\alpha,\beta}, A_{\alpha,\beta\gamma}, \dots) = 0 \quad p = 1, \dots, M$$

$(x_\mu$ and A_α are the arguments and functions respectively) admits a Lie-Bäcklund tangent transformation group generated by a Lie-Bäcklund operator

$$\begin{aligned} X &= f_\alpha \frac{\partial}{\partial A_\alpha} + (d_\nu f_\alpha) \frac{\partial}{\partial A_{\alpha,\nu}} + (d_\mu d_\nu f_\alpha) \frac{\partial}{\partial A_{\alpha,\nu\mu}} + \dots \\ f_\alpha &= f_\alpha(x_\mu, A_\beta, A_{\beta,\nu}, A_{\beta,\nu\mu}, \dots) \end{aligned} \tag{4}$$

† Present address: Tartu State University, Tähe 4, Tartu 202400, Estonian SSR, USSR.

(we write the corresponding tangent vector field of the group in the canonical form) if (and only if)

$$\begin{aligned} X\omega_p &= 0 \\ \omega_p &= 0 \\ d_i\omega_p &= 0 \\ d_jd_i\omega_p &= 0 \\ &\vdots \end{aligned}$$

(where $d_i\omega_p = 0, d_jd_i\omega_p = 0 \dots$ are the differential consequences of the initial system). For details of the theory of Lie-Bäcklund transformations, see Anderson and Ibragimov (1979) and Ibragimov (1983).

Thus, the Maxwell equation (1) admits the group with the Lie-Bäcklund operator (4) with

$$f_\mu = d_\mu\varphi \tag{5}$$

where $\varphi = \varphi(x_\nu, A_\alpha, A_{\alpha\beta}, \dots)$ is an arbitrary function. If $\varphi = \varphi(x)$, equation (5) defines the usual gauge transformation (2). Each U(1) gauge invariant system (depending on $F_{\mu\nu}$ only) evidently possesses the same property.

Now let us proceed to non-Abelian gauge theories and consider the Yang-Mills equation with an arbitrary simple gauge group G.

$$\begin{aligned} \partial_\mu G_{\mu\nu}^a + gf_{abc}A_\mu^b G_{\mu\nu}^c &= 0 \\ G_{\mu\nu}^a &= A_{\nu,\mu}^a - A_{\mu,\nu}^a + gf_{abc}A_\mu^b A_\nu^c \quad a, b, c = 1, \dots, N \end{aligned} \tag{6}$$

where N is the dimension of the group G and f_{abc} are the structure constants. The tensor f_{abc} is completely antisymmetric, due to the choosing of the invariant inner product of the basis generators orthonormal (in the adjoint representation):

$$(L_a, L_b) = K \text{Tr}(L_a L_b) = \delta_{ab}.$$

Using the invariance of the Yang-Mills equation (6) under gauge transformations (in the infinitesimal and finite form)

$$\begin{aligned} A_\mu^a &\rightarrow A_\mu^a - \frac{1}{g} \partial_\mu \omega^a + f_{abc} \omega^b A_\mu^c \\ A_\mu &\rightarrow UA_\mu U^{-1} + \frac{i}{g} U \partial_\mu U^{-1} \\ A_\mu &= A_\mu^a L_a \quad U = \exp(-i\omega^a L_a) \quad \omega^a = \omega^a(x) \end{aligned} \tag{7}$$

we introduce the dependence of the group parameters on all the variables of the extended space $x_\mu, A_\mu^a, A_{\mu,\alpha}^a, A_{\mu,\alpha\beta}^a, \dots$ (analogously to the Abelian case).

Then equations (7) change to

$$\begin{aligned} A_\mu^a &\rightarrow A_\mu^a - \frac{1}{g} d_\mu \omega^a + f_{abc} \omega^b A_\mu^c \\ A_\mu &\rightarrow UA_\mu U^{-1} + \frac{i}{g} d_\mu U^{-1} \\ \omega^a &= \omega^a(x_\mu, A_\mu^b, A_{\mu,\alpha}^b, \dots) \end{aligned} \tag{8}$$

(the invariance of equations (6) under transformations (8) can be checked easily). Thus, the Yang-Mills equation admits a group of Lie-Bäcklund transformations with the operator

$$\begin{aligned}
 X_\varphi &= f_{\mu\varphi}^a \frac{\partial}{\partial A_\mu^a} + (d_\alpha f_{\mu\varphi}^a) \frac{\partial}{\partial A_{\mu,\alpha}^a} + (d_\beta d_\alpha f_{\mu\varphi}^a) \frac{\partial}{\partial A_{\mu,\alpha\beta}^a} + \dots \\
 f_{\mu\varphi}^a &= -\frac{1}{g} d_\mu \varphi^a + f_{abc} \varphi^b A_\mu^c \\
 \varphi^a &= \varphi^a(x_\nu, A_\nu^b, A_{\nu,\alpha}^b, \dots).
 \end{aligned}
 \tag{9}$$

The commutation relation in the algebra of the Lie-Bäcklund operators is

$$\begin{aligned}
 [X_\varphi, X_\psi] &= X_\theta \\
 \theta^a &= X_\psi \psi^a - X_\varphi \varphi^a + f_{abc} \psi^b \varphi^c.
 \end{aligned}
 \tag{10}$$

Here the Jacobi identity

$$f_{abn} f_{ncd} + f_{bcn} f_{nad} + f_{can} f_{nbd} = 0$$

has been used. Note that the Lie-Bäcklund algebra (9) for equation (6) is non-trivial: it cannot be obtained from the Lie point symmetry group of the Yang-Mills equation by a simple prolongation (e.g., Ovsyannikov 1978). Really, besides the local gauge invariance (7) the Yang-Mills equation possesses the group of conformal transformations (Mack and Salam 1969, Schwartz 1982). Therefore, by a simple prolongation the group of point transformations of the Yang-Mills equation evidently does not lead to the group with operators (9). (The same conclusion is valid also for the Maxwell equation.)

Similar Lie-Bäcklund operators can also be constructed for other gauge theories with interaction between different fields. For example, let us consider the gauge invariant (with an arbitrary simple group G of dimension N) system of Yang-Mills fields coupled with the multiplet of the N scalar particles ϕ^a :

$$L = -\frac{1}{4} G_{\mu\nu}^a G_{\mu\nu}^a + \frac{1}{2} (D_\mu \phi^a)(D_\mu \phi^a) - V(\phi^2) \tag{11}$$

where the adjoint representation is chosen,

$$D_\mu \phi^a = \partial_\mu \phi^a + g f_{abc} A_\mu^b \phi^c \quad a, b, c = 1, \dots, N$$

and $V(\phi^2)$ is the G -invariant polynomial with respect to ϕ^a (for the Higgs model $V(\phi^2)$ is the polynomial of fourth order with the minimum at $\phi = v$ (e.g., Abers and Lee 1973)).

The local gauge transformations leaving Lagrangian (11) invariant, are given by expressions (7) with $(\omega^a = \omega^a(x))$

$$\begin{aligned}
 \phi^a &\rightarrow \phi^a + f_{abc} \omega^b \phi^c \\
 \phi^a &\rightarrow (U)_{ab} \phi^b.
 \end{aligned}$$

As earlier, let us allow for the dependence of the group parameters ω^a on all the variables of the extended space:

$$\omega^a = \omega^a(x_\mu, A_\mu^b, A_{\mu,\alpha}^b, \dots, \phi^b, \phi_{,\alpha}^b, \dots) \tag{12}$$

(the invariance of Lagrangian (11) holds as well).

The Lie-Bäcklund operator for system (11) has the form

$$\begin{aligned}
 X_\omega = & f_\mu^a \frac{\partial}{\partial A_\mu^a} + (d_\alpha f_\mu^a) \frac{\partial}{\partial A_{\mu,\alpha}^a} + \dots \\
 & + \bar{f}_\phi^a \frac{\partial}{\partial \phi^a} + (d_\alpha \bar{f}_\phi^a) \frac{\partial}{\partial \phi_{,\alpha}^a} + \dots
 \end{aligned}
 \tag{13}$$

where

$$f_\omega^a = -\frac{1}{g} d_\mu \omega^a + f_{abc} \omega^b A_\mu^c$$

$$\bar{f}_\omega^a = f_{abc} \omega^b \phi^c$$

and ω^a is an arbitrary function (12). As in the case of the pure Yang-Mills theory, the commutator of two Lie-Bäcklund fields has the same form (10) in space (12).

The authors are grateful to M Kóiv for valuable discussions.

References

- Abers E S and Lee B W 1973 *Phys. Rep.* **9C** 1-141
 Anderson R L and Ibragimov N H 1979 *Lie-Bäcklund Transformations in Applications* (Philadelphia: SIAM)
 Ibragimov N H 1983 *Transformation Groups in Mathematical Physics* (Moscow: Nauka) (in Russian)
 Mack G and Salam A 1969 *Ann. Phys., NY* **53** 174-202
 Ovsyannikov L V 1978 *Group Analysis of Differential Equations* (Moscow: Nauka) (Engl. transl: Ames W F
 1982 *Group Analysis of Differential Equations* (New York: Academic))
 Schwartz F 1982 *Lett. Math. Phys.* **6** 355-9