

## The $q$ -deformed boson realisation of the quantum group $SU(n)_q$ and its representations

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1989 J. Phys. A: Math. Gen. 22 L983

(<http://iopscience.iop.org/0305-4470/22/21/001>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 129.252.86.83

The article was downloaded on 31/05/2010 at 12:44

Please note that [terms and conditions apply](#).

LETTER TO THE EDITOR

The  $q$ -deformed boson realisation of the quantum group  $SU(n)_q$  and its representations

Chang-Pu Sun<sup>†‡§</sup> and Hong-Chen Fu<sup>‡</sup>

<sup>†</sup> CCAST (World Laboratory), P O Box 8730, Beijing, People's Republic of China

<sup>‡</sup> Physics Department, Northeast Normal University, Changchun, People's Republic of China||

<sup>§</sup> Theoretical Physics Division, Nankai Institute of Mathematics, People's Republic of China

Received 30 August 1989

**Abstract.** The  $q$ -deformed boson realisation of the quantum group  $SU(n)_q$  ( $(A_{n-1})_q$ ) is constructed and certain types of representations of  $SU(n)_q$  are obtained in the  $q$ -deformed Fock space by this boson realisation. The Jimbo representations of the quantum group  $SU(2)_q$  are given as an example in this letter.

Recently, as a new type of algebraic structure, the  $q$ -deformation of Lie algebras has been discovered by Jimbo [1-3] and Drinfeld [4]. It is mathematically a Hopf algebra and loosely called the quantum group. It appears when one tries to solve the famous quantum Yang-Baxter equation (QYBE) from different physical models, such as the exactly soluble models in statistical mechanics, integrable models in two-dimensional field theory, etc. The representations of the quantum group associated with its QYBE are of central importance in these problems. So it is important to study the representations of quantum groups.

The boson realisation method is a useful method for studying representations of groups. It has been used to construct indecomposable representations of Lie algebras [5], Lie superalgebras [6] and Loop algebras [7] and give inhomogeneous differential realisations of Lie algebras [8]. In this letter, by introducing the  $q$ -deformation of the Bose algebra, we generalise the usual boson realisation method to obtain explicit expressions for the representations of the quantum group  $SU(n)_q$  ( $(A_{n-1})_q$ ).

We first consider a  $q$ -deformation of the one-state Bose algebra (we call it  $q$  Bose algebra  $\mathcal{B}_q$ ) generated by operators  $b$ ,  $b^\dagger$ ,  $\mathbb{1}$  and  $N$  in Hilbert space that satisfy

$$\begin{aligned} bb^\dagger - q^{-1}b^\dagger b &= q^N & [B, b^\dagger] &= b^\dagger & [N, b] &= -b \\ [\mathbb{1}, x] &= 0 & \forall x &\in \mathcal{B}_q. \end{aligned} \tag{1}$$

When the parameter  $q \rightarrow 1$  (or  $q = e^\hbar$ ,  $\hbar \rightarrow 0$ )  $\mathcal{B}_q$  becomes the usual Bose algebra. It is easy to prove that

$$\begin{aligned} Nb^{\dagger n} &= b^\dagger N + nb^{\dagger n} \\ b^\dagger q^N &= q^{-1}q^N b^\dagger & bq^N &= qq^N b \\ bb^{\dagger n} &= q^{1-n}[n]q^N b^{\dagger n-1} + q^{-n}b^{\dagger n}b & [n] &\equiv (q^n - q^{-n})/(q - q^{-1}) \end{aligned} \tag{2}$$

|| Current mailing address of Chong-Pu Sun.

by induction from (1). The relations in (2) can naturally be extended to the case of the  $n$ -state Bose algebra  $\{b_i, b_i^\dagger, \mathbb{1}, N_i \mid i = 1, 2, \dots, n\}$ . For the  $n$ -state Bose algebra we define the  $q$ -analogous Fock space  $F_q$  (also called  $q$ -Fock space) with the basis

$$|m_i\rangle = |m_1, m_2, \dots, m_n\rangle = \left( \prod_{i=1}^n \frac{b_i^{\dagger m_i}}{\sqrt{[m_i]!}} \right) |0\rangle \quad [n]! \equiv [n][n-1] \dots [2][1] \quad (3)$$

where the vacuum state  $|0\rangle$  satisfies  $b_i|0\rangle = 0, N_i|0\rangle = 0 \ (i = 1, 2, \dots, n)$ . Then, it follows from (2) that

$$\begin{aligned} b_j^\dagger |m_j\rangle &= \sqrt{[m_j+1]} |m_j + \delta_j\rangle & b_j |m_j\rangle &= \sqrt{[m_j]} |m_j - \delta_j - \delta_{ij}\rangle \\ N_j |m_j\rangle &= m_j |m_j\rangle & j &= 1, 2, \dots, n. \end{aligned} \quad (4)$$

For given classical Lie algebra  $\mathcal{L}$  with the Chevalley basis  $\{h_i, e_i, f_i, i = 1, 2, \dots, l\}$  which satisfies

$$\begin{aligned} [h_i, h_j] &= 0 & [e_i, f_j] &= \delta_{ij} h_i \\ [h_i, e_j] &= A_{ij} e_j & [h_i, f_j] &= -A_{ij} e_j \end{aligned} \quad (5)$$

( $\mathbf{A}$  is the Cartan matrix of  $\mathcal{L}$ ). A  $q$ -deformed boson realisation (also called a  $q$ -boson realisation) of the quantum group  $\mathcal{L}_q$  associated with the Lie algebra  $\mathcal{L}$  is a mapping  $B$  of  $\mathcal{L}$  onto the operator algebra  $\mathcal{O}$  on the  $q$ -Fock space  $F_q$ , which satisfies

$$\begin{aligned} [\tilde{h}_i, \tilde{h}_j] &= 0 & [\tilde{e}_i, \tilde{f}_j] &= \delta_{ij} [\tilde{h}_i] \\ [\tilde{h}_i, \tilde{e}_j] &= A_{ij} \tilde{e}_j & [\tilde{h}_i, \tilde{f}_j] &= -A_{ij} \tilde{f}_j \\ \tilde{x} &\equiv B(x) & \forall x &\in \mathcal{L} \end{aligned} \quad (6)$$

where we define

$$[\hat{0}] = \frac{q^{\hat{0}} - q^{-\hat{0}}}{q - q^{-1}}$$

for any operator  $\hat{0}$ . The  $q$ -Boson realisation of  $\mathcal{L}_q$  generated by  $\{\tilde{h}_i, \tilde{e}_i, \tilde{f}_i, i = 1, 2, \dots, l\}$  can be regarded as a subalgebra of  $\mathcal{O}$ , i.e.  $\mathcal{L}_q \subset \mathcal{O}$ .

Consider the Lie algebra  $A_{n-1} = \text{SU}(n)$  with the Chevalley basis

$$h_j = E_{jj} - E_{j+1, j+1} \quad e_j = E_{jj+1} \quad f_j = E_{j+1, j} \quad j = 1, 2, \dots, n-1 \quad (7)$$

where  $E$  is a  $n \times n$  matrix with elements  $(E_{ij})_{\alpha\beta} = \delta_{i\alpha} \delta_{j\beta} \ (\alpha, \beta = 1, 2, \dots, n)$ . The  $q$ -boson realisation of its quantum group  $(A_{n-1})_q$  is given by

$$\tilde{h}_j = N_j - N_{j+1} \quad \tilde{e}_j = b_j^\dagger b_{j+1} \quad \tilde{f}_j = b_{j+1}^\dagger b_j \quad j = 1, 2, \dots, n-1. \quad (8)$$

By direct computation we check

$$\begin{aligned} [\tilde{h}_j, \tilde{h}_i] &= 0 & [\tilde{e}_i, \tilde{f}_j] &= \delta_{ij} [\tilde{h}_i] \\ [\tilde{h}_i, \tilde{e}_j] &= A_{ij} \tilde{e}_j & [\tilde{h}_i, \tilde{f}_j] &= -A_{ij} \tilde{f}_j \end{aligned} \quad (9)$$

where  $A_{ij} = 2\delta_{ij} - \delta_{ij+1} - \delta_{i, j-1}$  is a element of the cartan matrix  $\mathbf{A}$  of  $\text{SU}(n)$ .

Now, we construct the representations of  $(A_{n-1})_q$  from the Boson realisation (8) by defining the action of  $(A_{n-1})_q$  on  $F_q$

$$\Gamma(x)|u\rangle = x|u\rangle \quad \forall x \in (A_{n-1})_q \quad |u\rangle \in F_q$$

obtaining

$$\begin{aligned}
 \Gamma(\tilde{h}_j)|m_i\rangle &= (m_j - m_{j+1})|m\rangle \\
 \Gamma(\tilde{e}_j)|m\rangle &= ([m_{j+1}][m_j + 1])^{1/2}|m_i + \delta_{ij} - \delta_{i,j+1}\rangle \\
 \Gamma(\tilde{f}_j)|m\rangle &= ([m_j][m_{j+1} + 1])^{1/2}|m_i - \delta_{ij} + \delta_{i,j+1}\rangle.
 \end{aligned}
 \tag{10}$$


Because the sum  $\sum_{i=1}^n m_i$  remains the same under the action of  $\Gamma$ ,  $F_q$  is decomposed into the direct sum of all the invariant subspaces  $F_q^{[m]}$  ( $m = 0, 1, 2, \dots$ ):

$$\left\{ |m_i\rangle \left| \sum_{i=1}^n m_i = m \right. \right\}.$$

Each one of these subspaces  $F_q^{[m]}$  ( $m = 0, 1, \dots$ ) carries an irreducible representation  $\Gamma^{[m]}$  of  $(A_{n-1})_q$  with dimension

$$d_n^m = \frac{(n+m-1)!}{m!(n-1)!}.
 \tag{11}$$

When  $q \rightarrow 1$  ( $q = e, \hbar \rightarrow 0$ ), the representation  $\Gamma^{[m]}$  becomes a symmetrised representation of Lie algebra  $A_{n-1}$  labelled by the Young diagram



In fact, from (10) we can give explicit matrices of some representations of the quantum group  $(A_{n-1})_q$ . For example, when  $n = 3$  and  $M = 2$ , the six-dimensional representation of  $SU(3)_q = (A_2)_q$  on the basis  $\{|2, 0, 0\rangle, |0, 2, 0\rangle, |0, 0, 2\rangle, |1, 1, 0\rangle, |0, 11\rangle, |101\rangle\}$  is

$$\begin{aligned}
 \Gamma^{[2]}(\tilde{h}_1) &= \text{diag}[2, -2, 0, 0, -1, 1] \\
 \Gamma^{[2]}(\tilde{h}_2) &= \text{diag}[0, 2, -2, 1, 0, -1] \\
 \Gamma^{[2]}(\tilde{e}_1) &= \begin{pmatrix} 0 & 0 & 0 & \sqrt{[2]} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \sqrt{[2]} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix} & \Gamma^{[2]}(\tilde{e}_2) &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \sqrt{[2]} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & \sqrt{[2]} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} \\
 \Gamma^{[2]}(f_1) &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \sqrt{[2]} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ \sqrt{[2]} & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix} & \Gamma^{[2]}(\tilde{f}_2) &= \begin{pmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & \sqrt{[2]} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & \sqrt{[2]} & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \end{pmatrix}
 \end{aligned}
 \tag{12}$$

The Jimbo standard angular momentum representation of the quantum group  $SU(2)_q$  is only a special example of the above discussion for  $n = 1$ . Let  $J_+ = \tilde{e}_1$ ,  $J_- = \tilde{f}_1$  and  $J_3 = \frac{1}{2}h_1$ , it follows from (10) that

$$\begin{aligned}
 \Gamma(J_+)|n_1 n_2\rangle &= ([n_1 + 1][n_2]^{1/2})|n_1 + 1, n_2 - 1\rangle \\
 \Gamma(J_-)|n_1 n_2\rangle &= ([n_1][n_2 + 1]^{1/2})|n_1 - 1, n_2 + 1\rangle \\
 \Gamma(J_3)|n_1 n_2\rangle &= \frac{1}{2}(n_1 - n_2)|n_1 n_2\rangle.
 \end{aligned}
 \tag{13}$$

By defining angular momentum basis  $|\Psi_{jm}\rangle = |j+m, j-m\rangle$  for the representation, (13) is rewritten as

$$\Gamma(J_{\pm})|\Psi_{jm}\rangle = (j \pm m + 1)[j \mp m]^{1/2}|\Psi_{j, m \pm 1}\rangle \quad (14)$$

$$\Gamma(J_3)|\Psi_{jm}\rangle = m|\Psi_{jm}\rangle.$$

This is just the Jimbo representation, which becomes the standard irreducible representation of  $SU(2)$  when  $q \rightarrow 1$ .

The method used in this paper can be applied to  $C_{n-1}$  and also expected to study quantum groups of any classical Lie algebra by further generalisations.

## References

- [1] Jimbo M 1985 *Lett. Math. Phys.* **10** 63
- [2] Jimbo M 1985 *Lett. Math. Phys.* **11** 247
- [3] Jimbo M 1986 *Commun. Math. Phys.* **102** 537
- [4] Drinfel'd V G 1986 *Proc. ICM Berkeley* p 798
- [5] Sun C-P 1987 *J. Phys. A: Math. Gen.* **20** 4551
- [6] Sun C-P 1987 *J. Phys. A: Math. Gen.* **20** 5823
- [7] Sun C-P 1987 *J. Phys. A: Math. Gen.* **20** L1157
- [8] Fu H-C and Sun C-P 1989 *J. Math. Phys.* **30** in press