

General solutions of $R^{j1/2}(x)$ and quantum group structure

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1991 J. Phys. A: Math. Gen. 24 4955

(<http://iopscience.iop.org/0305-4470/24/21/009>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 129.252.86.83

The article was downloaded on 01/06/2010 at 13:59

Please note that [terms and conditions apply](#).

General solutions of $R^{j_1 j_2}(x)$ and quantum group structure

Mo-Lin Ge†, Xu-Feng Liu†, Chang-Pu Sun†‡ and Kang Xue†‡

† Department of Theoretical Physics, Nankai Institute of Mathematics, Tianjin, 300071, People's Republic of China

‡ Department of Physics, North-east Normal University, Changchun, 130024, People's Republic of China

Abstract. A general form of solutions of braid relations associated with $V^j \otimes V^{1/2}$ for $Su(2)$ is calculated. A Yang-Baxterization prescription is presented to generate the corresponding solutions of Yang-Baxter equations. We show that this general form of solutions gives rise to a new construction of quantum group structure even though the q -commutation relations are kept standard.

1. Introduction

It is known that for spin models the solutions of braid relations

$$S_{12}^{j_1 j_2} S_{23}^{j_1 j_3} S_{12}^{j_2 j_3} = S_{23}^{j_2 j_3} S_{12}^{j_1 j_3} S_{23}^{j_1 j_2} \tag{1.1}$$

can best be explicitly constructed by the quantum groups (QA) through the standard theory of Drinfeld and Reshetikhin [1-3], where j_1, j_2 and j_3 are the weight labels associated with the corresponding spin spaces. In the following, for convenience, we will call the solutions that can be obtained by the QG (QG) in the usual way the 'standard' ones, which can also be worked with the help of the fusion rule [4, 5].

In comparison to the statistical model for a vertex model the Yang-Baxter equation (YBE) can be written in the form [6, 7]

$$\sum_{\gamma, \mu'' \nu''} W(\mu, \alpha | \gamma, \mu') W'(\nu, \delta | \beta, \nu'') W''(\nu, \mu'' | \nu', \mu') = \sum_{\gamma, \mu'', \nu''} W''(\nu, \mu | \nu'', \mu'') W'(\mu'', \alpha | \gamma, \mu') W(\nu'', \gamma | \beta, \nu') \tag{1.2}$$

where W, W' and W'' are Boltzmann weights. It is well known that if equation (1.2) can be additively parametrized, then one can introduce

$$T_{\mu' \alpha}^{\beta \nu'}(x) = W(\mu', \alpha | \beta, \nu') = \begin{array}{c} \beta \quad \nu' \\ \diagdown \quad \diagup \\ \theta \\ \diagup \quad \diagdown \\ \mu' \quad \alpha \end{array} \tag{1.3}$$

where $x = \exp(-i\theta) = \exp(-u)$ and matrices T satisfy

$$\sum_{\gamma, \mu'', \nu''} T_{\nu'' \gamma}^{\beta \nu'}(x) T_{\mu'' \alpha}^{\gamma \mu'}(xy) T_{\nu \mu''}^{\nu'' \mu''}(y) = \sum_{\gamma, \mu'', \nu''} T_{\nu' \mu''}^{\nu' \mu''}(y) T_{\nu \gamma}^{\beta \nu'}(xy) T_{\mu \alpha}^{\gamma \mu'}(x) \tag{1.4}$$

which by redefining

$$(\check{R}^{j_1 j_2}(x))_{a(j_1) b(j_2)}^{c(j_2) d(j_1)} = T_{\beta' \beta'}^{\alpha \alpha'}(x) \tag{1.5}$$

is referred to as the incomplete YBE:

$$\check{R}_{12}^{j_1 j_2}(x) \check{R}_{23}^{j_1 j_3}(xy) \check{R}_{12}^{j_2 j_3}(y) = \check{R}_{23}^{j_2 j_3}(y) \check{R}_{12}^{j_1 j_3}(xy) \check{R}_{23}^{j_1 j_2}(x). \tag{1.6}$$

Noting that by considering the asymptotic behaviour

$$S^{j_1 j_2} = \lim_{x \rightarrow 0} \check{R}^{j_1 j_2}(x) \tag{1.7}$$

the YBE is reduced to the braid relation (1.1).

If we consider the model shown in figure 1, where the full lines denote spin j and the broken lines denote spin $\frac{1}{2}$, then equation (1.1) is reduced to

$$\begin{aligned} S_{12}^{j \frac{1}{2}} S_{23}^{\frac{1}{2} j} S_{12}^{\frac{1}{2} j} &= S_{23}^{\frac{1}{2} j} S_{12}^{j \frac{1}{2}} S_{23}^{j \frac{1}{2}} \\ S_{12}^{\frac{1}{2} j} S_{23}^{j \frac{1}{2}} S_{12}^{\frac{1}{2} j} &= S_{23}^{j \frac{1}{2}} S_{12}^{\frac{1}{2} j} S_{23}^{\frac{1}{2} j} \\ S_{12}^{j \frac{1}{2}} S_{23}^{\frac{1}{2} j} S_{12}^{\frac{1}{2} j} &= S_{23}^{j \frac{1}{2}} S_{12}^{\frac{1}{2} j} S_{23}^{j \frac{1}{2}} \end{aligned} \tag{1.8}$$

which satisfies the graph in figure 2. As usual the CP invariance should be satisfied,

$$(S^{j \frac{1}{2}})_{c(\frac{1}{2})d(j)}^{a(j)b(\frac{1}{2})} = (S^{\frac{1}{2} j})_{-d(j)-c(\frac{1}{2})}^{-b(\frac{1}{2})-a(j)} \tag{1.9}$$

under which only two equations in equation (1.8) are independent:

$$\begin{aligned} S_{12}^{j \frac{1}{2}} S_{23}^{\frac{1}{2} j} S_{12}^{\frac{1}{2} j} &= S_{23}^{\frac{1}{2} j} S_{12}^{j \frac{1}{2}} S_{23}^{j \frac{1}{2}} \\ S_{12}^{\frac{1}{2} j} S_{23}^{j \frac{1}{2}} S_{12}^{\frac{1}{2} j} &= S_{23}^{j \frac{1}{2}} S_{12}^{\frac{1}{2} j} S_{23}^{\frac{1}{2} j}. \end{aligned} \tag{1.10}$$

Hence, solving the model in figure 1 consists of solving equation (1.10). So far no general solutions of equation (1.10) have been derived. All of the known solutions are based on the standard construction with usual treatment by QG or fusion rule.




Figure 1.




Figure 2.

Motivated by the results in [8–10] where the solutions are beyond the standard construction, for the braid relation with the same spin spaces

$$S_{12} S_{23} S_{12} = S_{23} S_{12} S_{23}$$

we want to find a general form of solutions for equation (1.10) which contains the standard ones as special case.

After lengthy computation we reach the following results:

(i) General solutions of equation (1.1) can be explicitly derived.

In the general case they depend on $j+2$ (for $j=1, 2, \dots$) or $j+\frac{3}{2}$ (for $j=\frac{3}{2}, \frac{5}{2}, \dots$) free parameters, respectively, and cannot be related to the standard ones by a similar transformation.

(ii) The Yang-Baxterization prescription [10] can be performed to generate solutions of the YBE [11]:

$$\begin{aligned} \check{R}_{12}^{j \frac{1}{2}}(x) \check{R}_{23}^{j \frac{1}{2}}(xy) \check{R}_{12}^{j \frac{1}{2}}(y) &= \check{R}_{23}^{j \frac{1}{2}}(y) \check{R}_{12}^{j \frac{1}{2}}(xy) \check{R}_{23}^{j \frac{1}{2}}(x) \\ \check{R}_{12}^{\frac{1}{2} j}(x) \check{R}_{23}^{\frac{1}{2} j}(xy) \check{R}_{12}^{\frac{1}{2} j}(y) &= \check{R}_{23}^{\frac{1}{2} j}(y) \check{R}_{12}^{\frac{1}{2} j}(xy) \check{R}_{23}^{\frac{1}{2} j}(x) \\ \check{R}_{12}^{\frac{1}{2} j}(x) \check{R}_{23}^{\frac{1}{2} j}(xy) \check{R}_{12}^{\frac{1}{2} j}(y) &= \check{R}_{23}^{\frac{1}{2} j}(y) \check{R}_{12}^{\frac{1}{2} j}(xy) \check{R}_{23}^{\frac{1}{2} j}(x). \end{aligned} \tag{1.11}$$

The procedure of Yang-Baxterization is the consequence of CP invariance and is based on the Reshetikhin approach [2] and cannot be made by simply following [12].

(iii) The FRT approach [13] is used to give QG structure for a given $S^{j\frac{1}{2}}$. The general solutions of equation (1.1) give rise to the standard structure of the QG, which is not surprising. However, the centralizer representation (in this paper only $j_2 = \frac{1}{2}$ is taken into account)

$$R^{j_1 j_2} = P^{j_1 j_2} S^{j_1 j_2} \tag{1.12}$$

where $P^{j_1 j_2}(V^{j_1} \otimes V^{j_2}) = V^{j_2} \otimes V^{j_1}$, possess a different structure from the standard one, namely

$$R^{j_1 j_2} = R_{\text{standard}}^{j_1 j_2} + \Delta^{j_1 j_2} \tag{1.13}$$

where $R_{\text{standard}}^{j_1 j_2}$ can be expressed in terms of the basis $e_s(v^{j_1})$ and $e^s(v^{j_2})$ in the usual way [1-3, 12]. The additional terms are determined through recurrence relations. In other words, the 'exotic' property occurs in the different way of construction of $R^{j_1 j_2}$ with the basis rather than in the commutation relations.

2. General solutions for $S^{j\frac{1}{2}}$

We are looking for general solutions satisfying equations (1.8) and (1.9) and the weight conservation

$$a(j) + b(\frac{1}{2}) = c(\frac{1}{2}) + d(j). \tag{2.1}$$

Then equations (1.8), (1.9) and (2.1) constrain $S^{j\frac{1}{2}}$ and $S^{j\frac{1}{2}}$ to the forms

$$(S^{j\frac{1}{2}})_{c(\frac{1}{2})d(\frac{1}{2})}^{a(\frac{1}{2})b(\frac{1}{2})} = q|_{a=b=c=d} + \delta_d^a \delta_c^b |_{a \neq b} + w \delta_c^a \delta_d^b |_{a < b} \tag{2.2}$$

$$(S^{j\frac{1}{2}})_{c(\frac{1}{2})d(j)}^{a(j)b(\frac{1}{2})} = p^{(a(j),b(\frac{1}{2}))} \delta_{d(j)}^{a(j)} \delta_{c(\frac{1}{2})}^{b(\frac{1}{2})} + q^{(a(j),b(\frac{1}{2}))} \delta_{d(j)-2b(\frac{1}{2})}^{a(j)} \delta_{-c(\frac{1}{2})}^{b(\frac{1}{2})} \tag{2.3}$$

where $w = q - q^{-1}$ and the parameters $p^{(a(j),b(\frac{1}{2}))}$ and $q^{(a(j),b(\frac{1}{2}))}$ are to be determined. Obviously, for $S^{j\frac{1}{2}}$ only the standard solution is allowed, as shown by equation (2.2).

Substituting equations (2.2) and (2.3) into equation (1.8) and taking equation (1.9) into account we obtain the relations that the unknown parameters must satisfy:

$$\begin{aligned} q^{(a(j),-\frac{1}{2})} &= 0 & p^{(a(j),\frac{1}{2})} &= qp^{(a(j)-1,\frac{1}{2})} & p^{(a(j),-\frac{1}{2})} &= q^{-1} p^{(a(j)-1,-\frac{1}{2})} \\ q^{(-a(j),\frac{1}{2})} q^{(a(j)-1,\frac{1}{2})} |_{a(j) > -j} &- q^{(a(j),\frac{1}{2})} q^{(-a(j)-1,\frac{1}{2})} |_{a(j) < j} \\ &= q^{-1} w (p^{(a(j),\frac{1}{2})} p^{(-a(j),-\frac{1}{2})} - p^{(-a(j),\frac{1}{2})} p^{(a(j),-\frac{1}{2})}). \end{aligned} \tag{2.4}$$

Without loss of generality, taking

$$p^{(j,\frac{1}{2})} = q \quad p(-j, -\frac{1}{2}) = Q \tag{2.5}$$

where q and Q are arbitrary parameters, we then obtain the general solutions of equation (1.8) under equation (1.9):

$$\begin{aligned} p^{(a(j),\frac{1}{2})} &= q^{a(j)+1-j} & p^{(a(j),-\frac{1}{2})} &= q^{-a(j)-j} Q & q^{(a(j),-\frac{1}{2})} &= 0 \\ q^{(-a(j)-1,\frac{1}{2})} q^{(a(j),\frac{1}{2})} &= q^{-2j} Q w^2 [j - a(j)]_q [j + a(j) + 1]_q & & & (a(j) \neq j) \end{aligned} \tag{2.6}$$

$$w = q - q^{-1}$$

where

$$[n]_q = (q^n - q^{-n}) / (q - q^{-1}). \tag{2.7}$$

Thus $q^{(-a(j)-1, \frac{1}{2})}$ can be determined if $q^{(a(j), \frac{1}{2})}$ are regarded as free parameters. In the general solutions there will be $j+2$ or $j+\frac{3}{2}$ free parameters for $j=1, 2, \dots$ or $j=\frac{3}{2}, \frac{5}{2}, \dots$, respectively.

When

$$Q = q \quad q^{(-a(j)-1, \frac{1}{2})} = q^{(a(j), \frac{1}{2})} \tag{2.8}$$

the solutions (2.6) are reduced to the standard solutions, under which

$$\begin{aligned} q^{(a(j), -\frac{1}{2})} &= 0 & p^{(a(j), \frac{1}{2})} &= q^{a(j)+1-j} \\ p^{(a(j), -\frac{1}{2})} &= q^{-a(j)+1-j} \\ q^{(a(j), \frac{1}{2})} &= q^{-j+\frac{1}{2}} w\{[j-a(j)]_q [j+a(j)+1]_q\}^{\frac{1}{2}} \end{aligned} \tag{2.9}$$

which coincide with the results given in [1-3]. Actually, by direct calculation, up to an overall factor q^{-j+1} , equation (2.9) leads to the representation of the universal R -matrix:

$$\begin{aligned} (R^{1j})_{m_1, m_2}^{m'_1, m'_2} &= \delta_{m'_1+m'_2}^{m_1+m_2} \frac{(1-q)^{m'_1-m_1}}{[m'_1-m_1]_q!} q^{m_1 m'_2 + m_2 m'_1} \\ &\times \left(\frac{[\frac{1}{2}+m'_1]_q! [\frac{1}{2}-m_1]_q! [j-m'_2]_q! [j+m_2]_q!}{[\frac{1}{2}-m'_1]_q! [\frac{1}{2}+m_1]_q! [j+m'_2]_q! [j-m_2]_q!} \right)^{1/2} \end{aligned} \tag{2.10}$$

($a(j)$ is taken as m).

To illustrate we now give some examples. When $j=1$ we have the solutions

$$S^{1\frac{1}{2}} = \begin{bmatrix} q & & & & & \\ & 0 & q^{-2}Q & & & \\ & 1 & q_1 & & & \\ & & & 0 & q^{-1}Q & \\ & & & q^{-1} & q_2 & \\ & & & & & \end{bmatrix} \tag{2.11}$$

with

$$q_1 q_2 = Qw(1-q^{-4}) \tag{2.12}$$

and

$$\tilde{S}^{1\frac{1}{2}} = \begin{bmatrix} q & & & & & \\ & 0 & \tilde{Q} & & & \\ & 1 & q_1 & & & \\ & & & \tilde{q}_2 & q\tilde{Q} & \\ & & & q & 0 & \\ & & & & & \tilde{Q} \end{bmatrix} \tag{2.13}$$

with

$$\tilde{q}_2 = -q\tilde{q}_1. \tag{2.14}$$

Both solutions satisfy equations (1.8), (1.9) and (2.1).

We emphasize that the general solution possesses six distinct eigenvalues for $j=1$; for example, $q, Q, \frac{1}{2}(q_i \pm (q_i^2 + 4q^{-2}Q)^{1/2}, i=1, 2$. However, the standard one has only three distinct eigenvalues because of the reduction. Obviously they cannot be connected through a similar transformation. This feature holds for any j . The example with $j=\frac{3}{2}$ has been analysed explicitly. We omit this example in this paper.

3. Yang–Baxterization

As was pointed out in [10] the Yang–Baxterization prescription is developed on the basis of Jimbo’s loop extension theory [11], which is based on projectors and corresponding eigenvalues. It is worth noting that Yang–Baxterization for the present solutions cannot simply be done by following either [10] or [11], because in the present case the numbers of distinct eigenvalues and decomposed subspaces are no longer the same. The number of such subspaces is two, resulting in the decomposition

$$j \otimes \frac{1}{2} = j + \frac{1}{2} \oplus j - \frac{1}{2}. \tag{3.1}$$

Our scheme is very simple and is based on direct verification. For $S^{\frac{1}{2}}$ it is well known that

$$\check{R}^{\frac{1}{2}}(x) = x(S^{\frac{1}{2}})^{-1} - S^{\frac{1}{2}}. \tag{3.2}$$

Taking equation (3.1) into account we know Yang–Baxterization for $S^{j\frac{1}{2}}$ takes the form

$$\check{R}^{j\frac{1}{2}}(x) = A_1 x(S^{j\frac{1}{2}})^{-1} + A_2 S^{j\frac{1}{2}} \tag{3.3}$$

$$\check{R}^{j\frac{1}{2}}(x) = A_1 x(S^{j\frac{1}{2}})^{-1} + A_2 S^{j\frac{1}{2}} \tag{3.4}$$

where the CP invariance has been considered. Substituting equations (3.2)–(3.4) into equation (1.11) we find equation (1.11) is satisfied for any spectral parameter x if

$$A_1 = -Qq^{-2j}A_2. \tag{3.5}$$

For convenience we take $A_1 = q$ then

$$A_2 = -q^{2j+1}Q^{-1}. \tag{3.6}$$

In this manner the Yang–Baxterization of the general solutions leads to

$$\begin{aligned} \check{R}^{j\frac{1}{2}}(x) = & \sum_{a(j), b(\frac{1}{2})} p^{(a(j), b(\frac{1}{2}))}(x) E_{a(j), b(\frac{1}{2})} \otimes E_{b(\frac{1}{2}), a(j)} \\ & + \sum_{a(j), b(\frac{1}{2})} q^{(a(j), b(\frac{1}{2}))}(x) E_{a(j), -b(\frac{1}{2})} \otimes E_{b(\frac{1}{2}), a(j)+2b(\frac{1}{2})} \end{aligned} \tag{3.7}$$

where $(E_{a,b})_{cd} = \delta_{ac}\delta_{bd}$ and

$$p^{(a(j), b(\frac{1}{2}))}(x) = \begin{cases} q^{j+1}(xq^{-a(j)} - Q^{-1}q^{a(j)+1}) & \text{for } b(\frac{1}{2}) = \frac{1}{2} \\ q^j(xq^{a(j)} - Q^{-1}q^{-a(j)+1}) & \text{for } b(\frac{1}{2}) = -\frac{1}{2} \end{cases} \tag{3.8}$$

$$q^{(a(j), b(\frac{1}{2}))} = \begin{cases} -q^{2j+1}Q^{-1}q^{(a(j), \frac{1}{2})} & \text{for } b(\frac{1}{2}) = \frac{1}{2} \\ -q^{2j+1}xq^{(-a(j), \frac{1}{2})} & \text{for } b(\frac{1}{2}) = -\frac{1}{2}. \end{cases} \tag{3.9}$$

Equation (3.8) allows the standard solutions as a special case where

$$p^{(a(j), b(\frac{1}{2}))}(x) = \begin{cases} q^{j+1}(xq^{-a(j)} - q^{a(j)}) & \text{for } b(\frac{1}{2}) = \frac{1}{2} \\ q^j(xq^{a(j)} - q^{-a(j)}) & \text{for } b(\frac{1}{2}) = -\frac{1}{2} \end{cases} \tag{3.10}$$

$$q^{(a(j), b(\frac{1}{2}))}(x) = \begin{cases} -q^{j+\frac{1}{2}}w\{[j-a(j)]_q[j+a(j)+1]_q\}^{1/2} & \text{for } b(\frac{1}{2}) = \frac{1}{2} \\ -q^{j+\frac{1}{2}}w\{[j+a(j)]_q[j-a(j)+1]_q\}^{\frac{1}{2}}x & \text{for } b(\frac{1}{2}) = -\frac{1}{2}. \end{cases} \tag{3.11}$$

It is interesting to point out that by taking $x = -1$ the standard solutions of $R^{j\frac{1}{2}}(x)|_{x=-1}$ given by equations (3.10) and (3.11) exactly coincide with the results given in [14, 15]. This particular example ($j = 1$) had been given in [16].

4. Quantum group structure

Following the FRT approach [13] for a given $S^{j\frac{1}{2}}$ the corresponding quantum group can be set up. For convenience the indices $a(j)$ will henceforth be replaced by Greek letters and $b(\frac{1}{2})$ by Latin letters; we then have the solutions

$$(S^{j\frac{1}{2}})_{m\nu}^{\mu l} = p^{(\mu,l)} \delta_{\nu}^{\mu} \delta_m^l + q^{(\mu,l)} \delta_{\nu-2l}^{\mu} \delta_{-m}^l \tag{4.1}$$

$$(S^{j\frac{1}{2}})_{\nu m}^{\mu} = p^{(-\mu,-l)} \delta_{-\nu}^{-\mu} \delta_{-m}^{-l} + q^{(-\mu,-l)} \delta_{-\nu+2l}^{-\mu} \delta_m^{-l} \tag{4.2}$$

$$\mu, \nu = j, j-1, \dots, -j+1, -j \quad l, m = \pm\frac{1}{2} \tag{4.3}$$

where

$$p^{(\mu,\frac{1}{2})} = q^{\mu+1-j} \quad p^{(\mu,-\frac{1}{2})} = q^{-\mu-j} Q \quad q^{(\mu,-\frac{1}{2})} = 0 \tag{4.4}$$

$$q^{(-\mu-\frac{1}{2},\frac{1}{2})} q^{(\mu,\frac{1}{2})} = q^{-2j} Q w^2 [j-\mu]_q [j+\mu+1]_q \quad (\mu \neq \pm j). \tag{4.5}$$

The corresponding FRT relations read

$$\begin{aligned} S^{j\frac{1}{2}}(L_{\pm}^{\frac{1}{2}} \otimes L_{\pm}^j) &= (L_{\pm}^j \otimes L_{\pm}^{\frac{1}{2}}) S^{j\frac{1}{2}} \\ S^{j\frac{1}{2}}(L_{\pm}^j \otimes L_{\pm}^{\frac{1}{2}}) &= (L_{\pm}^{\frac{1}{2}} \otimes L_{\pm}^j) S^{j\frac{1}{2}} \\ S^{j\frac{1}{2}}(L_{+}^{\frac{1}{2}} \otimes L_{-}^j) &= (L_{-}^j \otimes L_{+}^{\frac{1}{2}}) S^{j\frac{1}{2}} \\ S^{j\frac{1}{2}}(L_{+}^j \otimes L_{-}^{\frac{1}{2}}) &= (L_{-}^{\frac{1}{2}} \otimes L_{+}^j) S^{j\frac{1}{2}}. \end{aligned} \tag{4.6}$$

After calculation we find

$$\begin{aligned} (L_{+})_{\nu}^{\mu} &= \alpha_{\nu}^{\mu} K_1^{-2\nu} K_2^{2(\mu-j)} (X^{+})^{\mu-\nu} \\ (L_{-})_{\nu}^{\mu} &= \beta_{\nu}^{\mu} K_1^{2\mu} K_2^{2(\nu-j-2\mu)} (X^{-})^{\nu-\mu} \\ (L_{+})_{\frac{1}{2}}^{\frac{1}{2}} &= K_1^{-1} K_2 \quad (L_{-})_{-\frac{1}{2}}^{-\frac{1}{2}} = K_1 K_2^{-1} \quad (L_{+})_{\frac{1}{2}}^{\frac{1}{2}} = K_1 K_2 X^{+} \\ (L_{-})_{\frac{1}{2}}^{\frac{1}{2}} &= K_1 K_2 \quad (L_{-})_{-\frac{1}{2}}^{-\frac{1}{2}} = K_1^{-1} K_2 \quad (L_{-})_{\frac{1}{2}}^{\frac{1}{2}} = K_1^{-1} K_2^3 X^{-} \\ (L_{+})_{\frac{1}{2}}^{-\frac{1}{2}} &= (L_{-})_{-\frac{1}{2}}^{\frac{1}{2}} = 0 \end{aligned} \tag{4.7}$$

where

$$\begin{aligned} \alpha_{\nu}^{\mu} &= 0 \quad \text{for } \mu < \nu \\ \beta_{\nu}^{\mu} &= 0 \quad \text{for } \mu > \nu \\ \alpha_{\mu}^{\mu} &= \beta_{\mu}^{\mu} = 1 \end{aligned}$$

$$\alpha_{\nu}^{\nu+n} = \prod_{l=1}^n (q^{(-\nu-l,\frac{1}{2})} q^{\nu+j} (Qq^{-1})^{-(l-1)/j}) / (q^l - q^{-l}) \quad (n = 1, 2, \dots, j-\nu) \tag{4.8}$$

$$\beta_{\nu}^{\nu-n} = \prod_{l=1}^n (q^{(\nu-l,\frac{1}{2})} q^{\nu+j+1} Q^{-1} q^{-2l+1} (Qq^{-1})^{(l-1)/j}) / (q^l - q^{-l}) \quad (n = 1, 2, \dots, j+\nu)$$

and the operators satisfy the relations

$$\begin{aligned} [K_1, K_2] &= 0 \quad K, X^{\pm} K_1^{-1} = (Qq^{2j-1})^{\pm 1/2j} X^{\pm} \\ K_2 X^{\pm} K_2^{-1} &= (Qq^{-1})^{\pm 1/2j} X^{\pm} \\ (Qq^{-1})^{-1/j} X^{+} X^{-} - (Qq^{-1})^{1/j} X^{-} X^{+} &= q^{-1} w K_2^{-4} (K_1^{-2} K_2^2 - K_1^2 K_2^{-2}). \end{aligned} \tag{4.9}$$

The coproduct Δ , co-unit ε and antipode γ are given by the following:

$$\begin{aligned}
 \Delta K_1 &= K_1 \otimes K_1 & \Delta K_2 &= K_2 \otimes K_2 \\
 \Delta X^+ &= K_1^{-2} \otimes X^+ + X^+ \otimes K_2^{-2} \\
 \Delta X^- &= X^- \otimes K_1^2 K_2^{-4} + K_2^{-2} \otimes X^- \\
 \varepsilon(K_1) &= \varepsilon(K_2) = 1 & \varepsilon(X^\pm) &= 0 \\
 \gamma(X^+) &= -Q^{-1} q K_1^2 K_2^2 X^+ & \gamma(X^-) &= -Qq^{-3} K_1^{-2} K_2^6 X^- \\
 \delta(K_1) &= K_1^{-1} & \gamma(K_2) &= K_2^{-1}.
 \end{aligned}
 \tag{4.10}$$

Similar to the discussion in [17] the commutation relations can be simplified by introducing

$$\tilde{X}^\pm = X^\pm K_2^2 \quad \mathcal{H} = K_1 K_2^{-1}. \tag{4.11}$$

We then have the usual commutation relations except a centre

$$\begin{aligned}
 K_2 \tilde{X}^\pm K_2^{-1} &= (Qq^{-1})^{\pm 1/2j} \tilde{X}^\pm & [\mathcal{H}, K_2] &= 0 \\
 \mathcal{H} \tilde{X}^\pm \mathcal{H}^{-1} &= q^{\pm 1} \tilde{X}^\pm & & \\
 [\tilde{X}^+, \tilde{X}^-] &= q^{-1} w (\mathcal{H}^{-2} - \mathcal{H}^2).
 \end{aligned}
 \tag{4.12}$$

It is not surprising that the above result has been reached since we cannot go beyond the basic structure of the QG. Actually, this result checks the validity of our solutions. We would like to emphasize that the derivation of such commutation relations does not mean that the QG structure is 'trivial'. In the next section we will point out that the construction of the R -matrix is no longer standard, i.e. it cannot be formed in the usual way in terms of basis, nor can it be related to any similar transformation of the standard basis.

5. Relationship between general and standard solutions

According to the standard construction of the R -matrix [1-3] the centralizer can be represented in terms of the basis of universal enveloping algebra. The representation in this case is

$$\begin{aligned}
 S_{\text{standard}}^{\frac{1}{2}j} &= q^{\rho^{(\frac{1}{2})(J_3)} \otimes \rho^{(j)(J_3)}/2} \sum_{n=0}^{\infty} \frac{(1-q^{-2})^n}{[n]_q!} \\
 &\times q^{(\frac{1}{2})n(n-1)} (q^{\rho^{(\frac{1}{2})(J_3)} \rho^{(\frac{1}{2})(J_+)} \otimes q^{-\rho^{(j)(J_3)} \rho^{(j)(J_-)}})^n
 \end{aligned}
 \tag{5.1}$$

where $\rho^{(j)}$ stands for the representation of $SU_q(2)$ and

$$\begin{aligned}
 \rho^{(j)}(J_\pm)_m^{m'} &= ([j \mp m]_q [j \pm m + 1]_q)^{1/2} \delta_{m+1}^{m'} \\
 \rho^{(j)}(J_3)_m^{m'} &= 2m \delta_m^{m'} \quad m, m' \in (j, j-1, \dots, -j).
 \end{aligned}
 \tag{5.2}$$

Following the idea q -weight conservation in [18] we write our solutions as

$$\begin{aligned}
 S^{\frac{1}{2}j} &= S_{\text{standard}}^{\frac{1}{2}j} + \Delta^{\frac{1}{2}j} \\
 S^{j\frac{1}{2}} &= S_{\text{standard}}^{j\frac{1}{2}} + \Delta^{j\frac{1}{2}} \\
 S^{\frac{1}{2}\frac{1}{2}} &= S_{\text{standard}}^{\frac{1}{2}\frac{1}{2}}.
 \end{aligned}
 \tag{5.3}$$

Let us determine Δ^{lj} and Δ^{jl} by substituting equation (5.3) into equation (1.1). The constraint relations yield

$$S_{12}^{lj}(\Delta_{13}^{lj}S_{23}^{lj} + S_{13}^{lj}\Delta_{23}^{lj} + \Delta_{13}^{lj}\Delta_{23}^{lj}) = (\Delta_{23}^{lj}S_{13}^{lj} + S_{23}^{lj}\Delta_{13}^{lj} + \Delta_{23}^{lj}\Delta_{13}^{lj})S_{12}^{lj} \tag{5.4}$$

and

$$\Delta_{12}^{lj}S_{13}^{lj}S_{23}^{lj} + S_{12}^{lj}S_{13}^{lj}\Delta_{23}^{lj} + \Delta_{12}^{lj}S_{13}^{lj}S_{23}^{lj} = \Delta_{23}^{lj}S_{13}^{lj}S_{12}^{lj} + S_{23}^{lj}S_{13}^{lj}\Delta_{12}^{lj} + \Delta_{23}^{lj}\Delta_{13}^{lj}\Delta_{12}^{lj}. \tag{5.5}$$

After calculation we find that Δ^{lj} and Δ^{jl} should be expressed through the q -generators as

$$\begin{aligned} \Delta^{lj} = & \sum_{k=0}^{2j} D_k \rho^{(l)}(J_- \rho^{(j)}(J_+) \otimes (\rho^{(j)}(J_-))^k (\rho^{(j)}(J_+))^k \\ & + \sum_{k=1}^{2j} B_k \rho^{(l)}(J_+) \otimes (\rho^{(j)}(J_+))^{k-1} (\rho^{(j)}(J_-))^k \\ & + \sum_{k=1}^{2j} C_k \rho^{(l)}(J_-) \otimes (\rho^{(j)}(J_+))^k (\rho^{(j)}(J_-))^{k-1}. \end{aligned} \tag{5.6}$$

Using

$$\begin{aligned} [(\rho^{(j)}(J_-))^k (\rho^{(j)}(J_+))^k]_m^{m'} &= \prod_{l=1}^k [j+m+l]_q [j-m+l+1]_q \delta_m^{m'} \\ [\rho^{(l)}(J_-) \rho^{(l)}(J_+) \otimes (\rho^{(j)}(J_-))^k (\rho^{(j)}(J_+))^k]_{m_1 m_2}^{m'_1 m'_2} & \\ &= \prod_{l=1}^k [j+m_2+l]_q [j-m_2-l+1]_q \delta_{-\frac{1}{2}}^{m'_1} \delta_{m_1}^{-\frac{1}{2}} \delta_{m_2}^{m'_2} \end{aligned} \tag{5.7}$$

we obtain the recurrence relations for the coefficients D_k and B_k ,

$$D_0 + \sum_{k=1}^{2j} D_k \left(\prod_{l=1}^k [j+m+l]_q [j-m-l+1]_q \right) = q^{m-1} (Q-q) \quad (m \neq j) \tag{5.8}$$

$$D_0 = q^{-j-1} (Q-q)$$

and

$$\begin{aligned} ([j+m]_q [j-m+1]_q)^{\frac{1}{2}} & \left[B_1 + \sum_{k=2}^{2j} \left(\prod_{l=2}^k B_k [j+m-l+1]_q [j-m+l]_q \right) \right] \\ & = q_{m-1} - q^{-\frac{1}{2}} w ([j+m]_q [j-m+1]_q)^{\frac{1}{2}} \quad (m \neq -j) \end{aligned} \tag{5.9}$$

as well as a sufficient choice

$$C_k = 0. \tag{5.10}$$

Equations (5.8)–(5.10) can be used to determine Δ^{jl} and Δ^{lj} in terms of the representations of the q -generators J_{\pm} and J_3 satisfying the standard QG, as shown by equation (4.12) (\tilde{X}^{\pm} identifies with J_{\pm} after rescaling).

It should be emphasized that since the general solution cannot be related to the standard one by a similar transformation and by virtue of equation (5.3), in general, our solution cannot be expressed by a new basis which is a similar transformation of the standard basis in the same manner as for the standard case. We have met the same situation for a new solution of S^{11} [17, 18].

If $C_k \neq 0$, the calculation is very complicated, and we give an example only. For equation (2.13) we find

$$\begin{aligned} D_0 &= \tilde{Q} - q^{-1}, D_1 = (1/[2]_q)(q-1)(\tilde{Q} - q^{-1}) \\ D_2 &= (1-q)(\tilde{Q} + 1) \\ B_1 &= (-1/[2]_q)[(1-q^{-2})(q^2 - q^{-2})]^{1/2}, B_2 = (1/[2]_q)\tilde{q}_1 \\ C_1 &= (1/[2]_q)\tilde{q}_2 \quad C_2 = -([2]_q)^{3/2}\tilde{q}_2 \\ \tilde{q}_2 &= -q\tilde{q}_1. \end{aligned}$$

In conclusion we have verified all of the results listed in the introduction.

Acknowledgment

This work was partly supported by NSF of the People's Republic of China.

References

[1] Drinfeld V G 1985 *Dokl. Akad. Nauk. CCCP* **283** 1060; 1986 *Proc. ICM, Berkeley, California, USA* pp 798-820

[2] Reshetikhin N Yu 1988 *Preprint LOMI E-4-1987*

[3] Kirillov A N and Reshetikhin N Yu 1989 *Infinite Dimensional Lie Algebras and Groups* ed V G Kac (Singapore: World Scientific)

[4] Kulish, P P, Reshetikhin N Yu and Sklyanin E K 1981 *Lett. Math. Phys.* **5** 393-403

[5] Jimbo M 1986 *Lett. Math. Phys.* **11** 247
Cherednik I V 1986 *Soviet Math. Dokl.* **33** 507-10

[6] Baxter R J 1982 *Exactly Solved Models in Statistical Mechanics* (London: Academic)

[7] de Vega H J 1990 *Int. J. Mod. Phys. B* **4** 735

[8] Lee H C, Couture M and Scheing N 1988 *Preprint* Chalk River, Canada

[9] Ge M L and Xue K 1990 *Phys. Lett.* **146A** 245
Cheng Y, Couture M, Ge M L and Xue K 1991 *Int. J. Mod. Phys. A* **6** 559
Ge M L and Xue K 1991 *J. Phys. A: Math. Gen.* **24** L895; 1991 *J. Math. Phys.* **32** 1301

[10] Ge M L, Wu Y S and Xue K 1991 *Int. J. Mod. Phys. A* **6** 3735

[11] We would like to emphasize that so far for $V^{1/2} \otimes V^{1/2}$ the general Yang-Baxterization has not been set up satisfactorily. In [10] an extension of Jimbo's theory (see [12]) is given which works only for $j_1 = j_2$. If the fusion rule does not work as in the present case one should rediscuss the problem

[12] Jimbo M 1985 *Lett. Math. Phys.* **10** 63

[13] Faddeev, L D, Reshetikhin N Yu and Takhtajan, A L 1989 *Algebra Analysis* **1** 178
Takhtajan L 1990 *Introduction to Quantum Group and Integrable Massive Models of Quantum Field Theory* ed Ge M L and Zhao B H (Singapore: World Scientific) pp 69-197

[14] Curtright and Zachos C 1989 *Preprint ANL-HEP-PR 89-105*

[15] Curtright T, Ghandour G I and Zachos C 1990 *Preprint ANL-HEP-PR-90-08*

[16] Ge M L, Li W and Xue K 1990 Yang-Baxterization and new solutions associated with $SU_q(2)$ tensorial Space $V^1 \otimes V^{1/2}$ *Preprint Nankai*

[17] Ge M L and Wu A C T 1991 Quantum groups constructed from the non-standard braid group representations in the Faddeev-Reshetikhin-Takhtajan approach *J. Phys. A: Math. Gen.* **24** L725 (where the separation like equation (4.12) for new solutions of A_n was firstly made)

[18] Ge M L, Sun C P and Xue K, 1990 *J. Phys. A: Math. Gen.* **23** L645
The general discussion on the application of weight conservation to find solutions of braid relations can be found in the following references:
Ge M L, Li Y Q and Xue K 1990 *J. Phys. A: Math. Gen.* **23** 605, 619
Ge M L, Gwa L H, Piao F and Xue K 1990 *J. Phys. A: Math. Gen.* **23** 2273
Ge M L, Wang L Y, Xue K and Wu Y S 1989 *Int. J. Mod. Phys. A* **4** 3351; 1990 *Int. J. Mod. Phys. A* **5** 1975
Ge M L, Wang L Y and Kong X P 1991 *J. Phys. A: Math. Gen.* **24** 569