

Some remarks on the visible points of a lattice

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1994 J. Phys. A: Math. Gen. 27 5041

(<http://iopscience.iop.org/0305-4470/27/14/030>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.68

The article was downloaded on 01/06/2010 at 23:20

Please note that [terms and conditions apply](#).

Corrigenda

Some remarks on the visible points of a lattice

Baake M, Grimm U and Warrington D H 1994 *J. Phys. A: Math. Gen.* 27 2669–74

In the stage of proof reading, we have apparently overlooked that equation (3) is missing. After equation (2), the last paragraph on the first page of the article should continue as follows:

This is because $\ell \cdot F_\Lambda = \{m_1 b_1 + \dots + m_n b_n | \gcd(m_1, \dots, m_n) = \ell\}$. On the other hand, we have

$$\Lambda = \{0\} \cup \bigcup_{\ell=1}^{\infty} \ell \cdot F_\Lambda \tag{3}$$

which is the correct version of equation (1) in [2]. Note that (3) is the union of pairwise disjoint sets.

A new finite-size scaling approach to random walks

Giacometti A and Nakanishi H 1994 *J. Phys. A: Math. Gen.* 27 2277–88

By mistake, instead of figure 7, figure 5 was printed again. The correct figure is reported below.




Figure 1. Computed phase diagram for the surface–bulk problem for the random walk on a 2×2 cell for the diluted case for bond ($p = 0.5$). The rule chosen was the *myopic ant* described in the text with periodic boundary conditions in the $\hat{2}$ direction (parallel to the surface), and open boundary conditions in the $\hat{1}$ direction, where only those walks which span were averaged. The symbols (Δ) and (\square) correspond to the bulk and surface singularities as explained in the text.