

Numerical diagonalization study of the trimer deposition-evaporation model in one dimension

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1994 J. Phys. A: Math. Gen. 27 L831

(<http://iopscience.iop.org/0305-4470/27/21/008>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.68

The article was downloaded on 01/06/2010 at 22:57

Please note that [terms and conditions apply](#).

LETTER TO THE EDITOR

Numerical diagonalization study of the trimer deposition–evaporation model in one dimension

Peter B Thomas†‡, M K Hari Menon§ and Deepak Dhar||

Theoretical Physics Group, Tata Institute of Fundamental Research, Homi Bhabha Road, Bombay 400005, India

Received 21 June 1994

Abstract. We study the model of deposition–evaporation of trimers on a line recently introduced by Barma *et al.* The stochastic matrix of the model can be written in the form of the Hamiltonian of a quantum spin- $\frac{1}{2}$ chain with three-spin couplings given by $H = \sum_i [(1 - \sigma_i^- \sigma_{i+1}^- \sigma_{i+2}^-) \sigma_i^+ \sigma_{i+1}^+ \sigma_{i+2}^+ + \text{hc}]$. We study, by exact numerical diagonalization of H , the variation of the gap in the eigenvalue spectrum with the system size for rings of size up to 30. For the sector corresponding to the initial condition in which all sites are empty, we find that the gap vanishes as L^{-z} where the gap exponent z is approximately 2.55 ± 0.15 . This model is equivalent to an interfacial roughening model where the dynamical variables at each site are matrices. From our estimate for the gap exponent we conclude that the model belongs to a new universality class, distinct from that studied by Kardar *et al.*

Many physical processes such as heterogeneous catalysis, chemical reactions on polymer chains, adsorption on solid surfaces, etc involve evaporation and deposition of reactants on a substrate. Recently Barma *et al* have introduced a simple model which shows that the excluded volume effect together with dissociation and re-combination of the reactants on the surface can give rise to very interesting dynamical behaviour. In their model they have studied a random deposition–evaporation process of k identical atoms (called k -mers, $k = 1, 2, 3 \dots$) on a surface [1, 2]. It has been shown that in one-dimension when $k \geq 3$ the phase space breaks up into an exponentially large number of dynamically disconnected sectors and the model has an infinite number of conserved quantities [3]. It is found that in this case the auto-correlation function in the steady state decays with time t as $t^{-1/4}$, $t^{-1/2}$, $t^{-0.6}$ or as $e^{-\sqrt{t}}$, depending on the initial condition. The behaviour of the auto-correlation function for different initial conditions is understood in terms of the random walk of the substrings which constitutes what is called an irreducible string [4, 5]. However, for the steady state corresponding to the empty configuration as the initial condition, this analysis does not apply. In this case, for the trimer model, Monte Carlo simulations show power-law decay of the autocorrelation function with an approximate value for the exponent 0.6 [5]. A theoretical understanding of this exponent is still lacking. Thus our main motivation is to understand the dynamics of trimer model in this sector. We have done a study of the

† Present address: Department of Physics, Brookhaven National Laboratory, Upton, NY 11973, USA.

‡ E-mail address: thomas@cmth.phy.bnl.gov

§ E-mail address: hari@theory.tifr.res.in

|| E-mail address: ddhar@theory.tifr.res.in

trimer model on a one-dimensional lattice, by exact diagonalization of the stochastic matrix in this sector.

In this letter we restrict ourselves to the study of trimers ($k = 3$) on a line ($d = 1$). We consider a ring of L sites. At each site i is a dynamical variable n_i which takes the values 0 or 1, depending on whether the site is occupied or not. A configuration $\{n_i\}$ evolves stochastically in time by Markovian dynamics as follows: any three adjacent empty sites can become occupied with a rate ϵ and any three adjacent occupied sites can become empty with a rate ϵ' .

If $P(C, t)$ is the probability that the ring has configuration C at time t , then $P(C, t)$ satisfies the master equation

$$\frac{\partial}{\partial t} P(C, t) = \sum_{C'} W_{CC'} P(C', t) \quad (1)$$

where the transition rate matrix \hat{W} for the case $\epsilon = \epsilon'$ can be written as

$$\hat{W} = \epsilon \sum_{i=1}^L [(1 - \sigma_i^- \sigma_{i+1}^- \sigma_{i+2}^-) \sigma_i^+ \sigma_{i+1}^+ \sigma_{i+2}^+ + \text{hc}] \quad (2)$$

where σ_i^- and σ_i^+ are the Pauli annihilation and creation operators at site i .

Since \hat{W} is a stochastic matrix where the transition rates satisfy detailed balance, all its eigenvalues are real and non-positive. The infinite number of conservation laws of this Hamiltonian can be encoded into a single conservation law of the irreducible string [3]. For any configuration the irreducible string is defined as follows: from the L -bit string of 0s and 1s representing the configuration, we recursively delete any consecutive occurrence of three 0s or 1s until no further deletions are possible. The irreducible string is conserved under dynamics and can be used to label uniquely each of the dynamically disconnected sectors. There is a large degeneracy for the eigenvalue 0, reflecting the large number of conservation laws in the model. An example of an eigenvector with zero eigenvalue is any configuration which has no three adjacent 0s or 1s. Such a state cannot evolve in time. The number of such configurations has been shown to vary as μ^L for large L , where μ is the golden mean $(\sqrt{5} + 1)/2$ [1, 2].

We can exactly diagonalize \hat{W} in some almost totally jammed sectors. For example, if the sector corresponds to an irreducible string of length $L - 3$, then it is easy to see that the corresponding stochastic matrix, in general, has a size of $O(L^2)$. Under dynamics the position of the reducible block on the ring changes and its motion can be described as a random walk [1, 2]. In this case it can be shown that the mean square displacement increases linearly with time. This corresponds to a dynamical exponent of $z = 2$. Sectors with irreducible string length $L - 6$ correspond to diffusion of two interacting random walkers. In this case the size of the stochastic matrix will be of $O(L^3)$. When the two walkers are next to each other, they stay there longer, which corresponds to an attractive interaction. The dynamical exponent will also be 2 in this case.

The most interesting sector corresponds to the case when the length of the irreducible string (l) is very small compared to L . In this case Monte Carlo simulations [5] have shown that the attractive interaction between these 'random walkers' gives rise to a sub-diffusive behaviour, with the dynamical exponent $z > 2$. In this paper, we estimate this exponent by diagonalizing numerically the stochastic matrix for small systems and assuming finite-size scaling.

For numerical diagonalization it is desirable to reduce the size of the matrix as much as possible by making use of the known symmetries and conservation laws of the model. For periodic boundary conditions, and for the special case of deposition and evaporation

rates equal ($\epsilon = \epsilon'$), in addition to the conservation law of the irreducible string, one can make use of the three symmetries of the system namely translation, reflection and flip, to reduce the size of the matrix by about a factor of $2L$. Let \hat{T} , \hat{P} and \hat{F} be the operators corresponding to these symmetries. They are defined by

$$\begin{aligned}\hat{T}|n_1, n_2, \dots, n_i, \dots, n_L\rangle &= |n_2, n_3, \dots, n_{i+1}, \dots, n_L, n_1\rangle \\ \hat{P}|n_1, n_2, \dots, n_i, \dots, n_L\rangle &= |n_L, n_{L-1}, \dots, n_i, \dots, n_1\rangle \\ \hat{F}|n_1, n_2, \dots, n_i, \dots, n_L\rangle &= |\bar{n}_1, \bar{n}_2, \dots, \bar{n}_i, \dots, \bar{n}_L\rangle \quad \text{where } \bar{n}_i = 1 - n_i.\end{aligned}\tag{3}$$

Here $|n_1, n_2, \dots, n_i, \dots, n_L\rangle$ is a vector in the Hilbert space representing the configuration $\{n_i\}$. These operators satisfy the following algebra:

$$[\hat{T}, \hat{F}] = [\hat{P}, \hat{F}] = 0 \quad \hat{T}^L = \hat{P}^2 = \hat{F}^2 = 1 \quad \hat{T}\hat{P} = \hat{P}\hat{T}^{-1}.\tag{4}$$

Note that \hat{T} and \hat{P} do not commute. The three operators which simultaneously commute with \hat{W} and with each other are \hat{F} , \hat{P} and $(\hat{T} + \hat{T}^{-1})$. Let their corresponding eigenvalues be f , p and $2\cos(k)$, respectively, where $f = \pm 1$, $p = \pm 1$ and $k = 2n\pi/L$; $n = 0, 1, \dots, L-1$. The simultaneous eigenvectors of these three operators are of the form

$$\begin{aligned}|k, f, p, +\rangle &= (1 + f\hat{F})(1 + p\hat{P}) \sum_{r=1}^L T^r \cos(kr)|C\rangle \\ |k, f, p, -\rangle &= (1 + f\hat{F})(1 + p\hat{P}) \sum_{r=1}^L T^r \sin(kr)|C\rangle\end{aligned}\tag{5}$$

where $|C\rangle$ is any of the vectors $|\{n_i\}\rangle$.

We have used the states (5) as the basis for the stochastic matrix. For the null sector, the matrix splits into $2L$ blocks, corresponding to combinations of the two eigenvalues of \hat{F} and the L eigenvalues of \hat{T} . Of these, due to a Kramers-type degeneracy in the eigenvalues for the momentum values k and $2\pi - k$, we can fix p to always be equal to unity, and sweep over only half of the allowed momentum values. For lattice lengths which are not multiples of three, there is an additional degeneracy in the eigenvalues for $f = 1$ and -1 , since these states and their flipped counterparts are not connected by the dynamics. Since the size of the null sector $\sim (27/4)^{L/3} L^{-3/2}$ [3], the size of each block $\sim (27/4)^{L/3} L^{-5/2}$. For any lattice length, each block of the matrix is real and sparse, since all rows or columns have at most L non-zero entries.

The difference between the largest and the second largest eigenvalue of the complete matrix is proportional to the inverse relaxation time. The largest eigenvalue is zero and it lies in the block $k = 0$, $f = 1$. To find the second largest eigenvalue of the full matrix, we have computed numerically the largest eigenvalue in all the other blocks, and the second-largest eigenvalue in the $k = 0$, $f = 1$ block. Simple iteration of the eigenvector after suitably shifting all the eigenvalues, converged sufficiently fast for these blocks. This method preserves the sparseness of the blocks, which is necessary to keep the memory requirement of the program as low as possible. For the $k = 0$, $f = 1$ block, we computed the second largest eigenvalue, by ensuring orthogonality of the iterated vector to the eigenvector corresponding to the zero eigenvalue.

We have computed these eigenvalues for lattice sizes ranging from $L = 3-30$. When L is a multiple of 3, the irreducible string in the null sector has length zero and in this case we have diagonalized the stochastic matrix for both the $f = 1$ and -1 case. For the case $f = -1$ the smallest eigenvalue occurs for $k = 2\pi(1 - 1/L)/3$, and for the case $f = 1$ it occurs for $k = 2\pi/3$. When $L = 3n + 1$ and $3n + 2$, where n is an integer, the irreducible




Figure 1. A plot of the effective exponent z_L versus $1/L$, where L is the length of the lattice.

Table 1.

| Length of the lattice | $f = -1$ | | |
|-----------------------|-------------|------------------|----------|
| | Matrix size | λ_{\min} | z_L |
| 3 | 1 | -6.000 00 | |
| 6 | 1 | -2.000 00 | 1.584 92 |
| 9 | 2 | -0.871 13 | 2.049 77 |
| 12 | 10 | -0.438 76 | 2.384 00 |
| 15 | 35 | -0.260 65 | 2.333 75 |
| 18 | 170 | -0.169 32 | 2.366 07 |
| 21 | 815 | -0.117 44 | 2.373 73 |
| 24 | 4 176 | -0.085 45 | 2.379 29 |
| 27 | 21 872 | -0.064 55 | 2.383 33 |
| 30 | 118 175 | -0.050 20 | 2.386 72 |

string in the sector where the initial state is all empty has length 1 and 2, respectively. In this case, as explained earlier, the eigenvalues for $f = 1$ and -1 are degenerate. We have estimated the gap exponent z for each of these four sets of data, by assuming the scaling relation $\lambda \sim L^{-z}$. We define the effective exponent

$$z_L = \frac{\log[\lambda_{L-3}/\lambda_L]}{\log[L/(L-3)]} \tag{6}$$

The sizes of the matrices, eigenvalues and estimate of the dynamical exponent z_L are shown in the tables below. The z_L values are also plotted as a function of $1/L$ in figure 1.

It is clear from an inspection of these tables that while the convergence in each sector is reasonably good, there is a large difference between them if we compare them between different sectors. To see if this could be due to the presence of a correction to the asymptotic scaling form, we have tried to incorporate various forms for the correction into the scaling. But none of these fit the data well, and at the same time decrease the discrepancy in z between different sectors. This can be seen from the fact that the effective values of z_L do not show a significant tendency to converge to a single value as L increases, for the largest sizes reached in our study. This is somewhat surprising as in previous studies very good convergence has been found for much smaller L (~ 15) [8, 9].

One possible explanation for this is that different sectors have different gap exponents.

Table 2.

| Length of the lattice | $f = -1$ | | |
|--------------------------|-------------|------------------|----------|
| | Matrix size | λ_{\min} | z_L |
| 6 | 1 | -2.000 00 | |
| 9 | 2 | -1.255 53 | 1.148 28 |
| 12 | 10 | -0.674 12 | 2.161 80 |
| 15 | 35 | -0.442 17 | 1.883 75 |
| 18 | 173 | -0.295 77 | 2.213 07 |
| 21 | 811 | -0.208 03 | 2.282 76 |
| 24 | 4 186 | -0.152 13 | 2.343 60 |
| 27 | 21 874 | -0.114 85 | 2.386 37 |
| 30 | 118 175 | -0.089 03 | 2.417 25 |

Table 3.

| Length of the lattice | Matrix size | λ_{\min} | z_L |
|--------------------------|-------------|------------------|----------|
| 4 | 1 | -1.000 00 | |
| 7 | 4 | -0.230 08 | 2.625 57 |
| 10 | 17 | -0.092 77 | 2.546 55 |
| 13 | 84 | -0.047 54 | 2.546 25 |
| 16 | 428 | -0.028 02 | 2.548 29 |
| 19 | 2 305 | -0.018 06 | 2.555 71 |
| 22 | 12 744 | -0.012 40 | 2.564 72 |
| 25 | 72 311 | -0.008 92 | 2.574 33 |

Table 4.

| Length of the lattice | Matrix size | λ_{\min} | z_L |
|--------------------------|-------------|------------------|----------|
| 5 | 1 | -1.000 00 | |
| 8 | 4 | -0.284 76 | 2.672 55 |
| 11 | 21 | -0.119 43 | 2.728 55 |
| 14 | 103 | -0.062 15 | 2.708 57 |
| 17 | 553 | -0.036 78 | 2.701 22 |
| 20 | 3 014 | -0.023 72 | 2.698 57 |
| 23 | 16 985 | -0.016 27 | 2.699 33 |
| 26 | 97 419 | -0.011 68 | 2.701 93 |

This is quite intriguing, but somewhat unlikely. The possible reason behind this could be the existence of an infinite number of conserved quantities in the model. It is hoped that further studies will clarify this point.

However, from our data it can be concluded that the gap exponent for all these sectors fall within the range $z = 2.55 \pm 0.15$. To get a more precise estimate for z one needs further study either of larger size lattices, or by Monte Carlo simulations, or analytical methods.

In figure 2 we have shown a plot of λ versus k (dispersion curve) for three different lattice sizes. This is related to the spectrum of the excitations of the quantum Hamiltonian \hat{W} . It is seen that the spectrum for different sizes is qualitatively similar, but shows a complicated, yet not understood structure as a function of k . We have also studied the




Figure 2. Dispersion curve of the quantum Hamiltonian corresponding to the trimer model (equation (2)).

same model for the case of unequal deposition–evaporation rates (in this case there is no flip symmetry). The range of estimated value of z is the same as that for equal deposition–evaporation rates.

The stochastic evolution of the trimer model can be mapped to the stochastic dynamics of a string, both ends fixed to the same point, by defining a matrix variable U_i at each site [3]. This matrix U_i has information about the length of the irreducible string corresponding to the substring from site 1 of the lattice up to site i . Under the dynamics the length of this irreducible string changes, and is related to the change in the matrix variables U_i . Thus this model corresponds to a generalization of the KPZ model where the scalar height variables are replaced by matrix variables. It is well known that $z = 1.5$ for the KPZ model [6]. Our results show that this model falls under a new universality class. It is also different from the model studied recently by Doherty *et al* which is also a generalization of the KPZ equation to n component variables. In their model, the dynamical exponent $z = \frac{3}{2}$ in one dimension, independent of n , though in higher dimensions it depends on n [7].

References

- [1] Barma M, Grynberg M D and Stinchcombe R B 1993 *Phys. Rev. Lett.* **70** 1033
- [2] Stinchcombe R B, Grynberg M D and Barma M 1993 *Phys. Rev. E* **47** 4018
- [3] Dhar D and Barma M 1993 *Pramana* **41** L193
- [4] Barma M and Dhar D 1993 *Proc. Solid State Physics Symp. C* **36** 489
- [5] Barma M and Dhar D 1994 *Phys. Rev. Lett.* at press
- [6] Kardar M, Parisi G and Zhang Y C 1986 *Phys. Rev. Lett.* **56** 889
- [7] Doherty J P, Moore M A, Kim J M and Bray A J 1994 *Phys. Rev. Lett.* **72** 2041
- [8] Kinzel W and Yeomans J M 1981 *J. Phys. A: Math. Gen.* **14** L163
- [9] Uzelac K and Jullien R 1990 *J. Phys. A: Math. Gen.* **23** 4525