

Lie group weight multiplicities from conformal field theory

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1995 J. Phys. A: Math. Gen. 28 2617

(<http://iopscience.iop.org/0305-4470/28/9/020>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.68

The article was downloaded on 02/06/2010 at 01:41

Please note that [terms and conditions apply](#).

Lie group weight multiplicities from conformal field theory

T Gannon†§, C Jakovljević† and M A Walton‡||

† Institut des Hautes Etudes Scientifiques, 91440 Bures-sur-Yvette, France

‡ Physics Department, University of Lethbridge, Lethbridge, Alberta, Canada T1K 3M4

Received 4 November 1994

Abstract. Dominant-weight multiplicities of simple Lie groups are expressed in terms of the modular matrices of Wess–Zumino–Witten conformal field theories, and related objects. Symmetries of the modular matrices give rise to new relations among multiplicities. At least for some Lie groups, these new relations are strong enough to completely fix all multiplicities.

1. Introduction

Wess–Zumino–Witten (WZW) models form an important class of conformal field theories (see [1], for example). They realize a current algebra equivalent to an affine Kac–Moody algebra. Included as a subalgebra is a semi-simple Lie algebra, the global symmetry algebra of the theory. Not surprisingly then, the theory of compact Lie groups and their Lie algebras has been very useful in elucidating the properties of WZW models.

Here we present an example where the reverse is true: WZW conformal field theories tell us something useful about Lie groups and their algebras. Weight multiplicities are crucial numbers in the representation theory of Lie groups (see [2, 3], for example). We will express the dominant-weight multiplicities of unitary highest-weight representations of Lie groups in terms of matrices relevant to WZW models. We also show that our expressions give rise to new relations between the multiplicities.

In order to explain our results, we must first discuss the connection between modular transformations and the Lie symmetry algebra of a WZW model. Primary fields of WZW models are in one-to-one correspondence with unitary highest-weight representations of affine Kac–Moody algebras \hat{g} [4, 5]. The partition function of a WZW model on a torus is a sesquilinear combination of characters of affine Kac–Moody representations [5]. These characters transform among themselves under the action of the modular group of the torus [6]. Remarkably, the ratios of elements of the modular S -matrix equal characters of the Lie symmetry algebra g evaluated at special points (see equation (3.5) below). This happy fact, discovered by Kac and Peterson [6], is the starting point of our work, and has been much exploited elsewhere ([7–9], for example).

Perhaps the relation between modular matrices and semi-simple Lie algebra characters is not so surprising, if other facts are taken into account. Consider the number of couplings between three primary fields of a WZW model, the so-called fusion coefficient. Since a WZW model has a Lie symmetry algebra, this fusion coefficient is less than or equal to the corresponding tensor product coefficient of the Lie algebra [5, 10]. Products of Lie

§ E-mail: gannon@ihes.fr

|| Supported in part by NSERC (e-mail: walton@hg.uleth.ca).

algebra characters decompose into integer linear combinations of characters, according to these coefficients. Verlinde realized that the modular transformation matrix S of a conformal field theory could tell us the fusion coefficients: products of certain ratios of elements of the modular matrix S decompose into integer linear combinations of these same ratios, with the coefficients being the fusion coefficients [11]. It is reasonable that the only way such ratios could satisfy these properties is for them to coincide with the Lie algebra characters, but evaluated at special points.

The outline of this paper is as follows. In section 2 the work of Patera and Sharp [12] is reviewed, so that it can be adapted to the use of Kac–Peterson modular matrices in section 3. There we write a new expression for the dominant-weight multiplicities of semi-simple Lie algebras. The symmetries of the Kac–Peterson modular S -matrix, and the even Weyl sums $E^{(n)}$ we introduce in section 3, are written down in section 4. The relations between the multiplicities that follow are also given. Section 5 contains some simple explicit examples of the new relations between multiplicities, and section 6 is our conclusion.

2. Lie group multiplicities from Lie characters

Define the Weyl orbit sum

$$E_\lambda(\sigma) := \frac{|W(\lambda - \rho)|}{|W|} \sum_{w \in W} \sigma^{w(\lambda - \rho)}. \tag{2.1}$$

Here E stands for an even Weyl sum, $|W|$ is the order of the Weyl group of g , and $|W\mu|$ is the order of the Weyl orbit of $\mu \in P_+ := \{\sum_{i=0}^r \mu_i \omega^i \mid \mu_i \in \mathbb{Z}_{\geq 0}\}$. In equation (2.1) $\lambda \in P_{++} := \{\sum_{i=0}^r \lambda_i \omega^i \mid \lambda_i \in \mathbb{Z}_{> 0}\}$, and ρ is the half-sum of positive roots: $\rho = \sum_{\alpha > 0} \alpha / 2 = \sum_{i=1}^r \omega^i$, where the ω^i are the fundamental weights of g . The somewhat abusive notation is $\sigma^\mu = \exp[-i\mu \cdot \sigma]$, with σ any weight, so that $\sigma^\lambda \sigma^\mu = \sigma^{\lambda + \mu}$.

An odd Weyl sum is the so-called discriminant

$$O_\lambda(\sigma) := \sum_{w \in W} \det(w) \sigma^{w\lambda} \tag{2.2}$$

where $\det(w)$ is the sign of the Weyl group element w . Let H^j be the elements of the Cartan subalgebra of g . Weyl’s famous character formula for the trace $\chi_\lambda(\sigma)$ of $\exp[-i \sum_{j=1}^r H^j \sigma_j]$ in the representation of highest-weight $\lambda - \rho$ is

$$\chi_\lambda(\sigma) = O_\lambda(\sigma) / O_\rho(\sigma). \tag{2.3}$$

The character, being an even Weyl function, can be expanded in terms of the even functions E_μ [13]:

$$\chi_\lambda(\sigma) = \sum_{\mu \in P_{++}} m_\lambda^\mu E_\mu(\sigma). \tag{2.4}$$

The non-negative integers m_λ^μ are the dominant-weight multiplicities: m_λ^μ denotes the multiplicity of the weight $\mu - \rho$ in the representation with highest-weight $\lambda - \rho$. Note that $m_\lambda^\lambda = 1$ for all $\lambda \in P_{++}$. We can consider the m_λ^μ to be the elements of an infinite matrix M . If $\lambda - \mu \notin \mathbb{Z}_{\geq 0} \{\alpha_1, \dots, \alpha_r\}$, where α_i are the simple roots, then $m_\lambda^\mu = 0$. Therefore M is a lower triangular matrix, provided the weights are ordered appropriately.

Not only can the Weyl character be expanded in terms of the Weyl orbit sums E_λ , but the reverse is also true:

$$E_\lambda(\sigma) = \sum_{\mu \in P_{++}} \ell_\lambda^\mu \chi_\mu(\sigma). \tag{2.5}$$

The coefficients ℓ_λ^μ are easily shown to be integers, but are in general *not* non-negative. However if L is the matrix with elements ℓ_λ^μ , then clearly $M = L^{-1}$. So, if the triangular matrix L can be calculated, it is a simple matter to invert it to obtain the dominant-weight multiplicities [12].

Patera and Sharp [12] also point out that the equations above allow the calculation of the ℓ_λ^μ for fixed λ using the Weyl group. This corresponds to the following formula:

$$\ell_\lambda^\mu = \frac{|W(\lambda - \rho)|}{|W|} \sum_{w, x \in W} \det(w) \delta_{w\rho + x(\lambda - \rho)}^\mu \tag{2.6}$$

that can be derived from the defining relation for the ℓ .

3. Lie group multiplicities from wzw modular matrices

Define

$$O_\lambda^{(n)}(\sigma) := F_n \sum_{w \in W} \det(w) \sigma_{(n)}^{w\lambda} = S_{\lambda\sigma}^{(n)} \tag{3.1}$$

and

$$E_\lambda^{(n)}(\sigma) := \frac{|W(\lambda - \rho)|}{|W|} \sum_{w \in W} \sigma_{(n)}^{w(\lambda - \rho)} \tag{3.2}$$

with

$$\sigma_{(n)}^\mu := \exp[-2\pi i \mu \cdot \sigma / n] \quad F_n := \frac{i^{|\Delta_+|}}{n^{r/2} \sqrt{|M^*/M|}} \tag{3.3}$$

where $|\Delta_+|$ is the number of positive roots of g , and M here is the weight lattice. The matrix $S^{(n)}$ in (3.1) is the Kac–Peterson modular matrix of WZW models, corresponding to the affine algebra \hat{g} at level $k = n - h^\vee$, where h^\vee is the dual Coxeter number of g . Define

$$P_+^n := \left\{ \sum_{i=0}^r \lambda_i \omega^i \mid \lambda_i \in \mathbb{Z}_{\geq}, \sum_{i=0}^r a_i^\vee \lambda_i = n \right\} \tag{3.4}$$

and P_{++}^n similarly, except with \mathbb{Z}_{\geq} replaced with $\mathbb{Z}_{>}$. The positive integers a_i^\vee in (3.4) are the *co-marks* [14] of \hat{g} . We read from (3.1) that

$$S_{\lambda\sigma}^{(n)} / S_{\rho\sigma}^{(n)} = O_\lambda^{(n)}(\sigma) / O_\rho^{(n)}(\sigma) =: \chi_\lambda^{(n)}(\sigma) \tag{3.5}$$

where

$$\chi_\lambda^{(n)}(\sigma) = \chi_\lambda(2\pi \sigma / n). \tag{3.6}$$

This is the ‘happy fact’ referred to in the introduction. M and L are lower triangular; therefore whenever $\lambda \in P_{++}^n$, this remarkable relation implies

$$\chi_\lambda^{(n)}(\sigma) = \sum_{\mu \in P_{++}^n} m_\lambda^\mu E_\mu^{(n)}(\sigma) \tag{3.7}$$

and

$$E_\lambda^{(n)}(\sigma) = \sum_{\mu \in P_{++}^n} \ell_\lambda^\mu \chi_\mu^{(n)}(\sigma). \tag{3.8}$$

Equation (3.8) also holds for $\lambda \in P_{++} \cap P_+^n$. Using the unitarity of the modular S -matrix

$$\sum_{\nu \in P_{++}^n} O_\lambda^{(n)}(\nu) O_\mu^{(n)*}(\nu) = \sum_{\nu \in P_{++}^n} S_{\lambda\nu}^{(n)} S_{\nu\mu}^{(n)*} = \delta_\lambda^\mu \tag{3.9}$$

we arrive at

$$\ell_\lambda^\mu = \sum_{\sigma \in P_{++}^n} E_\lambda^{(n)}(\sigma) O_\rho^{(n)}(\sigma) O_\mu^{(n)*}(\sigma) = \sum_{\sigma \in P_{++}^n} E_\lambda^{(n)}(\sigma) S_{\rho\sigma}^{(n)} S_{\mu\sigma}^{(n)*} \tag{3.10}$$

valid whenever both $\lambda \in P_{++} \cap P_+^n$ and $\mu \in P_{++}^n$. Equation (3.10) can be generalized to

$$\sum_{\sigma \in P_{++}^n} \ell_\lambda^\sigma N_{\mu\sigma}^{(n)\nu} = \sum_{\sigma \in P_{++}^n} E_\lambda^{(n)}(\sigma) S_{\mu\sigma}^{(n)} S_{\nu\sigma}^{(n)*} \tag{3.11}$$

where $N_{\mu\sigma}^{(n)\nu}$ are the WZW fusion rules, which we may take to be defined by Verlinde’s formula [11]:

$$N_{\lambda\mu}^{(n)\nu} = \sum_{\sigma \in P_{++}^n} \chi_\lambda^{(n)}(\sigma) S_{\mu\sigma}^{(n)} S_{\nu\sigma}^{(n)*}. \tag{3.12}$$

Of course we also get directly from (3.7) that

$$N_{\lambda\mu}^{(n)\nu} = \sum_{\sigma, \gamma \in P_{++}^n} m_\lambda^\gamma E_\gamma^{(n)}(\sigma) S_{\mu\sigma}^{(n)} S_{\nu\sigma}^{(n)*}. \tag{3.13}$$

Because L is lower triangular, a simple argument gives

$$m_\lambda^\mu = (L^{(n)-1})_\lambda^\mu \tag{3.14}$$

for all $\lambda, \mu \in P_{++}^n$, where $L^{(n)}$ is defined to be the sublattice of L obtained by restricting it to the set P_{++}^n . Thus equations like (3.10) provide a simple method of calculating dominant-weight multiplicities m_λ^μ . Moreover, if we find a permutation π of P_{++}^n which commutes with both $S^{(n)}$ and $E^{(n)}$, then it will be an exact symmetry of both ℓ and m . More generally, if $S^{(n)}$ and $E^{(n)}$ both transform ‘nicely’ under a permutation π of P_{++}^n , then we can expect to derive new relations for ℓ and m . This is the motivation for the following section.

4. New relations between multiplicities

In this section, we will show that symmetries of the Kac–Peterson modular matrices $S^{(n)}$ of WZW models give rise to new relations between finite-dimensional Lie algebra multiplicities.

The most obvious symmetry concerns the affine Weyl group \widehat{W} of \widehat{g} . We know [14] that it is isomorphic to the semi-direct product of the (finite) Weyl group W with the group of translations in the co-root lattice Q^\vee . We also know that the \widehat{W} -orbit of any weight intersects P_+^n in exactly one point. More precisely, let $\lambda \in M$ be some weight. Then an element α exists in the co-root lattice of g , and some $w \in W$, such that

$$[\lambda] := w(\lambda + n\alpha) \in P_+^n. \tag{4.1}$$

We will use this observation throughout this section. $[\lambda]$ is uniquely determined by λ (and n), but w will be only if $[\lambda] \in P_{++}^n$. Define $\epsilon(\lambda) := 0$ if $[\lambda] \notin P_{++}^n$, and $\epsilon(\lambda) := \det(w)$ otherwise, where $w \in W$ satisfies (4.1).

We read directly from (3.1) and (3.2), respectively, that

$$S_{\lambda\mu}^{(n)} = \epsilon(\lambda) S_{[\lambda]\mu}^{(n)} = \epsilon(\mu) S_{\lambda[\mu]}^{(n)} \tag{4.2}$$

$$E_\lambda^{(n)}(\sigma) = E_\lambda^{(n)}([\sigma]) = \frac{|W(\lambda - \rho)|}{|W([\lambda - \rho])|} E_{[\lambda - \rho] + \rho}^{(n)}(\sigma). \tag{4.3}$$

By the argument which gave us equation (3.10), we find that for any $\lambda \in P_{++}^n$, $\mu \in P_{++}^n$

$$\sum_{\substack{\nu \in P_{++}^n \\ [\nu] = \mu}} \epsilon(\nu) \ell_\lambda^\nu = \sum_{\sigma \in P_{++}^n} E_\lambda^{(n)}(\sigma) S_{\rho\sigma}^{(n)} S_{\mu\sigma}^{(n)*}. \tag{4.4}$$

Thus for any $\mu \in P_{++}^n$, and any $\lambda \in P_{++}^n$ with $[\lambda - \rho] + \rho \in P_{++}^n$, we get the truncation

$$\ell_{[\lambda - \rho] + \rho}^\mu = \frac{|W([\lambda - \rho])|}{|W(\lambda - \rho)|} \sum_{\substack{\nu \in P_{++}^n \\ [\nu] = \mu}} \epsilon(\nu) \ell_\lambda^\nu. \tag{4.5}$$

Roughly speaking, equation (4.5) states that if we know the ℓ for ‘large’ weights, then we know them for ‘small’ ones. Incidentally, if $[\lambda - \rho] + \rho \notin P_{++}^n$, then equation (4.5) holds if we replace its LHS with a sum similar to that of its RHS. Similar comments hold below if $\pi_A(\lambda) \notin P_+^n$ in (4.13), or $\pi_\mu(\lambda) \notin P_+^n$ in (4.17).

Next, consider the symmetries involving the outer automorphisms of affine Lie algebras \widehat{g} , or equivalently, the automorphisms of the extended Coxeter-Dynkin diagrams of g . If an outer automorphism is also a symmetry of the unextended Coxeter-Dynkin diagram of g , i.e. it fixes the extended node, then it is well known to be an exact symmetry C (a conjugation) of both the ℓ and the m :

$$m_{C\lambda}^{C\mu} = m_\lambda^\mu \quad \ell_{C\lambda}^{C\mu} = \ell_\lambda^\mu. \tag{4.6}$$

Here we are interested instead in those automorphisms which are not conjugations. Denote such an automorphism by A , and the fundamental weights of \widehat{g} by ω^i , with $i = 0, 1, 2, \dots, r$. There is one of these automorphisms $A = A_i$ for every node of the extended diagram with mark $a_i = 1$; A_i will send ω^0 to ω^i . Since

$$A(\lambda - n\omega^0) = w_A(\lambda - n\omega^0) \tag{4.7}$$

with w_A an element of the Weyl group W of g , for all $\lambda \in P_+^n$, we have [15]

$$S_{A\lambda, \sigma}^{(n)} = S_{\lambda\sigma}^{(n)} \exp[-2\pi i(A\omega^0) \cdot \sigma] \det(w_A) = S_{\lambda\sigma}^{(n)} \exp[-2\pi i(A\omega^0) \cdot (\sigma - \rho)]. \tag{4.8}$$

Similarly

$$E_\lambda^{(n)}(A\sigma) = E_\lambda^{(n)}(\sigma) \exp[-2\pi i(A\omega^0) \cdot (\lambda - \rho)] \tag{4.9}$$

$$E_{A\lambda}^{(n)}(\sigma) = E_{\pi_A\lambda}^{(n)}(\sigma) \frac{|W(A\lambda - \rho)|}{|W(\pi_A\lambda - \rho)|} \exp[-2\pi i(A\omega^0) \cdot \sigma] \tag{4.10}$$

where π_A denotes the one-to-one map from P_{++}^n to P_{++} given by

$$\pi_A(\lambda) := [\lambda - w_A^{-1}\rho] + \rho. \tag{4.11}$$

For fixed g, w_A and hence the map π_A is readily obtained from (4.7)— π_A will be the identity only when A is. For example, for $g = su(r + 1)$ and $A = A_j$ satisfying $A\omega^j = \omega^{j+j}$, we get

$$(w_A\lambda)_i = \begin{cases} \lambda_{i-j} & \text{if } i \neq j \\ -\lambda_1 - \dots - \lambda_r & \text{if } i = j \end{cases} \tag{4.12}$$

$\det(w_A) = (-1)^{rj}$, and $\pi_A(\lambda) = \lambda + (r + 1)\omega^{r+1-j}$.

From our main result (3.10), we immediately find

$$\ell_{A\lambda}^{A\mu} = \ell_{\pi_A(\lambda)}^\mu \frac{|W(A\lambda - \rho)|}{|W(\pi_A\lambda - \rho)|} \det(w_A) \tag{4.13}$$

for any $\lambda, \mu \in P_{++}^n$, provided $\pi_A(\lambda) \in P_{++}^n$. Unfortunately π_A will only be a permutation of P_{++}^n in the trivial case when $A = \text{id}$, so it is not easy to see what equation (4.13) directly implies for the m_λ^μ .

There are also Galois symmetries of the Kac–Peterson modular matrix $S^{(n)}$, first discovered in [16, 17] (and generalized to all rational conformal field theories in [18]). The $S_{\lambda\mu}^{(n)}/F_n$ and $E_\lambda^{(n)}(\sigma)$ are polynomials with rational coefficients in a primitive (nN) th root of unity, where $N = |M^*/M|^{\frac{1}{2}}$, M here being the weight lattice of g . So, any polynomial relation involving them and rational numbers only, will also be satisfied if this primitive (nN) th root of unity is replaced by another.

Let a be an integer coprime to nN , and let $a(S^{(n)})$ denote the Kac–Peterson matrix after the primitive (nN) th root of unity is replaced by its a th power (ignoring here the irrelevant factor F_n). For such a , and for $\lambda \in P_{++}^n$, recall the quantities $[a\lambda] \in P_{++}^n$ and $\epsilon(a\lambda) \in \{\pm 1\}$ defined around (4.1). For each a coprime to nN , the map $\lambda \mapsto [a\lambda]$ is a permutation of P_{++}^n . From the form of the matrix $S^{(n)}$, it is easy to find

$$a(S_{\lambda\mu}^{(n)}) = \epsilon(a\lambda) S_{[a\lambda], \mu}^{(n)} = \epsilon(a\mu) S_{\lambda, [a\mu]}^{(n)}. \tag{4.14}$$

In a similar fashion, we find

$$a(E_\lambda^{(n)}(\sigma)) = \frac{|W(\lambda - \rho)|}{|W(\pi_A\lambda - \rho)|} E_{\pi_A(\lambda)}^{(n)}(\sigma) = E_\lambda^{(n)}([a\sigma]) \tag{4.15}$$

where π_a denotes the one-to-one map from P_{++}^n to P_{++} defined by

$$\pi_a(\lambda) := [a\lambda - a\rho] + \rho. \tag{4.16}$$

A little work yields

$$\ell_\lambda^\mu = \epsilon(a\mu)\epsilon(a\rho) \frac{|W(\lambda - \rho)|}{|W(\pi_a\lambda - \rho)|} \sum_{\nu \in P_{++}^n} \ell_{\pi_a(\lambda)}^\nu N_{\nu, [a\rho]}^{(n) [a\mu]} \tag{4.17}$$

whenever $\mu \in P_{++}^n$ and $\lambda, \pi_a(\lambda) \in P_{++} \cap P_{++}^n$. Here we have used the Verlinde formula [11] given in (3.12) for the fusion coefficients $N_{\lambda\mu}^{(n) \nu}$. Let $N_\lambda^{(n)}$ denote the fusion matrix defined by $(N_\lambda^{(n)})_\mu^\nu := N_{\lambda\mu}^{(n) \nu}$. The matrix $N_{[a\rho]}^{(n)}$ will always be invertible [18], so equation (4.17) tells us that for any fixed $\lambda \in P_{++}$, the values $\ell_{\pi_a(\lambda)}^-$ will be known once the ℓ_λ^- are known, and conversely, provided a and n satisfy the usual conditions. Equation (4.17) can also be interpreted as an expression for the fusion matrices $N_{[a\rho]}^{(n)}$ in terms of the ℓ and the m .

It is again difficult to express this Galois symmetry directly at the level of the multiplicities m_λ^μ themselves. But if π_a is a permutation of P_{++}^n , we get from (4.17) that

$$m_\lambda^\mu = \epsilon(a\mu)\epsilon(a\rho) \frac{|W(\pi_a\lambda - \rho)|}{|W(\lambda - \rho)|} \sum_{\nu \in P_{++}^n} (N_{[a\rho]}^{(n-1)})_{[a\lambda]}^\nu m_{\nu}^{\pi_a(\mu)}. \tag{4.18}$$

A special case of (4.17) occurs when $[a\rho] = \rho$. (More generally, a similar simplification happens whenever $[a\rho] = A\rho$ for some outer automorphism A .) Then $\epsilon(a\rho) = \epsilon(a\lambda)$ for all $\lambda \in P_{++}^n$ (apply equation (4.14) with $\mu = \rho$, together with the fact that $S_{\rho\nu}^{(n)} > 0$ for all $\nu \in P_{++}^n$). Equation (4.17) reduces to

$$\ell_\lambda^\mu = \frac{|W(\lambda - \rho)|}{|W(\pi_a\lambda - \rho)|} \ell_{\pi_a(\lambda)}^{[a\mu]}. \tag{4.19}$$

For example this happens whenever $a = -1$, and we get an example of (4.6).

Similarly, suppose $[a\mu] = A\rho$, for some outer automorphism A . Then provided $\lambda, \pi_a(\lambda) \in P_{++} \cap P_{++}^n$, equation (4.17) reduces to

$$\ell_\lambda^\mu = \epsilon(a\mu)\epsilon(a\rho) \frac{|W(\lambda - \rho)|}{|W(\pi_a\lambda - \rho)|} \ell_{\pi_a\lambda}^{A[a\rho]}. \tag{4.20}$$

Another noteworthy special case of (4.17) involves those weights λ' with the property that $\ell_{\lambda'}^\mu = \delta_{\lambda'}^\mu$ for all $\mu \in P_{++}$, i.e. those λ' for which $\lambda' - \rho$ is a miniscule weight. For $su(r + 1)$, they are the fundamental weights $\lambda' = \omega^i + \rho$. Then for any $\lambda \in P_{++}^{(n)}$ with $\pi_a(\lambda) = \lambda'$

$$\ell_\lambda^\mu = \epsilon(a\mu)\epsilon(a\rho) \frac{|W(\lambda - \rho)|}{|W(\lambda' - \rho)|} N_{\lambda', [a\rho]}^{(n) [a\mu]} \tag{4.21}$$

if, as usual, a is coprime to nN and $\lambda', \mu \in P_{++}^n$. The fusions involving the fundamental weights $\omega^i + \rho$ are easy to compute, so the RHS of (4.21) can be explicitly evaluated in all cases. One of the reasons equations (4.17)–(4.21) could be interesting is because they suggest the rank-level duality that WZW fusions satisfy [19] could appear in some way in the ℓ and the m .

5. Examples: the case of $g = su(3)$

For concrete illustrations of our results, we will focus on the example of $g = su(3)$. The comments in this section can be extended to any $su(r + 1)$ without difficulty.

Consider first equation (4.13). It reduces to

$$\ell_{(\lambda_1, \lambda_2)}^{(\mu_1, \mu_2)} = \ell_{(n-\lambda_1-\lambda_2+3, \lambda_2)}^{(n-\mu_1-\mu_2, \mu_2)} \quad \forall n \in \mathbb{Z}_> \tag{5.1}$$

provided only that: $\mu_1 + \mu_2 < n$; $\lambda_2, \mu_1, \mu_2 \geq 1$; and either $\lambda_1 + \lambda_2 < n, \lambda_1 \geq 3$, or $\lambda_1 + \lambda_2 \leq n, \lambda_1 > 3$.

Equation (5.1) is very powerful. For example, suppose $\lambda_1 \geq 3$. Put $n = \lambda_1 + \lambda_2 + 1$. Then equation (5.1) reduces ℓ_λ^μ to $\ell_{\lambda'}^{\mu'}$, where $\lambda' = (4, \lambda_2)$. In fact, if both $\lambda_1, \lambda_2 \geq 3$ and $\mu \in P_{++}$, from (5.1) (once we compute the values of $\ell_{(4,4)}^-$) we obtain

$$\ell_\lambda^\mu = \begin{cases} +1 & \text{if } \mu \in \{\lambda, (\lambda_1 - 3, \lambda_2), (\lambda_1, \lambda_2 - 3)\} \\ -1 & \text{if } \mu \in \{(\lambda_1 - 2, \lambda_2 + 1), (\lambda_1 + 1, \lambda_2 - 2), (\lambda_1 - 2, \lambda_2 - 2)\} \\ 0 & \text{otherwise.} \end{cases} \tag{5.2}$$

In fact, using equation (4.5), we find that equation (5.2) gives the correct value of ℓ_λ^μ for any $\lambda, \mu \in P_{++}$ (whether or not $\lambda_1 \geq 3$), with one exception: $\ell_{(2,2)}^{(1,1)} = -2 \neq 0$,

Next, let us turn to the Galois symmetries. It is not difficult to see that $[\alpha\rho] = \rho$ only for $a \equiv \pm 1 \pmod n$, in which case $\pi_a(\lambda) = \pi_A(\lambda)$ for some outer automorphism A . Thus for $g = su(3)$, equation (4.19) only gives information also obtainable from (5.1).

However, equations (4.17), (4.20), (4.21) are very powerful here. To give one explicit example, consider $\lambda = (4, 4), a = 5, n = 8$. Then $\pi_a\lambda = (2, 2)$. The only possible non-zero elements of $\ell_{(2,2)}^-$ are $\ell_{(2,2)}^{(2,2)} = 1$ and $\ell_{(2,2)}^{(1,1)}$. We get from (4.17) that, for example,

$$\begin{aligned} \ell_{(4,4)}^{(3,3)} &= 0 & \ell_{(4,4)}^{(1,1)} &= N_{(2,2),(3,3)}^{(8)} + \ell_{(2,2)}^{(1,1)} & \ell_{(4,4)}^{(2,2)} &= -N_{(2,2),(3,3)}^{(8)} \\ \ell_{(4,4)}^{(5,2)} &= -N_{(2,2),(3,3)}^{(8)} & \ell_{(4,4)}^{(1,4)} &= N_{(2,2),(3,3)}^{(8)} \end{aligned} \tag{5.3}$$

Note that the relations (4.5), (4.13) and (4.17), together with the selection rule ' $\ell_\lambda^\mu \neq 0$ requires $\lambda - \mu \in \mathbb{Z}_{\geq}\{\alpha_1, \dots, \alpha_r\}$ ', and the normalization $\ell_\lambda^\lambda = 1$, easily determine all values of ℓ_λ^μ and hence m_λ^μ . In particular, we have seen that equations (4.5) and (4.13) determine all ℓ_λ^μ provided the values $\ell_{(4,4)}^-$ are known. Equation (5.3) fixes all values of $\ell_{(4,4)}^-$ except $\ell_{(4,4)}^{(1,1)}$, but by equation (5.1) we find $\ell_{(4,4)}^{(1,1)} = \ell_{(4,4)}^{(7,1)} = 0$. A similar conclusion should apply to any $su(r + 1)$.

6. Conclusion

Our main result is expression (3.10) for the inverse of the matrix of dominant-weight multiplicities, in terms of the Kac–Peterson modular matrix $S^{(n)}$ and the even Weyl orbit sums $E_\lambda^{(n)}(\sigma)$. Surely the even Weyl orbit sums will find other uses in the study of WZW models and affine Kac–Moody algebras.

Also obtained were relations (4.5), (4.13) and (4.17) between the ‘inverse multiplicities’ ℓ_λ^μ , that are consequences of the symmetries of the Kac–Peterson matrices and the $E_\lambda^{(n)}(\sigma)$. Relations between the dominant-weight multiplicities m_λ^μ follow in certain cases, as

equation (4.18) shows. These relations would be difficult to understand from the point of view of Lie groups and their semi-simple Lie algebras only, but are quite natural in WZW models, or in their affine Kac–Moody current algebras. The new relations arise because the semi-simple symmetry algebra of a WZW model is a special subalgebra of its affine current algebra. The affine Weyl and outer automorphism symmetries are especially natural in the Kac–Moody context. Perhaps the Galois symmetries can be better understood in terms of affine Kac–Moody algebras.

The debt we owe to the previous work of Patera and Sharp [12] is obvious when comparing sections 2 and 3. We should also mention a paper by Moody and Patera [20]. In it class functions of an arbitrary semi-simple compact Lie group are decomposed into sums of irreducible characters. The method uses the characters of elements of finite order (EFOs) of the group to approximate the characters of arbitrary elements. These characters of the EFOs have a marked similarity to the Kac–Peterson ratios $\chi_\lambda^{(n)}(\sigma)$. Moody and Patera also use modular arithmetic to ensure that the approximation leads to the correct answers, if a suitable set of EFOs is chosen. In a similar way, we automatically recover the exact weight multiplicities from the Kac–Peterson ratios, which are the Lie algebra characters evaluated at special points.

Further comparison with [20] is clearly warranted. Perhaps it will help us toward a better understanding of WZW models, one that approaches the current understanding of semi-simple compact Lie groups and their algebras.

Acknowledgments

TG and MW thank the IHES for hospitality, and MW also thanks R T Sharp and P Mathieu for discussions.

References

- [1] Goddard P and Olive D 1986 *Int. J. Mod. Phys. A* **1** 303
Fuchs J 1992 *Affine Lie Algebras and Quantum Groups* (Cambridge: Cambridge University Press)
- [2] Fulton W and Harris J 1991 *Representation Theory—A First Course* (Berlin: Springer)
- [3] Bremner M R, Moody R V and Patera J 1985 *Tables of Dominant Weight Multiplicities for Representations of Simple Lie Algebras* (New York: Dekker)
- [4] Witten E 1984 *Nucl. Phys.* **92** 455
Knizhnik V and Zamolodchikov A B 1984 *Nucl. Phys. B* **247** 83
- [5] Gepner D and Witten E 1986 *Nucl. Phys. B* **278** 493
- [6] Kac V G and Peterson D 1984 *Adv. Math.* **53** 125
- [7] Walton M A 1990 *Phys. Lett.* **241B** 365; 1990 *Nucl. Phys. B* **340** 777
- [8] Fuchs J 1991 *Comm. Math. Phys.* **136** 345
- [9] Gannon T 1994 The classification of $su(m)_k$ automorphism invariants *IHES preprint hep-th/9408119*
- [10] Kirillov A N, Mathieu P, Sénéchal D and Walton M A 1993 *Nucl. Phys. B* **391** 257
- [11] Verlinde E 1988 *Nucl. Phys. B* **300** 360
- [12] Patera J and Sharp R T 1989 *J. Phys. A: Math. Gen.* **22** 2329
- [13] Bourbaki N 1968 *Groupes et algèbres de Lie* (Paris: Hermann) chs IV–VI
- [14] Kac V G 1990 *Infinite Dimensional Lie Algebras* (Cambridge: Cambridge University Press) 3rd edn
- [15] Bernard D 1987 *Nucl. Phys. B* **288** 626
- [16] Gannon T 1993 *Nucl. Phys. B* **396** 708
- [17] Ruelle Ph, Thiran E and Weyers J 1993 *Nucl. Phys. B* **402** 693
- [18] Coste A and Gannon T 1994 *Phys. Lett.* **323B** 316
- [19] Goodman F and Wenzl H 1990 *Adv. Math.* **82** 244
Cummins C J 1991 *J. Phys. A: Math. Gen.* **24** 391
Mlawer E, Naculich S, Riggs H and Schnitzer H 1991 *Nucl. Phys. B* **352** 863
- [20] Moody R V and Patera J 1987 *Math. Comput.* **48** 799