

On a one-parameter family of q -exponential functions

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1996 J. Phys. A: Math. Gen. 29 L223

(<http://iopscience.iop.org/0305-4470/29/10/001>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.70

The article was downloaded on 02/06/2010 at 03:51

Please note that [terms and conditions apply](#).

LETTER TO THE EDITOR

On a one-parameter family of q -exponential functions

N M Atakishiyev^{†‡}

IIMAS–UNAM, Apartado Postal 139-B, 62191 Cuernavaca, Morelos, Mexico

Received 29 February 1996

Abstract. We examine the properties of a family of q -exponential functions, which depend on an extra parameter α . These functions have a well defined meaning for both the $0 < |q| < 1$ and $|q| > 1$ cases if only $\alpha \in [0, 1]$. It is shown that any two members of this family with different values of the parameter α are related to each other by a Fourier–Gauss transformation.

The one-parameter family of q -exponential functions

$$E_q^{(\alpha)}(z) = \sum_{n=0}^{\infty} \frac{q^{\alpha n^2/2}}{(q; q)_n} z^n \tag{1}$$

with $\alpha \in \mathfrak{R}$ has been considered in [1]. The q -shifted factorial $(q; q)_n$ in (1) is defined as $(z; q)_0 = 1$ and $(z; q)_n = \prod_{j=0}^{n-1} (1 - zq^j)$, $n = 1, 2, 3, \dots$. Consequently, in the limit when $q \rightarrow 1$ we have

$$\lim_{q \rightarrow 1} E_q^{(\alpha)}((1 - q)z) = e^z. \tag{2}$$

The possibility of introducing such a family of q -exponential functions has been already mentioned in [2]. Exton defines it as

$$E(q, \lambda; x) = \sum_{n=0}^{\infty} \frac{x^n q^{\lambda n(n-1)}}{[n; q]!} \tag{3}$$

where the symbol $[n; q]!$ denotes the product $\prod_{j=1}^n [j; q]$, with $[0; q] = 1$ and the bracket notation

$$[j; q] = \frac{1 - q^j}{1 - q}. \tag{4}$$

Since $(q; q)_n = (1 - q)^n [n; q]!$ by definition, it is obvious that the relationship between these two notations is

$$E(q, \lambda; x) = E_q^{(2\lambda)}(q^{-\lambda}(1 - q)x). \tag{5}$$

Two particular cases of this family with $\alpha = 0$ and $\alpha = 1$ are well known: they are the q -exponential function

$$e_q(z) = E_q^{(0)}(z) = \sum_{n=0}^{\infty} \frac{z^n}{(q, q)_n} \tag{6}$$

[†] Permanent address: Institute of Physics, Azerbaijan Academy of Sciences, Baku 370143, Azerbaijan. Visiting scientist at IIMAS–UNAM/Cuernavaca with Catedra Patrimonial, CONACYT, Mexico.

[‡] E-mail: natig@ce.ifisicam.unam.mx

and its reciprocal

$$E_q(z) = e_q^{-1}(z) = E_q^{(1)}(-q^{-\frac{1}{2}}z) = \sum_{n=0}^{\infty} \frac{q^{n(n-1)/2}}{(q; q)_n} (-z)^n \tag{7}$$

respectively [3]. Another particular example of (1) corresponds to the value $\alpha = \frac{1}{2}$ and is

$$E_q^{(1/2)}(z) = \mathcal{E}_q(-; 0, z) = \varepsilon_q(z) \tag{8}$$

where $\mathcal{E}_q(z; a, b)$ is a two-parameter q -exponential function, introduced in [4]. Exton denotes this q -exponential function as [2]

$$E(q, x) = E_q^{(1/2)}(q^{-1/4}(1 - q)x) \tag{9}$$

and he emphasizes that it ‘was originally considered in connection with a particular q -generalization of the circular functions which exhibit properties of q -orthogonality’.

By analogy with the exponentiating of Lie algebras into Lie groups, one may consider q -exponentials of the generators of a q -algebra and express their matrix elements in representation space in terms of q -special functions. In this way one manages to interpret algebraically the properties of these q -special functions through symmetry techniques [1]. Furthermore, as we show below, members of the family (1) with different values of the parameter α turn out to be Fourier–Gauss transforms of (6) and (7). In other words, they are of central importance for constructing Fourier–Gauss transforms of a number of q -special functions (cf [5, 6]). Therefore we wish to study here some additional properties of the q -exponential functions (1).

We start with the observation that by the ratio test the infinite series in (1) is convergent for $0 < |q| < 1$ and arbitrary complex z only if the parameter α is positive: $0 < \alpha < \infty$. The case $\alpha = 0$ is a little bit more involved, but $E_q^{(0)}(z) = e_q(z)$ and the properties of the q -exponential function (6) are well studied [2, 3, 7].

The series (1) converges also for $1 < |q| < \infty$ and arbitrary complex z , provided that $-\infty < \alpha < 1$. Thus, the q -exponential functions (1) are well defined for both $0 < |q| < 1$ and $1 < |q| < \infty$, if the parameter α belongs to the line segment $[0, 1]$. Observe that the inversion formula

$$(q^{-1}; q^{-1})_n = (-1)^n q^{-n(n+1)/2} (q; q)_n \tag{10}$$

leads to the relation

$$E_{q^{-1}}^{(\alpha)}(z) = E_q^{(1-\alpha)}(-q^{1/2}z). \tag{11}$$

When $\alpha = 0$, (11) reproduces the known relation [2, 7]

$$e_{q^{-1}}(z) = E_q(qz) \tag{12}$$

between the q -exponential functions (6) and (7) for $0 < |q| < 1$ and $1 < |q| < \infty$ (or vice versa), respectively.

There are two types of Fourier–Gauss transforms for the q -exponential functions (1), depending on whether the parameter α belongs to either the interval $[0, \frac{1}{2}]$, or $[\frac{1}{2}, 1]$. Let us consider these cases in turn.

(a) When $0 \leq \alpha \leq \frac{1}{2}$, it is not hard to show that

$$E_q^{(\alpha+\frac{1}{2})}(t e^{-\kappa x}) e^{-x^2/2} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{ixy-y^2/2} E_q^{(\alpha)}(t e^{i\kappa y}) dy \tag{13}$$

where $q = \exp(-2\kappa^2)$. Indeed, to evaluate the right-hand side of (13) one only needs to use the definition (1) with $z = t e^{i\kappa y}$ and to integrate this sum termwise by the Fourier transform

$$\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{ixy-y^2/2} dy = e^{-x^2/2} \tag{14}$$

for the Gauss exponential function $\exp(-x^2/2)$. Important particular cases of (13) are

$$\varepsilon_q(t e^{-\kappa x}) e^{-x^2/2} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{ixy-y^2/2} e_q(t e^{i\kappa y}) dy \tag{15}$$

and

$$E_q(-q^{1/2} t e^{-\kappa x}) e^{-x^2/2} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{ixy-y^2/2} \varepsilon_q(t e^{i\kappa y}) dy. \tag{16}$$

They correspond to the values 0 and $\frac{1}{2}$ of the parameter α , respectively.

(b) In like manner, for $\frac{1}{2} \leq \alpha \leq 1$ we have

$$E_q^{(\alpha-1/2)}(t e^{i\kappa x}) e^{-x^2/2} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{ixy-y^2/2} E_q^{(\alpha)}(t e^{i\kappa y}) dy. \tag{17}$$

In particular, when $\alpha = \frac{1}{2}$ from (17) follows the inverse Fourier transformation with respect to (15)

$$e_q(t e^{i\kappa x}) e^{-x^2/2} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{ixy-y^2/2} \varepsilon_q(t e^{i\kappa y}) dy \tag{15'}$$

whereas the value $\alpha = 1$ yields the inverse to (16), i.e.

$$\varepsilon_q(t e^{i\kappa x}) e^{-x^2/2} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{ixy-y^2/2} E_q(-q^{1/2} t e^{i\kappa y}) dy. \tag{16'}$$

Actually, the Fourier–Gauss transforms (13) and (17) may be written in the unified form

$$E_q^{(\alpha+v^2/2)}(t e^{-v\kappa x}) e^{-x^2/2} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{ixy-y^2/2} E_q^{(\alpha)}(t e^{i v \kappa y}) dy \tag{18}$$

provided that $-\alpha \leq v^2/2 \leq 1 - \alpha$. This is easy to prove in exactly the same way as (13), or (17). When $0 \leq \alpha \leq \frac{1}{2}$ and $v = 1$ from (18) one obtains (15) and when $\frac{1}{2} \leq \alpha \leq 1$ and $v = -i$ from (18) follows (17). Also, for $\alpha = 0$ and $v = \sqrt{2}$ the Fourier–Gauss transform (18) gives the following relation between the q -exponential functions $e_q(z)$ and $E_q(z)$ (cf (16))

$$E_q(-q^{1/2} t e^{-\sqrt{2}\kappa x}) e^{-x^2/2} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{ixy-y^2/2} e_q(t e^{i\sqrt{2}\kappa y}) dy \tag{19}$$

which is a particular case of Ramanujan’s integral with a complex parameter [8–10].

Finally, we would like to touch upon two intimately interrelated features of the q -exponential functions (1). One of them is an explicit form of reciprocal q -exponential functions for any parameter α from the interval $[0, 1]$. Unfortunately, we know such reciprocals for the boundary values $\alpha = 0$ and $\alpha = 1$ only. The formal solution of this problem is known: for any $\alpha \in [0, 1]$ the function reciprocal to (1) is represented by the infinite series

$$1/E_q^{(\alpha)}(z) = \sum_{n=0}^{\infty} c_n^{(\alpha)}(q) z^n \tag{20}$$

with the coefficients $c_0^{(\alpha)}(q) = 1$ and

$$c_n^{(\alpha)}(q) = (-1)^n \begin{vmatrix} a_1^{(\alpha)}(q) & 1 & 0 & \dots & 0 \\ a_2^{(\alpha)}(q) & a_1^{(\alpha)}(q) & 1 & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n-1}^{(\alpha)}(q) & a_{n-2}^{(\alpha)}(q) & a_{n-3}^{(\alpha)}(q) & \dots & 1 \\ a_n^{(\alpha)}(q) & a_{n-1}^{(\alpha)}(q) & a_{n-2}^{(\alpha)}(q) & \dots & a_1^{(\alpha)}(q) \end{vmatrix} \quad n = 1, 2, \dots \tag{21}$$

where $a_n^{(\alpha)}(q) = q^{\alpha n^2/2}(q; q)_n^{-1}$ are the corresponding coefficients in the expansion (1) for $E_q^{(\alpha)}(z)$.

Another interesting point is the possibility of representing $E_q^{(\alpha)}(z)$ as an infinite product. Again, we know that in the particular cases of $\alpha = 0$ and $\alpha = 1$ the affirmative answer to this question is given by Euler's formulae [3]

$$e_q(z) = (z; q)_\infty^{-1} \quad E_q(z) = (z; q)_\infty \tag{22}$$

for the q -exponential functions (6) and (7). Observe that Euler's formulae (22) follow from the functional relations

$$e_q(qz) = (1 - z)e_q(z) \quad E_q(z) = (1 - z)E_q(qz) \tag{23}$$

and the side conditions $e_q(0) = E_q(0) = 1$. One may try to combine one of the relations (23) with either (15), or (16), respectively, in order to get the corresponding representation at least for the parameter $\alpha = \frac{1}{2}$. However, in both cases this results in the functional relation (cf [11])

$$\varepsilon_q(qz) = \varepsilon_q(z) - q^{1/4}z\varepsilon_q(q^{1/2}z) \tag{24}$$

with $z = t e^{-\kappa x}$ and $z = t e^{i\kappa x}$, respectively. Actually, this type of functional relation holds for arbitrary z and $\alpha \in [0, 1]$ (not for $\alpha = \frac{1}{2}$ only!) and has the form

$$E_q^{(\alpha)}(qz) = E_q^{(\alpha)}(z) - q^{\alpha/2}zE_q^{(\alpha)}(q^\alpha z). \tag{25}$$

The validity of (25) can be readily verified by using the definition (1). When $\alpha = 0$ and $\alpha = 1$, from (25) follow the functional relations (23) for the q -exponential functions $e_q(z)$ and $E_q(z)$, respectively.

The relation (25) is equivalent to a q -difference equation. Indeed, in terms of the Jackson q -difference operator [12]

$$\Delta f(z) = \frac{f(z) - f(qz)}{z(1 - q)} \tag{26}$$

it may be represented as

$$\Delta E_q^{(\alpha)}(z) = \frac{q^{\alpha/2}}{1 - q} E_q^{(\alpha)}(q^\alpha z). \tag{25'}$$

Note also that applying the functional relations (23) n times in succession leads to

$$e_q(q^n z) = (z; q)_n e_q(z) \quad E_q(z) = (z; q)_n E_q(q^n z) \tag{27}$$

respectively. These simple formulae turn out to be very useful in proving the orthogonality of the classical q -polynomials with respect to measures, containing q -exponential functions $e_q(z)$ and $E_q(z)$ [13–16]. In an analogous manner, from (25) it follows at once that

$$E_q^{(\alpha)}(q^n z) = \sum_{k=0}^n \begin{bmatrix} n \\ k \end{bmatrix}_q (-z)^k q^{[(\alpha+1)k-1]k/2} E_q^{(\alpha)}(q^{\alpha k} z) \tag{28}$$

where

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}}$$

are the q -binomial coefficients. This relation is easily verified by induction, upon using the following property of the q -binomial coefficients [3]

$$\begin{bmatrix} n \\ k \end{bmatrix}_q + q^{n-k+1} \begin{bmatrix} n \\ k-1 \end{bmatrix}_q = \begin{bmatrix} n+1 \\ k \end{bmatrix}_q.$$

Discussions with A B Balantekin, F Leyvraz, and K B Wolf are gratefully acknowledged. This work is partially supported by the UNAM-DGAPA project IN106595.

References

- [1] Floreanini R, LeTourneau J and Vinet L 1995 *J. Phys. A: Math. Gen.* **28** L287–93
- [2] Exton H 1983 *q-Hypergeometric Functions and Applications* (Chichester: Ellis Horwood)
- [3] Gasper G and Rahman M 1990 *Basic Hypergeometric Functions* (Cambridge: Cambridge University Press)
- [4] Ismail M E H and Zhang R 1994 *Adv. Math.* **109** 1–33
- [5] Atakishiyev N M and Nagiyev Sh M 1994 *Theor. Math. Phys.* **98** 162–8; *J. Phys. A: Math. Gen.* **27** L611–15
- [6] Atakishiyev N M and Feinsilver P 1995 On the coherent states for the q -Hermite polynomials and related Fourier transformation *Preprint* 37 IIMAS-UNAM, México (1996 *J. Phys. A: Math. Gen.* to appear)
- [7] Atakishiyev N M 1996 A Ramanujan-type measure for the Al-Salam and Ismail biorthogonal rational functions *Proc. Workshop on Symmetries and Integrability of Difference Equations (Estérel, Canada, 22–29 May, 1994)* CRM Proceedings and Lecture Notes, Université de Montréal
- [8] Ramanujan S 1927 *Collected Papers* ed G H Hardy *et al* (Cambridge: Cambridge University Press) (reprinted 1959 (New York: Chelsea))
- [9] Askey R 1982 *Proc. Am. Math. Soc.* **85** 192–4
- [10] Atakishiyev N M and Feinsilver P 1996 Two Ramanujan's integrals with a complex parameter *Proc. IVth Wigner Symp. (Guadalajara, Mexico, 7–11 August, 1995)* (Singapore: World Scientific)
- [11] Nelson C A and Gartley M G 1994 *J. Phys. A: Math. Gen.* **27** 3857–81
- [12] Jackson F H 1908 *Trans. R. Soc. Edin.* **46** 253–81
- [13] Andrews G E and Askey R A 1984 *Lecture Notes in Mathematics* vol 1171 (Berlin: Springer) pp 36–63
- [14] Askey R A and Wilson J A 1985 *Mem. Am. Math. Soc.* **54** 1–55
- [15] Berg C and Ismail M E H 1996 q -Hermite polynomials and classical orthogonal polynomials *Can. J. Math.* to appear
- [16] Atakishiyev N M 1995 Ramanujan-type continuous measures for classical q -polynomials *Preprint* CRM–2254, Centre de Recherches Mathématiques, Université de Montréal (*Teor. Mat. Fiz.* to appear)