

## Strong chaos without the butterfly effect in dynamical systems with feedback

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1996 J. Phys. A: Math. Gen. 29 2291

(<http://iopscience.iop.org/0305-4470/29/10/009>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.70

The article was downloaded on 02/06/2010 at 03:51

Please note that [terms and conditions apply](#).

## Strong chaos without the butterfly effect in dynamical systems with feedback

Guido Boffetta<sup>†</sup>, Giovanni Paladin<sup>‡</sup> and Angelo Vulpiani<sup>§</sup>

<sup>†</sup> Istituto de Fisica Generale, Università di Torino, Via P. Giuria 1 I-10125 Torino, Italy

<sup>‡</sup> Dipartimento de Fisica, Università dell'Aquila, Via Vetoio I-67100 Coppito, L'Aquila, Italy

<sup>§</sup> Dipartimento de Fisica, Università di Roma 'La Sapienza' P.le A Moro 2 I-00185 Rome, Italy

Received 9 June 1995, in final form 11 December 1995

**Abstract.** We discuss the predictability of a system that drives a chaotic system with a positive Lyapunov exponent. In the absence of feedback, the driver is regular and fully predictable. With a small feedback of strength  $\epsilon$ , the state of the driver can be predicted up to a time diverging with a power of  $\epsilon^{-1}$ , although the total system is strongly chaotic. The exponential amplification of the uncertainty on the initial conditions of the driver coexists with very long predictability times as illustrated in a model of coupled maps and of three point vortices in a disc.

It is commonly believed that a sensible dependence on an initial condition makes forecasting impossible even in systems with a few degrees of freedom. This is the so-called butterfly effect discovered by Lorenz in a numerical simulation of a model of convection with three degrees of freedom (Lorenz 1963). Using his words, 'A butterfly moving its wings over Brazil might cause the formation of a tornado in Texas'.

In general, a dynamical system is considered chaotic when there is an exponential amplification of an infinitesimal perturbation  $\delta_0$  on the initial conditions, with a mean time rate given by the inverse of the maximum Lyapunov exponent  $\lambda$  (Benettin *et al* 1976). Indeed, such a deterministic system is expected to be predictable on times  $t \ll \lambda^{-1}$  and to behave like a random system on larger times. The purpose of this paper is to show that there exists a wide class of dynamical systems where a large value of the Lyapunov exponent does not imply a short predictability time on a physically relevant part of the system. In this respect, one can speak of strong chaos  $\lambda > 0$  without a butterfly effect.

In particular, we discuss the predictability of a conservative system that drives a strongly chaotic system with positive maximum Lyapunov exponent  $\lambda_0$ . In the absence of feedback the driving system is regular and completely predictable. A small feedback of strength  $\epsilon$  still allows us to predict the future of the driving system up to a very long predictability time  $T_p$  that diverges with  $\epsilon$ . The Lyapunov exponent of the total system is  $\lambda_{\text{tot}} \approx \lambda_0$ , and so there is a regime of strong chaos for all  $\epsilon$  values. The absence of the butterfly effect stems from saturation effects in the evolution laws for the growth of an uncertainty on the driven system.

To be explicit, let us consider a system with evolution given by two sets of equations

$$\frac{d\xi}{dt} = F(\xi) + \epsilon h(\eta) \quad (1a)$$

$$\frac{d\eta}{dt} = G(\xi, \eta) \quad (1b)$$

with  $\xi \in R^n$  and  $\eta \in R^m$ . The variables  $\eta$  are thus driven by a subsystem represented by the variables  $\xi$ , with a weak feedback of order  $\epsilon$ . A typical physical example is given by an asteroid moving in the gravitational field generated by two celestial bodies of much larger mass such as Jupiter and the Sun (Sussman and Wisdom 1992). Usually the feedback is neglected, and one considers the restricted three-body problem, i.e. the  $\eta$  variables passively driven. However, a finer description should take into account even the influence of the asteroid on the evolution of the other two bodies, i.e. an 'active' driving where  $\epsilon \neq 0$ . This situation appears in many other phenomena, such as the active advection of a contaminant in a fluid, a simple example which will be discussed in this paper.

The main properties of the system in the absence of feedback are the following:

(i) the driver is an independent dynamical system that exhibits a regular evolution with zero Lyapunov exponent;

(ii) the driven subsystem is chaotic with a positive maximum Lyapunov exponent, say  $\lambda_0$ . In other words, the behaviour of the driver ( $\xi$  variables) is completely predictable.

However, as soon as  $\epsilon \neq 0$ , one should consider the total system which is obviously chaotic with a Lyapunov exponent  $\lambda_{\text{tot}}$  that, a small correction of order  $\epsilon$ , is given by the Lyapunov exponent  $\lambda_0$  of the chaotic driven subsystem. This means that there is an exponential amplification of a small uncertainty on the knowledge of the initial conditions even in the driver. This is an amazing result, since it is natural to expect that it is possible to forecast the behaviour of the driver for very long times as  $\epsilon \rightarrow 0$ . Actually, intuition is correct while the Lyapunov analysis gives completely wrong hints on the predictability problem contrary to what was commonly believed. The famous butterfly effect of Lorenz seems not to forbid the possibility of predicting the future of a part of the system. The paradox stems from saturation effects in the evolution for the growth of the uncertainty. To fix notation and definitions, let us consider the evolution of the total system

$$\frac{dx_i}{dt} = f_i(x) \quad \text{where } x = (\xi, \eta) \in R^{n+m}. \quad (2)$$

The uncertainty on its state is  $\Delta(t) = x(t) - x'(t)$  where  $x$  and  $x'$  are trajectories starting from close initial conditions, i.e.  $|x(0) - x'(0)| = \delta_0$ . In the limit  $\delta_0 \rightarrow 0$ ,  $\Delta$  can be confused with the tangent vector  $z$  whose evolution equations are

$$\frac{dz}{dt} = J(t)z \quad \text{where } J_{ik}(t) = \left. \frac{df_i}{dx_j} \right|_{x(t)}. \quad (3)$$

The maximum Lyapunov exponent is then defined as the exponential rate of the uncertainty growth,

$$\lambda = \lim_{t \rightarrow \infty} \lim_{\delta_0 \rightarrow 0} \frac{1}{t} \ln \left( \frac{|\Delta(t)|}{\delta_0} \right) = \lim_{t \rightarrow \infty} \frac{1}{t} \ln |z(t)|. \quad (4)$$

It is worth stressing that the full equations for the evolution of an uncertainty are nonlinear:

$$\frac{d\Delta}{dt} = J(t)\Delta + O(\Delta^2) \quad (5)$$

so that the two limits in (4) cannot be interchanged.

The predictability of the system is defined in terms of the allowed maximal ignorance on the state of the system, a tolerance parameter  $\Delta_{\text{max}}$  which must be fixed according to the requirements of the observer. The predictability time is thus

$$T_p = \sup_t \{t \text{ such that } |\Delta(t')| \leq \Delta_{\text{max}} \text{ for } t' \leq t\}. \quad (6)$$

If  $\Delta_{\max} \ll 1$ , equation (5) is well approximated by (3) and the predictability time can be roughly identified with the inverse Lyapunov exponent, since

$$T_p \sim \frac{1}{\lambda} \ln \left( \frac{\Delta_{\max}}{\delta_0} \right) \tag{7}$$

and the dependence on the initial error  $\delta_0$  and on the tolerance parameter  $\Delta_{\max}$  is only logarithmic and can be safely ignored for many practical purposes.

Suppose we are now only interested in the uncertainty  $\Delta^{(\xi)}$  in the driver system. When  $\epsilon = 0$ , the Lyapunov exponent of the driver  $\lambda_\xi = 0$  so that it is fully predictable (we have in general,  $T_p^{(\xi)} \sim \delta_0^{-\beta}$ , the exponent  $\beta$  depending on the particular system) while the driven system has a Lyapunov exponent  $\lambda_0 > 0$ . However, for any  $\epsilon \neq 0$  the two subsystems are coupled, and the global Lyapunov exponent is expected to be

$$\lambda = \lambda_0 + O(\epsilon). \tag{8}$$

A direct application of (7) would give

$$T_p^{(\xi)} \sim T_p \sim \frac{1}{\lambda_0}. \tag{9}$$

One thus obtains a singular limit,  $\lim_{\epsilon \rightarrow 0} T_p^{(\xi)}(\epsilon) \neq T_p^{(\xi)}(\epsilon = 0)$ . The troubles stem from the identification between uncertainty  $\Delta(t)$  and tangent vector  $z(t)$ , which is not correct on long time scales.

It is convenient to illustrate the problem in a simpler context as the main qualitative aspects of equations (1a) and (1b) can be reproduced considering only the feedback effect in two coupled maps of the type

$$\xi_{t+1} = L\xi_t + \epsilon h(\eta_t) \tag{10a}$$

$$\eta_{t+1} = G(\eta_t) \tag{10b}$$

where the time  $t$  is an integer variable,  $\xi \in R^2$ ,  $h = (h_1, h_2)$  is a vector, function of the variable  $\eta \in R^1$  whose evolution is ruled by a chaotic one-dimensional map  $G$ , and  $L$  is the linear operator corresponding to a rotation of an arbitrary angle  $\theta$ ,

$$L = \begin{pmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{pmatrix}.$$

When  $\epsilon = 0$ , one is left with two independent systems, one of them regular and of Hamiltonian type, the other fully chaotic.

These maps provide a simple, maybe the simplest, example of a system with two different temporal regimes:

- (i) short times where  $\delta_0 \exp(\lambda_0 t) \ll 1$  so that it is correct to ignore the nonlinear terms in (5), so that  $\Delta \sim z$ ;
- (ii) long times where one should consider the full nonlinear equation (5) for the uncertainty growth.

From the observer's point of view, both these regimes might be interesting, according to his particular requirements. If one is interested in forecasting the very fine details of the systems, the tolerance threshold  $\Delta_{\max}$  could be quite small, hence  $T_p \sim \lambda_0^{-1}$ . In general, however, a system is considered unpredictable when the uncertainty is rather large (say discrimination between sun and rain in meteorology) and regime (ii) is the relevant one. In that case, nonlinear effects in (5) cannot be neglected and in order to give an analytical estimate of the predictability time we can use a stochastic model of the deterministic

equations. Indeed the chaotic feedback on the evolution of the ‘driver’ system can be simulated by a random vector  $w$ , i.e.

$$\xi_{t+1} = L\xi_t + \epsilon w_t. \quad (11)$$

The uncertainty  $\Delta_t^{(\xi)}$  is then given by the difference of two trajectories  $\xi_t$  and  $\xi_t'$  originated by nearby initial conditions and evolves according to the stochastic map

$$\Delta_{t+1}^{(\xi)} = L\Delta_t^{(\xi)} + \epsilon W_t \quad (12)$$

where  $W_t = w_t' - w_t$ . For short times  $t \ll \lambda_0^{-1} |\ln \delta_0|$  one cannot consider the ‘noises’  $w_t$  and  $w_t'$  as uncorrelated so that the uncertainty on the driver grows exponentially under the influence on the deterministic chaos given by the feedback,  $|\Delta_t^{(\xi)}| \sim |\Delta_t| \sim \exp(\lambda t)$ , with  $\lambda = \lambda_0 + O(\epsilon)$ . For long times  $t \gg \lambda_0^{-1} |\ln \delta_0|$ , the random variables are practically uncorrelated so that their difference  $W_t$  still acts as a noisy term. As a consequence the growth of the uncertainty is diffusive, since the formal solution of (12) is

$$\Delta_t^{(\xi)} = L^t \left( \delta_0 + \epsilon \sum_{\tau=0}^{t-1} L^{-\tau} W_\tau \right) \quad (13)$$

and noting that  $L^t$  is a unitary transformation, from (13) one can derive the bound

$$|\Delta_t^{(\xi)}| \leq \epsilon \left| \sum_{\tau=0}^{t-1} W_\tau \right| \sim \epsilon t^{1/2} \quad (14)$$

where we have used the estimate  $|\sum_{\tau=0}^{t-1} W_\tau| \sim t^{1/2}$  given by standard arguments borrowed from the central limit theorem. In conclusion, for our model maps (10a) and (10b), the predictability time on the driver diverges like

$$T_p^{(\xi)} \sim \epsilon^{-2} \quad (15)$$

although there is a regime of strong chaos since the total Lyapunov exponent  $\lambda = \lambda_0 + O(\epsilon)$  does not vanish with the strength of the feedback  $\epsilon$ .


It is important to stress that the particular power of the diffusive law in a realistic model can be different from that of a random walk, since the deterministic chaos of the feedback could be better represented by random variables with appropriate correlations. The qualitative behaviour exhibited by the stochastic model for the uncertainty growth (exponential followed by a power law) can be tested in a direct numerical simulation of the coupled maps (10a) and (10b), where we choose the linear vector function for the feedback,

$$h(\eta) = (\eta, \eta) \quad (16a)$$

and the logistic map at the Ulam point for the driving system,

$$G(\eta) = 4\eta(1 - \eta) \quad (16b)$$

with the Lyapunov exponent  $\lambda_0 = \ln 2$ . Figure 1 shows the behaviours of the uncertainty  $|\Delta^{(\xi)}|$  for the driver, starting with an error  $\delta_0$  on the initial condition  $\eta_0$  and no error on the driver. At the beginning both  $|\Delta^{(\eta)}|$  and  $|\Delta^{(\xi)}|$  grow exponentially. However, the phase space available to the variable  $\eta$  is finite, so that  $|\Delta^{(\eta)}|$  is bounded by a maximum value  $\Delta_M \sim O(1)$ . It will be attained at the time  $t = t^* \sim \lambda_0^{-1} \ln(\Delta_M/\delta_0)$ , when the uncertainty on the driver system is  $|\Delta^{(\xi)}| \sim \epsilon \Delta_M$ , and so much lower than the threshold. At larger times  $t > t^*$ , the uncertainty on the driven system remains practically constant and  $|\Delta^{(\xi)}|$  increases with a diffusive law of type (14) according to the mechanism described by the stochastic map (12).



**Figure 1.** Growth of the uncertainty  $|\Delta^{(\epsilon)}|$  of the driver system in the coupled maps (10a) and (10b) as a function of time  $t$ , where the rotation angle  $\theta = 0.82099$ , the feedback strength  $\epsilon = 10^{-5}$  and the error on the initial condition of the driven system (10b)  $\delta_0 = 10^{-10}$ . Broken curve: exponential regime  $\Delta^{(\epsilon)}(t) = \epsilon\delta_0 \exp(\lambda_0 t)$  where  $\lambda_0 = \ln 2$ . Full line:  $\Delta^{(\epsilon)}(t) = \epsilon t^{1/2}$ .

We have also studied a more realistic model of two coupled standard maps in action-angle variables  $I$  and  $\theta$ ,

$$\begin{aligned}
 I_{t+1}^{(1)} &= I_t^{(1)} - \epsilon \sin(\theta_t^{(1)} + \theta_t^{(2)}) \\
 \theta_{t+1}^{(1)} &= \theta_t^{(1)} + I_{t+1}^{(1)} \\
 I_{t+1}^{(2)} &= I_t^{(2)} - K \sin(\theta_t^{(1)} + \theta_t^{(2)}) \\
 \theta_{t+1}^{(2)} &= \theta_t^{(2)} + I_{t+1}^{(2)}
 \end{aligned}
 \tag{17}$$

where  $K \gg \epsilon$  is a control parameter of order unity, such that the system  $(I^{(2)}, \theta^{(2)})$  is chaotic when  $\epsilon = 0$ . We do not discuss in detail the results for these coupled maps, since they are qualitatively similar to those obtained for the simplified model (10).

We now consider an application to a physical phenomenon, the motion of an ensemble of point vortices in a fluid. It is a classical problem in fluid mechanics, formally similar to the planetary motion in a gravitational field. Both the systems are Hamiltonian with long-range interactions. The main qualitative differences are that the Hamiltonian for point vortices does not contain a kinetic term and the motion is confined on the two-dimensional plane. The phase space for a collection of  $N$  vortices thus has  $2N$  dimensions, related to the physical coordinates.

Dynamical properties of point vortex systems have been studied by several authors interested in their chaotic motion and connection with two-dimensional turbulence (see Aref 1983 for a review). The Hamiltonian theory for vortex motion inside a bounded domain

was developed many years ago (Lin 1941) for several boundaries. Here we are interested in the motion in the unitary disc  $D$  for which the Hamiltonian takes the form

$$H = -\frac{1}{4\pi} \sum_{i>j} \Gamma_i \Gamma_j \log \left[ \frac{r_i^2 + r_j^2 - 2r_i r_j \cos \theta_{ij}}{1 + r_i^2 r_j^2 - 2r_i r_j \cos \theta_{ij}} \right] + \frac{1}{4\pi} \sum_{i=1}^N \Gamma_i^2 \log(1 - r_i^2) \quad (18)$$

where the  $\Gamma_i$  represent the circulation of the  $i$ th vortex of coordinates  $\mathbf{x}_i = (x_i = r_i \cos \theta_i, y_i = r_i \sin \theta_i)$  and  $\theta_{ij} = \theta_i - \theta_j$ . The canonical conjugated variables are the scaled coordinates  $(\Gamma_i x_i, y_i)$  and the phase space is thus  $N$  times the configuration space  $D$ . The Hamiltonian (18) is invariant under rotations in the configuration space, and thus the angular momentum is a second conserved quantity

$$L^2 = \sum_{i=1}^N \Gamma_i (x_i^2 + y_i^2). \quad (19)$$

By general results of Hamiltonian mechanics, a system of two point vortices is always integrable, but we should expect chaotic motion for  $N > 2$  vortices.

In the following we will consider  $N = 3$  vortices, two of them carrying fixed circulation  $\Gamma_1 = \Gamma_2 = 1$  and representing the driver, which without feedback is integrable. The third vortex, of circulation  $\Gamma_3 = \epsilon$ , represents the driven system which now makes the total system chaotic. In the limit  $\epsilon \rightarrow 0$  the third vortex becomes a passive particle (it is passively transported by the flow generated by the two unit vortices) and does not influence the integrable motion of the two vortices as for the three-body restricted problem in celestial mechanics. The restricted system is still chaotic, but the uncertainty is confined to the passive tracer, while the motion of the two vortices is, in general, quasi-periodic. This limit is one of the simplest examples of chaotic advection in two-dimensional flow and it will be studied in detail in another paper (Boffetta et al 1996).


For our particular problem of three vortices the Hamiltonian can be rewritten in the following standard perturbation form:

$$H = H_0(\mathbf{x}_1, \mathbf{x}_2) + \epsilon H_1(\mathbf{x}_1, \mathbf{x}_2, \mathbf{x}_3) + \epsilon^2 H_2(\mathbf{x}_3). \quad (20)$$

The first term  $H_0$  describes the dynamics of the two unit vortices (and leads to integrable motion for  $\epsilon = 0$ ); the term  $H_1$  represents the interaction with the small vortex of circulation  $\epsilon$  and the last term is due to the interaction of the third vortex with its own image. The  $O(\epsilon)$  term is thus the perturbation to the integrable system  $H_0$  and we are reduced to the general framework described above if we identify  $\xi = (\mathbf{x}_1, \mathbf{x}_2)$  and  $\eta = \mathbf{x}_3$ . The only difference is that now the  $\eta$  dynamics is not chaotic by itself, but chaoticity is induced by the interaction with the integrable system  $\xi$ .

We now describe a typical simulation of error growth in the point vortex model which reproduces the effects obtained with the coupled maps model. We fix the value of the coupling constant (circulation of the third vortex)  $\epsilon = 10^{-6}$  and the initial conditions for the vortex positions are chosen in order to obtain chaotic motion with a global Lyapunov exponent  $\lambda \sim 0.041$ . The initial uncertainty on the coordinates of the small vortex is  $\Delta^{(\eta)}(0) = 10^{-3}$  while we assume we know the initial position of the two large vortices with a precision of  $\Delta^{(\xi)}(0) = 10^{-8}$ . The saturation value for the uncertainty is proportional to the disc radius, here  $\Delta_M \sim 1$ .

We let the system evolve according to the Hamiltonian dynamics (18) for quite a long time and we computed, at each time, the maximum value reached by the uncertainty (we used the maximum because in this system uncertainty shows strong oscillations: this is a memory of the quasi-periodic behaviour for  $\epsilon = 0$ ). This represents the worst situation for making predictions. The upper scatter plot in figure 2 shows the time evolution of the



**Figure 2.** Uncertainty growth for the point vortex model. Cross: maximum of the uncertainty  $\Delta^{(n)}(t)$  on the third vortex. Diamond: maximum error  $\Delta^{(\xi)}(t)$  on the driving system of two vortices. Dotted curve: exponential regime  $\Delta^{(\xi)}(t) \sim \exp(\gamma t)$  with  $\gamma = 0.064$ . Broken curve: power-law regime  $\Delta^{(\xi)}(t) \sim \epsilon t^{1/\alpha}$  with  $\alpha^{-1} = 0.88$ .

uncertainty for the driven system  $\Delta^{(n)}(t)$ . We can recognize a short ( $t < 100$ ) exponential growth until the nonlinear effect becomes important. At large times ( $t > t^* \sim 300$ ) the uncertainty saturates to its maximum value  $\Delta_M$ . The lower scatter plot represents the uncertainty for the driver system of two vortices,  $\Delta^{(\xi)}(t)$ . We can easily recognize the two expected limiting behaviours represented by the two curves. For small times, the error grows exponentially,  $\Delta^{(\xi)}(t) \sim \epsilon \gamma^t$  where  $\gamma \sim 0.064$  is close to the global Lyapunov exponent. For long times, the power-law behaviour is recovered,  $\Delta^{(\xi)}(t) \sim \epsilon t^{1/\alpha}$  with  $\alpha^{-1} \sim 0.88$ .

In conclusion, we must stress that all our results can be generalized in a straightforward way to a weakly chaotic driver with a maximum Lyapunov exponent  $\lambda_d \ll \lambda_0$ . In fact, the driver might be either conservative or dissipative. The important point is that the dynamics of the driver have a much longer characteristic time than the driven system so that, for an observer interested in the predictability problem, the two systems can be practically decoupled. The Lyapunov analysis, although mathematically correct, does not capture the physically relevant features of the phenomenon, and the exponential dependence on initial conditions does not affect the possibility of forecasting the future of the driver on a very long time scale. This is still true in systems with many different time scales instead of only two, such as fully developed turbulence, as suggested a long time ago on phenomenological grounds by Lorenz (see Lorenz 1969, Leith and Kraichnan 1972, Lilly 1973). In this case the inverse Lyapunov exponent is not related to the predictability time on the large length scale motion (Aurell *et al* 1996).

## Acknowledgments

We are grateful for financial support to INFN through *Iniziativa Specifica F13* and to Istituto Nazionale Fisica della Materia. GB thanks the ‘Istituto di Cosmogeofisica del CNR’, Torino, for hospitality.



**References**

- Aref H 1983 *Ann. Rev. Fluid Mech.* **15** 345  
Aurell E, Boffetta G, Crisanti A, Paladin G and Vulpiani A 1996 *Phys. Rev. E* at press  
Benettin G, Galgani L and Strelcyn J M 1976 *Phys. Rev. A* **14** 2338  
Boffetta G, Celani A and Franzese P 1996 *J. Phys. A: Math. Gen.* **29** at press  
Leith C E and Kraichnan R H 1972 *J. Atmos. Sci.* **29** 1041  
Lilly D K 1973 *Dynamic Meteorology* ed P Morel (Boston, MA: Reidel) p 353  
Lin C C 1941 *Proc. Natl Acad. Sci., US* **27** 570  
Lorenz E N 1963 *J. Atmos. Sci.* **20** 130  
Lorenz E N 1969 *Tellus* **21** 3  
Sussmann G J and Wisdom J 1992 *Nature* **257** 56