

Reply to comments on 'A new efficient method for calculating perturbation energies using functions which are not quadratically integrable'

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1996 J. Phys. A: Math. Gen. 29 6467

(<http://iopscience.iop.org/0305-4470/29/19/031>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.70

The article was downloaded on 02/06/2010 at 04:01

Please note that [terms and conditions apply](#).

COMMENT

## Reply to comments on ‘A new efficient method for calculating perturbation energies using functions which are not quadratically integrable’

L Skála<sup>†</sup> and J Čížek<sup>‡</sup>

<sup>†</sup> Charles University, Faculty of Mathematics and Physics, Ke Karlovu 3, 121 16 Prague 2, Czech Republic

<sup>‡</sup> University of Waterloo, Department of Applied Mathematics, Waterloo, Ontario N2L 3G1, Canada

Received 28 May 1996

**Abstract.** In a previous paper (Skála and Čížek 1996) a new approach to calculating perturbation energies of bound states based on the use of functions which are not quadratically integrable was suggested. This method has been analysed in a comment by Guardiola and Ros (1996). The present comment is the reply to this previous comment. The aim is to give a simple and straightforward proof of the method and clarify some other points important for its application.

Following [1] and [2], the Hamiltonian, wavefunction and energy in the perturbation theory are assumed in the usual forms

$$H = H_0 + \lambda H_1 \quad (1)$$

$$\psi = \psi_0 + \lambda \psi_1 + \lambda^2 \psi_2 + \dots \quad (2)$$

and

$$E = E_0 + \lambda E_1 + \lambda^2 E_2 + \dots \quad (3)$$

Using these assumptions in the Schrödinger equation  $H\psi = E\psi$ , the well known equations for  $E_n$  and  $\psi_n$  are obtained:

$$H_0\psi_0 = E_0\psi_0 \quad (4)$$

and

$$H_0\psi_n + H_1\psi_{n-1} = \sum_{i=0}^n E_i\psi_{n-i} \quad n = 1, 2, \dots \quad (5)$$

The perturbation energies  $E_n$  in this equation are determined from the corresponding boundary conditions.

Assuming the validity of the normalization and orthogonality conditions

$$\langle \psi_0 | \psi_n \rangle = \delta_{0,n} \quad n = 0, 1, \dots \quad (6)$$

the well known formula for  $E_n$

$$E_n = \langle \psi_0 | H_1 | \psi_{n-1} \rangle \quad n = 1, 2, \dots \quad (7)$$

can be easily derived.

In [1], we suggested another formula for the perturbation energy

$$E_n = -\frac{\psi_n(0, x_0)}{F(x_0)} \quad n = 1, 2, \dots \quad (8)$$

in which the validity of equation (6) is not assumed. Here,  $\psi_n(E_n, x)$  denotes the solution of equation (5) for the energy  $E_n$  and  $x_0$  is the point where the boundary conditions

$$\psi_n(x_0) = 0 \quad n = 0, 1, \dots \quad (9)$$

are applied.  $F(x)$  is a universal function which can be calculated from the equation

$$F(x) = \psi_1(1, x) - \psi_1(0, x). \quad (10)$$

We note that  $E_n$  in  $\psi_n(E_n, x)$  is not the value of the perturbation energy, but a parameter which is used during the integration of equation (5) for a given  $n$ . In consequence, functions  $\psi_n(E_n, x)$  are not in general quadratically integrable. It applies also for  $E_n = 1$  and  $E_n = 0$ .

Equation (8) is a very simple and interesting result showing that functions which are not quadratically integrable can be used for the calculation of the perturbation energies of quadratically integrable states.

In [1], detailed discussion of the validity of equations (8) and (10) was not given. This has been noted in [2] and a rather complicated proof of these equations has been elaborated. Discussion of the boundary conditions in the numerical solution of equation (5) has not been performed in [1] and [2]. The aim of this comment is to show that equations (8) and (10) follow very simply from equation (5) if these conditions are taken in the usual form. We would also like to comment on some remarks regarding the numerical properties of our method.

Now we assume that  $E_i$  and  $\psi_i(x)$ ,  $i = 0, \dots, n-1$  are the correct perturbation energies and wavefunctions.

Following [2], we denote

$$\sum_{i=1}^{n-1} E_i \psi_{n-i}(x) - H_1 \psi_{n-1}(x) = \mathcal{F}(x). \quad (11)$$

Then, equation (5) becomes

$$(H_0 - E_0)\psi_n(E_n, x) = E_n \psi_0(x) + \mathcal{F}(x) \quad n = 1, 2, \dots \quad (12)$$

Subtracting this equation for a general value of the parameter  $E_n$  and for  $E_n = 0$  we get

$$(H_0 - E_0)(\psi_n(E_n, x) - \psi_n(0, x)) = E_n \psi_0(x). \quad (13)$$

For  $E_n = 1$ , this equation yields

$$(H_0 - E_0)(\psi_n(1, x) - \psi_n(0, x)) = \psi_0(x). \quad (14)$$

Using the last expression for  $\psi_0(x)$  in equation (13) we get

$$(H_0 - E_0)(\psi_n(E_n, x) - \psi_n(0, x)) = E_n(H_0 - E_0)(\psi_n(1, x) - \psi_n(0, x)). \quad (15)$$

We see from this equation that, in general,

$$\psi_n(E_n, x) - \psi_n(0, x) = E_n(\psi_n(1, x) - \psi_n(0, x)) + G_n(E_n)\psi_0(x) \quad (16)$$

where  $G_n(E_n)$  is a function of  $E_n$ .

To determine  $G_n(E_n)$ , we assume the same boundary conditions for  $\psi_n$  as in the usual numerical solution of equation (5). We suppose that the integration of equation (5) is performed on a finite but sufficiently large interval  $(x_s, x_0)$ . Here,  $x_s$  ( $x_s < x_0$ ) is the

starting point of the integration and  $x_0$  is the final point where the boundary conditions (9) are applied. At the starting point  $x_s$ , we keep constant the value of the wavefunction  $\psi_n(E_n, x_s)$  for all the values of the parameter  $E_n$

$$\psi_n(E_n, x_s) = c_n \quad n = 0, 1, \dots \tag{17}$$

Here,  $c_n$  is a small number different from zero. To define  $\psi_n(E_n, x)$  uniquely, we make similar assumption also about its derivative

$$\left. \frac{\partial \psi_n(E_n, x)}{\partial x} \right|_{x=x_s} = d_n, \quad n = 0, 1, \dots \tag{18}$$

where  $d_n$  is a small number or zero. Values of the constants  $c_n$  and  $d_n$  are not critical. The value of the perturbation energy  $E_n$  is determined by the boundary condition (9).

Using the boundary condition (17) in equation (16) we get

$$G_n(E_n)\psi_0(E_0, x_s) = 0. \tag{19}$$

Taking into account the fact that  $\psi_0(E_0, x_s) = c_0 \neq 0$  we see that

$$G_n(E_n) = 0 \tag{20}$$

and

$$\psi_n(E_n, x) - \psi_n(0, x) = E_n(\psi_n(1, x) - \psi_n(0, x)). \tag{21}$$

Let us assume now that  $E_n$  is the perturbation energy determined by the boundary condition  $\psi_n(E_n, x_0) = 0$ . Then, equation (21) gives for  $x = x_0$

$$E_n = -\frac{\psi_n(0, x_0)}{\psi_n(1, x_0) - \psi_n(0, x_0)} \quad n = 1, 2, \dots \tag{22}$$

Writing equation (14) for general  $n$  and for  $n = 1$  we get the result similar to equation (16)

$$\psi_n(1, x) - \psi_n(0, x) = \psi_1(1, x) - \psi_1(0, x) + g_n\psi_0(x) \tag{23}$$

where  $g_n$  is a constant. It is obvious that  $g_n = 0$  for the same reason as  $G_n(E_n) = 0$  and

$$\psi_n(1, x) - \psi_n(0, x) = \psi_1(1, x) - \psi_1(0, x). \tag{24}$$

We see that the difference  $\psi_n(1, x) - \psi_n(0, x)$  is a universal function of  $x$  denoted as  $F(x)$  in [1]

$$\psi_n(1, x) - \psi_n(0, x) = F(x) \quad n = 1, 2, \dots \tag{25}$$

With this function, equation (22) can be written in the form of equation (8). Equation (11) of [1]

$$\frac{\partial \psi_n(E_n, x)}{\partial E_n} = F(x) \tag{26}$$

is a trivial consequence of equations (21) and (25).

We note that the perturbation  $H_1$  does not appear in (13). Therefore,  $F(x)$  is a universal function not only for all orders  $n = 1, 2, \dots$  but also for all the perturbations  $H_1$  assuming that the zero-order Hamiltonian  $H_0$  does not change and the boundary conditions for  $\psi_n$  have the form of equations (9), (17) and (18).

During our proof of equation (8), the validity of the normalization and orthogonality conditions (6) was not assumed. Therefore, equation (8) is more general than equation (7). It is remarkable that, in contrast to the variational methods or usual versions of the perturbation theory such as (7), no matrix elements have to be calculated in equation (8). To determine  $E_n$  from this equation, only one function value  $\psi_n(0, x_0)$  and  $F(x_0)$  must be known.

Since the validity of the normalization and orthogonality conditions (6) is not required in our method, the wavefunctions  $\psi_n$  calculated numerically from equation (5) can in general contain an arbitrary component of  $\psi_0$ . It is obvious that it leads to the renormalization of the  $\psi_0$  component in the resulting wavefunction  $\psi$ . If required, the orthogonality conditions (6) can be obeyed by the usual orthogonalization scheme.

Finally, we note that the calculation of eigenfunctions for eigenvalues taken as parameters is a part of standard algorithms for the calculation of eigenvalues of differential equations. In these algorithms, differential equations are numerically integrated and estimates of the eigenvalues are changed until the boundary conditions are obeyed. Such 'shooting' methods represent a standard numerical procedure and have no special problems with instabilities and overflows. In [1] and in this comment, we have discussed a special case of differential equations, namely, the perturbation equations. From this point of view, the remark in [2], according to which the method using functions that are not normalizable is prone to instabilities and overflows, seems to be out of place.

Finally, let us repeat [1] that the two most important applications are the strong coupling perturbation theory [3–11] as well as a rigorous mathematical theory of the upper and lower bounds [12] which is more effective than the inner projection technique [13].

## References

- [1] Skála L and Čížek J 1996 *J. Phys. A: Math. Gen.* **29** L129
- [2] Guardiola R and Ros J 1996 *J. Phys. A: Math. Gen.* **29** 6461
- [3] Turbiner A V and Ushveridze A G 1988 *J. Math. Phys.* **29** 2053
- [4] Fernández F M 1992 *Phys. Lett.* **166A** 173
- [5] Guardiola R, Solís M A and Ros J 1992 *Il Nuovo Cimento B* **107** 713
- [6] Fernández F M and Guardiola R 1993 *J. Phys. A: Math. Gen.* **26** 7169
- [7] Janke W and Kleinert H 1995 *Phys. Rev. Lett.* **75** 2787
- [8] Weniger E J 1996 *Ann. Phys., NY* **246** 133
- [9] Weniger E J Construction of the strong coupling expansion for the ground state energy of the quartic, sextic and octic anharmonic oscillator via a renormalized strong coupling expansion *Phys. Rev. Lett.* submitted
- [10] Čížek J and Skála L Strong coupling perturbation theory for quartic oscillator and large-order behaviour of the coefficients in preparation
- [11] Kapsa V, Čížek J and Skála L Strong coupling perturbation theory for the sextic and octic oscillator and large-order behaviour of the coefficients in preparation
- [12] Čížek J and Skála L Functions, which are not quadratically integrable, and mathematically rigorous upper and lower bounds for the eigenvalues of simple systems in preparation
- [13] Vinette F and Čížek J 1991 *J. Math. Phys.* **32** 3392