

Correlation length and average loop length of the fully packed loop model

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1996 J. Phys. A: Math. Gen. 29 7041

(<http://iopscience.iop.org/0305-4470/29/22/008>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.70

The article was downloaded on 02/06/2010 at 04:04

Please note that [terms and conditions apply](#).

Correlation length and average loop length of the fully packed loop model

Anton Kast†

Department of Physics, University of California at Berkeley, Berkeley, CA 94720, USA

Received 16 April 1996, in final form 9 August 1996

Abstract. The fully packed loop model of closed paths covering the honeycomb lattice is studied through its identification with the $sl_q(3)$ integrable lattice model. Some known results from the Bethe ansatz solution of this model are reviewed. The free energy, correlation length, and the ensemble average loop length are given explicitly for the many-loop phase. The results are compared with the known result for the model's surface tension. A perturbative formalism is introduced and used to verify results.

1. Review of the FPL model

The fully packed loop (FPL) model is a statistical model where the ensemble is the set of all combinations of closed paths on the honeycomb lattice that visit every vertex and do not intersect. The Boltzmann weight of such a filling set of paths is just the exponential of the number of paths, i.e. the energy of a configuration is the number of closed loops used to cover the lattice. An example of a fully packed configuration of loops on this lattice is shown in figure 1. The partition function for this model may be represented as

$$Z_{FPL}(n) = \sum_C n^{P(C)} \quad (1.1)$$

where the sum is over all C , the coverings of the vertices of the hexagonal lattice by closed nonintersecting paths, $P(C)$ is the number of paths in the covering C , and n is a generalized activity.

The partition function (1.1) is apparently the generating function for the numbers of ways to cover the hexagonal lattice by any number of closed paths. Its calculation in the thermodynamic limit is an interesting combinatorial problem.

This model is equivalent to a colouring problem solved by Baxter in 1970 [1]. The loop form of the problem was introduced by Reshetikhin [2] and by Blöte and Nienhuis [3]. It was originally studied for its interest as the low-temperature limit of the $O(n)$ vector lattice models [4, 5]. In this limit, the dimensionality of vectors n is just the activity n in equation (1.1). It has been applied to random-walk problems [6], and more recently its operator spectrum and corresponding scaling exponents have been deduced [6–8].

The FPL model is solvable because of an identification with the integrable lattice model associated to the quantum group $sl_q(3)$ [9]. This integrable model is a vertex model on the square lattice where each link of the lattice can be in one of three states, and the vertex weights are given by the R -matrix for $sl_q(3)$ as in figure 2. The R -matrix depends on a

† E-mail address: anton@physics.berkeley.edu




Figure 1. An example of a fully packed loop configuration. With periodic boundary conditions, this configuration has 11 loops.

$$\begin{array}{c}
 \text{b} \\
 | \\
 \text{i} - \text{---} - \text{j} \\
 | \\
 \text{a}
 \end{array}
 \quad = \quad R_{ab}^{ij}(\gamma, \theta)$$

$$\begin{aligned}
 &= \delta_b^i \delta_a^j \frac{\sinh(\theta)}{\sinh(\theta+\gamma)} + \\
 &\delta_a^i \delta_b^j \left[(1 - \delta_{ab}) \frac{\sinh(\gamma)}{\sinh(\theta+\gamma)} e^{\Delta_{ab}\theta} + \delta_{ab} \left(1 - \frac{\sinh(\theta)}{\sinh(\theta+\gamma)} \right) \right]
 \end{aligned}$$

Figure 2. The definition of vertex weights in the integrable lattice model associated to $sl_q(d+1)$. γ and θ are parameters of the model. States on edges are labelled by roman indices with $(d+1)$ possible values. In the formula, $\Delta_{ab} = \frac{a-b}{|a-b|} + 2\frac{b-a}{d+1}$.

deformation parameter $\gamma = \log(q)$, as well as a spectral parameter θ typical of integrable theories. Denoting the $sl_q(3)$ partition function as $Z_{sl_q(3)}(\gamma, \theta)$, the precise identification is

$$Z_{FPL}(e^\gamma + e^{-\gamma}) = (e^\gamma - e^{-\gamma})^{-N} \lim_{\theta \rightarrow -\gamma} (e^\theta e^\gamma - e^{-\theta} e^{-\gamma}) Z_{sl_q(3)}(\gamma, \theta) \quad (1.2)$$

where N is the volume of the lattice (the number of hexagonal faces). Since the model is integrable, much exact information can be derived. In particular, the model's Bethe equations have been constructed and solved.

One of the more important results that have been derived in this way is the existence of a phase transition in the model (1.1) at $n = 2$ [1, 2]. At larger n , larger numbers of loops are favoured and at smaller n configurations with fewer loops are favoured. It has been conjectured that this transition is between a large- n phase where the average loop length is finite and a small- n phase where this average is infinite.

In section 3 a simple relation between the free energy of the FPL model and the ensemble average length of loops is derived. From the known solution to the Bethe equations of the $sl_q(3)$ integrable lattice model, the free energy is identified and used to graph the exact value of the average loop length as a function of n .

Identifying $n = e^\gamma + e^{-\gamma}$, it becomes apparent that $n > 2$ corresponds to the integrable model for γ real and $0 < n < 2$ corresponds to γ purely imaginary. The former phase is known [9] to be massive, in the sense that there is a gap in the spectrum of eigenvalues of the transfer matrix between the leading eigenvalue and the next-leading eigenvalue. By standard arguments [11], this implies a finite correlation length. The gap tends to zero as γ goes to zero, showing that $n = 2$ is a critical point of the model (1.1).

In section 4 this correlation length is studied by considering the spectrum of eigenvalues of this transfer matrix. The spectrum may be deduced directly from the model's Bethe equations. In the case that the transfer matrix is symmetric and therefore has real eigenvalues, the correlation length is related to the maximum eigenvalue Λ_{\max} and next-

leading eigenvalue Λ_1 by

$$\xi^{-1} = \log \frac{\Lambda_{\max}}{\Lambda_1}. \tag{1.3}$$

2. Review of the $sl_q(3)$ integrable model

The R -matrix of the $sl_q(3)$ quantum group is a $3^2 \times 3^2$ matrix that may be interpreted as a matrix of Boltzmann weights of a vertex model on the square lattice as shown in figure 2. Since this matrix satisfies the Yang–Baxter equation, the associated transfer matrix commutes with itself evaluated at differing values of the spectral parameter and the model is exactly solvable by a recursive set of two nested Bethe ansätze [9].

Eigenvalues of the transfer matrix in finite size N are parametrized by roots λ_j^k , $k = 1, 2$ and $j = 1 \dots p_k$ of a system of Bethe equations. The eigenvalue $\Lambda_N^{p_1 p_2}$ is given in terms of these roots by

$$\Lambda_N^{p_1, p_2} = \prod_{j=1}^{p_1} \frac{\sinh(i\lambda_j^1 - \theta + \frac{1}{2}\gamma)}{\sinh(i\lambda_j^1 - \theta - \frac{1}{2}\gamma)} + \left[\frac{\sinh(\theta)}{\sinh(\theta + \gamma)} \right]^N \prod_{j=1}^{p_1} \frac{\sinh(i\lambda_j^1 - \theta - \frac{3}{2}\gamma)}{\sinh(i\lambda_j^1 - \theta - \frac{1}{2}\gamma)} \Lambda_{p_1}^{p_2} \tag{2.1}$$

$$\Lambda_{p_1}^{p_2} = \prod_{j=1}^{p_2} \frac{\sinh(i\lambda_j^2 - \theta)}{\sinh(i\lambda_j^2 - \theta - \gamma)} + \prod_{j=1}^{p_1} \frac{\sinh(i\lambda_j^1 - \theta - \frac{1}{2}\gamma)}{\sinh(i\lambda_j^1 - \theta - \frac{3}{2}\gamma)} \prod_{j=1}^{p_2} \frac{\sinh(i\lambda_j^2 - \theta - 2\gamma)}{\sinh(i\lambda_j^2 - \theta - \gamma)} \tag{2.2}$$

where $\Lambda_{p_2}^{p_1}$ is a convenient intermediate symbol. In the thermodynamic limit the roots are described by distributions $\rho^k(\lambda)$ on the interval $[-\pi/2, \pi/2]$. The distribution corresponding to the maximum eigenvalue and which yields the free energy is

$$\rho_{\max}^k(\lambda) = \sum_{m=-\infty}^{\infty} e^{2im\lambda} \frac{1}{\pi} \frac{\sinh[(3-k)m\gamma]}{\sinh[3m\gamma]}. \tag{2.3}$$

For eigenvalues near the maximum eigenvalue, the changes in this distribution are parametrized by the locations of holes θ_h^q , $q = 1, 2$, $h = 1 \dots N_q$, according to

$$\rho^1(\lambda) - \rho_{\max}^1(\lambda) = \sum_{m=-\infty}^{\infty} e^{2im\lambda} \sum_{q=1}^2 \left(-\frac{1}{\pi} \right) e^{|m\gamma|} \frac{\sinh[(3-q)m\gamma]}{\sinh[3m\gamma]} \sum_{h=1}^{N_q} e^{-2im\theta_h^q} \tag{2.4}$$

$$\rho^2(\lambda) - \rho_{\max}^2(\lambda) = \sum_{m=-\infty}^{\infty} e^{2im\lambda} \sum_{q=1}^2 \left(-\frac{1}{\pi} \right) e^{|m\gamma|} \frac{\sinh[qm\gamma]}{\sinh[3m\gamma]} \sum_{h=1}^{N_q} e^{-2im\theta_h^q}. \tag{2.5}$$

The numbers of holes are constrained to satisfy the relation,

$$\sum_{q=1}^2 qN_q = \text{multiple of } 3. \tag{2.6}$$

In the range of parameters $\theta > -\gamma/2$ the bracketed quantity in equation (2.1) vanishes in the thermodynamic limit. In this case the formulae (2.1), (2.4) may be combined to yield a compact formula for eigenvalues,

$$\log \Lambda = \int_{-\pi/2}^{\pi/2} \rho^1(\lambda) \log \left[\frac{\sinh(i\lambda - \theta + \frac{1}{2}\gamma)}{\sinh(i\lambda - \theta - \frac{1}{2}\gamma)} \right] d\lambda. \tag{2.7}$$

3. Free energy and average loop length

The formulae of the preceding section in the case $\theta = -\gamma$ yield directly the free energy density of the model (1.1) for the $n > 2$ phase as the logarithm of the maximum eigenvalue of the transfer matrix, rescaled by the factor of equation (1.2). This free energy was actually derived in 1970 by Baxter [1] as the solution to a weighted three-colouring problem on the honeycomb lattice. The free energy density of the FPL model in the $n > 2$ phase is

$$\begin{aligned} F_{FPL}(n) &\equiv \lim_{N \rightarrow \infty} \frac{1}{N} \log Z_{FPL}(n) \\ &= \log \left\{ q^{1/3} \prod_{m=1}^{\infty} \frac{(1 - q^{-6m+2})^2}{(1 - q^{-6m+4})(1 - q^{-6m})} \right\} \end{aligned} \quad (3.1)$$

where $n = q + q^{-1}$, and $q = e^\gamma > 1$. This function has an essential singularity at $q = 1$. The free energy for $n < 2$ and with periodic boundary conditions is given in integral form in [6].

It is interesting to note that for both phases, the free energy density gives the ensemble average length of loops. Since a configuration C on a lattice of N faces has $2N$ occupied links, the total length of loops is always $2N$. The average loop length of configuration C is therefore $2N/P(C)$. If we define the ensemble average loop length $L_N(n)$ by

$$L_N(n) = \frac{1}{Z_{FPL}(n)} \sum_C \frac{2N}{P(C)} n^{P(C)} \quad (3.2)$$

then from inspection of equation (1.1) it is clear that

$$\frac{d}{dn} [L_N(n) Z_{FPL}(n)] = \frac{2N}{n} Z_{FPL}(n). \quad (3.3)$$

The general solution to this equation can be written up to quadrature by direct integration:

$$L_N(n) = \frac{1}{Z_{FPL}(n)} \int_C^n \frac{2N}{n'} Z_{FPL}(n') dn' \quad (3.4)$$

where the lower limit of integration is an undetermined constant. In terms of the free energy density $F_{FPL} = (1/N) \log Z_{FPL}$, this becomes

$$L_N(n) = 2N e^{-NF_{FPL}(n)} \int_C^n \frac{e^{NF_{FPL}(n')}}{n'} dn'. \quad (3.5)$$

The integral in equation (3.5) can be evaluated by steepest descent. The result is

$$L_N(n) = \frac{2N}{n} \exp \left(-Nn \frac{dF_{FPL}}{dn}(n) \right) \left[\frac{\exp \left[Nn' \frac{dF_{FPL}}{dn}(n') \right]}{N \frac{dF_{FPL}}{dn}(n')} \right]_C^n. \quad (3.6)$$

The constant of integration may now be determined from the known value of $L_N(n)$ at $n \rightarrow \infty$. As will be shown in section 5, in this limit $Z_N(n) \simeq 3n^{N/3}$, $\frac{dF_{FPL}}{dn}(n) \simeq 1/3n$, and $L_N(n) = 6$. These imply that $C = -\infty$, so in the thermodynamic limit

$$L_N(n) = \frac{2}{n \frac{dF_{FPL}}{dn}}. \quad (3.7)$$

In this calculation we have neglected corrections of the order of $1/N$ to $L_N(n)$.

A graph of the ensemble average loop length versus n in the large- n phase is shown in figure 3. This verifies the conjecture of [2] that loop length diverges at the critical point.




Figure 3. This is a graph of the average loop length of the FPL model versus n , the fugacity of loops. The critical point is at $n = 2$.

4. Correlation length

To obtain the correlation length, we must compute the expression (2.7) for the minimal hole distribution. When $d = 2$, there are two choices for the numbers N_q which parametrize excited states in equation (2.4). Either $N_1 = 3$ and $N_2 = 0$, or $N_1 = 1$ and $N_2 = 1$. In each case, the eigenvalue gap is minimized for holes at $\theta_h^q = \frac{\pi}{2}$ where the sum in equation (2.4) after integration in (2.7) is oscillatory. The transfer matrix of the model is symmetric at the point $\theta = -\frac{1}{2}\gamma$, and its eigenvalues are then real. After setting θ to this value, the correlation length of the model is given by equation (1.3).

Considering the case $N_1 = 3$ and $N_2 = 0$, we denote the next-leading eigenvalue for this hole distribution as Λ_{30} . The equation (2.7) together with the formula for densities (2.4) gives

$$\log \frac{\Lambda_{30}}{\Lambda_{\max}} = \sum_{m=-\infty}^{\infty} \phi_m \left(-\frac{1}{\pi}\right) e^{|m\gamma|} 3(-1)^m \frac{\sinh(2m\gamma)}{\sinh(3m\gamma)} \tag{4.1}$$

where ϕ_m are the integrals over roots,

$$\phi_m \equiv \int_{-\pi/2}^{\pi/2} e^{2im\lambda} \log \left[\frac{\sinh(i\lambda + \frac{1}{2}\gamma - \theta)}{\sinh(i\lambda - \frac{1}{2}\gamma - \theta)} \right] d\lambda. \tag{4.2}$$

The integral in equation (4.2) can easily be performed by contour integration. After introducing the variables $q = e^\gamma$ and $z = e^\theta$, the result for $-\frac{1}{2}\gamma < \theta < 0$ is

$$\int_{-\pi/2}^{\pi/2} e^{2im\lambda} \log \left[\frac{\sinh(i\lambda + \frac{1}{2}\gamma - \theta)}{\sinh(i\lambda - \frac{1}{2}\gamma - \theta)} \right] d\lambda = \begin{cases} \frac{\pi}{m} [1 - (z^2 q^{-1})^m] & m > 0 \\ -2\pi \log z & m = 0 \\ \frac{\pi}{m} [1 - (z^2 q)^m] & m < 0. \end{cases} \tag{4.3}$$

Substituting this result into equation (4.1) gives

$$\log \frac{\Lambda_{30}}{\Lambda_{\max}} = 2 \log z + 3 \sum_{m>0} \frac{(-1)^m}{m} (z^{2m} - z^{-2m}) \left(\frac{q^{2m} - q^{-2m}}{q^{3m} - q^{-3m}} \right). \tag{4.4}$$

After expanding the demoninator of the summand in a power series in q^{-1} , this may be resummed to the form,

$$\log \frac{\Lambda_{30}}{\Lambda_{\max}} = 2 \log z - 3 \sum_{m \geq 0} \log \left[\frac{(1 + z^2 q^{-1} q^{-6m})(1 + z^{-2} q^{-5} q^{-6m})}{(1 + z^{-2} q^{-1} q^{-6m})(1 + z^2 q^{-5} q^{-6m})} \right]. \quad (4.5)$$

This form is now convergent at the symmetric point, $\theta = -\frac{1}{2}\gamma$ or equivalently $z^2 = q^{-1}$. We may therefore evaluate it there to obtain the correlation length according to equation (1.3),

$$\xi^{-1} = 3 \log \left\{ q^{1/3} \prod_{m > 0} \frac{(1 + q^2 q^{-6m})(1 + q^4 q^{-6m})}{(1 + q^{-6m})(1 + q^6 q^{-6m})} \right\}. \quad (4.6)$$

This is the desired result, the correlation length of the FPL model where $n = q + q^{-1}$ and $q > 1$, or equivalently $q = +\sqrt{n^2 - 4}$.

The other possible choice of holes, $N_1 = 1$ and $N_2 = 1$ may be computed in the same way to give

$$\log \frac{\Lambda_{\max}}{\Lambda_{11}} = \log \left\{ q \prod_{m > 0} \frac{(1 + q^2 q^{-6m})(1 + q^3 q^{-6m})(1 + q^3 q^{-6m})(1 + q^4 q^{-6m})}{(1 + q^{-6m})(1 + q q^{-6m})(1 + q^5 q^{-6m})(1 + q^6 q^{-6m})} \right\}. \quad (4.7)$$

This quantity is greater than (4.6) for all $q > 1$, so it is not the inverse correlation length. For large q , the inequality may be seen by considering the limiting forms of expressions (4.6) and (4.7). Rigorously, the multiplicands in (4.7) may be seen to be greater than those in (4.6) term by term in m .

5. Perturbative analysis

The FPL model has a natural large- n expansion which allows simple perturbative verifications of results.

When n is large, the dominant configurations are those with large numbers of loops. The perturbative procedure is to approximate the sum over states by including the configurations with the highest numbers of loops.

On a hexagonal lattice with number of faces N a multiple of three, there are three configurations with the maximum possible number of loops. In these states, one out of every three faces has a small loop around it and these small loops lie on a triangular lattice. A sample is shown in figure 4. These three configurations differ by translations and each has $N/3$ loops.

The smallest change in the number of loops that can be made is to introduce a defect somewhere in one of the maximal configurations, as shown in figure 5. There are $2N/3$ different such defects that can be introduced and each reduces the number of loops by two.

Introducing defects in this way, we can reach all possible configurations. To see that this is so, we can represent a configuration by labelling the links on the lattice that do not contain part of a path. One of every three links is unoccupied, and every vertex touches one unoccupied link. These unoccupied links form a dimer configuration for the vertices of the lattice. If we draw rhombuses around every dimer and interpret the resulting picture as the projection of the edges of a stack of cubes, we see that a FPL configuration is equivalent to a stack of cubes. Such an identification is shown in figure 6.

In this new representation, the action of inserting a defect is just the action of adding or removing a cube. This identification is exhibited in figure 7. The result then follows that since every stack of cubes can be made by adding or removing cubes, every FPL configuration can be made from one of the maximal ones by inserting some combination of defects.




Figure 4. A sample from a configuration with the maximum number of loops.




Figure 5. A configuration with two fewer than the maximum number of loops.




Figure 6. An example of the identification of FPL configurations and stacks of cubes. One rhombus is drawn centred on each unoccupied link.




Figure 7. In the cube representation, introducing a defect is adding or removing a cube.

To obtain an approximation for the free energy, consider first the maximal state shown in figure 4. For a lattice of N faces, this configuration has $(N/3)$ loops. There are three such configurations corresponding to the three-fold translational degeneracy of the state. To lowest order then $Z_{FPL} = 3n^{N/3}[1 + O(n^{-1})]$. This result was used in section 3 to determine the asymptotics of the average loop length.

Allowing defects, there are $(2N/3)$ locations for a defect and each defect reduces the number of loops by two. Defects may be applied in any number and in any combination, giving the usual sum over disconnected diagrams. We can write this as the exponential of the connected diagram (one defect) and we will be correct except for the effects of excluded volumes which begin with two-defect connected diagrams and are therefore higher order. To the next order, $Z_{FPL} = 3n^{N/3} \exp[(2N/3)n^{-2}] \exp[O(n^{-4})]$.

Perturbatively calculating the FPL free energy, we see that

$$F_{FPL}(n) = \frac{1}{3} \log(n) + \frac{2n^{-2}}{3} + O(n^{-4}) \tag{5.1}$$

in conformity with Baxter's result shown in equation (3.1).

This point of view incidentally leads to a simple expression for the entropy density of the FPL configurations at $n = 1$. At this point, all configurations are weighted equally and Z_{FPL} is just the number of configurations, or the exponential of the entropy. Then calculating the partition function is just the problem of counting the number of coverings of the honeycomb lattice by paths, which is the number of different possible stacks of

cubes, which is the old combinatorial problem of counting plane partitions. Elser [12] has calculated the asymptotics of plane partitions for large arrays of numbers.

The result applied to this case is entirely dependent on the shape of the boundary, even in the thermodynamic limit. While any FPL configuration is equivalent to a dimer configuration in the bulk, at boundaries there may be vertices with less than three neighbours and so it is possible for an FPL configuration to have no corresponding dimer configuration. For a lattice of N faces and the boundary condition that configurations must be equivalent to a dimer configuration, the maximum entropy is obtained for a hexagon-shaped boundary and in that case the partition function is asymptotically

$$Z_{FPL}(1) = \exp[N(\frac{3}{2} \log 3 - 2 \log 2)]. \quad (5.2)$$

6. Comparison with surface tension

The ground state of the $sl_q(d)$ integrable lattice model is $(d + 1)$ -fold degenerate. This implies the existence of a notion of interfacial tension $S(\gamma)$ away from the critical point between regions of differing antiferromagnetic polarization. By considering finite-size corrections, de Vega [10] has derived transcendental equations for this interfacial tension.

The asymptotic behaviour of the interfacial tension for $\gamma \rightarrow \infty$ was extracted by deVega, and the result is

$$S(\gamma) = \frac{d}{d+1} \gamma + O(1). \quad (6.1)$$

In the case of the FPL model, $d = 2$, $n = e^\gamma + e^{-\gamma}$, and

$$S(n) = \frac{2}{3} \log(n) + O(1). \quad (6.2)$$

This result may be compared with a perturbative calculation. Consider the sum over FPL states at large- n with the constraint that boundary conditions are fixed to cause frustration in the bulk, as in figure 8. The configuration in that figure has the maximum number of loops possible and is the analogue of the configuration shown in figure 4. Denoting the sum over defects in this configuration by Z'_{FPL} , the interfacial tension is defined to be the change in free energy per unit length of the interface:

$$\frac{Z'_{FPL}}{Z_{FPL}} \sim e^{-LS} \quad (6.3)$$

where L is the vertical size of the lattice.

For a lattice of N faces, the maximum number of loops possible in the presence of the constraint is $(N/3) - (2L/3)$ instead of $(N/3)$. The maximal state in the presence of the constraint is now $3 \times 2^{2L/3}$ -fold degenerate, because there are $(2L/3)$ locations near the




Figure 8. An interface separating two regions of differing polarization.

interface where defects may be freely introduced without changing the number of loops. To lowest order therefore,

$$Z_{FPL} \simeq 3n^{N/3} \quad (6.4)$$

$$Z'_{FPL} \simeq 3 \times 2^{2L/3} n^{(N-2L)/3}. \quad (6.5)$$

Reading off the exponents, we have from equation (6.3) the result that

$$S(\gamma) = \frac{2}{3} \log(n) + O(1). \quad (6.6)$$

Equation (6.6) is apparently consistent with the large- γ asymptotics derived in [10].

Acknowledgment

The author is grateful to Professor Nikolai Reshetikhin for many helpful conversations.

References

- [1] Baxter R H 1970 *J. Math. Phys.* **11** 784
- [2] Reshetikhin N Yu 1991 *J. Phys. A: Math. Gen.* **24** 2387
- [3] Blöte H W J and Nienhuis B 1994 *Phys. Rev. Lett.* **72** 1372
- [4] Nienhuis B 1982 *Phys. Rev. Lett.* **49** 1062
- [5] Domany E, Mukamel D, Nienhuis B and Schwimmer A 1981 *Nucl. Phys. B* **190** 279
- [6] Batchelor M T, Suzuki J and Yung C M 1994 *Phys. Rev. Lett.* **73** 2646
- [7] Kondev J and Henley C L 1994 *Phys. Rev. Lett.* **73** 2786
- [8] Kondev J, de Gier J and Nienhuis B 1996 *J. Phys. A: Math. Gen.* submitted
- [9] Babelon O, de Vega H J and Viallet C M 1982 *Nucl. Phys. B* **200** 266; 1983 *Nucl. Phys. B* **220** 283
- [10] de Vega H J 1987 *J. Phys. A: Math. Gen.* **20** 6023
- [11] Baxter R J 1982 *Exactly Solved Models in Statistical Mechanics* (New York: Academic)
- [12] Elser V 1984 *PhD Thesis* University of California, Berkeley