

Temperature evolution and bifurcations of metastable states in mean-field spin glasses, with connections with structural glasses

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1997 J. Phys. A: Math. Gen. 30 5593

(<http://iopscience.iop.org/0305-4470/30/16/006>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.108

The article was downloaded on 02/06/2010 at 05:50

Please note that [terms and conditions apply](#).

Temperature evolution and bifurcations of metastable states in mean-field spin glasses, with connections with structural glasses

Alain Barrat^{†§}, Silvio Franz^{†||} and Giorgio Parisi^{‡¶}

[†] International Center for Theoretical Physics, Strada Costiera 11, PO Box 586, 34100 Trieste, Italy

[‡] Università di Roma 'La Sapienza', Piazzale A Moro 2, 00185 Rome, Italy

Received 19 March 1997

Abstract. The correlations of the free-energy landscape of mean-field spin glasses at different temperatures are investigated, concentrating on models with a first-order freezing transition. Using a 'potential function' we follow the metastable states of the model in temperature, and discuss the possibility of level crossing (which we do not find) and multifurcation (which we find). The dynamics at a given temperature starting from an equilibrium configuration at a different temperature is also discussed. In the presence of multifurcation, we find that the equilibrium is never achieved, leading to an aging behaviour at slower energy levels than usual aging. The relevance of the observed mechanisms for real structural glasses is discussed, and some numerical simulations of a soft sphere model of glass are presented.

1. Introduction

Free-energy landscapes in high-dimensional spaces have been used for a long time as metaphors for describing the physics of complex systems as glasses and spin glasses, and also proteins and evolutionary fitness landscapes [1]. The basic idea of this approach is that complex system dynamics can be viewed as a search for optima in a rough hypersurface. Although free-energy surfaces can in principle be defined for large classes of finite-dimensional models, the actual construction of such functions has been achieved only in the case of long-range disordered systems (mean-field spin glasses), which relevance for finite-dimensional spin glasses has been a subject of a long debate, growing evidence [2] pointing in the direction that mean-field theory is a good starting point to describe finite-dimensional physics. Thouless *et al* [3] (TAP) showed that stable and metastable states of long-range spin glasses are associated to minima of a suitable free energy, which is a random function in a N ($\rightarrow \infty$) dimensional space. The complex phenomenology of equilibrium spin glasses can be read as a set of propositions about the low-lying minima of the TAP free energy, and the barriers separating them.

Due to the random character of the TAP free-energy function, analytic statements on the structure of the stationary point have an inescapable statistical character. Various techniques have been invented to study the minima of TAP free energy, and the structure of the stable

[§] E-mail address: barrat@ictp.trieste.it

^{||} E-mail address: franz@ictp.trieste.it

[¶] E-mail address: parisi@roma1.infn.it

and metastable minima for the fixed external parameter is known in great detail, and gives a coherent picture of the glassy transition. Two classes of models are known, according to the order of the freezing transition. In models such as the Sherrington–Kirkpatrick models, which display a second-order phase transition, the metastable states of the TAP free energy do not play an important physical role. A second-order transition suggests a mechanism of bifurcation (or rather multifurcation) of the paramagnetic minimum as T_c is crossed. In off-equilibrium dynamics it is found that all the extensive quantities tend to their equilibrium values for large times. Whereas these mean-field models, with continuous transition, seem to apply for the description of real spin glasses, a second class of models, such as, for example the Potts glass, show a first-order freezing transition, and seem closer to describing the physics of structural glasses. Indeed, for these models a purely dynamical transition occurs where the relaxation time diverges, while the static thermodynamic quantities show singularities only at a lower temperature. Below the dynamical transition temperature, metastable states dominate the physics, and, dynamically, the extensive quantities do not tend to their equilibrium values if a random initial condition is chosen. Statically, the partition function is dominated by metastable states (between the statical and the dynamical transitions, by an exponentially large number of mutually inaccessible states) that the system is unable to dynamically reach.

The scenario in which the barriers between metastable states are infinite, and where a quenched system never reaches any of these states, is clearly linked to the mean-field approximation. For finite systems, metastable states have a finite lifetime, and the system should be able to find them in a finite time. The time, and the states the system is able to find, can depend, for example on the cooling rate. A modified scenario would include ‘activated processes’ and suppress the divergence of relaxation times at the dynamical transition, replacing it by a rapid increase (with divergence only at the static transition).

This picture, as was already advocated in [4–6] and more recently in [7], could be relevant for real glasses: indeed, the glass transition temperature is also a purely dynamical quantity, defined by the fact that the relaxation time reaches a certain value, the existence of a static transition at a lower temperature still being a subject of debate. Below this temperature, the system remains out of equilibrium for all available timescales, and static quantities are not reached. In the same way, the aforementioned scenario would yield a glass transition (corresponding to a large but finite value of the relaxation time) occurring above the static transition, and a dynamical evolution resulting from a mixture of mean-field-like dynamics and activated processes.

In this context, the relevance of mean-field studies depends on the various timescales involved: if the barriers between metastable states are low, activated processes are fast and will dominate the evolution; if in contrast, the energy barriers are finite but large, there will exist time windows in which the mean-field scenario will hold.

To address this question, we therefore have to gain knowledge of the metastable states, both statically and dynamically, for the mean-field models, and to compare the emerging picture and dynamical scenarios with the real world, or at least with numerical simulations.

If the structure of the TAP minima for a fixed temperature is rather well known [8, 9], a much less coherent picture is available for the correlation of the free-energy landscapes for different temperatures. In the Sherrington–Kirkpatrick model, the study in [10, 11] revealed ‘chaotic temperature dependence’ of the low-lying states. States of equilibrium at different temperatures are, no matter how small the temperature differences, as uncorrelated as they can. In some other models the chaotic property is absent. A clear example is the spherical p -spin model, where the homogeneity of the Hamiltonian implies that the order of the free-energy minima does not depend on temperature, so that, in the whole low-temperature

phase, the statics is given by the same low-lying states. Some general conclusions about the fate of TAP minima under temperature changes can be drawn on the basis of the smoothness of the TAP free energy as a function of temperature. For example, absolutely stable minima cannot disappear or multifurcate for an infinitesimal change of the temperature, and the generation of new stationary points has to pass by a marginally stable situation.

Two issues appear to be relevant for the description of correlations of the landscapes for different temperatures: level bifurcations and level crossing. Therefore, in this paper we try to gain some generic insights into these topics, by addressing the issue of following the TAP states in temperature for a spherical model which displays first-order glassy transition. Different from the p -spin model, the Hamiltonian does not have the homogeneity which prevents the chaoticity with respect to temperature changes. We therefore expect that its behaviour is generic in the class of models with first-order transition. The analysis is performed with the aid of a recent method where the metastable states are associated to local minima of some macroscopic ‘potential’ function of the spin-glass order parameter. The basic idea of this approach is that the free-energy manifold can be probed introducing an external field pointing in the direction of some typical equilibrium configuration [12–14]. In section 2, we review the construction of the potential of [12] and, using a more general ansatz [15], we extend the discussion to some properties not mentioned there. In particular, the extension of the ansatz allows us to discuss, in section 3, the possibility of multifurcation of the metastable states that we follow in temperature, a possibility not considered in [12].

In section 4, we use another powerful approach, a dynamical one, with appropriate initial conditions: the dynamics of a system thermalized at a certain temperature, and then brought at another, also allows us to explore the phase space of the system [12, 16]. We show that the two methods yield the same results, as was suggested, but not proved for the general case, in [12, 16], and use moreover this dynamical study to tackle another relevant issue: the dynamical behaviour of the systems when a TAP solution bifurcates†. In particular, the problem of whether the system will fall into one of the new valleys or will be unable to decide where to fall and age forever.

After having described these mechanisms for the considered mean-field models, in section 5 we tentatively compare them with the case of real glasses, via numerical simulations of a soft sphere glass. In particular, the dynamical mean-field approach of thermalized initial conditions can be thought of as a previous very slow cooling to a certain temperature, followed by a rapid change. The study of the energy reached with various cooling rates in the simulated system shows the relevance of the mean-field scenario, in the available time window.

2. Studied models; the potential; previous results

The class of models we consider is defined by N (real) spins $\mathbf{s} = \{s_1, \dots, s_N\}$ interacting through a Hamiltonian $H(\mathbf{s})$ and a global (spherical) constraint $\sum_i s_i^2 = N$. The Hamiltonian is random, Gaussian, with correlations

$$\overline{H(\mathbf{s})H(\mathbf{s}')} = Nf(q_{\mathbf{s}\mathbf{s}'}) \quad (1)$$

where $q_{\mathbf{s}\mathbf{s}'} = 1/N \sum_i s_i s'_i$ is the overlap between the configurations \mathbf{s} and \mathbf{s}' . If f is a polynomial function, the Hamiltonian can be presented as a linear combination of terms of

† In [12], the ansatz used did not allow us to see the multifurcation, and the TAP solution seemed instead to disappear, leaving the issue open of the correspondence between the dynamical and the statical approaches.

the type

$$H_p(\mathbf{s}) = - \sum_{1 \leq i_1 < i_2 < \dots < i_p \leq N} J_{i_1 i_2 \dots i_p} s_{i_1} s_{i_2} \dots s_{i_p} \quad (2)$$

with Gaussian-independent couplings $J_{i_1 i_2 \dots i_p}$, with zero mean and variance $p!/(2N^{p-1})$. It is easy to see that $\overline{H_p(\mathbf{s})H_{p'}(\mathbf{s}')} = \delta_{p,p'} q_{ss'}^p/2$. As we will see in the following, the purely monomial case, the so-called p -spin model, has remarkably simple properties under temperature changes [9] thanks to the homogeneity of the Hamiltonian under contemporary rescaling of all the variables. In order to study the generic behaviour it is therefore necessary to consider the inhomogeneous Hamiltonian, giving rise to non-monomial correlation functions. The specific form of the function $f(q)$ we will use in our examples will be mainly $f(q) = \frac{1}{2}(q^3 + q^4)$, however the results will be generic for the inhomogeneous Hamiltonian verifying the condition that $f''(q)^{-3/2} f'''(q)$ is monotonically decreasing with q for all q (this ensures that the transition is a discontinuous one).

In that case the statics of the model is described by:

- a high temperature phase, for $T > T_d$, where the dominant contribution is given by a paramagnetic state;
- a temperature range $T_d > T > T_s$ where the replica calculations yield a replica-symmetric result, which in fact corresponds to ergodicity breaking in an exponentially large number of states (finite complexity);
- a low-temperature phase, for $T_s > T$, with a one-step replica-symmetry breaking, corresponding to the predominance of the lowest TAP states, with zero complexity.

The relaxation dynamics from a random initial state, yields equilibrium dynamics in the paramagnetic state for $T > T_d$, while, for $T_d > T$, the aging phenomena appears [17] and the long time limit of the energy per spin is higher than the equilibrium value.

In this section we review the construction of the potential function [12], and we expose some new results coming from a replica-symmetry breaking ansatz [15], which clarify some of the ‘mysteries’ left open in [12].

2.1. Construction of the potential

One of the characteristics of spin glasses, due to their random character, is that the different equilibrium states cannot be selected by an external field uncorrelated with the landscape defined by the Hamiltonian. The basic idea underlying the potential function is to use an external field pointing in the direction of a particular equilibrium configuration [13, 12]. So, if σ_i denotes a typical equilibrium configuration at a temperature T' one can define the partition function

$$Z[T, \epsilon, \sigma] = \sum_{\mathbf{S}} e^{-\beta H[\mathbf{S}] + \epsilon \mathbf{S} \sigma}. \quad (3)$$

Besides being self-averaging with respect to the distribution of the quenched Hamiltonian, the free energy $\Gamma[T, T', \epsilon] = -T/N \log Z[T, \epsilon, \sigma]$ is also independent on the particular configuration σ we choose, and therefore coincides with its average over the probability distribution $\exp(-\beta' H[\sigma])/Z[T']$. We define the ‘potential’ as the Legendre transform of Γ :

$$V[T, T', q_{12}] = \min_{\epsilon} \Gamma[T, T', \epsilon] + \epsilon q_{12} - F[T]. \quad (4)$$

From the Legendre transform we have subtracted F the free energy at temperature T in order to have $V[T, T', 0] = 0$. Defined in this way, the potential has the meaning of the

free energy cost to keep a system at temperature T at a fixed overlap q_{12} from a generic configuration of equilibrium at a different temperature T' .

V is also self-averaging with respect to the quenched disorder distribution, which we denote by an overline. The basic object we need to evaluate is then

$$\overline{\frac{1}{Z[T']} \sum_{\sigma} e^{-\beta' H[\sigma]} \log \left(\sum_S e^{-\beta H[S] + \epsilon S \sigma} \right)}. \quad (5)$$

It is explained in detail in [12] that in order to perform the averages it is possible to use a double analytic continuation from integer values of the parameters m and n , used to represent $\log Z[T, \epsilon, \sigma]$ as $\lim_{m \rightarrow 0} (Z[T, \epsilon, \sigma]^m - 1)/m$, and $1/Z[T']$ as $\lim_{n \rightarrow 0} Z^{n-1}$. There are then n replicas σ_a ($a = 1, \dots, n$), and m replicas S_{α} ($\alpha = 1, \dots, m$). The ‘external field’ term is an interaction term of all the replicas S_{α} with one privileged replica, say, σ_1 . Three overlap matrices turn out to be relevant for the description of the physics of the model: $Q_{ab}^* = \frac{1}{N} \sum_i \langle \sigma_i^a \sigma_i^b \rangle$ describing the overlap statistics of the replicas at equilibrium at temperature T' , $P_{a,\alpha} = \frac{1}{N} \sum_i \langle \sigma_i^a S_i^{\alpha} \rangle$ describing the overlaps among the replicas at temperature T' and the replicas at temperature T , and finally $Q^{\alpha\beta} = \frac{1}{N} \sum_i \langle S_i^{\alpha} S_i^{\beta} \rangle$ describing the overlaps between replicas at temperature T . As is physically clear, it is found that the structure of the matrix Q_{ab}^* is not affected at the leading order by the presence of the replicas S^{α} . In this paper we will restrict ourselves to the temperature range $T \geq T_S$, where $Q_{ab}^* = \delta_{ab}$. In this regime it is sensible to assume $P_{a,\alpha} = \delta_{a,1} q_{12}$ for all α . The structure of the matrix $Q_{\alpha\beta}$ is more subtle. Assuming a single-state picture in [12] the form $Q_{\alpha\beta} = \delta_{\alpha\beta} + q(1 - \delta_{\alpha\beta})$ was taken. But it turns out also to be necessary to consider the possibility that ergodicity is broken for the system in a ‘field’, with consequent replica symmetry breaking in $Q_{\alpha\beta}$. The most general ansatz we shall need is the ‘one step’ form (see, e.g. [8]), characterized by the parameters (q_0, q_1, x) . With this ansatz it is easy to find that the potential, as a function of all the order parameters, is [15]:

$$V(q_{12}) = -\frac{1}{2\beta} \left\{ 2\beta\beta' f(q_{12}) - \beta^2((1-x)f(q_1) + xf(q_0)) + \frac{x-1}{x} \ln(1-q_1) - \frac{1}{x} \ln(1-(1-x)q_1 - xq_0) + \frac{q_0 - q_{12}^2}{1 - (1-x)q_1 - xq_0} \right\} \quad (6)$$

where V has to be maximized with respect to q_0 , q_1 and x . These saddle-point equations read:

$$\begin{aligned} q_{12}^2 &= q_0 - \beta^2 f'(q_0)(1 - (1-x)q_1 - xq_0)^2 \\ \beta^2(f'(q_1) - f'(q_0))(1-x) &= (1-x) \frac{q_1 - q_0}{(1-q_1)(1 - (1-x)q_1 - xq_0)} \\ \beta^2(f(q_1) - f(q_0)) + \frac{1}{x^2} \ln \left(\frac{1-q_1}{1 - (1-x)q_1 - xq_0} \right) + \beta^2 \frac{(1-q_1)}{x} f'(q_1) & \\ -\beta^2 \frac{f'(q_0)}{x} (1 - (1-x)q_1 - xq_0) &= 0. \end{aligned} \quad (7)$$

A numerical resolution allows us to construct the curve $V(q_{12})$ †.

In general, this curve can be divided into three regions. There are small and large q_{12} regions (outside the interval A–B in figure 1) where replica symmetry holds. In between, the symmetry is broken. In the large q_{12} region, the solution is $q_1 = q_0$ testifying ergodicity

† In [12], the form $Q_{\alpha\beta} = \delta_{\alpha\beta} + q(1 - \delta_{\alpha\beta})$ yielded simplified equations, corresponding to $q_0 = q_1$ in (7). The resulting potential will be denoted as the ‘replica symmetric’ potential.




Figure 1. Potential V as a function of q_{12} , for the $p = 3 + p = 4$, $\beta = 1.25$, $\beta' = 1.243$; full curve: replica symmetric solution; dotted curve: RSB solution, from A ($x = 1$) to B (where q_0 and q_1 merge).

in a single state. In the point B a de Almeida Thouless instability develops. The replica-symmetry breaking region is interpreted as usual ergodicity breaking with dominance of small number of valleys for typical samples. At the point A one finds $x = 1$, and the restoring of replica symmetry implies in fact a number of valleys exponentially large $\mathcal{N} \sim e^{N\Sigma(q_{12})}$. In this region (between $q_{12} = 0$ and A), $x = 1$, the Edwards–Anderson parameter inside the valleys is obtained as the value of q_1 from the second equation of (7) divided by $(1 - x)$ in $x = 1$, and is depicted with crosses in figure 2. The complexity $\Sigma(q_{12})$ can be calculated as in the usual case as $\frac{\partial V}{\partial x}|_{x=1}$, and is depicted in figure 3. For $q_{12} = 0$, where there is no effective constraint, the second replica is at equilibrium at T and we find the total complexity at temperature T , and the equilibrium Edwards–Anderson parameter at T .

The global situation is displayed for a typical example in figures 1 and 2.

2.2. Minima of the potential

The qualitative features of the potential are largely independent of the form of the function f . Let us briefly discuss the case of equal temperatures $\beta = \beta'$ [12]. The potential always has an absolute minimum for $q_{12} = 0$, corresponding, as previously mentioned, to the second replica being at equilibrium. Another minimum appears for a non-zero value of q_{12} for temperatures below T_d (see figure 4) (at $T = T' = T_d$, it is a horizontal flex). This relative minimum corresponds to having both replicas in the same state, with $q_{12} = q = q_{EA}$. Since the number of equilibrium states at temperature T is $\exp(N\Sigma(T))$ by the definition of the complexity (or configurational entropy) Σ , the free-energy cost of having this situation is

$$V_{\text{relative minimum}}(q_{EA}) = T\Sigma(T) = -\frac{\beta}{2}f(q_{EA}) - \frac{q_{EA}}{2\beta} - \frac{1}{2\beta}\ln(1 - q_{EA}). \quad (8)$$

For the p -spin model, the case of different temperatures β and β' , has also been treated in some details [12]. The relative minimum, which still exists for $0 < T < T_{\text{final}}(T')$, and $T' < T_d$, remains in the replica-symmetric region of the potential, and can be clearly




Figure 2. $q_0(q_{12}), q_1(q_{12})$ as a function of q_{12} , for the $p = 3 + p = 4$, $\beta = 1.25$, $\beta' = 1.243$ (lines); diamonds: RS solution, given by inserting $q_0 = q_1$ in (7); crosses: continuation of $q_1(q_{12})$ in the first RS region, with $\lim_{q_{12} \rightarrow 0} q_1(q_{12}) = q_{EA}(\beta)$.




Figure 3. $\Sigma(q_{12})$ for the $p = 3 + p = 4$, $\beta = 1.25$, $\beta' = 1.243$; for $q_{12} \rightarrow 0$ we recover the complexity at β . At A the complexity goes to zero, corresponding to the entrance in the RSB region of the potential.

interpreted. Indeed, the homogeneity of the Hamiltonian allows us to write the TAP free energy in a simple form [9, 12, 18, 16]:

$$f_{\text{TAP}}(\{m_i\}) = q^{\frac{p}{2}} E^0(\{\hat{s}_i\}) - \frac{T}{2} \ln(1 - q) - \frac{1}{4T} [(p - 1)q^p - pq^{p-1} + 1] \quad (9)$$

where we have written $m_i = \langle s_i \rangle = \sqrt{q} \hat{s}_i$, with $\sum_i \hat{s}_i^2 = N$, and the angular energy (zero-temperature energy) is:

$$E^0(\{\hat{s}_i\}) \equiv -\frac{1}{N} \sum_{1 \leq i_1 < \dots < i_p \leq N} J_{i_1, \dots, i_p} \hat{s}_{i_1} \cdots \hat{s}_{i_p}. \quad (10)$$




Figure 4. Potential V as a function of q_{12} , for the $p = 3$ p -spin model, for $\beta = \beta'$, from top to bottom $\beta = 1.6$, $\beta = \beta_d \approx 1.633$, $\beta = 1.7$; here $\beta_s = 1.706$.

The order in free energy of the solutions of the TAP equations does not depend on temperature, nor can a solution bifurcate as the temperature is changed. All these solutions can be easily parametrized [9, 12, 18, 16] and followed with temperature.

It is then easy to show that the properties of the primary minimum (value of $q_0 = q_1$, and energy) are precisely the properties of the TAP states of equilibrium at T' (characterized by a zero-temperature energy E'_0) followed at T (parameter q and energy), with

$$V_{\text{primary}} = T\Sigma(T') + F_{\text{TAP}}(T, E'_0) - F(T). \quad (11)$$

This situation corresponds therefore to having the second replica in a TAP state of equilibrium at T' followed at T †. This situation was also ascertained by the study of the dynamics of a system thermalized at T' , and whose temperature was then changed to T [12, 16, 19]: the dynamics obtained is indeed of equilibrium in these particular TAP states, chosen by the thermalization at T' and followed when the temperature of the system is changed.

In the case where f is different from a monomial, i.e. of an inhomogeneous Hamiltonian, many points remained unclear. In particular, the TAP free energy cannot be parametrized in such a simple form, and it is not granted that the TAP solutions keep their order in free energy when the temperature changes. Moreover, the role of the breaking of replica symmetry was not studied. In the p -spin model, as a consequence of the absence of bifurcation of the solutions, the minimum of the potential is always in the replica-symmetric region, and the inclusion of replica-symmetry breaking effects does not affect the discussion of the metastable states, except for eliminating the spurious secondary minimum found in [12], whose meaning was not clear.

For an inhomogeneous Hamiltonian, by studying the potential, including RSB effects, and the dynamics with thermalized initial conditions, we will show that the potential still allows us to determine the characteristics of TAP states, as long as the minimum is in the replica-symmetric region. We will associate the entrance of the minimum in the RSB region

† Also note that, for $\beta' = \beta_d$, the minimum in fact becomes a horizontal flex of the potential, with the energy and parameter q_{12} equal to those obtained in off-equilibrium dynamics.

of the potential with bifurcations, and show that, in this case, the dynamics with thermalized initial conditions gives rise to a particular form of aging.

3. Potential for an inhomogeneous Hamiltonian

Let us turn to the detailed study of the potential in the case of an inhomogeneous Hamiltonian. The numerical examples will be given for $f(q) = (q^3 + q^4)/2$ for simplicity, but the analysis is independent of this particular form.

We first note that, like for the homogeneous case, a minimum with $q_{12} \neq 0$ only exists for $T' < T_d$. Moreover, we will limit ourselves to $T' > T_s$. For $T = T'$, the primary minimum is still in the replica-symmetric part of the potential. If T is raised, this minimum stays in the RS region, and disappears at a certain temperature $T_{\text{final}}(T')$, which verifies $T_{\text{final}}(T_d) = T_d$.

As T is lowered however, the endpoint of the RSB region (where $q_0 = q_1 \equiv q$) gets closer to the minimum, and finally reaches it at $T_{\text{rsb}}(T') = \frac{1}{\beta_{\text{rsb}}}$ given by

$$\begin{aligned} \beta_{\text{rsb}}^2 f''(q)(1-q)^2 &= 1 \\ \beta_{\text{rsb}}^2 (1-q)^2 f'(q) &= q - \tilde{p}^2 \\ \beta_{\text{rsb}} \beta' f'(\tilde{p}) &= \frac{\tilde{p}}{1-q} \end{aligned} \quad (12)$$

where \tilde{p} is the value of q_{12} in the minimum. For even lower temperatures, the minimum is within the RSB region.

$T_{\text{rsb}}(T')$ reaches zero for some T' (see figure 5); for lower T' , the minimum is always in the RS region.

An example of the situation $T > T_{\text{rsb}}(T')$ is displayed in figure 1, while the limiting case $T = T_{\text{rsb}}(T')$ and a case where $T < T_{\text{rsb}}(T')$ are shown in figure 6.

For temperatures $T_{\text{final}}(T') > T > T_{\text{rsb}}(T')$ the primary minimum can be interpreted as the state of equilibrium at temperature T' followed down at temperature T . Indeed, if we




Figure 5. $T_{\text{final}}(T')$ (full curve) and $T_{\text{rsb}}(T')$ (broken curve) for $f(q) = \frac{1}{2}(q^3 + q^4)$; $T_d \approx 0.805166$. Note that at low enough temperatures, the states never bifurcate. The vertical line corresponds to $T' = T_d$, i.e. the temperature of the appearance of the minimum: along this line the potential displays an horizontal flex.




Figure 6. Potential for $p = 3 + p = 4$, for $\beta' = 1.243$ and $\beta = 1.4625$ (top), $\beta = 3$ (bottom); full curves=RS solution, broken curves=RSB solution. For $\beta = 1.4625$ the endpoint of the RSB solution coincides with the minimum of the potential; for $\beta = 3$ the minimum has disappeared from the RS solution, while it still exists for the RSB curve.

consider the TAP states with values of the energy and of the parameter q equal to those of the primary minimum, and if we compute their free energy f_{TAP} , we obtain (see the appendix):

$$V_{\text{primary}} = f_{\text{TAP}} - F_{\text{RS}}(T). \quad (13)$$

Following the computation of [18, 20], it is also possible to obtain the number of TAP solutions with fixed parameter q and energy E_{TAP} , and, writing it in the form

$$\exp(NS(q, E_{\text{TAP}}, T)) \quad (14)$$

we have checked numerically the identity:

$$S(q^{pr}, E_{\text{primary}}, T) = \Sigma(T'). \quad (15)$$




Figure 7. Total free energy.

Therefore, the number of equilibrium TAP solutions at T' ($\exp(N\Sigma(T'))$) is equal to the number of TAP solutions at T with the energy and the parameter q of the primary minimum. This fact, together with (13), shows that the state of equilibrium at T' has been followed at T , and is a stable state with free-energy cost

$$V_{\text{primary}} = T\Sigma(T') + F_{\text{TAP}}(E_{\text{primary}}, q^{pr}, \beta) - F_{\text{RS}}(T) \quad (16)$$

(where the total, i.e. with the complexity term, free energy of the TAP states is $F_{\text{TAP}}(E_{\text{TAP}}, q, \beta) = f_{\text{TAP}}(E_{\text{TAP}}, q, \beta) - TS(q, E_{\text{TAP}}, T)$). In the next section we will show how these states can be followed dynamically, by choosing appropriate initial conditions.

In contrast, for $T < T_{\text{rsb}}(T')$ the primary minimum is in the region of the potential which displays replica symmetry breaking[†]. The obvious interpretation for this is that at $T_{\text{rsb}}(T')$, the metastable states multifurcate, according to the usual pattern known from the physics of the Sherrington–Kirkpatrick model.

Let us now address the problem of level crossing: the p -spin model seems very particular, in that the order in free energy of the TAP states does not depend on temperature. For $T < T_s$, the statics are given by the lowest TAP states, therefore there are high correlations between equilibrium states at different temperatures. On the other hand, for temperatures between T_s and T_d , equilibrium measures at different temperatures are given by different bunches of TAP states; therefore the overlap between equilibrium states at different temperatures is zero, but the TAP states can be followed at other temperatures, and their order in free energy (without the complexity term) remains the same.

For the case of an inhomogeneous Hamiltonian, we also show that, as long as we consider TAP states giving the equilibrium measure at temperatures higher than T_s , we have no crossing in the free energies f_{TAP} : indeed, if we note $f(T, T')$ the free energy of one TAP state of equilibrium at T' , followed at T , we have

$$f(T, T') = V_{\text{primary}}(T, T') + F(T) \quad (17)$$

[†] Note that $T_{\text{rsb}}(T_d) = T_d$, and that, for $T' = T_d$, $T < T_d$, the minimum is in fact a horizontal flex, like for the homogeneous case, except that it lies within the RSB region of the potential. Besides, the energy in this point is equal to the dynamical energy at T .

and thus we obtain

$$\frac{\partial}{\partial T'} f(T, T') = \frac{f(q_{12}^{pr})}{T'^2}. \quad (18)$$

This quantity is always positive, so, if we have two temperatures $T_d > T'_1 > T'_2 > T_s$, at any temperature T for which we can follow the states giving equilibrium at T'_1 and T'_2 , the order

$$f(T, T'_1) > f(T, T'_2) \quad (19)$$

is conserved. Of course, this is not the case if we consider the full free energy, with the complexity term, i.e. $F(T, T') = f(T, T') - T\Sigma(T')$. We then have that each curve $F(T, T')$ as a function of T is tangent to the curve $F_{RS}(T)$ at the point $(T', F(T', T') = F_{RS}(T'))$.

Hence, this global situation, with the replica-symmetric free energy as the envelope of the curves giving the total TAP free energies, whereas the curves giving the TAP free energies without the complexity term do not cross, seems very generic between the static and dynamic transitions.

Considering the case of $T' = T_s$, the lowest T' for which we are allowed to use the simple ansatz $Q_{ab}^* = \delta_{ab}$, we find a different situation. In fact, we find that if $T < T_s$ the value of the potential in the primary minimum is slightly higher than 0, with $q_{EA}(T) = q_1(q_{12} \rightarrow 0) > q^{pr} > q_{12}^{pr}$. This is in contrast to the case of the p -spin model, for which the states of equilibrium at T_s , followed at T , are still of equilibrium at T : these are the lowest TAP states, and they dominate the equilibrium measure for temperatures ranging from 0 to T_s . In this case we obtain $V_{\text{primary}} = V(0)$, and $q^{pr} = q_1(q_{12} \rightarrow 0)$. Here, in contrast, the difference between the quantities at $q_{12} = 0$ and at the primary minimum show that the states of equilibrium at T_s are no longer of equilibrium at $T < T_s$. Therefore, chaos is present in temperature. For a detailed study of the $T' < T_s$ region, we would however need to take into account the RSB effects on the first replica, which would yield another form for the potential, and we will not do it here.

4. Dynamics

We now address the problem of the dynamics of the system at T , starting from thermalized initial conditions at T' . In the case of the p -spin model, it was shown that such a procedure allows us to dynamically reach the states described by the minimum of the potential, i.e. to follow dynamically the TAP states. As usual, we study the Langevin relaxation dynamics of the model, given by

$$\frac{ds_i(t)}{dt} = -\frac{\partial H}{\partial s_i} - \mu(t)s_i(t) + \eta_i(t) \quad (20)$$

where the η_i are Gaussian thermal noises with $\langle \eta_i(t)\eta_j(t') \rangle = 2T\delta_{ij}\delta(t-t')$, and $\mu(t)$ has to be computed self-consistently in order to implement the spherical constraint $\sum_i s_i^2 = N$.

In the infinite N limit, we can obtain the dynamical equations for the correlation and response functions ($C(t, t') = \frac{1}{N} \sum_i \overline{\langle s_i(t)s_i(t') \rangle}$, $r(t, t') = \frac{1}{N} \sum_i \overline{\langle \frac{\partial s_i(t)}{\partial \eta_i(t')} \rangle}$) [12, 22], that for

$t > t'$ read:

$$\begin{aligned} \frac{\partial r(t, t')}{\partial t} &= -\mu(t)r(t, t') + \int_{t'}^t ds f''(C(t, s))r(t, s)r(s, t') \\ \frac{\partial C(t, t')}{\partial t} &= -\mu(t)C(t, t') + \int_0^{t'} ds f'(C(t, s))r(t', s) \\ &\quad + \int_0^t ds f''(C(t, s))r(t, s)C(s, t') + \frac{1}{T'} f'(C(t, 0))C(t', 0) \end{aligned} \quad (21)$$

complemented by the equation that enforces the spherical condition

$$\begin{aligned} \mu(t) &= \int_0^t ds f'(C(t, s))r(t, s) + \int_0^t ds f''(C(t, s))r(t, s)C(s, t) \\ &\quad + T + \frac{1}{T'} f'(C(t, 0))C(t, 0). \end{aligned} \quad (22)$$

In [12], it was noted that a numerical integration of (21) for a particular choice of the temperatures, after some transient led to equilibrium with time translation invariance (TTI) and validity of the fluctuation dissipation theorem (FDT). However, no systematic study was undertaken.

As long as the primary minimum of the potential is in the replica-symmetric region it is reasonable to take as an ansatz, that indeed an equilibrium regime is reached after a short transient. We therefore deal with the functions $C_{as}(\tau)$, $r_{as}(\tau)$ related by FDT, with the introduction of the limiting quantities \tilde{p} and q :

$$\begin{aligned} C(t, t') &= C_{as}(t - t'); \quad r(t, t') = r_{as}(t - t'); \quad r_{as}(\tau) = -\beta \frac{\partial}{\partial \tau} C_{as}(\tau) \\ \lim_{t \rightarrow \infty} C(t, 0) &= \tilde{p}; \quad \lim_{\tau \rightarrow \infty} C_{as}(\tau) = q. \end{aligned} \quad (23)$$

This ansatz yields the same equations for \tilde{p} and q as those for q_{12} and q (33) specifying the extremum of the potential in the RS region [12]. Besides, it coincides very well with the results of a numerical integration of equations (21). We can therefore conclude that the dynamics takes place in a TAP state, of equilibrium at T' , in which the system was put at T' by thermalization, followed dynamically at the new temperature T . This behaviour is exactly the same as that for the p -spin model [16].

For $T < T_{\text{rsb}}(T')$ another ansatz has to be chosen. In particular, since for low enough T the minimum disappears from the RS potential, the dynamical equations for \tilde{p} and q have no more solutions. We therefore propose an ansatz similar to the one used in the aging dynamics of such models [17], except that the motion will be confined in the vicinity of the initial state. We assume then that:

- for finite time separations $\tau = t - t'$, with τ/t going to zero, the equilibrium properties are valid, which means that we deal with the functions $C_{\text{FDT}}(\tau)$ and $r_{\text{FDT}}(\tau)$, related by FDT. We note that $\lim_{\tau \rightarrow \infty} C_{\text{FDT}}(\tau) = q_1$;

- an aging regime is present: for t and t' going to infinity, without $(t - t')/t \rightarrow 0$, TTI is violated, and the FDT is replaced by the quasi-FDT

$$x \frac{\partial C(t, t')}{\partial t'} = \text{Tr}(t, t') \quad (24)$$

with constant $x \neq 1$. In this regime, we have the limits $\lim_{t'/t \rightarrow 1} C(t, t') = q_1$, $\lim_{t'/t \rightarrow 0} C(t, t') = q_0$;

- we have moreover to introduce the quantity $\lim_{t \rightarrow \infty} C(t, 0) = \tilde{p}$, which tells us how much the system remembers its initial conditions.

As it happens in the random initial condition case, the parameters q_1, q_0, \tilde{p}, x can be determined from the asymptotic analysis of equation (21) without fully solving the dynamics. The hypothesis of the existence of an aging regime, and the continuity of the response function implies the equation

$$\beta^2 f''(q_1)(1 - q_1)^2 = 1 \quad (25)$$

which coincides with the ‘marginal stability condition’ of the statics [23, 17]. The other three equations

$$\begin{aligned} \frac{q_1}{\beta(1 - q_1)} &= \beta f'(q_1)(1 - q_1) + \beta x(q_1 f'(q_1) - q_0 f'(q_0)) + \beta' \tilde{p} f'(\tilde{p}) \\ \frac{\tilde{p}}{\beta(1 - q_1)} &= \beta \tilde{p} x(f'(q_1) - f'(q_0)) + \beta' f'(\tilde{p}) \\ \frac{q_0}{\beta(1 - q_1)} &= \beta f'(q_0)(1 - q_1) + \beta q_0 x(f'(q_1) - f'(q_0)) + \beta' \tilde{p} f'(\tilde{p}) \end{aligned} \quad (26)$$

can be shown to be equivalent to the vanishing of the derivatives of the potential function 4 with respect to q_1, q_0 and q_{12} . In terms of these parameters the asymptotic energy is given by:

$$E = -\beta' f(\tilde{p}) - \beta(f(1) - f(q_1)) - \beta x(f(q_1) - f(q_0)). \quad (27)$$

The usual aging behaviour with $q_0 = \tilde{p} = 0$ is of course the solution of these equations. This corresponds to forgetting the initial conditions, and happens when T' is higher than the dynamical transition temperature. However, for $T' < T_d$, this solution, besides contrasting with the statical picture of the model, would be internally contradictory: indeed, at T_{rsb} the energy in the primary minimum is lower than the dynamical energy at the same temperature. Therefore, such a solution, which would yield an asymptotic energy equal to the dynamical one, would lead to a higher energy for a lower temperature! Finally, the numerical integration of the dynamical equations shows that the behaviour of the dynamical quantities is very different from the case of infinite T' , and that $C(t, 0)$ does not seem to decrease to zero. These facts lead to the conclusion that we must prefer the solution with non-zero q_0 and \tilde{p} . The aging therefore takes place in a restricted phase-space region. However, for $T < T_{\text{rsb}}(T')$ the dynamic internal energy is higher than the static one, similarly to what happens starting from random initial conditions.

Let us also note that the asymptotic energy (27) in the case of thermalized initial conditions is lower than the dynamical energy after a quench, showing that this procedure allows us to reach states with lower energies. An immediate consequence is the importance of the way in which the final temperature is reached.

5. A comparison with real glasses

5.1. General considerations

In the studied mean-field models, we found that, below the dynamical transition T_D , we could define a whole spectrum of internal energies for the system at temperature T , depending on the way the system has been put at its final temperature:

- the equilibrium energy $E_{\text{eq}}(T)$, which is done by the usual Boltzmann Gibbs formula;
 - the dynamical energy, corresponding to the energy of a system which is quenched to the final temperature from a temperature higher than T_D ;
 - the energies $E(T', T)$, obtained for a system at equilibrium at T' and then put at T .
- Depending on T' and T , the system can be at equilibrium or exhibit aging dynamics.

These energies can be consistently computed using the explicit form of the dynamics. It is also possible to compute them by using the appropriate statistical prescription which does not make explicit reference to the dynamics.

At this point the reader may ask how much all these findings are relevant for the real world. Metastable states with an *infinite* life do not exist in short-range finite-dimensional models and their presence in mean-field models is a clear artifact of the approximation. The would-be infinite life metastable states of the mean field theory do decay through some activated processes (whose detailed properties have not yet been fully clarified). If the mean-field picture is relevant for the real world the timescale of the activated processes should be large enough that there is a time window in which the behaviour predicted by the mean-field theory can be observed.

Given our lack of command on the activated processes, we cannot treat this question analytically and we have to resort to numerical simulations. We will consider a simple system, one of the prototypes of glass forming systems, known to have a glass transition at a given temperature T_G .

We will see that we can also define various energies:

- the equilibrium energy $E_{\text{eq}}(T)$;
- the slow cooling energy $E_S(T)$, which is obtained by the limit to infinite cooling time of the energy of a system which starts at temperatures greater than the dynamical transition;
- the fast cooling energy $E_F(T)$, which is obtained by the limit to infinite cooling rate of the energy of a system which is quenched to the final temperature from a temperature higher than T_D .

Between $E_S(T)$ and $E_F(T)$, various cooling rates will yield various asymptotic energies.

We will see that if we cool the system fast to a temperature near or below T_G (we have investigated up to temperatures equal to $0.25T_G$) the energy as function of the time may be represented by the following form

$$E_F(t) = E_F + At^{-\mu} + O(t^{-2\mu}) \quad (28)$$

where the exponent μ is in the range 0.5–0.7 and weakly depends on the temperature. The previous formula well represent the data for time in the window 10^2 – 10^5 time units (i.e. one Monte Carlo sweep).

In a similar way, we can represent the data for the slow-cooled energy as a function of time with a similar form in the same time window:

$$E_S(t) = E_S + At^{-\mu} + O(t^{-2\mu}) \quad (29)$$

where the exponent μ is compatible to be equal to the one used in equation (28).

The two functions E_F and E_S are different from one another below T_G and their difference vanishes when we approach T_G . It is clear that the two previous formulae can be valid only in a limited time window; asymptotically the two energies $E_F(t)$ and $E_S(t)$ must go to the same limit (i.e. the equilibrium value of the energy). This is likely to happen on a much longer scale. Here we want to stress the presence of a time window in which the prediction of a theory based on the existence of metastable states can be tested.

Before showing the results of the numerical simulations, we will give some details of the model we consider.

5.2. The Hamiltonian

The model we consider is the following. We have taken a mixture of soft particles of different sizes. Half of the particles are of type *A*, half of type *B* and the interaction among

the particle is given by the Hamiltonian:

$$H = \sum_{i < k} \left(\frac{\sigma(i) + \sigma(k)}{|\mathbf{x}_i - \mathbf{x}_k|} \right)^{12} \quad (30)$$

where the radius (σ) depends on the type of particles. This model has been carefully studied in the past [24–29]. It is known that a choice of the radius, such that $\sigma_B/\sigma_A = 1.2$, strongly inhibits crystallization and that the systems go into a glassy phase when it is cooled. Using the same conventions as the previous investigators we consider particles of an average diameter of 1, more precisely we set

$$\frac{\sigma_A^3 + 2(\sigma_A + \sigma_B)^3 + \sigma_B^3}{4} = 1. \quad (31)$$

Due to the simple scaling behaviour of the potential, the thermodynamic quantities depend only on the quantity $T^{3/4}/\rho$, T and ρ being respectively the temperature and the density. For definiteness we have taken $\rho = 1$. It is usual to introduce the quantity $\Gamma^4 \equiv \beta$. The glass transition is known to happen around $\Gamma = 1.45$ (i.e. for $T \approx 0.226$) [25].

5.3. Numerical results

Our simulations are done using a Monte Carlo algorithm, which is more easy to deal with than molecular dynamics, if we change the temperature in an abrupt way. Each particle is shifted by a random amount at each step, and the size of the shift is fixed by the condition that the average acceptance rate of the proposal change is about 0.4. Particles are placed in a cubic box with periodic boundary conditions. In our simulations we have considered a relatively small number of particles, i.e. $N = 66$. Note that for all the simulations, the system is always out of equilibrium and exhibits aging: the ergodic time is far beyond reach [27, 28, 30].

We start by placing the particles at random and quench the system by putting it at its final temperature (i.e. infinite cooling rate). The typical value of the energy density of the initial configuration is very high ($O(10^5)$) due to the singular form of the potential and it takes a few iterations to arrive at a more reasonable value. We show the data as a function of Monte Carlo time t in figure 8 for $\Gamma = 1.8$ ($T \approx 0.095$).

In the slow cooling approach we also start by placing the particles at random at the beginning. We divide the cooling time into five equal intervals: in the first interval we have $\Gamma = 1$, in the second interval $\Gamma = 1.2$, ... and the fifth interval $\Gamma = 1.8$. The data are taken for each temperature only in the second half of the corresponding interval. The results, as a function of the time spent at each temperature, i.e. of the inverse of the cooling rate, are shown in figure 8 for $\Gamma = 1.8$. We clearly see that the two curves E_F and E_S definitely extrapolate to a different value. The extrapolated values of the energy as function of the temperature can be seen in figure 10 using the fast and the slow cooling method in the region $\Gamma > 1.4$ ($T < 0.26$). The data are not shown at higher temperatures, because the two methods give the same result. Other procedures to investigate the dependence of the cooling rate involve a similar cooling from $\Gamma = 1$ to $\Gamma = 1.8$ in a total time t_{cooling} , with times $t_{\text{cooling}}/4$ spent at $\Gamma = 1, 1.2, 1.4, 1.6$, and then the study of the evolution of the energy at $\Gamma = 1.8$. The long time limit of the energy lies then between E_F (for $t_{\text{cooling}} \rightarrow 0$) and E_S (for $t_{\text{cooling}} \rightarrow \infty$). In figure 9 we show in the evolution of the energy at $\Gamma = 1.8$ for various cooling rates. The effects are quite small, so it is necessary to compare reasonably different rates. For the available times, the energy of the system depends on the cooling it has followed: the energy is lower for slower cooling procedures.




Figure 8. ‘Fast’ curve: energy as a function of $t^{-0.5}$ where t is the time spent at $\Gamma = 1.8$ after a rapid quench; ‘slow’ curve: energy reached at the end of the time spent at $\Gamma = 1.8$, as a function of $t^{-0.5}$, where t is the time spent at each temperature during the gradual quench process. We see that slower cooling yields lower energies. The continuation to $t^{-0.5} \rightarrow 0$ corresponds to an infinitely slow cooling.




Figure 9. Evolution of the energy at $\Gamma = 1.8$ ($T \approx 0.095$), as a function of time, for various cooling rates; the horizontal lines correspond to E_F and E_S . The lower curves correspond to slower coolings.

Moreover, it is worth noting that the value of μ is, roughly speaking, independent of the temperature [30]. This phenomenon happens in the only model of mean-field theory where the exponent has been computed [31] and this is a strong indication that the approach to equilibrium in this region is not dominated by activated processes, but more (roughly speaking) by entropic barriers: the barriers between metastable states could include both energetic and entropic effects [32]. Moreover, this shows the possible relevance of the scenario detailed in the preceding paragraph (aging similar to usual, but at lower energies), and of some intuitive mean-field scenarios [33].




Figure 10. Extrapolations of the energies E_F and E_S at large times, i.e. asymptotic energies after a quench or an infinitely slow cooling, for various temperatures.

It would be interesting to be able to simulate the thermalization at a certain temperature, followed by a quench at a lower temperature, like in mean-field models. Unfortunately, the available time window does not allow us to reach thermalization at temperatures lower than the dynamic transition.

Another possibility would be to cool the system very slowly to a certain value T' , such that its energy is lower than $E_S(T)$ for a certain T ($T > T'$), and then to bring the system back to T , to see whether the obtained energy is still lower than $E_S(T)$. Such investigations are however beyond the scope of this short study.

6. Summary and conclusions

In this paper we have investigated the behaviour in temperature of the metastable states of long-range spin glasses with first-order freezing transition. We have shown that the metastable states can be followed up and down in temperature, from the temperature where they are dominating the partition function. Going up in temperature, one finds some temperature where the states disappear, merging with some maxima. Going down in temperature, the states never disappear, although in some range of T' multifurcation is found. We also studied the dynamics at temperature $T < T_d$, following a quench from equilibrium at temperature T' . If $T' > T_d$ we find no difference from the usual aging behaviour [17] that follows a quench from infinite temperature. For $T' < T_d$ we have found two possibilities. If $T > T_{rsb}(T')$ the original valley has ‘deformed’ but not bifurcated and the system is able to equilibrate inside it. In the complementary interval $T < T_{rsb}(T')$ the landscape has changed drastically as the original valley has bifurcated. The system is then unable to thermalize and falls in an aging regime, while remaining confined in the vicinity of the initial data. Besides, this dynamical study shows that the aging after a slow quench (in the mean-field case, the case of thermalized initial conditions at T' , can be thought of as a situation after an infinitely slow quench) allows us to reach a situation where the behaviour is qualitatively similar to the one after a rapid quench (i.e. aging corresponding to a slow touring of the phase space), but within a phase-space region with lower energies. Therefore, at a given temperature T , the possibilities are not only of aging at a relatively high energy, after a sudden quench, or of equilibrium dynamics after an infinitely slow quench, but also

of aging at intermediate energies, depending on the route from a high temperature to T .

In the last section, we tried to emphasize the possible relevance of such mean-field scenarios for finite dimensions, where it has been advocated that metastable states may still exist, but with a finite lifetime: coming from a high-temperature phase, the system may be able to find these states in a finite time, and the resulting aging behaviour when decreasing the temperature could be a mixture of jumps between states and periods of wandering when states bifurcate.

Indeed, the numerical study of section 5 shows indications that, at least in the explored time window, for a soft sphere model of glass exhibiting aging, the dynamics is not dominated by activated processes. Depending on the cooling rate from the high-temperature phase, various energies can be reached. Since the system is finite, it should however reach equilibrium in a finite time (the energy should reach the equilibrium energy, whatever the route to the final temperature may be) but, these simulations show that, even for a relatively small system, this finite time is very large, and therefore that mean-field conclusions can be of importance in the real world.

Acknowledgments

It is a pleasure to thank A Cavagna, I Giardina and M Virasoro for interesting discussions.

Appendix

We consider the case when the primary minimum of the potential is in the RS region: $q_0 = q_1 \equiv q$. Then, for fixed T' , we compute the value of q_{12} and q for this minimum, q_{12}^{pr} and q^{pr} . The saddle-point equations for q_0 , q_1 , x reduce to

$$\beta^2 f'(q^{pr}) = \frac{q^{pr} - q_{12}^{pr2}}{(1 - q^{pr})^2} \quad (32)$$

and the equation $\frac{\partial V}{\partial q_{12}} = 0$ is

$$\beta \beta' f'(q_{12}^{pr}) = \frac{q_{12}^{pr}}{1 - q^{pr}}. \quad (33)$$

The value of the potential is

$$V_{\text{primary}} = -\beta' f(q_{12}^{pr}) + \frac{\beta}{2} f(q^{pr}) - \frac{\beta}{2} (1 - q^{pr}) f'(q^{pr}) - \frac{1}{2\beta} \ln(1 - q^{pr}). \quad (34)$$

The energy of the second replica, in this minimum, is

$$E_{\text{primary}} = \frac{\partial}{\partial \beta} (\beta V + \beta F(T)) \quad (35)$$

which yields

$$E_{\text{primary}} = -\beta' f(q_{12}^{pr}) + \beta f(q^{pr}) - \beta f(1). \quad (36)$$

On the other hand, we can write the TAP free energy as:

$$f_{\text{TAP}}(H, q, \beta) = H - \frac{1}{2\beta} \ln(1 - q) - \frac{\beta}{2} (f(1) - f(q) - (1 - q)f'(q)) \quad (37)$$

where $q = \frac{1}{N} \sum_i m_i^2$, and H is the value taken by the Hamiltonian $H(\{m_i\})$, so the energy of a TAP state m_i is

$$E_{\text{TAP}} = \frac{\partial \beta f}{\partial \beta} = H - \beta (f(1) - f(q) - (1 - q)f'(q)). \quad (38)$$

Then, taking

$$H_{pr} = E_{\text{primary}} + \beta(f(1) - f(q^{pr}) - (1 - q^{pr})f'(q^{pr})) \quad (39)$$

we obtain immediately that

$$V_{\text{primary}} = f_{\text{TAP}}(H_{pr}, q^{pr}, \beta) - F_{\text{RS}}(T). \quad (40)$$

This means that V_{primary} is the free-energy cost of having the second replica in a TAP state with parameter q^{pr} and energy E_{primary} at inverse temperature β .

References

- [1] Sherrington D 1996 Landscape paradigms in physics and biology: introduction and overview *Preprint cond-mat/9608088 Physica D* to appear and references therein
- [2] For a recent review see Marinari E, Parisi G and Ruiz-Lorenzo J 1997 Numerical simulations of spin glass systems *Preprint cond-mat/9701016* contribution to the volume *Spin Glasses and Random Fields* ed P Young
- [3] Thouless D J, Anderson P W and Palmer R G 1977 *Phil. Mag.* **35** 593
- [4] Kirkpatrick T R and Thirumalai D 1988 *Phys. Rev. B* **37** 5342
Kirkpatrick T R and Wolynes P G 1987 *Phys. Rev. B* **36** 8552
Thirumalai D and Kirkpatrick T R 1988 *Phys. Rev. B* **38** 4881
- [5] Kirkpatrick T R and Thirumalai D 1987 *Phys. Rev. B* **36** 5388
- [6] Kirkpatrick T R, Thirumalai D and Wolynes P G 1989 *Phys. Rev. A* **40** 1045
- [7] Parisi G 1994 *Nuovo Cimento II* **16** 939
- [8] Mézard M, Parisi G and Virasoro M A 1987 *Spin Glass Theory and Beyond* (Singapore: World Scientific)
- [9] Kurchan J, Parisi G and Virasoro M A 1993 *J. Physique* **3** 1819
- [10] Kondor I 1989 *J. Phys. A: Math. Gen.* **22** L163
Kondor I and Vegsö A 1993 *J. Phys. A: Math. Gen.* **26** L641
- [11] Franz S and Ney-Nifle M 1995 *J. Phys. A: Math. Gen.* **28** 2499
- [12] Franz S and Parisi G 1995 *J. Physique* **5** 1401
- [13] Monasson R 1995 *Phys. Rev. Lett.* **75** 2847
- [14] Franz S and Parisi G 1997 Phase diagram of glassy systems in an external field *Preprint cond-mat/9701033*
- [15] Cavagna A, Giardinà I and Parisi G 1997 *J. Phys. A: Math. Gen.* **30** 4449
- [16] Barrat A, Burioni R and Mézard M 1996 *J. Phys. A: Math. Gen.* **29** L81
- [17] Cugliandolo L F and Kurchan J 1993 *Phys. Rev. Lett.* **71** 173
- [18] Crisanti A and Sommers H J 1995 *J. Physique* **5** 805
- [19] Barrat A 1997 The p -spin spherical spin glass model *Preprint cond-mat/9701031*
- [20] Rieger H 1992 *Phys. Rev. B* **46** 14 655
- [21] Franz S and Mézard M 1994 *Physica* **210A** 48
- [22] Houghton A, Jain S and Young A P 1983 *Phys. Rev. B* **28** 290
- [23] Horner H 1987 *Z. Phys. B* **66** 175
Crisanti A, Horner H and Sommers H J 1993 *Z. Phys. B* **92** 257
De Santis E, Parisi G and Ritort F 1995 *J. Phys. A: Math. Gen.* **28** 327
- [24] Bernu B, Hansen J-P, Hitawari Y and Pastore G 1987 *Phys. Rev. A* **36** 4891
- [25] Barrat J-L, Roux J-N and Hansen J-P *Chem. Phys.* **149** 197
- [26] Hansen J-P and Yip S 1995 *Trans. Theor. Stat. Phys.* **24** 1149
- [27] Lancaster D and Parisi G 1997 A study of activated processes in soft sphere glass *Preprint cond-mat/9701045*
- [28] Parisi G 1997 Short time aging in binary glasses *Preprint cond-mat/9701015*
- [29] Parisi G 1997 On the mean field approach to glassy systems *Preprint cond-mat/9701034*
- [30] Parisi G 1997 Numerical indications for the existence of a thermodynamic transition in binary glasses *Preprint cond-mat/9701100*
- [31] Franz S, Marinari E and Parisi G 1995 *J. Phys. A: Math. Gen.* **28** 5437
- [32] Ritort F 1995 *Phys. Rev. Lett.* **75** 1190
Franz S and Ritort F 1996 *Europhys. Lett.* **31** 507
Franz S and Ritort F 1996 *J. Stat. Phys.* **85** 131
- [33] Kurchan J and Laloux L 1996 *J. Phys. A: Math. Gen.* **29** 1929