

High-order equations of motion in quantum mechanics and Galilean relativity

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1997 J. Phys. A: Math. Gen. 30 L131

(<http://iopscience.iop.org/0305-4470/30/6/004>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.112

The article was downloaded on 02/06/2010 at 06:13

Please note that [terms and conditions apply](#).

LETTER TO THE EDITOR

High-order equations of motion in quantum mechanics and Galilean relativity

W I Fushchych†§¶ and Z I Symenoh‡||

† Department of Mathematics, UMIST, PO Box 88, Manchester M60 1QD, UK

‡ Institute of Mathematics of the National Academy of Sciences of Ukraine, 3 Tereshchenkivs'ka Street, 252004 Kyiv, Ukraine

Received 1 October 1996, in final form 3 December 1996

Abstract. Linear partial differential equations of arbitrary order invariant under the Galilei transformations are described. Symmetry classification of potentials for these equations in two-dimensional space is carried out. High-order nonlinear partial differential equations invariant under the Galilei, extended Galilei and full Galilei algebras are studied.

Non-relativistic quantum mechanics is based on the equation

$$L\Psi \equiv (S + V)\Psi = 0 \quad (1)$$

where $S = p_0 - p_a^2/2m$, $p_0 = i\partial/\partial x_0 = i\partial/\partial t$, $p_a = -i\partial/\partial x_a$, $V = V(\mathbf{x}, \Psi^*\Psi)$. In the case where V is a function only of \mathbf{x} , equation (1) coincides with the standard linear Schrödinger equation.

The fundamental property of (1) (in the case $V = 0$) is the fact that this equation is compatible with the Galilean relativity principle. In other words, equation (1) ($V = 0$) is invariant under the Galilei group $G(1, 3)$. The Lie algebra $AG(1, 3) = \langle P_0, P_a, J_{ab}, G_a \rangle$ of the Galilei group is generated (see, e.g., [1, 2]) by the operators

$$\begin{aligned} P_0 &= p_0 & P_a &= p_a \\ J_{ab} &= x_a p_b - x_b p_a & a \neq b, a, b &= 1, 2, 3 \\ G_a &= t p_a - m x_a. \end{aligned} \quad (2)$$

The operators $\langle G_a \rangle$ generate the standard Galilei transformations

$$t \rightarrow t' = t \quad x_a \rightarrow x'_a = x_a + v_a t.$$

Definition 1. We say that the equation of type (1) is compatible with the Galilei principle of relativity if it is invariant under the operators $\langle P_0, P_a, J_{ab}, G_a \rangle$.

Let X be one of the operators $\langle P_0, P_a, J_{ab}, G_a \rangle$.

§ E-mail address: mcbtwif@fs1.ma.umist.ac.uk

¶ On research leave from the Institute of Mathematics of the National Academy of Sciences of Ukraine, 3 Tereshchenkivs'ka Street, 252004 Kyiv, Ukraine; E-mail address: math@nonlinear.kiev.ua or nonlin@apmat.freenet.kiev.ua

|| E-mail address: zoya@apmat.freenet.kiev.ua

Definition 2. Equation (1) is invariant under the operator X if the following condition is true:

$$\left. \begin{matrix} X L\Psi \\ (2) \end{matrix} \right|_{L\Psi=0} = 0 \tag{3}$$

where $X_{(2)}$ is the second Lie prolongation of the operator X [1–4].

The equation of type (3) is a Lie condition of invariance of the equation under the Lie algebra. In our case, it is the condition of invariance under the algebra $AG(1, 3)$.

Theorem 1. ([1, 2, 5]) Among linear equations of the first order in t and of the second order in the space variables x there exists the unique equation (1) ($V = \lambda = \text{constant}$) invariant under the algebra $AG(1, 3)$ with the basic elements (2).

Conclusion. We can regard the theorem formulated above as a method of deriving the Schrödinger equation from the Galilei principle of relativity [5, 6].

In the present paper, we give the answer to the following question: Do there exist equations not equivalent to the Schrödinger equation for which the Galilei principle of relativity is true?

In [6, 7], the following generalization of the Schrödinger equation was proposed,

$$(\lambda_1 S + \lambda_2 S^2 + \dots + \lambda_n S^n + V)\Psi = 0 \tag{4}$$

where $S^2 = SS, \dots, S^n = S^{n-1}S, \lambda_1, \lambda_2, \dots, \lambda_n$ are arbitrary parameters.

If $V = 0$, equation (4), as well as equation (1), is invariant under the algebra $AG(1, 3)$, i.e. this equation is compatible with the Galilei principle of relativity. Is this equation unique among high-order linear equations? In what follows, we find a positive answer for this question.

More precisely, we solve the following problems.

(i) We describe all linear equations of arbitrary order invariant under the algebra $AG(1, 3)$.

(ii) We describe the maximal (in Lie sense) symmetry of equation (4) in the two-dimensional space (t, x) .

(iii) We describe nonlinear equations of type (4) invariant under the algebra $AG(1, 3)$, the extended Galilei algebra $AG_1(1, 3) = \langle AG(1, 3), D \rangle$, and the full Galilei algebra $AG_2(1, 3) = \langle AG_1(1, 3), A \rangle$. D and A are the dilation and projective operators, respectively.

(i) For solving the above problems we use the method described in [1, 2, 5–7].

Theorem 2. A: Among linear partial differential equations (PDE) of arbitrary even order $2n$

$$L\Psi = 0$$

$$L = A + B^\mu \partial_\mu + C^{\mu\nu} \partial_{\mu\nu} + D^{\mu\nu\sigma} \partial_{\mu\nu\sigma} + \dots + E^{\overbrace{\mu\nu\sigma\dots\kappa}^{2n}} \partial_{\underbrace{\mu\nu\sigma\dots\kappa}_{2n}} \tag{5}$$

there exists the unique equation

$$(\lambda_1 S + \lambda_2 S^2 + \dots + \lambda_n S^n)\Psi = \lambda\Psi \tag{6}$$

invariant under the algebra $AG(1, 3)$.

B: There are no linear PDE of arbitrary odd order $2n + 1$

$$L\Psi = 0$$

$$L = A + B^\mu \partial_\mu + C^{\mu\nu} \partial_{\mu\nu} + D^{\mu\nu\sigma} \partial_{\mu\nu\sigma} + \dots + \underbrace{E^{\mu\nu\sigma\dots\kappa}}_{2n} \partial_{\mu\nu\sigma\dots\kappa} + \underbrace{G^{\mu\nu\sigma\dots\kappa\rho}}_{2n+1} \partial_{\mu\nu\sigma\dots\kappa\rho} \quad (7)$$

with one non-zero coefficient of the highest derivatives at least, invariant under $AG(1, 3)$.

Here, $A, B^\mu, C^{\mu\nu}, D^{\mu\nu\sigma}, \dots, \underbrace{E^{\mu\nu\sigma\dots\kappa}}_{2n}, \underbrace{G^{\mu\nu\sigma\dots\kappa\rho}}_{2n+1}$ are arbitrary functions of t and \mathbf{x} ; $\lambda_1, \lambda_2, \dots, \lambda_n, \lambda$ are arbitrary constants, $\lambda_n \neq 0$; $\partial_\mu \equiv \partial/\partial x_\mu$, $\partial_{\mu\nu} \equiv \partial^2/\partial x_\mu \partial x_\nu, \dots (\mu, \nu, \dots, \rho = 0, 3)$.

Proof. The scheme and idea of the proof of the theorem is very simple but the concrete realization is not simple. We describe in more detail the proof of part A. Part B is proved in the same way as the first part of the theorem.

According to the Lie method [1, 3, 4], we find the $2n$ th prolongations of the operators (2) and consider the system of determining equations

$$\left. \begin{matrix} X \\ (2n) \end{matrix} L\Psi \right|_{L\Psi=0} = 0 \quad \forall X \in AG(1, 3). \quad (8)$$

Writing equations (8) in the explicit form and equating coefficients for equal derivatives, we solve the system of partial differential equations to obtain functions $A, B^\mu, C^{\mu\nu}, D^{\mu\nu\sigma}, \dots, \underbrace{E^{\mu\nu\sigma\dots\kappa}}_{2n}$.

Invariance of equation (5) under the operators P_0, P_a results in the fact that functions $A, B^\mu, C^{\mu\nu}, D^{\mu\nu\sigma}, \dots, \underbrace{E^{\mu\nu\sigma\dots\kappa}}_{2n}$ do not depend on t and \mathbf{x} , i.e. these coefficients are arbitrary constants. In other words, our PDE has the form $L\Psi \equiv Q^{(1)}(p_0, p_a)\Psi = 0$, where $Q^{(1)}$ is a polynomial in (p_0, p_a) with constant coefficients.

After taking into account the invariance under the operators J_{ab} , we find that the equation has the form $L\Psi \equiv Q^{(2)}(p_0, p_a^2)\Psi = 0$, where $Q^{(2)}$ is a polynomial in (p_0, p_a^2) . After considering the invariance under the Galilei operators G_a , we obtain that the equation has the form $L\Psi \equiv Q^{(3)}(p_0 - (1/2m)p_a^2)\Psi = 0$, where $Q^{(3)}$ is a polynomial in $(p_0 - (1/2m)p_a^2)$. In other words, the equation has the form (6). The theorem is proved. \square

Consequence. Among fourth-order linear PDEs there exists the unique equation invariant under the algebra $AG(1, 3)$ with basic operators (2). This equation has the form

$$(\lambda_1 S + \lambda_2 S^2)\Psi = \lambda \Psi$$

where $\lambda_2 \neq 0$.

(ii) Now, we consider equation (4) in two dimensions t, x and carry out symmetry classification of the potentials $V = V(x)$ of this equation, i.e. we find all functions $V = V(x)$ admitting an extension of symmetry of (4). The following statement is true.

Theorem 3. The two-dimensional equation (4) with $\lambda_n \neq 0, n \neq 1$ is invariant under the following algebras:

- (1) $\langle P_0, I \rangle$, iff $V(x)$ is an arbitrary differentiable function;
- (2) $AG(1, 1) = \langle P_0, P_1, G, I \rangle$, iff $V = \text{constant}$;

(3) $AG_2(1, 1) = \langle \tilde{P}_0, P_1, G, D, A, I \rangle$, iff $V = V_1 = \text{constant}$ the following equalities are true:

$$\frac{\lambda_k}{\lambda_n} = \binom{n}{k} \left(\frac{V_1}{\lambda_n} \right)^{(n-k)/n} \quad k = 1, \dots, n - 1; \quad (9)$$

(4) $\langle \tilde{P}_0, D, A, I \rangle$, iff $V = V_1 + C/x^{2n}$, V_1, C are constants and (9) are true. $\binom{n}{k}$ are the binomial coefficients.

The operators in theorem 3 have the following representation:

$$\begin{aligned} P_0 &= p_0 & P_1 &= p_1 & G &= tp_1 - mx \\ \tilde{P}_0 &= \tilde{p}_0 = P_0 + \sqrt[n]{V_1/\lambda_n} & D &= 2t\tilde{p}_0 - xp_1 - (i/2)(2n - 3) \\ A &= t^2\tilde{p}_0 - tD - (1/2)mx^2 \end{aligned} \quad (10)$$

I is the unit operator.

Consequence. The $2n$ th-order PDE

$$(S^n + V(x))\Psi = 0$$

is invariant under the following algebras:

- (1) $\langle P_0, I \rangle$, iff $V(x)$ is an arbitrary differentiable function;
- (2) $AG(1, 1) = \langle P_0, P_1, G, I \rangle$, iff $V = \text{constant}$;
- (3) $AG_2(1, 1) = \langle P_0, P_1, G, D, A, I \rangle$, iff $V = 0$;
- (4) $\langle P_0, D, A, I \rangle$, iff $V = C/x^{2n}$ where C is an arbitrary constant.

The above operators have representation (10) with $V_1 = 0$.

Note that symmetry classification of potentials for the fourth-order PDE of the form

$$(\lambda_1 S + \lambda_2 S^2 + V(x))\Psi = 0$$

was carried out in [8]. In this case, symmetry operators have representation (10) with $V_1 = \lambda_1^2/4\lambda_2$ and $n = 2$.

(iii) Now, let us consider nonlinear PDEs of type (4) in $(r + 1)$ -dimensional space:

$$S^n \Psi + F(\Psi \Psi^*) \Psi = 0 \quad (11)$$

where Ψ^* is complex conjugated function, n is an arbitrary integer power and F is an arbitrary complex function of $\Psi \Psi^*$

We study symmetry classification of (11), i.e. we find all functions $F(\Psi \Psi^*)$ which admit an extension of symmetry of equation (11).

Theorem 4. Equation (11) is invariant under the following algebras:

- (1) $\langle P_0, P_a, J_{ab}, G_a, Q_1 \rangle$, iff F is an arbitrary differentiable function;
- (2) $\langle P_0, P_a, J_{ab}, G_a, Q_1, Q_2 \rangle$, iff $F = \text{constant} \neq 0$;
- (3) $\langle P_0, P_a, J_{ab}, G_a, Q_1, D \rangle$, iff $F = C(\Psi \Psi^*)^k$, $k \neq 0$;
- (4) $\langle P_0, P_a, J_{ab}, G_a, Q_1, D, A \rangle$, iff $F = C(\Psi \Psi^*)^{(2n)/(r+2-2n)}$;
- (5) $\langle P_0, P_a, J_{ab}, G_a, Q_1, Q_2, D, A \rangle$, iff $F = 0$.

Here, indices a, b are from 1 to r , $a \neq b$, k is an arbitrary number ($k \neq 0$), and the above operators have the following representation:

$$\begin{aligned} P_0 &= p_0 & P_a &= p_a & J_{ab} &= x_a p_b - x_b p_a & G_a &= t \partial_{x_a} + i m x_a Q_1 \\ Q_1 &= \Psi \partial_\Psi - \Psi^* \partial_{\Psi^*} & Q_2 &= \Psi \partial_\Psi + \Psi^* \partial_{\Psi^*} \\ \tilde{D} &= 2t \partial_t + x^c \partial_{x_c} - (n/k) Q_2 & D &= 2t \partial_t + x^c \partial_{x_c} - \frac{r+2-2n}{2} Q_2 \\ A &= t^2 \partial_t + t x^c \partial_{x_c} + (i/2) m x^c x_c Q_1 - \frac{r+2-2n}{2} t Q_2 \end{aligned}$$

where summation from 1 to r over the repeated indices c is understood.

Thus, in the present paper, we have described the unique linear PDE of arbitrary even order which is invariant under the Galilei group. We have investigated the exhaustive symmetry classification of potentials $V(x)$ of (4) and functions $F(\Psi\Psi^*)$ of the nonlinear equation (11), i.e. we have pointed out all functions admitting an extension of the invariance algebra.

The authors would like to thank an anonymous reviewer for kindness and helpful suggestions. The paper is partly supported by INTAS, Royal Society, and the International Soros Science Education Program (grant No PSU061097).

References

- [1] Fushchych W, Shtelen W and Serov N 1993 *Symmetry Analysis and Exact Solutions of Equations of Nonlinear Mathematical Physics* (Dordrecht: Kluwer) p 436
- [2] Fushchich W I and Nikitin A G 1994 *Symmetry of Equations of Quantum Mechanics* (New York: Allerton) p 465
- [3] Ovsyannikov L V 1982 *Group analysis of differential equations* (New York: Academic) p 400
- [4] Olver P 1986 *Application of Lie Groups to Differential Equations* (New York: Springer) p 497
- [5] Fushchych W I and Cherniha R M 1985 The Galilean relativistic principle and nonlinear partial differential equations *J. Phys. A: Math. Gen.* **18** 3491–503
- [6] Fushchych W I 1981 Symmetry in problems of mathematical physics *Algebra-Theoretic Analysis in Mathematical Physics* (Kiev: Institute of Mathematics, Ukrainian Academy of Sciences) pp 6–28 (in Russian)
- [7] Fushchych W 1995 *Ansatz'95, J. Nonlin. Math. Phys.* **2** 216–35
- [8] Symenoh Z I 1995 Symmetry classification of potentials for the generalized Schrödinger equation *Dop. Nat. Acad. Nauk Ukraïny* **3** 20–2