

Integrable multiparametric quantum spin chains

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1998 J. Phys. A: Math. Gen. 31 687

(<http://iopscience.iop.org/0305-4470/31/2/024>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.122

The article was downloaded on 02/06/2010 at 06:52

Please note that [terms and conditions apply](#).

Integrable multiparametric quantum spin chains

Angela Foerster[†]||, Jon Links[‡]¶ and Itzhak Roditi[§]+

[†] Instituto de Física da UFRGS, Av. Bento Gonçalves 9500, Porto Alegre, RS - Brazil

[‡] Department of Mathematics, University of Queensland, Queensland, 4072, Australia

[§] Theory Division, CERN, CH-1211 Geneva 23, Switzerland

Received 1 August 1997, in final form 13 October 1997

Abstract. Using Reshetikhin's construction for multiparametric quantum algebras we obtain the associated multiparametric quantum spin chains. We show that under certain restrictions these models can be mapped to quantum spin chains with twisted boundary conditions. We illustrate how this general formalism applies to construct multiparametric versions of the supersymmetric t - J and U models.

1. Introduction

The advent of quantum algebras [1, 2] precipitated many new results in the area of integrable models. Likewise their supersymmetric counterparts quantum superalgebras [3–6] facilitate systematic treatments of integrable models which accommodate both bosonic and fermionic degrees of freedom. An important subclass are those which may be interpreted as describing systems of correlated electrons for their obvious physical applications in condensed matter physics. Among these are the supersymmetric (SUSY) t - J model [7, 8] and supersymmetric generalizations of the Hubbard model [9, 10]. Particularly, it would be beneficial if these models could provide some knowledge of the transitions between the metallic, insulating and superconducting phases. It has long been known that an insight into these properties may be gained by studying the effects of the boundary conditions on such models (see, e.g., [11, 12]). Subsequently several authors have studied electronic models with twisted boundary conditions [16–19].

Integrable models with twisted boundary conditions may be formulated within the framework of quantum (super)algebras. This notion of twisting is more general than that usually treated in electronic models whereby a twisted boundary condition is thought of as the introduction of a phase factor into the periodicity of the model. Here our twisted boundary conditions correspond to more general transformations of the local states and have their origins in the underlying symmetry of the model [20]. For a given R -matrix $R(u)$ any matrix M satisfying $[M_1 M_2, R(u)] = 0$ allows an integrable model to be constructed with the trace over the auxiliary space weighted by M . This is what we will refer to as a twisted boundary condition.

The work of Reshetikhin [21] on multiparametric quantum (super)algebras permits a natural way to construct integrable models also dependent upon additional free parameters.

|| E-mail: angela@if.ufrgs.br

¶ E-mail: jrl@maths.uq.edu.au

+ E-mail: Itzhak.Roditi@cern.ch; permanent address: Centro Brasileiro de Pesquisas Físicas - CBPF, Rua Dr Xavier Sigaud 150, 22290-180, Rio de Janeiro, RJ - Brazil.

In fact the model of Perk and Schultz [22] may be formulated within this framework [23]. We will make it apparent that there exists under suitable conditions a integrability preserving mapping between these multiparametric models and those with twisted boundary conditions. Some particular cases have been studied previously [24–27]. We show that these methods may be applied to any model with an underlying quantum (super)algebra symmetry. By establishing this correspondence between models with twisted boundary conditions and multiparametric models, it is reasonable to expect that the multiparametric solutions may provide suitable test models for describing the various phases associated with correlated electron systems.

As examples we will consider the cases of the SUSY t - J model [8] and the SUSY U model [10]. As well as giving the Hamiltonians for the integrable multiparametric generalizations of these two models, we have also determined the corresponding Bethe ansatz equations which provide the starting point for an investigation into their thermodynamics.

The paper is organized as follows. In section 2 we present the general construction of multiparametric spin chains and their relation to models with twisted boundary conditions. In section 3 we illustrate how our general formalism applies to construct multiparametric versions of the SUSY t - J and U models. The Bethe ansatz equations of the models are also obtained. A summary of our main results is presented in section 4.

2. General construction

Using Reshetikhin's construction [21] for multiparametric quantum algebras, it is a straightforward matter to obtain the associated multiparametric quantum spin chain. Here we demonstrate that under appropriate constraints these models may be transformed to quantum spin chains with twisted boundary conditions; i.e. the additional parameters arising from Reshetikhin's construction may be mapped to the boundary.

Let (A, Δ, R) denote a quasitriangular Hopf (super)algebra where Δ and R denote the co-product and R -matrix, respectively. Suppose that there exists an element $F \in A \otimes A$ such that

$$\begin{aligned} (\Delta \otimes I)(F) &= F_{13}F_{23} & (I \otimes \Delta)(F) &= F_{13}F_{12} \\ F_{12}F_{13}F_{23} &= F_{23}F_{13}F_{12} & F_{12}F_{21} &= I. \end{aligned} \quad (1)$$

Theorem 1 of [21] states that (A, Δ^F, R^F) is also a quasitriangular Hopf algebra with co-product and R -matrix respectively given by

$$\Delta^F = F_{12}\Delta F_{21} \quad R^F = F_{21}RF_{21}. \quad (2)$$

In the case where (A, Δ, R) is an affine quantum (super)algebra from [21] we have that F can be chosen to be

$$F = \exp \sum_{i < j} (H_i \otimes H_j - H_j \otimes H_i) \phi_{ij} \quad (3)$$

where $\{H_i\}$ is a basis for the Cartan subalgebra of the affine quantum (super)algebra and the ϕ_{ij} , $i < j$ are arbitrary complex parameters. For our purposes we will extend the Cartan subalgebra by an additional central extension (not the usual central charge) H_0 which will act as a scalar multiple of the identity operator in any representation.

Suppose that π is a loop representation of the affine quantum superalgebra. We let $R(u)$, $R^F(u)$ be the (super)matrix representatives of R and R^F , respectively, which both

satisfy the Yang–Baxter equation

$$R_{12}(u - v)R_{13}(u)R_{23}(v) = R_{23}(v)R_{13}(u)R_{12}(u - v).$$

For the supersymmetric case it is necessary to impose the multiplication rule

$$(a \otimes b)(c \otimes d) = (-1)^{|b||c|}(ac \otimes bd)$$

for homogeneous supermatrices a, b, c, d where $[a] = 0$ if a is even and $[a] = 1$ if a is odd [28]. However, with an appropriate redefinition of its matrix elements $R(u)$ satisfies the usual (non-graded) Yang–Baxter equation (see, e.g., [29]). We adopt this latter convention throughout.

If $R(u)|_{u=0} = P$ with P the permutation operator, then $R^F(u)|_{u=0} = P$ as a result of (1). We may construct the transfer matrix

$$\begin{aligned} t^F(u) &= \text{str}_0(\pi^{\otimes(N+1)}(I \otimes \Delta_N^F)R_{01}^F) \\ &= \text{str}_0(R_{0N}^F(u)R_{0(N-1)}^F(u) \cdots R_{01}^F(u)) \end{aligned} \tag{4}$$

where Δ_N^F is defined recursively through

$$\begin{aligned} \Delta_N^F &= (I \otimes I \cdots \otimes \Delta^F) \Delta_{N-1}^F \\ &= (\Delta^F \otimes I \cdots \otimes I) \Delta_{N-1}^F. \end{aligned} \tag{5}$$

The subscripts 0 and 1, 2, ..., N denote the auxiliary and quantum spaces, respectively, and str_0 is the supertrace over the zeroth space. From the Yang–Baxter equation it follows that the multiparametric transfer matrices $t^F(u)$ form a commuting family. The associated multiparametric spin chain Hamiltonian is given by

$$\begin{aligned} H^F &= (t^F(u))^{-1} \frac{d}{du} t^F(u) \Big|_{u=0} \\ &= \sum_{i=1}^{N-1} h_{i,i+1}^F + h_{N1}^F \end{aligned} \tag{6}$$

with

$$h^F = \frac{d}{du} P R^F(u) \Big|_{u=0}.$$

Through the use of equation (1) we may alternatively write

$$t^F(u) = \text{str}_0(\pi^{\otimes(N+1)}(I \otimes J_N)[(I \otimes \Delta_N)(F_{10}R_{01}F_{10})(I \otimes J_N)^{-1})$$

with

$$\begin{aligned} J_N &= G_{N-1}G_{N-2} \cdots G_1 \\ G_i &= F_{iN}F_{i(N-1)} \cdots F_{i(i+1)}. \end{aligned} \tag{7}$$

We now define a new transfer matrix

$$\begin{aligned} t(u) &= J_N^{-1}t^F(u)J_N \\ &= \text{str}_0(\pi^{\otimes(N+1)}(I \otimes \Delta_N)(F_{10}R_{01}F_{10})) \end{aligned} \tag{8}$$

where we have employed the convention to let F denote both the algebraic object and its (super)matrix representative. Through further use of equation (1) we may show that

$$t(u) = \text{str}_0(F_{10}F_{20} \cdots F_{N0}R_{0N}(u)R_{0(N-1)}(u) \cdots R_{01}(u)F_{10} \cdots F_{N0})$$

and the associated Hamiltonian is given by

$$\begin{aligned} H &= (t(u))^{-1} \frac{d}{du} t(u) \Big|_{u=0} \\ &= \sum_{i=1}^{N-1} h_{i,i+1} + (F_{N(N-1)} \cdots F_{N1})^2 h_{N1} (F_{1N} \cdots F_{(N-1)N})^2 \end{aligned} \quad (9)$$

where

$$h = \frac{d}{du} P R(u) \Big|_{u=0} .$$

The above Hamiltonian describes a closed system where instead of the usual periodic boundary conditions we now have a more general type of boundary condition. The boundary term in the above Hamiltonian is a global operator; i.e. it acts non-trivially on all sites. However, we can in fact interpret this term as a local operator which couples only the sites labelled 1 and N . It can be shown that the boundary term commutes with the local observables $h_{i,i+1}$ for $i \neq 1, N-1$. This situation is analogous to the closed quantum (super)algebra invariant chains discussed in [30].

From the above construction we may also obtain models with twisted boundary conditions by an appropriate choice of F . Recall that we extend the Cartan subalgebra by the central element H_0 . Let this element act as cI in the representation π where c is some complex number. If we now choose $\phi_{ij} = 0$ for $i \neq 0$ in the expression (3) the matrix F factorizes as $F = M_1^{-1} M_2$ with

$$M = \exp\left(\sum_{i=1}^l c\phi_{0i} H_i\right)$$

and l is the rank of the underlying quantum (super)algebra $U_q(g)$. Using the fact that the R -matrix satisfies

$$[R(u), I \otimes H_i + H_i \otimes I] = 0 \quad i = 1, 2, \dots, l$$

tells us that

$$[R(u), M_1 M_2] = 0.$$

In this case the Hamiltonian (9) reduces to

$$H = \sum_{i=1}^{N-1} h_{i,i+1} + M_1^{2N} h_{N1} M_1^{-2N} \quad (10)$$

which is precisely the form for a system with twisted boundary conditions (see [20]).

3. Examples

In this section we illustrate how our formalism applies to the construction of a multiparametric version of the SUSY t - J model [7] and the SUSY U model [10]. Both models are $gl(2/1)$ invariant and their formulation through the quantum inverse scattering method can be found in [8] and [31], respectively. The first model describes electrons with nearest-neighbour hopping and spin exchange interaction on a chain, while the second can be considered an extension of the Hubbard model with additional pair-hopping and bond-charge interaction terms. These models are of interest because of their possible connection with high- T_c superconductivity. In order to turn our discussion more general, we will in fact handle with their anisotropic or q -deformed versions [32–34]. Of course, in the rational limit $q \rightarrow 1$ all results reduce to their corresponding isotropic ones.

3.1. The supersymmetric t - J model

We begin by introducing the multiparametric $U_q(gl(2/1))$ R -matrix, which in terms of a generic spectral parameter x and a deformation parameter q reads

$$R^F(x) = \begin{pmatrix} a & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & t_1^2 b & 0 & c_- & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & t_2^2 b & 0 & 0 & 0 & c_- & 0 & 0 \\ 0 & c_+ & 0 & b/t_1^2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & a & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & t_3^2 b & 0 & c_- & 0 \\ 0 & 0 & c_+ & 0 & 0 & 0 & b/t_2^2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & c_+ & 0 & b/t_3^2 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & w \end{pmatrix} \quad (11)$$

where

$$\begin{aligned} a &= xq - \frac{1}{xq} & b &= x - \frac{1}{x} \\ c_+ &= x \left(q - \frac{1}{q} \right) & c_- &= \frac{1}{x} \left(q - \frac{1}{q} \right) & w &= -\frac{x}{q} + \frac{q}{x} \end{aligned} \quad (12)$$

and t_1, t_2, t_3 are independent parameters written in terms of $\phi_{01}, \phi_{02}, \phi_{12}$ as

$$\begin{aligned} t_1 &= \exp(-2\phi_{01} + \phi_{02} + \phi_{12}) \\ t_2 &= \exp(-\phi_{01} + \phi_{02} + \phi_{12}) \\ t_3 &= \exp(\phi_{01} - \phi_{12}). \end{aligned} \quad (13)$$

The above matrix has already been presented by Perk and Schultz [22] in their study of a multicomponent generalization of the six-vertex model.

Next we construct the transfer matrix $t^F(x)$ according to equation (4) from which we find the associated multiparametric Hamiltonian (see equation (6)) on a one-dimensional periodic lattice:

$$H^F = \sum_{i=1}^{N-1} h_{i,i+1}^F + h_{N1}^F$$

where

$$\begin{aligned} h_{i,i+1}^F &= - \left(\frac{1}{t_2^2} c_{i\uparrow}^\dagger c_{i+1\uparrow} + t_2^2 c_{i+1\uparrow}^\dagger c_{i\uparrow} + \frac{1}{t_3^2} c_{i\downarrow}^\dagger c_{i+1\downarrow} + t_3^2 c_{i+1\downarrow}^\dagger c_{i\downarrow} \right) - 2 \left[\frac{1}{t_1^2} S_i^+ S_{i+1}^- \right. \\ &\quad \left. + t_1^2 S_i^- S_{i+1}^+ + 2 \cos \gamma \left(S_i^z S_{i+1}^z - \frac{n_i n_{i+1}}{4} \right) \right] + i \sin(\gamma)(n_i - n_{i+1}) \\ &\quad - i \sin(\gamma)(n_i S_{i+1}^z - S_i^z n_{i+1}) - \cos \gamma n_i + 2 \cos \gamma. \end{aligned} \quad (14)$$

Here the $c_{i\pm}^{(\dagger)}$ are spin-up or spin-down annihilation (creation) operators, the S_i spin matrices, the n_i 's occupation numbers of electrons at lattice site i and γ is the anisotropy parameter ($q = e^{i\gamma}$). A similar version of a multiparametric SUSY t - J model has already been discussed in [35]. Notice that here it emerges systematically from our general construction. By setting $t_1, t_2, t_3 \rightarrow 1$ in equation (14), the usual terms of the anisotropic SUSY t - J model [32] can be recovered.

The Hamiltonian (14) can be exactly solved through the algebraic nested Bethe ansatz method. This procedure is carried out in two steps and the Bethe ansatz equations are given by

$$\begin{aligned}
 t_1^{2(N-M_2)} t_2^{2M_2} t_3^{-2M_2} \left(\frac{a(x_k^{(1)})}{b(x_k^{(1)})} \right)^N \prod_{i=1}^{M_1} \frac{a(x_i^{(1)}/x_k^{(1)}) b(x_k^{(1)}/x_i^{(1)})}{b(x_i^{(1)}/x_k^{(1)}) a(x_k^{(1)}/x_i^{(1)})} \prod_{j=1}^{M_2} \frac{b(x_j^{(2)}/x_k^{(1)})}{a(x_j^{(2)}/x_k^{(1)})} &= -1 \\
 k = 1, \dots, M_1 & \\
 t_1^{-2(N-M_1)} t_2^{2(N-M_1)} t_3^{2M_1} (-1)^{M_2} \prod_{i=1}^{M_1} \frac{a(x_k^{(2)}/x_i^{(1)})}{b(x_k^{(2)}/x_i^{(1)})} \prod_{j=1}^{M_2} \frac{a(x_j^{(2)}/x_k^{(2)}) b(x_k^{(2)}/x_j^{(2)})}{b(x_j^{(2)}/x_k^{(2)}) w(x_k^{(2)}/x_j^{(2)})} &= 1 \\
 k = 1, \dots, M_2 &
 \end{aligned} \tag{15}$$

where $x_k^{(m)}$ ($m = 1, 2$; $k = 1, \dots, M_m$) denote the Bethe ansatz parameters, N is the number of lattice sites, M_1 is the number of holes plus down spins and M_2 is the number of holes. We see from equation (15) that the additional parameters t_1, t_2, t_3 have the meaning of external fields (see, e.g., [36, 37])

Following the approach presented in the previous section, we perform the transformation (8) and then set $t_3 = t_2/t_1$ (or $\phi_{12} = 0$, see equation (13)) in order to find the anisotropic SUSY t - J model with twisted boundary conditions (10)

$$H = \sum_{i=1}^{N-1} h_{i,i+1} + h_{N1}$$

where $h_{i,i+1} = \lim_{\{t_1, t_2, t_3 \rightarrow 1\}} h_{i,i+1}^F$ and

$$\begin{aligned}
 h_{N,1} = & - \left[t_2^{2N} c_{N\uparrow}^\dagger c_{1\uparrow} + \frac{1}{t_2^{2N}} c_{1\uparrow}^\dagger c_{N\uparrow} + \left(\frac{t_2}{t_1} \right)^{2N} c_{N\downarrow}^\dagger c_{1\downarrow} + \left(\frac{t_1}{t_2} \right)^{2N} c_{1\downarrow}^\dagger c_{N\downarrow} \right] - 2 \left[t_1^{2N} S_N^+ S_1^- \right. \\
 & + \frac{1}{t_1^{2N}} S_N^- S_1^+ + 2 \cos \gamma \left(S_N^z S_1^z - \frac{n_N n_1}{4} \right) \left. \right] + i \sin(\gamma) (n_N - n_1) \\
 & - i \sin(\gamma) (n_N S_1^z - S_N^z n_1) - \cos \gamma n_N + 2 \cos \gamma.
 \end{aligned} \tag{16}$$

3.2. The supersymmetric U model

Let us now construct a multiparametric version of the anisotropic SUSY U model, which has recently been proposed as a new integrable model for correlated electrons (see [33, 34] for more details).

We begin by recalling the trigonometric R-matrix associated with the one parameter family of four-dimensional representations of $U_q(gl(2/1))$

$$\begin{aligned}
 R(x) &= P \check{R}(x) \\
 \check{R}(x) &= \frac{q^x - q^{2\alpha}}{1 - q^{x+2\alpha}} P_1 + P_2 + \frac{1 - q^{x+2\alpha+2}}{q^x - q^{2\alpha+2}} P_3.
 \end{aligned} \tag{17}$$

Here x and q are respectively the spectral and deformation parameters, and α is a free parameter which arises from the underlying representation. P is the permutation operator and $P_i, i = 1, 2, 3$ are projectors whose explicit form can be found in [34].

We find the corresponding multiparametric R -matrix

$$R^F(x) = \begin{pmatrix} * & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & t_1^2 * & 0 & 0 & * & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & t_2^2 * & 0 & 0 & 0 & 0 & 0 & * & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & t_1^2 t_2^2 * & 0 & 0 & t_1 t_2 t_3 * & 0 & 0 & \frac{t_1 t_2}{t_3} * & 0 & 0 & * & 0 & 0 \\ 0 & * & 0 & 0 & \frac{*}{t_1} & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & * & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & t_1 t_2 t_3 * & 0 & 0 & t_3^2 * & 0 & 0 & * & 0 & 0 & \frac{t_3}{t_1 t_2} * & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & t_1^2 t_3^2 * & 0 & 0 & 0 & 0 & 0 & * & 0 \\ 0 & 0 & * & 0 & 0 & 0 & 0 & 0 & \frac{*}{t_2} & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & \frac{t_1 t_2}{t_3} * & 0 & 0 & * & 0 & 0 & \frac{*}{t_3} & 0 & 0 & \frac{*}{t_1 t_2 t_3} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & * & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & * \frac{t_2^2}{t_3} & 0 & 0 & * \\ 0 & 0 & 0 & * & 0 & 0 & \frac{t_3}{t_1 t_2} * & 0 & 0 & \frac{*}{t_1 t_2 t_3} & 0 & 0 & \frac{*}{t_1^2 t_2} & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & * & 0 & 0 & 0 & 0 & 0 & \frac{*}{t_1^2 t_3} & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & * & 0 & 0 & * \frac{t_3^2}{t_2} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & * \end{pmatrix} \tag{18}$$

where t_1, t_2 and t_3 are independent parameters also given by equation (13) and ‘*’ denote the elements of the R -matrix (17), which can be obtained from the projectors given in [34]. We do not write them explicitly here since we will not need them later. Notice that here, in contrast to the previous case (see equation (11)), the new parameters t_1, t_2, t_3 occupy also non-diagonal positions. This is a peculiarity of higher representations and can also be verified for other higher spin models (e.g., the spin-1 XXZ chain). In fact, our prescription for the element F of the multiparametric R matrix is particularly interesting in these cases, where it is not obvious how to construct R^F .

Next we construct the transfer matrix $t^F(x)$ according to equation (4) from which we find the multiparametric version of the anisotropic SUSY U model (see equation (6))

$$H^F = \sum_{i=1}^{N-1} h_{i,i+1}^F + h_{N1}^F$$

where

$$\begin{aligned} h_{i,i+1}^F = & -\xi c_{i\uparrow}^+ c_{i+1\uparrow} [-\eta^{-1} v^+]^{n_{i\downarrow}} [-\eta^{-1} v^-]^{n_{i+1\downarrow}} - \text{hc} \\ & - \rho c_{i\downarrow}^+ c_{i+1\downarrow} [-\eta v^-]^{n_{i\uparrow}} [-\eta v^+]^{n_{i+1\uparrow}} - \text{hc} + [\alpha]_q^{-1} \xi \rho c_{i\downarrow}^+ c_{i\uparrow}^+ c_{i+1\uparrow} c_{i+1\downarrow} + \text{hc} \\ & + [\alpha]_q^{-1} (n_{i\uparrow} n_{i\downarrow} + n_{i+1\uparrow} n_{i+1\downarrow}) + q^{\alpha+1} (n_{i\uparrow} + n_{i\downarrow} - 1) \\ & + q^{-\alpha-1} (n_{i+1\uparrow} + n_{i+1\downarrow} - 1) \end{aligned} \tag{19}$$

and $\xi = t_3^2/t_2^2$, $\eta = t_1 t_3/t_2$, $\rho = 1/t_1^2 t_3^2$, $v^\pm = \text{sgn}(\alpha) q^{\pm 1/2} ([\alpha + 1]_q / [\alpha]_q)^{1/2}$. In the hc terms one should notice that the parameters $t_i \rightarrow t_i^{-1}$, $i = 1, 2, 3$. Equation (19) is a generalization of the anisotropic SUSY U model [34], which can be recovered in the limit $t_1, t_2, t_3 \rightarrow 1$.

This model can be exactly solved by means of the algebraic nested Bethe ansatz method and the Bethe ansatz equations are given by

$$\begin{aligned} t_1^{2(N-N_2)} t_2^{2N_2} t_3^{-2N_2} \left(\frac{x_k^{(1)} q^{\alpha+1} - q^{-1}}{x_k^{(1)} q^{-\alpha} - 1} \right)^N &= \prod_{j \neq k}^{N_2} \frac{x_k^{(1)} - x_j^{(2)} q^2}{q(x_k^{(1)} - x_j^{(2)})} & k = 1, \dots, N_1 \\ t_1^{-2(N-N_1)} t_2^{2(N-N_1)} t_3^{2N_1} \prod_i^{N_1} \frac{x_i^{(1)} - x_k^{(2)} q^2}{q(x_i^{(1)} - x_k^{(2)})} &= \prod_{j \neq k}^{N_2} \frac{-x_j^{(2)} + x_k^{(2)} q^2}{x_k^{(2)} - x_j^{(2)} q^2} & k = 1, \dots, N_2 \end{aligned} \quad (20)$$

where $x_k^{(m)}$ ($m = 1, 2$; $k = 1, \dots, N_m$) are the Bethe ansatz parameters, N_1 is the total number of spins and N_2 is the number of spins down.

According to the approach presented in the previous section, we perform the transformation (8) and then choose $t_3 = t_2/t_1$ in order to find the anisotropic SUSY U model with twisted boundary conditions, which yields

$$H = \sum_{i=1}^{N-1} h_{i,i+1} + h_{N1}$$

where $h_{i,i+1}$ denotes the local terms of the anisotropic SUSY U model [34] and

$$\begin{aligned} h_{N,1} &= -t_1^{2N} c_{N\uparrow}^+ c_{1\uparrow} (-v^+)^{n_{N\downarrow}} (-v^-)^{n_{1\downarrow}} - \text{hc} - t_2^{2N} c_{N\downarrow}^+ c_{1\downarrow} (-v^-)^{n_{N\uparrow}} (-v^+)^{n_{1\uparrow}} - \text{hc} \\ &+ [\alpha]_q^{-1} (t_1 t_2)^{-2N} c_{N\downarrow}^+ c_{N\uparrow}^+ c_{1\uparrow} c_{1\downarrow} + \text{hc} + [\alpha]_q^{-1} (n_{N\uparrow} n_{N\downarrow} + n_{1\uparrow} n_{1\downarrow}) \\ &+ q^{\alpha+1} (n_{N\uparrow} + n_{N\downarrow} - 1) + q^{-\alpha-1} (n_{1\uparrow} + n_{1\downarrow} - 1). \end{aligned} \quad (21)$$

4. Conclusions

In this paper we have demonstrated a correspondence between multiparametric spin chains and models with twisted boundary conditions in the expectation that this connection will provide further insight into the description of the phase transitions of such integrable systems. Our approach can be applied to any model with an underlying quantum (super)algebra symmetry. We are particularly interested in models which describe systems of correlated electrons and have studied the SUSY t - J and U models as examples.

Another important class of integrable models are those associated with the Temperley–Lieb algebra. In [38] Zhang proposes a systematic method to generate multiparametric extensions of these models. It is possible to adapt the techniques employed in this paper to establish a mapping from models based on the Temperley–Lieb algebra with twisted boundary conditions to associated multiparametric generalizations. With respect to correlated electron systems, an example based on the Temperley–Lieb algebra has been described in [39, 40] to which this procedure can be applied.

Acknowledgments

The authors wish to thank J H H Perk for illuminating discussions held at the International Workshop on Statistical Mechanics and Integrable Systems at Coolangatta, Australia, in July

1997. We also thank D Arnaudon for helpful comments. AF would like to thank CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico) for financial support. JL is supported by an Australian Research Council Postdoctoral Fellowship. IR thanks CERN and the Instituto de Física da UFRGS for their hospitality and CNPq for financial support.

References

- [1] Jimbo M 1985 *Lett. Math. Phys.* **10** 63
- [2] Drinfeld V G 1986 *Proc. Int. Congress of Mathematicians* (Berkeley) p 798
- [3] Bracken A J, Gould M D and Zhang R B 1990 *Mod. Phys. Lett.* **5A** 831
- [4] Yamane H 1991 *Proc. Japanese Acad. A* **67** 108
- [5] Khoroshkin S M and Tolstoy V N 1991 *Commun. Math. Phys.* **141** 599
- [6] Scheunert M 1993 *J. Math. Phys.* **34** 3780
- [7] Anderson P W 1987 *Science* **235** 1196
Zhang F C and Rice T M 1988 *Phys. Rev. B* **37** 3759
- [8] Essler F H L and Korepin V E 1992 *Phys. Rev. B* **46** 9147
Foerster A and Karowski M 1992 *Phys. Rev. B* **46** 9234
Foerster A and Karowski M 1993 *Nucl. Phys. B* **396** 611
- [9] Essler F H L, Korepin V E and Schoutens K 1992 *Phys. Rev. Lett.* **68** 2960
Essler F H L, Korepin V E and Schoutens K 1993 *Phys. Rev. Lett.* **68** 70 73
- [10] Bracken A J, Gould M D, Links J R and Zhang Y-Z 1995 *Phys. Rev. Lett.* **74** 2769
- [11] Byers N and Yang C N 1961 *Phys. Rev. Lett.* **7** 46
- [12] Kohn W 1964 *Phys. Rev.* **133** A171
- [13] Yang C N 1962 *Rev. Mod. Phys.* **34** 696
- [14] Yang C N 1989 *Phys. Rev. Lett.* **63** 2144
- [15] Yang C N and Zhang S 1990 *Mod. Phys. Lett.* **4B** 759
- [16] Shastry B S and Sutherland B 1990 *Phys. Rev. Lett.* **65** 243
- [17] Bares P A, Carmelo J M P, Ferrer J and Horsch P 1992 *Phys. Rev. B* **46** 14624
- [18] Arrachea L, Aligia A A and Gagliano E 1996 *Phys. Rev. Lett.* **76** 4396
- [19] Bedürftig G, Essler F H L and Frahm H 1997 *Nucl. Phys. B* **498** 697
- [20] de Vega H J 1984 *Nucl. Phys. B* **240** 495
Yung C M and Batchelor M T 1995 *Nucl. Phys.* **446** [FS] 461
- [21] Reshetikhin N 1990 *Lett. Math. Phys.* **20** 331
- [22] Perk J H H and Schultz C L 1981 *Phys. Lett.* **84A** 407
Schultz C L 1983 *Physica* **122A** 71
- [23] Okado M and Yamane H 1990 *ICM-90 Satellite Conf. Proc.* ed M Kashiwara and T Miwa (Berlin: Springer) p 289
- [24] Alcaraz F C, Barber M N and Batchelor M T 1988 *Ann. Phys., NY* **182** 280
- [25] Pasquier V and Saleur H 1990 *Nucl. Phys. B* **316** 523
- [26] R-Monteiro M, Roditi I, Rodrigues L M C S, Sciuto S 1995 *Phys. Lett.* **354B** 389
- [27] Foerster A, Roditi I and Rodrigues L M C S 1996 *Mod. Phys. Lett.* **11A** 987
- [28] Kulish P P and Sklyanin E K 1982 *J. Sov. Maths.* **19** 1596
- [29] Delius G W, Gould M D, Links J R and Zhang Y-Z 1995 *Int. J. Mod. Phys. A* **10** 3259
- [30] Links J and Foerster A 1997 *J. Phys. A: Math. Gen.* **30** 2483
- [31] Hibberd K E, Gould M D, Links J R 1996 *Phys. Rev. B* **54** 8430
Ramos P B and Martins M J 1996 *Nucl. Phys. B* **474** [FS] 678
- [32] Foerster A and Karowski M 1993 *Nucl. Phys. B* **408** 512
- [33] Bariev R Z, Klümper A and Zittartz J 1995 *Europhys. Lett.* **32** 85
- [34] Gould M D, Hibberd K E, Links J R and Zhang Y-Z 1996 *Phys. Lett.* **212A** 156
Hibberd K E, Gould M D, Links J R 1996 *J. Phys. A: Math. Gen.* **29** 8053
- [35] Arnaudon D, Chryssomalakos C and Frappat L 1995 *J. Math. Phys.* **36** 5262
- [36] de Vega H J and Lopes E 1991 *Phys. Rev. Lett.* **67** 489
- [37] Hlavaty L 1985 Solution to the YBE corresponding to the XXZ models in an external magnetic field *Dubna preprint* E5-85-959
- [38] Zhang R B 1991 *J. Phys. A: Math. Gen.* **24** L535
- [39] Links J 1996 *J. Phys. A: Math. Gen.* **29** L69
- [40] Foerster A, Links J and Roditi I 1997 *Mod. Phys. Lett.* **12A** 1035