

Duality and the modular group in the quantum Hall effect

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1999 J. Phys. A: Math. Gen. 32 L243

(<http://iopscience.iop.org/0305-4470/32/21/101>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.105

The article was downloaded on 02/06/2010 at 07:32

Please note that [terms and conditions apply](#).

LETTER TO THE EDITOR

Duality and the modular group in the quantum Hall effect

Brian P Dolan

Department of Mathematical Physics, National University of Ireland, Maynooth, Republic of Ireland

and

Dublin Institute for Advanced Studies, 10 Burlington Road, Dublin, Republic of Ireland

E-mail: bdolan@thphys.may.ie

Received 22 December 1998

Abstract. We explore the consequences of introducing a complex conductivity into the quantum Hall effect. This leads naturally to an action of the modular group on the upper-half complex conductivity plane. Assuming that the action of a certain subgroup, compatible with the law of corresponding states, commutes with the renormalization group flow, we derive many properties of both the integer and fractional quantum Hall effects including: universality; the selection rule $|p_1q_2 - p_2q_1| = 1$ for transitions between quantum Hall states characterized by filling factors $\nu_1 = p_1/q_1$ and $\nu_2 = p_2/q_2$; critical values of the conductivity tensor; and Farey sequences of transitions. Extra assumptions about the form of the renormalization group flow lead to the semicircle rule for transitions between Hall plateaux.

The purpose of this letter is to explore the consequences of the proposal, made in [1] and examined further in [2, 3], that the hierarchical structure of the zero-temperature integer and fractional quantum Hall effects can be interpreted in terms of the properties of a subgroup of the modular group, $Sl(2, \mathbb{Z}) := \Gamma(1)$ —specifically the subgroup which consists of elements of $\Gamma(1)$ whose bottom left entry is even, sometimes denoted $\Gamma_0(2)$ in the mathematical literature. This group acts on the upper-half complex plane, parametrized by the complex conductivity, $\sigma = \sigma_{xy} + i\sigma_{xx}$, in units of $\frac{e^2}{h}$, and is generated by two operations, $T : \sigma \rightarrow \sigma + 1$ and $X : \sigma \rightarrow \frac{\sigma}{2\sigma+1}$. If $\gamma = \begin{pmatrix} a & b \\ 2c & d \end{pmatrix} \in \Gamma_0(2)$, with a, b, c , and $d \in \mathbb{Z}$ and $ad - 2cb = 1$, then $\gamma(\sigma) = \frac{a\sigma+b}{2c\sigma+d}$. Thus $T = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ and $X = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}$. Some consequences of this assumption for the phase diagram in the σ -plane were examined in [2] and in the second of these references the author notes that there is a connection with the work of Kivelson *et al* [4], but remarks that the comparison between [2] and [4] is not immediate. One of the aims of this paper is to explore the relation between these two approaches.

Following [1–3], it will be assumed that the phase diagram for the quantum Hall effect can be generated by the action of $\Gamma_0(2)$ on the upper-half σ plane. This immediately implies the ‘law of corresponding states’ of [4] and [5]. At Hall plateaux we have $\sigma_{xx} = 0$ and $\sigma_{xy} = s$ where s is a ratio of two mutually prime integers, with odd denominator (note that s is being used here to label the quantum phases and is denoted by s_{xy} in [4]). Plateaux can be related to each other by repeated action of T and X . At the centre of the plateaux, the filling factor, ν , is equal to the ratio $s = p/q$ and $T : \nu \rightarrow \nu + 1$ is the Landau level addition transformation of [4]

while $X : \nu \rightarrow \frac{\nu}{2\nu+1}$ is the flux attachment transformation. The particle–hole transformation $\nu \rightarrow 1 - \nu$, can be realized as the outer automorphism $\sigma \rightarrow 1 - \bar{\sigma}$ acting on the upper-half plane, where $\bar{\sigma} = \sigma_{xy} - i\sigma_{xx}$ (it will be assumed throughout that the electron spins are split, for the spin-degenerate case one must rescale $\sigma \rightarrow 2\sigma$).

The upper-half σ -plane can be completely covered by copies of a single ‘tile’, or fundamental region (see e.g. [6]), related to each other by elements of $\Gamma_0(2)$. The fundamental region has cusps at 0 and 1, linked by a semicircle of unit diameter, and consists of a vertical strip of unit width constructed above this semicircle. By assumption, all allowed quantum Hall transitions are images of the transition $\nu = 0 \rightarrow \nu = 1$ under some $\gamma \in \Gamma_0(2)$, and hence also linked by a semicircle.

Each such semicircle has a special point, in addition to the endpoints, which is a fixed point of $\Gamma_0(2)$ in the following sense. The point $\sigma^* = \frac{1+i}{2}$ is left fixed by $\gamma^* = \begin{pmatrix} 1 & -1 \\ 2 & -1 \end{pmatrix}$. Similarly the points obtained from σ^* by the other elements of $\Gamma_0(2)$, $\sigma_\gamma^* := \gamma(\sigma^*)$, are left fixed by $\gamma\gamma^*\gamma^{-1}$. It can be shown that the imaginary part, $\text{Im}(\sigma_\gamma^*) \leq \frac{1}{2}$ or $\text{Im}(\sigma_\gamma^*) = \infty, \forall \gamma$. The points σ_γ^* can be interpreted as critical points for the transition $\gamma(0) \leftrightarrow \gamma(1)$ if we further assume that the action of $\Gamma_0(2)$ commutes with the renormalization group (RG) flow. For if σ_γ^* were not a RG fixed point, we could move to an infinitesimally close point $\phi(\sigma_\gamma^*) \neq \sigma_\gamma^*$ with a RG transformation, ϕ . Demanding $\gamma \circ \phi(\sigma_\gamma^*) = \phi \circ \gamma(\sigma_\gamma^*) = \phi(\sigma_\gamma^*)$ then implies that $\phi(\sigma_\gamma^*)$ is also left invariant by γ . But the fixed points of $\Gamma_0(2)$ are isolated, so there is no other fixed point infinitesimally close to σ_γ^* . Hence $\phi(\sigma_\gamma^*) = \sigma_\gamma^*$ and σ_γ^* must be a RG fixed point. The end points of the arches, at $\sigma = \nu$ with $\nu = p/q$ rational, are also fixed points of $\Gamma_0(2)$. For q odd these are stable Hall states. Note, however, that a fixed point of the RG need not necessarily be a fixed point of $\Gamma_0(2)$ —but there is no experimental evidence of such extraneous fixed points of the RG.

Thus the fixed points of $\Gamma_0(2)$ must be fixed points of the RG, i.e. critical points. This




Figure 1. The phase structure in the upper-half complex σ plane. The solid curves represent phase boundaries and the dotted curves allowed transitions. Points where dotted and solid curves cross are critical points.




Figure 2. A magnified view of the phase structure in the upper-half complex σ plane.

leads to the topology of the flow diagram of [2], reproduced here in figures 1 and 2 where solid curves represent phase boundaries and dashed curves represent quantum Hall transitions. This implies the flow diagram proposed in [7], with its experimental support [8] and is also compatible with the phase diagram derived in [4]. That $\sigma^* = \frac{1+i}{2}$ is a critical point for the lowest Landau level was argued in [9]. Phase boundaries and transitions are represented by semicircles in the figures, but this is not forced on us by the $\Gamma_0(2)$ symmetry. They could be distorted from this geometry, provided that all phase boundaries are copies of a distorted ‘fundamental’ phase boundary (running from $\frac{1}{2}$ to $\frac{1}{2} + i\infty$) under the action of $\Gamma_0(2)$. Similarly, the dashed transition trajectories must all be copies of a distortion of the ‘fundamental’ arch spanning 0 and 1. Note, however, that the *fixed points are immovable*. A useful aspect of the semicircular arches used in the figures is that the intersection of any solid phase boundary with a dashed transition is a fixed point of $\Gamma_0(2)$, as are the end points of the arches (which are rational numbers or points at $\sigma = r + i\infty$ for integral or half-integral r). Any distortion from semicircular geometry must leave the endpoints and intersections of phase boundaries and transition trajectories pinned at the fixed points of $\Gamma_0(2)$.

As in [4], the phase diagram generated by $\Gamma_0(2)$ determines which transitions are allowed and which are not. Thus, for example, $s : \frac{1}{3} \rightarrow 0$ is allowed while $s : \frac{1}{3} \rightarrow \frac{1}{7}$ is not. All allowed transitions are generated by acting on the arch passing through $\sigma = 0$ and $\sigma = 1$ by some, $\gamma \in \Gamma_0(2)$. This allows the derivation a selection rule for a transition $s_1 = p_1/q_1 \rightarrow s_2 = p_2/q_2$, where q_1 and q_2 are odd, and the pairs p_i and q_i ($i = 1, 2$) are relatively prime (for brevity we shall not always distinguish below between s , labelling the

quantum Hall phase, and ν , the filling factor, except where necessary for comparison with [4]—on the real axis, when $\sigma_{xx} = 0$, they are the same). We shall see that a transition is allowed if and only if $p_1q_2 - p_2q_1 = \pm 1$.

From the above assumptions we have (relabelling if necessary) $\nu_1 = \gamma(0)$, $\nu_2 = \gamma(1)$. Thus $\frac{p_1}{q_1} = \frac{b}{d}$ and $\frac{p_2}{q_2} = \frac{a+b}{2c+d}$ where $\gamma = \begin{pmatrix} a & b \\ 2c & d \end{pmatrix} \in \Gamma_0(2)$. Since $ad - 2cb = 1$, b and d are mutually prime, by an elementary result of number theory, hence (taking plus signs without loss of generality) $b = p_1$, $d = q_1$. Similarly $(a+b)d - (2c+d)b = 1$ implies that $a+b$ and $2c+d$ are mutually prime, hence $a+b = p_2$ and $2c+d = q_2$. Thus $\gamma = \begin{pmatrix} p_2 - p_1 & p_1 \\ q_2 - q_1 & q_1 \end{pmatrix}$ and the condition $\det \gamma = 1$ then requires $p_2q_1 - p_1q_2 = 1$. The only possible exception to this rule would be a transition from $\nu = n \rightarrow \nu = m$ ($n, m \in \mathbb{Z}$), which could occur by going first from $\sigma = n$ to $\sigma = n + i\infty$ and then in to $\sigma = m$ from $\sigma = m + i\infty$. In a real experiment the maximum value of $|\sigma|$ would presumably be finite, due to impurities.

One can determine sequences of allowed transitions as follows. Suppose $\nu_0 = p_0/q_0$, with q_0 odd, is an allowed state, with p_0 and q_0 relatively prime. Consider the sequence $\nu_n = \frac{kn+p_0}{ln+q_0} := \frac{p_n}{q_n}$ for $n, k, l \in \mathbb{Z}$, where l is even (so that q_n is odd). Then $p_{n+1}q_n - p_nq_{n+1} = \pm 1 \Leftrightarrow kq_0 - lp_0 = \pm 1$. Thus a transition $\nu_{n+1} \rightarrow \nu_n$ is allowed provided $|kq_0 - lp_0| = 1$. In this way we can, for example, generate the three sequences

$$\begin{aligned} \frac{1}{3} &\rightarrow \frac{2}{5} \rightarrow \frac{3}{7} \rightarrow \frac{4}{9} \rightarrow \frac{5}{11} \rightarrow \frac{6}{13} \rightarrow \dots \\ \dots &\rightarrow \frac{7}{13} \rightarrow \frac{6}{11} \rightarrow \frac{5}{9} \rightarrow \frac{4}{7} \rightarrow \frac{3}{5} \rightarrow \frac{2}{3} \rightarrow 1 \\ \frac{2}{3} &\rightarrow \frac{5}{7} \rightarrow \frac{8}{11} \rightarrow \frac{11}{15} \rightarrow \dots \end{aligned} \quad (1)$$

plus higher sequences obtained by adding an integer to each term in these sequences. Such sequences are called Farey sequences and their relevance to the quantum Hall effect was examined in [10]. Note that a given experiment may jump from one sequence to another. Thus

$$\dots \rightarrow \frac{3}{5} \rightarrow \frac{2}{3} \rightarrow \frac{5}{7} \rightarrow \dots$$

is observed in [11].

Each transition contains a critical point given by $\gamma(\sigma^*)$. Thus if $\gamma = \begin{pmatrix} a & b \\ 2c & d \end{pmatrix}$, the critical point is at

$$\sigma_\gamma^* = \frac{2ac + 2bc + ad + 2bd + i}{2d^2 + 4cd + 4c^2} = \frac{(p_1q_1 + p_2q_2) + i}{(q_1^2 + q_2^2)} \quad (2)$$

when the transition goes from $\nu_1 = \gamma(0) = b/d = p_1/q_1$ to $\nu_2 = \gamma(1) = \frac{a+b}{2c+d} = p_2/q_2$. The parameters of γ can be related to physical parameters as follows. Following [5], let η be the effective charge of the quasi-holes of a Hall state, $e^* = \eta$, θ the statistical parameter (i.e. the phase of the two quasi-particle wavefunction changes by $\pi\theta$ when the positions of the two particles are exchanged) and s be the Hall state, with magic filling factor $\nu = s$. Then the critical conductivity for a transition from $s = \nu$ to $s' = \nu - \eta^2/\theta$ is given by equation (26) of [5] (in dimensionless units)

$$\sigma_{xx} = \frac{\eta^2}{1 + \theta^2} \quad \sigma_{xy} = s - \theta \frac{\eta^2}{1 + \theta^2}. \quad (3)$$

Equating these with the critical values in equation (2), there are two possibilities, depending on whether $\nu = \gamma(1)$ or $\gamma(0)$:

$$(i) \quad \nu = \frac{a+b}{2c+d} = \frac{p_2}{q_2} \quad \theta = \frac{d}{2c+d} = \frac{q_1}{q_2} \quad \eta^2 = \frac{1}{(2c+d)^2} = \frac{1}{q_2^2} \quad (4)$$

$$(ii) \quad \nu = \frac{b}{c} = \frac{p_1}{p_2} \quad \theta = -\frac{(2c+d)}{d} = -\frac{q_2}{q_1} \quad \eta^2 = \frac{1}{d^2} = \frac{1}{q_1^2}. \quad (5)$$

Table 1. Some examples of allowed transitions. The matrix γ maps the points $\sigma = 0$ and $\sigma = 1$ to the transition indicated in the leftmost column. Some representative experimental support (not exhaustive) is also indicated. The last two columns assume the semicircle law (ρ is the resistivity).

Transition $v_1 \rightarrow v_2$	γ	Critical conductivity	Critical resistivity	σ at σ_{xx}^{Max}	ρ at ρ_{xx}^{Max}
$n + 1 \rightarrow n$	$\begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix}$	$\frac{(2n+1)+i}{2}$	$\frac{(2n+1)+i}{2n^2+2n+1}$ ^(a)	$\frac{(2n+1)+i}{2}$	$\frac{(2n+1)+i}{2n(n+1)}$ ^(b)
$\frac{1}{2n+1} \rightarrow 0$	$\begin{pmatrix} 1 & 0 \\ 2n & 1 \end{pmatrix}$	$\frac{(2n+1)+i}{2(2n^2+2n+1)}$ ^(c)	$(2n + 1) + i$ ^(d)	$\frac{1+i}{2(2n+1)}$	$(2n + 1) + i\infty$
$\frac{n}{2n+1} \rightarrow \frac{n+1}{2n+3}$	$\begin{pmatrix} 1 & n \\ 2 & 2n + 1 \end{pmatrix}$	$\frac{(4n^2+6n+3)+i}{2(4n^2+8n+5)}$	$\frac{(4n^2+6n+3)+i}{2n^2+2n+1}$	$\frac{(4n^2+6n+1)+i}{2(2n+1)(2n+3)}$	$\frac{(4n^2+6n+1)+i}{2n(n+1)}$ ^(e)
$\frac{3n+2}{4n+3} \rightarrow \frac{3n+5}{4n+7}$	$\begin{pmatrix} 3 & 3n + 2 \\ 4 & 4n + 3 \end{pmatrix}$	$\frac{(24n^2+58n+41)+i}{2(16n^2+40n+29)}$	$\frac{(24n^2+58n+41)+i}{18n^2+42n+29}$	$\frac{(24n^2+58n+29)+i}{2(4n+3)(4n+7)}$	$\frac{(24n^2+58n+29)+i}{2(3n+2)(3n+5)}$ ^(f)

(a) These points all lie on the semicircle $\rho = i - e^{i\theta}, 0 \leq \theta \leq \pi$. For $n = 1$ see [21].
 (b) Assumes $n \neq 0$.
 (c) These points all lie on the semicircle $\sigma = \frac{1}{2}(i - e^{i\theta}), 0 \leq \theta \leq \pi$.
 (d) For $n = 0$ see [22] and [23], for $n = 1$ see [22] and [24], for $n = 2$ see [25].
 (e) Assumes $n \neq 0$. For $n = 1, \dots, 5$ and $n = -3, \dots, -7$ see [14].
 (f) For $n = 0$ see [14].

In both cases we reproduce the result, that $\eta = \pm 1/q$, [12] and [13]. Note in passing that the transition from bosonic to fermionic conductivities given by equation (14) of [4] is implemented by the action of an element of $\Gamma(1)$ which is not in $\Gamma_0(2)$. Thus $\sigma = \gamma(\sigma^{(b)})$ where $\sigma^{(b)} = \sigma_{xy}^{(b)} + i\sigma_{xx}^{(b)}$ is the complex conductivity of the bosonic Chern–Simons action and $\gamma = \frac{1}{\eta} \begin{pmatrix} \eta^2 - \theta s & s \\ -\theta & 1 \end{pmatrix}$. The above discussion gives the explicit connection between the Chern–Simons analysis of [4] and the group theory analysis of [2].

We make a final comment about the ‘semicircle’ law of [14–16]. By assumption, each quantum Hall transition can be obtained from the one between 0 and 1, passing through $\sigma^* = \frac{1+i}{2}$, by the action of some element of $\Gamma_0(2)$. Since $\Gamma_0(2)$ maps semicircles built on the real axis into other such semicircles we can deduce the ‘semicircle law’ of [14–16] by making one extra assumption—that the ‘fundamental’ arch between 0 and 1 is a semicircle. This implies that *all* other transitions are semicircles and allows predictions to be made of the maximum values of σ_{xx} and ρ_{xx} in any allowed transition, $v_1 \rightarrow v_2$, as well as the values of σ_{xy} and ρ_{xy} at which they occur. Thus the maximum value of σ_{xx} is at $\sigma_{xx}^{max} = \frac{v_1 - v_2}{2}$, where $\sigma_{xy} = \frac{v_1 + v_2}{2}$ (where $v_1 > v_2$). In general, this does not coincide with the critical value $\sigma_{\gamma}^* = \gamma(\sigma^*)$, except for the integer transitions (table 1).

The maximum value of ρ_{xx} is found by constructing the semicircle through $\frac{1}{v_1}$ and $\frac{1}{v_2}$ (provided neither vanishes). Thus $\rho_{xx}^{max} = \frac{1}{2}(\frac{1}{v_2} - \frac{1}{v_1})$, where $\rho_{xy} = \frac{1}{2}(\frac{1}{v_2} + \frac{1}{v_1})$. Some representative examples are shown in table 1.

To summarize, assuming (as in [3]) that the phase and flow diagram for the upper-half complex conductivity plane can be generated from an action of $\Gamma_0(2)$ which commutes with the RG, one deduces: (i) that all critical points are given by $\gamma(\sigma^*)$, where $\sigma^* = \frac{1+i}{2}$, with $\gamma \in \Gamma_0(2)$; (ii) the phase diagram of [4, 2, 8]; (iii) the laws of corresponding states [4, 5]; and (iv) the selection rule $|p_1q_2 - p_2q_1| = 1$, dictating which transitions are allowed and which are forbidden. Lastly, the semicircle law of [14–16] is compatible with, but not implied by, $\Gamma_0(2)$.

It should be noted that the full modular group does *not* provide the correct phase structure, since it would imply further critical points at the images of $\sigma = i$ and $\sigma = \frac{1+i\sqrt{3}}{2}$, under

$\gamma \in \Gamma(1)$, which are not observed experimentally. The appearance of $\Gamma_0(2)$ is due to the extension of Kramers–Wannier duality $\sigma_{xx} \rightarrow 1/\sigma_{xx}$ to the whole complex plane. It was argued in [17] that this extension leads naturally to $\Gamma(1)$ acting on the upper-half complex plane, for a coupled clock model. This was applied to the quantum Hall effect in [18, 19]. It appears to have been noted first in [2] that the subgroup $\Gamma_0(2)$ has the special property of preserving the parity of the denominator for rational $\nu = p/q$. The subgroup $\Gamma(2)$, consisting of all elements of $\Gamma(1)$ which are congruent to the identity, mod 2, was also considered in [2] and has been further investigated in [20]. Note, however, that there is no element of $\Gamma(2)$ which leaves $\sigma^* = \frac{1+i}{2}$ fixed, indeed there is no element of $\Gamma(2)$ which leaves *any* σ with $\infty > \text{Im}(\sigma) > 0$ fixed.

It is a pleasure to thank Jan Pawłowski for discussions about the RG flow of the quantum Hall effect.

References

- [1] Lütken C A and Ross G G 1993 *Phys. Rev. B* **48** 2500
- [2] Lütken C A 1993 *Nucl. Phys. B* **396** 670
Lütken C A 1993 *J. Phys. A: Math. Gen.* **26** L811–17
- [3] Burgess C P and Lütken C A 1997 *Nucl. Phys. B* **500** 367
- [4] Kivelson S, Lee D-H and Zhang S-C 1992 *Phys. Rev. B* **46** 2223
- [5] Lee D-H, Kivelson S and Zhang S-C 1992 *Phys. Rev. Lett.* **68** 2386
- [6] Rankin R A 1977 *Modular Forms and Functions* (Cambridge: Cambridge University Press)
- [7] Khmel'nitskii D E 1983 *Pis'ma Zh. Eksp. Teor. Fiz.* **38** 454 (Engl. transl. 1983 *JETP Lett.* **38** 552)
- [8] Wei H P, Chang A M, Tsui D C, Pruisken A M M and Razeghi M 1986 *Surf. Sci.* **170** 238
- [9] Huo Y, Hetzel R E and Bhatt R N 1993 *Phys. Rev. Lett.* **70** 481
- [10] Zang J and Birman J L 1993 *Phys. Rev. B* **47** 16 305
- [11] Willett R, Eisenstein J P, Störmer H L, Tsui D C, Gossard A C and English J H 1987 *Phys. Rev. Lett.* **59** 1776
- [12] Laughlin R B 1983 *Phys. Rev. Lett.* **50** 1395
Haldane F D M 1983 *Phys. Rev. Lett.* **51** 605
- [13] Clark R G, Mallett J R, Haynes S R, Harris J J and Foxon C T 1988 *Phys. Rev. Lett.* **60** 1747
- [14] Dykhne A M and Ruzin I M 1994 *Phys. Rev. B* **50** 2369
- [15] Ruzin I and Feng S 1995 *Phys. Rev. Lett.* **74** 154
- [16] Ruzin I M, Cooper N R and Halperin B I 1996 *Phys. Rev. B* **53** 1558
- [17] Cardy J L and Rabinovici E 1982 *Nucl. Phys. B* **205** 1
Cardy J L 1982 *Nucl. Phys. B* **205** 17
- [18] Shapere A and Wilczek F 1989 *Nucl. Phys. B* **320** 669
- [19] Lütken C A and Ross G G 1992 *Phys. Rev. B* **45** 11 837
- [20] Georgelin Y and Wallet J-C 1997 *Phys. Lett. A* **224** 303
Georgelin Y, Masson T and Wallet J-C 1997 *J. Phys. A: Math. Gen.* **30** 5065
- [21] Shahar D, Tsui D C, Shayegan M, Shimshoni E and Sondhi S L 1997 *Phys. Rev. Lett.* **79** 479
- [22] Shahar D, Tsui D C, Shayegan M, Bhatt R N and Cunningham J E 1995 *Phys. Rev. Lett.* **74** 4511
Shahar D, Tsui D C, Shayegan M, Cunningham J E, Shimshoni E and Sondhi S L 1997 *Solid State Commun.* **102** 817
(Shahar D, Tsui D C, Shayegan M, Cunningham J E, Shimshoni E and Sondhi S L 1996 *Preprint cond-mat/9607127*)
- [23] Hilke M, Shahar D, Song S H, Tsui D C, Xie Y H and Monroe D 1997 *Phys. Rev. B* **56** R15 545
- [24] Wong L W, Jiang H W, Trivedi N and Palm E 1995 *Phys. Rev. B* **51** 18 033
- [25] Jiang H W, Willett R L, Stormer H L, Tsui D C, Pfeiffer L N and West K W 1990 *Phys. Rev. Lett.* **65** 633