

$T = 0$ partition functions for Potts antiferromagnets on square lattice strips with (twisted) periodic boundary conditions

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

1999 J. Phys. A: Math. Gen. 32 L489

(<http://iopscience.iop.org/0305-4470/32/46/102>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.111

The article was downloaded on 02/06/2010 at 07:49

Please note that [terms and conditions apply](#).

LETTER TO THE EDITOR

 $T = 0$ partition functions for Potts antiferromagnets on square lattice strips with (twisted) periodic boundary conditionsNorman Biggs[†] and Robert Shrock[‡][†] Centre for Discrete and Applicable Mathematics, London School of Economics, London WC2A 2AE, UK[‡] Institute for Theoretical Physics, State University of New York, Stony Brook, NY 11794-3840, USA

E-mail: n.l.biggs@lse.ac.uk and robert.shrock@sunysb.edu

Received 27 July 1999

Abstract. We present exact calculations of the zero-temperature partition function for the q -state Potts antiferromagnet (or, equivalently, the chromatic polynomial) for two families of arbitrarily long strip graphs of the square lattice with periodic boundary conditions in the transverse direction and (i) periodic and (ii) twisted periodic boundary conditions in the longitudinal direction, so that the strip graphs are embedded on (i) a torus and (ii) a Klein bottle. In the limit of infinite length, we calculate the exponent of the entropy, $W(q)$, show it to be the same for (i) and (ii), and determine its analytic structure.

The chromatic polynomial $P(G, q)$ counts the number of ways that one can colour a graph G with q colours such that no two adjacent vertices have the same colour [1] (for reviews, see [2–4]). The least positive integer q for which $P(G, q)$ is nonzero is the chromatic number, $\chi(G)$. Besides its intrinsic mathematical interest, the chromatic polynomial has an important connection with statistical mechanics since it is the zero-temperature partition function of the q -state Potts antiferromagnet (AF) [5, 6] on G : $P(G, q) = Z(G, q, T = 0)_{PAF}$. The Potts AF exhibits nonzero ground state entropy $S_0 \neq 0$ (without frustration) for sufficiently large q on a given lattice graph and is thus an exception to the third law of thermodynamics. This is equivalent to a ground state degeneracy per site $W > 1$, since $S_0 = k_B \ln W$. Denoting the number of vertices of G as $n = v(G)$ and assuming a reasonable definition of $\{G\} = \lim_{n \rightarrow \infty} G$, we have $W(\{G\}, q) = \lim_{n \rightarrow \infty} P(G, q)^{1/n}$. Since $P(G, q)$ is a polynomial, one can generalize q from \mathbb{Z}_+ to \mathbb{C} . The zeros of $P(G, q)$ in the complex q plane are called chromatic zeros; a subset of these may form an accumulation set in the $n \rightarrow \infty$ limit, denoted \mathcal{B} , which is the continuous locus of points where $W(\{G\}, q)$ is nonanalytic [7, 8, 16] (\mathcal{B} may be null, and W may also be nonanalytic at certain discrete points). The maximal region in the complex q plane to which one can analytically continue the function $W(\{G\}, q)$ from physical values where there is nonzero ground state entropy is denoted R_1 . The maximal value of q where \mathcal{B} intersects the (positive) real axis is labelled $q_c(\{G\})$.

§ At certain special points q_s (typically $q_s = 0, 1, \dots, \chi(G)$), one has the noncommutativity of limits $\lim_{q \rightarrow q_s} \lim_{n \rightarrow \infty} P(G, q)^{1/n} \neq \lim_{n \rightarrow \infty} \lim_{q \rightarrow q_s} P(G, q)^{1/n}$, and hence it is necessary to specify the order of the limits in the definition of $W(\{G\}, q_s)$ [7]. We use the first order of limits here; this has the advantage of removing certain isolated discontinuities in W .

We consider strips of the square lattice with arbitrary length $L_x = m$ vertices and fixed width L_y vertices (with the longitudinal and transverse directions taken to be \hat{x} and \hat{y}). The chromatic polynomials for the cyclic and Möbius strip graphs of the square lattice were calculated for $L_y = 2$ in [9] (see also [10–12]) and for $L_y = 3$ in [13, 14]. After studies of the chromatic zeros for $L_y = 2$ in [9, 15, 16], W and \mathcal{B} were determined for this case in [7] and for $L_y = 3$ in [13]. An important question concerns the effect of boundary conditions (BCs), and hence graph topology, on P , W , and \mathcal{B} . We use the symbols FBC_y and PBC_y for free and periodic transverse boundary conditions and FBC_x , PBC_x , and TPBC_x for free, periodic, and twisted periodic longitudinal boundary conditions. The term ‘twisted’ means that the longitudinal ends of the strip are identified with reversed orientation. These strip graphs can be embedded on surfaces with the following topologies†: (i) (FBC_y , FBC_x): strip; (ii) (PBC_y , FBC_x): cylindrical; (iii) (FBC_y , PBC_x): cylindrical (denoted cyclic here); (iv) (FBC_y , TPBC_x): Möbius; (v) (PBC_y , PBC_x): torus; and (vi) (PBC_y , TPBC_x): Klein bottle‡. Here we present and analyse chromatic polynomials for the strip graph of the square lattice with $L_y = 3$ (i.e. cross sections forming triangles) and boundary conditions of type (v) and (vi): torus and Klein bottle. We recall that unlike graphs of type (i)–(v), the Klein bottle graph (vi) cannot be embedded without self-intersection in \mathbb{R}^3 . For $L_x = m \geq 2$ where they are well defined, the $L_y = 3$ torus and Klein bottle graphs have $n = L_x L_y$ vertices, $e = 2n$ edges, the same girth g (length of minimum closed circuit) and number k_g of circuits of length g , and the respective chromatic numbers $\chi = 3$ and $\chi = 4$.

We label a particular type of strip graph as G_s and the specific graph of length $L_x = m$ repeated subgraph units, e.g. columns of squares in the case of the square strip, as $(G_s)_m$. If one thinks of the graph as embedded on a rectangular strip of paper, with its upper and lower sides glued together and its longitudinal ends glued with direct or reversed orientation, then L_x is the length of this strip of paper in subgraph units. Writing

$$P((G_s)_m, q) = \sum_{j=0}^{n-1} (-1)^j h_{n-j} q^{n-j} \quad (1)$$

and using the results that [3, 25] $h_{n-j} = \binom{e}{j}$ for $0 \leq j < g - 1$ (whence $h_n = 1$ and $h_{n-1} = e$) and $h_{n-(g-1)} = \binom{e}{g-1} - k_g$, it follows that for m greater than the above-mentioned minimal value, these h_j are the same for the torus and Klein bottle of each type G_s . For a given G_s , as m increases, the h_{n-j} for the torus and Klein bottle graphs become equal for larger j .

A generic form for chromatic polynomials for recursively defined families of graphs, of which strip graphs G_s are special cases, is

$$P((G_s)_m, q) = \sum_{j=1}^{N_\lambda} c_j(q) (\lambda_j(q))^m \quad (2)$$

where $c_j(q)$ and the N_λ terms $\lambda_j(q)$ depend on the type of strip graph G_s but are independent of m .

For an $L_y = 3$, $L_x = m$ strip with (PBC_y , FBC_x) one has [20] $P(sq(L_y = 3)_m, \text{PBC}_y, \text{FBC}_x, q) = q(q-1)(q-2)(q^3 - 6q^2 + 14q - 13)^{m-1}$, whence

$$W(sq(L_y = 3), \text{PBC}_y, \text{FBC}_x, q) = (q^3 - 6q^2 + 14q - 13)^{1/3} \quad (3)$$

with $\mathcal{B} = \emptyset$.

† For the triangular lattice with cylindrical BCs, W and \mathcal{B} were calculated in [17]. Other calculations of P , W , and \mathcal{B} have been performed for strips having BCs of type (i) [18–20], (ii) [20], (iii) [21, 22], (iv) [14, 21].

‡ These BCs can all be implemented in a manner that is uniform in the length L_x ; the case (vii) (TPBC_y , TPBC_x) with the topology of the projective plane requires different identifications as L_x varies and will not be considered here. For connections between topology and graph properties, see e.g. [23, 24].

In order to calculate P , one may use recursive methods based on iterative use of deletion-contraction theorems [9, 10, 18] or a colouring compatibility matrix method described in [12, 26]. For the $L_y = 3$ torus (t) graphs, one finds

$$P(sq(L_y = 3)_m, PBC_y, PBC_x, q) = \sum_{j=1}^8 c_{t,j}(\lambda_{t,j})^m \tag{4}$$

where

$$\lambda_{t,1} = -1 \quad c_{t,1} = q^3 - 6q^2 + 8q - 1 \tag{5}$$

$$\lambda_{t,2} = q^3 - 6q^2 + 14q - 13 \quad c_{t,2} = 1 \tag{6}$$

$$\lambda_{t,3} = q - 1 \quad c_{t,3} = \frac{(q-1)(q-2)}{2} \tag{7}$$

$$\lambda_{t,4} = q - 4 \quad c_{t,4} = (q-1)(q-2) \tag{8}$$

$$\lambda_{t,5} = q - 2 \quad c_{t,5} = q(q-3) \tag{9}$$

$$\lambda_{t,6} = q - 5 \quad c_{t,6} = \frac{q(q-3)}{2} \tag{10}$$

$$\lambda_{t,7} = -(q^2 - 7q + 13) \quad c_{t,7} = q - 1 \tag{11}$$

$$\lambda_{t,8} = -(q-2)^2 \quad c_{t,8} = 2(q-1). \tag{12}$$

For the $L_y = 3$ Klein (K) bottle graphs one finds

$$P(sq(L_y = 3)_m, PBC_y, TPBC_x, q) = \sum_{j=1}^5 c_{K,j}(\lambda_{K,j})^m \tag{13}$$

where

$$\lambda_{K,1} = \lambda_{t,1} = -1 \quad c_{K,1} = -(q-1) \tag{14}$$

$$\lambda_{K,2} = \lambda_{t,2} = q^3 - 6q^2 + 14q - 13 \quad c_{K,2} = c_{t,2} = 1 \tag{15}$$

$$\lambda_{K,3} = \lambda_{t,3} = q - 1 \quad c_{K,3} = -c_{t,3} = -\frac{(q-1)(q-2)}{2} \tag{16}$$

$$\lambda_{K,4} = \lambda_{t,6} = q - 5 \quad c_{K,4} = c_{t,6} = \frac{q(q-3)}{2} \tag{17}$$

$$\lambda_{K,5} = \lambda_{t,7} = -(q^2 - 7q + 13) \quad c_{K,5} = c_{t,7} = q - 1. \tag{18}$$

The terms $\lambda_{t,j}$, $j = 4, 5, 8$ do not enter in equation (13). We contrast this with earlier findings. For a given strip, N_λ was found to be larger for (FBC_y, PBC_x) than (FBC_y, FBC_x) [13, 18, 21]. For the $L_y = 3$ square lattice strip case, $N_\lambda = 2$ for (FBC_y, FBC_x) but $N_\lambda = 1$ for (PBC_y, FBC_x) , because of the special feature that the cross sections were complete graphs, K_p with $p = 3$ and hence the intersection theorem led to a factorized, monomial form for P . It was found [13, 14] that for a given type of lattice strip, N_λ is the same for the $(FBC_y, PBC_x) =$ cyclic and $(FBC_y, TPBC_x) =$ Möbius topologies, although the c_j were, in general, different. The present results show that reversal of orientation in the identification of opposite ends of a strip can lead to a change in N_λ †.

Let $C = \sum_{j=1}^{N_\lambda} c_j$. We find $C = P(K_3, q) = q(q-1)(q-2)$ for the $L_y = 3$ torus graphs and $C = 0$ for the $L_y = 3$ Klein bottle graphs. The zero results from the special constraints introduced by the boundary conditions and is analogous to the fact that $C = 0$ for the $L_y = 2$ Möbius square strip [9] and its homeomorphic expansions [21]. However, not all Möbius strip

† A different sort of change in P , accompanied by a change in B , can be obtained if one considers a homogeneous recursive family and the same family with a finite inhomogeneous subgraph inserted, e.g., the ‘rope ladder’ graphs of [16] or two such subgraphs forming ends, namely, the $J(\Pi H)I$ strip graphs in [18].




Figure 1. Chromatic zeros for the $L_y = 3$, $L_x = m = 10$ torus graph.

graphs have $C = 0$; for example, for the $L_y = 3$ Möbius strips of the square and kagomé lattices, $C = q(q - 1)$ and $C = q$, respectively [14].

Chromatic zeros for the $L_y = 3$, $m = 10$ torus graph are shown in figure 1; with this value of m , the chromatic zeros for the Klein bottle graph are quite similar. The locus \mathcal{B} and the W functions are the same for the torus and Klein bottle graph families. We find $q_c = 3$, which, interestingly, is the same value as for the infinite 2D square lattice. The locus \mathcal{B} has support for $\text{Re}(q) \geq 0$ and separates the q plane into three regions. The outermost one, region R_1 , extends to infinite $|q|$ and includes the intervals $q \geq 3$ and $q \leq 0$ on the real q axis. Region R_2 includes the real interval $2 \leq q \leq 3$ and extends upward and downward to the complex conjugate triple points on \mathcal{B} at q_t and q_t^* , where $q_t \simeq 2.5 + 1.4i$. Region R_3 is the innermost one and includes the real interval $0 \leq q \leq 2$. The boundary between R_2 and R_3 curves to the right as one increases $|\text{Im}(q)|$, extending from $q = 2$ upward to q_t and downward to q_t^* . As is evident in figure 1, the density of chromatic zeros on the R_1 – R_3 boundary near $q = 0$ and on the R_3 – R_2 boundary is somewhat smaller than on the right-hand part of the R_2 – R_1 boundary.

In region R_1 , $\lambda_{t,2} = \lambda_{K,2}$ is the dominant λ_j , so

$$W = (q^3 - 6q^2 + 14q - 13)^{1/3} \quad q \in R_1. \quad (19)$$

The fact that this is the same as W for the (PBC_y, FBC_x) case, equation (3), is a general result. The importance of the PBC_y is evident from the fact that for the same width of three squares, the strip with (FBC_y, FBC_x) yields a different W [18].

In region R_2 , $\lambda_{t,6} = \lambda_{K,4}$ is dominant, so

$$|W| = |q - 5|^{1/3} \quad q \in R_2 \quad (20)$$

(in regions other than R_1 , only $|W|$ can be determined unambiguously [7]). In region R_3 , $\lambda_{t,7} = \lambda_{K,5}$ is dominant, so

$$|W| = |q^2 - 7q + 13|^{1/3} \quad q \in R_3. \quad (21)$$

The outer boundary separating R_1 from the inner two regions is oblate, extending out to a maximum of about $|\text{Im}(q)| \simeq 2.5$ for $\text{Re}(q) \simeq 1.5$ (and passing through $q = 0$ and 3). For all the three points, $q = 0, 2, 3$, where \mathcal{B} crosses the real q axis, it does so vertically. The present

results are in accord with the inference [13, 18] that for a recursive graph with regular lattice structure, a necessary and sufficient condition for \mathcal{B} to separate the q plane into two or more regions is that it contains a global circuit, i.e. a path along a lattice direction whose length goes to infinity as $n \rightarrow \infty$; here this is equivalent to PBC_x . The fact that \mathcal{B} is the same for these torus and Klein families means that none of $\lambda_{t,j}$, $j = 4, 5, 8$ is a dominant term.

In this letter we have reported the first calculations of the zero-temperature Potts antiferromagnet partition function (chromatic polynomial) and exponential of the entropy, W , for strips of the square lattice with width $L_y = 3$ and arbitrary length, having periodic transverse and periodic and twisted periodic longitudinal boundary conditions (torus and Klein bottle families of graphs). Our work elucidates the role that these boundary conditions and the associated topologies play; the torus and Klein bottle graphs have interestingly different chromatic polynomials, with different N_λ ($N_\lambda = 8$ and 5 , respectively), but the W functions and hence the boundaries \mathcal{B} are the same. The forms that we obtain for P are simpler than those found [13, 14] for the analogous $L_y = 3$ cyclic and Möbius square lattice strips, which had $N_\lambda = 10$ and some algebraic nonpolynomial λ_j , in contrast to the polynomial λ_j in equations (4)–(18). The resultant boundaries \mathcal{B} are also simpler, involving only three rather than seven, regions. Remarkably, the value $q_c = 3$ that we find for the torus and Klein bottle strips is the same as for the infinite square lattice.

The research of RS was supported in part by the US NSF grant PHY-97-22101.

References

- [1] Birkhoff G D 1912 *Ann. Math.* **14** 42
- [2] Read R C 1968 *J. Comb. Theory* **4** 52
- [3] Read R C and Tutte W T 1988 Chromatic polynomials *Selected Topics in Graph Theory* vol 3 (New York: Academic) p 15
- [4] Biggs N L 1993 *Algebraic Graph Theory* (Cambridge: Cambridge University Press)
- [5] Potts R B 1952 *Proc. Camb. Phil. Soc.* **48** 106
- [6] Wu F Y 1982 *Rev. Mod. Phys.* **54** 235
- [7] Shrock R and Tsai S-H 1997 *Phys. Rev. E* **55** 5165
- [8] Beraha S, Kahane J and Weiss N 1980 *J. Comb. Theory B* **28** 52
- [9] Biggs N L, Damerell R M and Sands D A 1972 *J. Comb. Theory B* **12** 123
- [10] Biggs, N L and Meredith G H 1976 *J. Comb. Theory B* **20** 5
- [11] Biggs N L 1977 *Bull. London Math. Soc.* **9** 54
- [12] Biggs N L 1999 *LSE Report* LSE-CDAM-99-03 to appear
- [13] Shrock R and Tsai S-H 1999 *Phys. Rev. E* at press
Shrock R and Tsai S-H 1999 *Physica A* at press
- [14] Shrock R 1999 *Phys. Lett. A* at press
- [15] Read R C 1988 *Proc. 5th Caribbean Conf. on Combinatorics and Computing (Barbados)*
- [16] Read R C and Royle G F 1991 *Graph Theory, Combinatorics, and Applications* vol 2 (New York: Wiley) p 1009
- [17] Baxter R J 1987 *J. Phys. A: Math. Gen.* **20** 5241
- [18] Roček M, Shrock R and Tsai S-H 1998 *Physica A* **252** 505
Roček M, Shrock R and Tsai S-H 1998 *Physica A* **259** 367
- [19] Shrock R and Tsai S-H 1998 *Physica A* **259** 315
- [20] Shrock R and Tsai S-H 1998 *Phys. Rev. E* **58** 4332
(Shrock R and Tsai S-H 1998 *Preprint* cond-mat/9808057)
- [21] Shrock R and Tsai S-H 1999 *J. Phys. A: Math. Gen. Lett.* **32** L195
- [22] Shrock R and Tsai S-H 1999 *J. Phys. A: Math. Gen.* **32** 5053
- [23] Saaty T L and Kainen P C 1977 *The Four-Colour Problem* (New York: McGraw-Hill) p 45
- [24] White A T and Beineke L W 1978 *Selected Topics in Graph Theory* ed L W Beineke and R J Wilson (New York: Academic) p 15
- [25] Meredith G H 1972 *J. Comb. Theory B* **13** 14
- [26] Biggs N L 1999 *LSE Report* LSE-CDAM-99-05