

Commuting difference operators with elliptic coefficients from Baxter's vacuum vectors

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2000 J. Phys. A: Math. Gen. 33 3825

(<http://iopscience.iop.org/0305-4470/33/20/308>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.118

The article was downloaded on 02/06/2010 at 08:09

Please note that [terms and conditions apply](#).

Commuting difference operators with elliptic coefficients from Baxter's vacuum vectors

A Zabrodin

Joint Institute of Chemical Physics, Kosygina str. 4, 117334, Moscow, Russia and ITEP, 117259, Moscow, Russia

Received 13 January 2000

Abstract. For quantum integrable models with elliptic R -matrix, we construct the Baxter Q -operator in infinite-dimensional representations of the algebra of observables.

1. Introduction

The method of Q -operator, first proposed by Baxter in his solution of the 8-vertex model and the spin- $\frac{1}{2}$ XYZ chain [1], is perhaps the most powerful approach to quantum integrable models known today. As compared with the algebraic Bethe ansatz [2], one or another version of this approach, known also as the method of vacuum vectors, is indispensable for models which do not possess bare vacuum states. Among later striking developments we mention the works [3], where the Baxter method was applied to the chiral Potts model.

At that time, however, the Q -operator itself seemed to be a kind of virtual object, for its explicit form was not known. In Baxter's original work, the Q -operator played an auxiliary role since the local Hamiltonian to be diagonalized was made of the quantum transfer matrix $T(\lambda)$. The main properties of the Q -operator $Q(\lambda)$ are the commutativity $[Q(\lambda), Q(\mu)] = 0$, $[Q(\lambda), T(\mu)] = 0$ for all λ, μ , and the following T - Q relation:

$$T(\lambda)Q(\lambda) = a(\lambda)Q(\lambda + \eta) + b(\lambda)Q(\lambda - \eta). \quad (1.1)$$

Here $a(\lambda)$, $b(\lambda)$ are c -number functions depending on a particular model. Combined with (model-dependent) analytical properties of the $Q(\lambda)$, this relation allows one to obtain a system of Bethe equations and eventually to solve the diagonalization problem. In the absence of global bare vacuum states, Baxter's method to derive (1.1) was to consider continuous-parametric family of special states that factorize into products of *local* vacuum vectors. Such states have especially simple properties under action of $T(\lambda)$. Treating them as columns or rows of some matrix $Q(\lambda)$, one arrives, after a proper normalization, at (1.1). Some years ago this approach was extended [4] to the case of twisted boundary conditions. It remained obscure, however, whether the Q -matrix could play any other role than just a technical tool for deriving Bethe equations.

The situation changed only recently, when it was realized that in integrable lattice models with an infinite-dimensional Hilbert space at each site it is Q rather than T that is the generating function of local Hamiltonians. With this motivation in mind, a considerable progress in constructing the Q -operator for a number of integrable models and in understanding its nature was achieved.

The work [5] was perhaps the first in which the Baxter approach was adopted for integrable models with an infinite-dimensional quantum space. Specifically, in that paper the Q -operator for the quantum Toda chain was constructed as a certain integral operator with explicitly known kernel. Recently, this remarkable development was extended [6] to the integrable homogeneous XXX spin chain in a unitary discrete series representation of the non-compact group $SU(1, 1)$ (see also [7]).

For integrable models of XXZ type, the nature of the Q -operator was clarified in [8, 9], where it was given a representation theory meaning. Specifically, the Q -operator can be obtained from the universal R -matrix for the quantum affine algebra $U_q(\widehat{sl(2)})$, by taking the trace of a chain product of the universal R -matrices in a special q -oscillator representation in one of the spaces. The commutativity of the Q -operators, which emerges as a surprise in Baxter's approach, is now guaranteed by characteristic properties of the universal R -matrix. The T - Q relation, from this point of view, is a simple consequence of fusion rules for representations of the quantum affine algebra. In fact, two different commuting Q -operators, $Q^+(\lambda)$ and $Q^-(\lambda)$, can be introduced in this way. They are two linearly independent operator solutions to the T - Q relation (1.1) treated as a second-order difference equation in λ . As such, they obey a quantum Wronskian condition, which turns out to be even more instructive than the T - Q relation itself.

As for matters more customary for the theory of solitons, it was argued in [10, 11] that the Q -operator is a quantum analogue of the Bäcklund transformation. In other words, it defines an integrable quantum model with discrete time by setting $Q(\lambda)$ to be the time shift operator. For any observable O , one sets $O(t + \delta) = Q(\lambda)O(t)(Q(\lambda))^{-1}$, where the discrete time shift δ is a function of λ . Integrability implies that these discrete evolution operators commute for all δ .

However, all this was about models associated with rational or trigonometric solutions of the Yang–Baxter equation (R -matrices). General R -matrices with additive spectral parameter are parametrized via elliptic functions. The simplest elliptic R -matrix is

$$R(\lambda) = \sum_{a=0}^3 \frac{\theta_{a+1}(2\lambda + \eta|\tau)}{\theta_{a+1}(\eta|\tau)} \sigma_a \otimes \sigma_a \quad (1.2)$$

where $\theta_a(z|\tau)$ are Jacobi θ functions, σ_a are Pauli matrices, and σ_0 is the unit matrix. This R -matrix is associated with the 8-vertex model, being the matrix of local Boltzmann weights at a vertex. The transfer matrix of this model is the generating function of conserved quantities for the integrable XYZ spin- $\frac{1}{2}$ chain. By models of the XYZ type we mean the more general integrable models with elliptic R -matrix, for instance integrable anisotropic spin chains with higher spin. Integrable spin chains of XYZ type can be solved by the generalized algebraic Bethe ansatz [2, 12].

However, the quantum affine algebra approach is not applicable in the XYZ case since nothing similar to the universal elliptic R -matrix is known. Nevertheless, the elliptic dependence on the spectral parameter does not destroy the functional relations between transfer matrices and Q -operators, including the T - Q relation and the Wronskian condition [13, 14]. This suggests that everything that it is possible to do in the XXX and XXZ cases, might be possible *mutatis mutandis* for XYZ too.

In this paper we present a direct construction of the Q -operator for homogeneous N -site spin chains of the XYZ type. In lattice integrable models with elliptic R -matrix, the algebra of local observables is the Sklyanin algebra [15, 16]. A concrete model is defined by fixing a particular representation of this algebra. Similar to the $sl(2)$ case (models of XXX type), the representations are labelled by a continuous parameter which is called *spin*, by analogy with $sl(2)$. We allow the spin to take any complex value, so we work in an infinite-dimensional

representation of the algebra of observables. In other words, we deal with representations of what would be called a non-compact real form of the Sklyanin algebra. Such representations can be realized by difference operators, so our Q -operator is a difference operator in many variables.

Our strategy closely follows the original Baxter argument. First, we construct a family of local (infinite-dimensional) vacuum vectors that obey the celebrated ‘pair propagation through a vertex’ property. Next, their product over all sites of the chain is, up to normalization, the kernel of the Q -operator. Finally, one should prove the commutativity of the Q -operators. However, the practical realization of these ideas in the infinite-dimensional setting is by no means obvious. Technically, it is very different from what is customary in the 8-vertex model. The construction extensively uses such truly special functions as elliptic generalizations of the gamma function and basic hypergeometric functions, and highly non-trivial identities for them. For example, commutativity of the Q -operators relies on an elliptic version of the Bailey transformation formula for the very well poised basic hypergeometric series ${}_{10}W_9$. The appearance of such a sophisticated mathematics in integrable models is not so surprising. Recently, the elliptic series ${}_{10}W_9$ was shown [17] to offer a natural representation for ‘elliptic $6j$ symbols’, which are nothing else than solutions of the Yang–Baxter equation in a special basis.

It is now a challenge to reveal a representation theory meaning of the elliptic Q -operator. Hopefully, this would make its properties and their proofs obvious.

The paper is organized as follows. In section 2 we present the main result in its shortest possible form. Section 3 contains the necessary material on representations of the Sklyanin algebra. The construction of the Q -operator begins in section 4, where we find a family of local vacuum states for the quantum L-operator. In section 5 we investigate direct products of such states along the chain and define unnormalized ‘pre- Q -operators’, Q_L and Q_R , which do not yet commute. Section 6 is rather technical. It is devoted to the proof of the basic commutation relation for the pre- Q -operators, $Q_L(\lambda)Q_R(\mu) = Q_L(\mu)Q_R(\lambda)$, which entails commutativity of the Q -operators introduced in section 7. In section 8 we discuss more specific issues: (a) twisted boundary conditions, (b) the exceptional case of a ‘spin chain’ with only one site, $N = 1$, and (c) the trigonometric (XXZ type) degeneration. A few concluding remarks are given in section 9.

Appendix A contains necessary information on the special functions involved in the construction. In appendix B we give a direct proof of the intertwining relation between representations of spins ℓ and $-\ell - 1$.

2. The main result

To state the result, we recall the definition of the quantum transfer matrix $T(\lambda)$ for the XYZ spin chain with spin ℓ . The elliptic quantum L-operator is the matrix

$$L(\lambda) = \frac{1}{2} \begin{pmatrix} \theta_1(2\lambda)s_0 + \theta_4(2\lambda)s_3 & \theta_2(2\lambda)s_1 + \theta_3(2\lambda)s_2 \\ \theta_2(2\lambda)s_1 - \theta_3(2\lambda)s_2 & \theta_1(2\lambda)s_0 - \theta_4(2\lambda)s_3 \end{pmatrix} \quad (2.1)$$

with non-commutative matrix elements. Specifically, s_a are difference operators in a complex variable z :

$$s_a = \frac{\theta_{a+1}(2z - 2\ell\eta)}{\theta_1(2z)} e^{\eta\partial_z} - \frac{\theta_{a+1}(-2z - 2\ell\eta)}{\theta_1(2z)} e^{-\eta\partial_z} \quad (2.2)$$

introduced by Sklyanin [16]. Here $\theta_a(z) \equiv \theta_a(z|\tau)$ are Jacobi θ functions with the elliptic module τ , $\text{Im } \tau > 0$, ℓ is a complex number (the spin), and $\eta \in \mathbb{C}$ is a parameter which is assumed to belong to the fundamental parallelogram with vertices $0, 1, \tau, 1 + \tau$, and to be

incommensurate with 1, τ . Definitions and transformation properties of the θ functions are listed in appendix A.

Let us consider the matrix product of L-operators (2.1): $\overset{1}{L}(\lambda) \overset{2}{L}(\lambda) \dots \overset{N}{L}(\lambda)$, $N > 1$, where each operator matrix $\overset{i}{L}(\lambda)$ has the form (2.1) with s_a acting on its own variable z_i . Clearly, matrix elements of $\overset{i}{L}(\lambda)$ and $\overset{j}{L}(\lambda)$ commute for $i \neq j$. The trace of this product (the sum of two diagonal elements),

$$T(\lambda) = \text{tr}[\overset{1}{L}(\lambda) \overset{2}{L}(\lambda) \dots \overset{N}{L}(\lambda)] \tag{2.3}$$

is called the quantum transfer matrix of the spin chain.

Our main result is the explicit realization of the Q -operator for this spin chain:

$$Q(\lambda) = \Gamma^{-N}(2\lambda - 2\ell\eta) \underset{i=1}{\prod}^N q_{i,i+1}(\lambda) \underset{j=1}{\prod}^N e^{(\lambda+\ell\eta)\theta_{z_j}} \tag{2.4}$$

where

$$q_{i,i+1}(\lambda) = \frac{\Gamma(2z_i)\Gamma(z_i - z_{i+1} + 2\lambda)\Gamma(z_i + z_{i+1} + 2\lambda)}{\Gamma(2z_i + 2\lambda + 2\ell\eta)\Gamma(z_i - z_{i+1} - 2\ell\eta)\Gamma(z_i + z_{i+1} - 2\ell\eta)} \times {}_6W_5 \left(-\frac{z_i+\lambda}{\eta} - \ell; -\frac{\lambda}{\eta} - \ell, -\frac{z_{i+1}+z_i}{2\eta} - \ell, \frac{z_{i+1}-z_i}{2\eta} - \ell; e^{-2\eta\theta_{z_i}} \mid \eta, \tau \right) \tag{2.5}$$

(it is implied that $z_{N+1} = z_1$). Here $\Gamma(z) = \Gamma(z|\tau, 2\eta)$ is the elliptic gamma function [18, 19]:

$$\Gamma(z|\tau, 2\eta) = \prod_{k,m=0}^{\infty} \frac{1 - e^{2\pi i((k+1)\tau+2(m+1)\eta-z)}}{1 - e^{2\pi i(k\tau+2m\eta+z)}} = \exp \left(-i \sum_{k=1}^{\infty} \frac{\sin(\pi k(2z - 2\eta - \tau))}{2k \sin(2\pi k\eta) \sin(\pi k\tau)} \right)$$

and ${}_6W_5(\alpha_1; \alpha_4, \dots, \alpha_6; z|\eta, \tau)$ is the elliptic analogue of the very well poised hypergeometric series (with the operator argument):

$${}_6W_5(\alpha_1; \alpha_4, \dots, \alpha_6; z|\eta, \tau) = \sum_{k=0}^{\infty} \frac{[\alpha_1 + 2k][\alpha_1]_k}{[\alpha_1][1]_k} \left(\prod_{m=1}^3 \frac{[\alpha_{m+3}]_k}{[\alpha_1 - \alpha_{m+3} + 1]_k} \right) z^k$$

where $[\alpha]_k = \prod_{j=0}^{k-1} \theta_1(2\alpha\eta + 2j\eta)$. For more information on the special functions $\Gamma, {}_6W_5$ see appendix A. The normal ordering means that all operators $e^{-2\eta\theta_{z_j}}$ are moved to the right.

We prove the commutativity $[Q(\lambda), T(\mu)] = [Q(\lambda), Q(\mu)] = 0$, and the T - Q relation

$$T(\lambda)Q(\lambda) = \theta_1^N(2\lambda - 2\ell\eta)Q(\lambda + \eta) + \theta_1^N(2\lambda + 2\ell\eta)Q(\lambda - \eta). \tag{2.6}$$

Another solution to this operator difference equation has a similar form (see (7.8)).

The limiting case $\tau \rightarrow 0$, when elliptic functions degenerate to trigonometric ones, corresponds to the XXZ spin chain. The structure of the formulae (2.4), (2.5) remains the same while the special functions get simplified. The elliptic gamma function becomes the q -gamma function (or the q -exponential function) while the elliptic hypergeometric series ${}_6W_5$ turns into the little q -Jacobi function represented by the basic hypergeometric series ${}_2\phi_1$.

3. Representations of the Sklyanin algebra

The aim of this section is to give the necessary preliminaries on representations of the Sklyanin algebra. We begin with a few formulae related to the quantum L-operator with elliptic dependence on the spectral parameter.

Consider the quantum L-operator (2.1), where s_a are realized as in (2.2). The four operators s_a obey the commutation relations of the Sklyanin algebra[†]:

$$\begin{aligned} (-1)^{\alpha+1} I_{\alpha 0} s_\alpha s_0 &= I_{\beta\gamma} s_\beta s_\gamma - I_{\gamma\beta} s_\gamma s_\beta \\ (-1)^{\alpha+1} I_{\alpha 0} s_0 s_\alpha &= I_{\gamma\beta} s_\beta s_\gamma - I_{\beta\gamma} s_\gamma s_\beta \end{aligned} \tag{3.1}$$

with the structure constants $I_{ab} = \theta_{a+1}(0)\theta_{b+1}(2\eta)$. Here $a, b = 0, \dots, 3$ and $\{\alpha, \beta, \gamma\}$ stands for any cyclic permutation of $\{1, 2, 3\}$. The relations of the Sklyanin algebra are equivalent to the condition that the L-operator satisfies the ‘RLL = LLR’ relation with the elliptic R-matrix (1.2).

Plugging the difference operators (2.2) into (2.1), one can represent the L-operator in a factorized form, which is especially convenient for computations:

$$L(\lambda) = \theta_1(2\lambda_+) \mathfrak{V}^{-1}(\lambda_+, z) \begin{pmatrix} e^{\eta\partial_z} & 0 \\ 0 & e^{-\eta\partial_z} \end{pmatrix} V(\lambda_-, z) \mathfrak{V}. \tag{3.2}$$

Hereafter we use the convenient notation $\lambda_\pm \equiv \lambda \pm \ell\eta$, $\bar{\theta}_a(z) \equiv \theta_a(z|\frac{\tau}{2})$. The dots mean the normal ordering when the shift operators are moved to the right and

$$V(\lambda, z) = \begin{pmatrix} \bar{\theta}_4(z + \lambda) & \bar{\theta}_3(z + \lambda) \\ \bar{\theta}_4(z - \lambda) & \bar{\theta}_3(z - \lambda) \end{pmatrix}.$$

Note that

$$L(-\lambda + \frac{1}{2}(1 + \tau)) = -e^{-\pi i\tau + 4\pi i\lambda} L^\top(\lambda) \tag{3.3}$$

where L^\top means transposition in the two-dimensional space.

The parameter ℓ in (2.2) is called the spin of the representation. If necessary, we write $s_a = s_a^{(\ell)}$ to indicate the dependence on ℓ . When $\ell \in \frac{1}{2}\mathbb{Z}_+$, these operators have a finite-dimensional invariant subspace, namely the space $\Theta_{4\ell}^+$ of even θ functions of order 4ℓ (see appendix A). This is the representation space of the $(2\ell + 1)$ -dimensional irreducible representation (of series A) of the Sklyanin algebra. For example, at $\ell = \frac{1}{2}$ the functions $\bar{\theta}_4(z)$, $\bar{\theta}_3(z)$ form a basis in Θ_2^+ , and the generators s_a , with respect to this basis, are represented by 2×2 matrices $(-i)^{\delta_{a,2}}(\theta_{a+1}(\eta))^{-1}\sigma_a$. In this case, $L(\lambda) = R(\lambda - \frac{1}{2}\eta)$, where R is the 8-vertex model R-matrix (1.2).

As is proved in [20], the space $\Theta_{4\ell}^+$ for $\ell \in \frac{1}{2}\mathbb{Z}_+$ is annihilated by the operator

$$w_\ell = \sum_{k=0}^{2\ell+1} (-1)^k \begin{bmatrix} 2\ell + 1 \\ k \end{bmatrix} \frac{\theta_1(2z + 2(2\ell - 2k + 1)\eta)}{\prod_{j=0}^{2\ell+1} \theta_1(2z + 2(j - k)\eta)} e^{(2\ell - 2k + 1)\eta\partial_z}. \tag{3.4}$$

Here and below we use the ‘elliptic factorial’ and ‘elliptic binomial’ notation:

$$[j] \equiv \theta_1(2j\eta) \quad [n]! = \prod_{j=1}^n [j] \quad \begin{bmatrix} n \\ m \end{bmatrix} \equiv \frac{[n]!}{[m]![n - m]!}. \tag{3.5}$$

Another important property of the operator w_ℓ established in [20] is that w_ℓ intertwines representations of spin ℓ and of spin $-(\ell + 1)$:

$$s_a^{(-\ell-1)} w_\ell = w_\ell s_a^{(\ell)} \quad a = 0, \dots, 3. \tag{3.6}$$

The same intertwining relation can be written for the quantum L-operator (2.1). The operator w_ℓ is an elliptic analogue of $(d/dz)^{2\ell+1}$ in the following sense. In the case of the algebra sl_2 , the intertwining operator between representations of spins ℓ and $-\ell - 1$ (realized by differential operators in z) is $(d/dz)^{2\ell+1}$. It annihilates the linear space of polynomials of degree $\leq 2\ell$.

[†] The standard generators of the Sklyanin algebra [15], S_a , are related to ours as follows: $S_a = (i)^{\delta_{a,2}}\theta_{a+1}(\eta)s_a$.

The operator w_ℓ can be extended to arbitrary complex values of ℓ in which case it is represented by a half-infinite series in the shift operator $e^{2\eta\partial_z}$. The series is an elliptic analogue of the very well poised basic hypergeometric series with an operator argument. For its explicit form see (7.13). The intertwining relations (3.6) hold true in this more general case, too. The proof is outlined in appendix B.

In what follows we need infinite-dimensional representations of the Sklyanin algebra in some functional space. Operators (2.2) provide such a representation. However, they are not sufficient for our purpose since it is necessary to specify the functional space where these operators are going to act. It is not clear how to do that for continuous functions. On the other hand, the difference character of the operators (2.2) allows one to restrict them to *discontinuous* functions of the form described below.

Let $\delta(z)$ be the function equal to zero everywhere but at $z = 0$, where it equals 1: $\delta(z) = 0$, $z \neq 0$, $\delta(0) = 1$. Clearly, $z\delta(z) = 0$. The Sklyanin algebra realized as in (2.2) acts, for any $\nu \in \mathbb{C}$, in the space of functions of the form

$$f(z) = \sum_{k \in \mathbb{Z}} f_k \delta(z - \nu + 2k\eta) \quad f_k \in \mathbb{C}. \quad (3.7)$$

This space is isomorphic to the linear space of sequences $\{f_k\}_{k \in \mathbb{Z}}$. We call functions of the form (3.7) *combs*.

A comb is said to be finite from the right (respectively, from the left) if there exists $M \in \mathbb{Z}$ such that $f_k = 0$ as $k > M$ (respectively, $k < M$). Let \mathcal{C}^+ (respectively, \mathcal{C}^-) be the space of combs finite from the left (respectively, from the right).

We define the pairing

$$(F(z), \delta(z - a)) = F(a) \quad (3.8)$$

for any function $F(z)$, not necessarily of the form (3.7). In particular,

$$(\delta(z - a), \delta(z - b)) = \delta(a - b). \quad (3.9)$$

Formally, this pairing can be written as an integral:

$$(F(z), \delta(z - a)) = \int F(z) \delta(z - a) dz$$

(perhaps a q -integral symbol would be more appropriate). We stress that the integral here means nothing more than another notation for the pairing, especially convenient in the case of many variables. By linearity, the pairing can be extended to the whole space of combs. We note that the pairing between the spaces \mathcal{C}^+ and \mathcal{C}^- is well defined since the sum is always finite.

Combs are to be viewed as kernels of difference operators. By a difference operator in one variable we mean any expression of the form

$$D = \sum_{k \in \mathbb{Z}} c_k(z) e^{(\mu + 2k\eta)\partial_z} \quad \mu \in \mathbb{C}. \quad (3.10)$$

The comb

$$D(z, \zeta) = \sum_{k \in \mathbb{Z}} c_k(z) \delta(z - \zeta + \mu + 2k\eta) \quad (3.11)$$

regarded as a function of any one of the variables z, ζ , is the kernel of this difference operator in the following sense. Using the pairing introduced above, we can write

$$(Df)(z) = \int D(z, \zeta) f(\zeta) d\zeta = \sum_{k \in \mathbb{Z}} c_k(z) f(z + \mu + 2k\eta). \quad (3.12)$$

The kernel $D(z, \zeta)$ can be viewed as an infinite matrix with continuously numbered rows (z) and columns (ζ). Then the convolution with respect to the second argument of the kernel, as in (3.12), defines action of the operator from the left. The convolution with respect to the first argument defines the action from the right,

$$(fD)(z) = \int f(\zeta)D(\zeta, z) d\zeta \tag{3.13}$$

equivalent to the action of the transposed difference operator from the left:

$$D^t = \sum_{k \in \mathbb{Z}} e^{-(\mu+2k\eta)\partial_z} c_k(z) = \sum_{k \in \mathbb{Z}} c_k(z - \mu - 2k\eta) e^{-(\mu+2k\eta)\partial_z}. \tag{3.14}$$

The transposition t is the anti-automorphism of the algebra of difference operators such that $(c(z)e^{\alpha\partial_z})^t = e^{-\alpha\partial_z}c(z)$. In terms of the above pairing we can write $(f, Dg) = (D^t f, g)$.

The kernels of Sklyanin’s operators (2.2) are

$$s_a(z, z') = \frac{\theta_{a+1}(2z - 2\ell\eta)}{\theta_1(2z)} \delta(z - z' + \eta) - \frac{\theta_{a+1}(-2z - 2\ell\eta)}{\theta_1(2z)} \delta(z - z' - \eta). \tag{3.15}$$

Let us also note that $s_a^{(\ell)t} = -(-1)^{\delta_{a,0}}\theta_1(2z)s_a^{(-\ell-1)}(\theta_1(2z))^{-1}$.

4. Local vacuum vectors

We are going to adopt the Baxter method of vacuum vectors to the infinite-dimensional representations of the Sklyanin algebra described in the previous section.

Let us recall the general definition of vacuum vectors. Consider an arbitrary L-operator L with two-dimensional auxiliary space \mathbb{C}^2 , i.e. an arbitrary 2×2 operator-valued matrix whose matrix elements are operators $L_{11}, L_{12}, L_{21}, L_{22}$:

$$L = \begin{pmatrix} L_{11} & L_{12} \\ L_{21} & L_{22} \end{pmatrix}. \tag{4.1}$$

They act in a linear space \mathcal{H} which is called the *quantum space* of the L-operator.

For a moment, let ψ, ϕ denote vectors from \mathbb{C}^2 . Let $X \in \mathcal{H}$, then acting by the quantum L-operator on the tensor product $X \otimes \phi$, we, generally speaking, obtain a mixed state in the quantum space: $LX \otimes \phi = X_1 \otimes \phi_1 + X_2 \otimes \phi_2$. The special case of a pure state,

$$LX \otimes \phi = X' \otimes \psi \tag{4.2}$$

is of particular importance. Given an L-operator, Baxter’s recipe is to find all pairs X, ϕ such that (4.2) holds. Taking the scalar product with the vector ψ^\perp orthogonal to ψ , we get

$$(\psi^\perp L\phi)X = 0 \tag{4.3}$$

i.e. the operator $K = (\psi^\perp L\phi)$ (acting in the quantum space only) has a zero mode $X \in \mathcal{H}$. Suppose (4.2) (or (4.3)) holds with some vectors ϕ, ψ ; then the vector X is called a *vacuum vector* of the L-operator.

A few remarks are in order. The relation (4.2) (in the particular case $\mathcal{H} \cong \mathbb{C}^2$) was the starting point for Baxter in his solution of the 8-vertex and XYZ models [1]. This is what he called the ‘pair propagation through a vertex’ property. In the context of the quantum inverse scattering method [2] the equivalent condition (4.3) is more customary. It defines local vacua of the (gauge-transformed) L-operator. A generalization of that solution to the higher spin XYZ model was given by Takebe in [12]. An algebro-geometric approach to the equation (4.2) for finite-dimensional matrices L_{ik} was developed by Krichever [21].

For L-operators with elliptic spectral parameter it is convenient to pass to the elliptic parametrization of the components of the vectors ϕ, ψ . We introduce the two-component vector

$$|\zeta\rangle = \begin{pmatrix} \bar{\theta}_4(\zeta) \\ \bar{\theta}_3(\zeta) \end{pmatrix}$$

and the corresponding covector $\langle\zeta| = (\bar{\theta}_4(\zeta), \bar{\theta}_3(\zeta))$ (recall that $\bar{\theta}_a(\zeta) = \theta_a(\zeta|\frac{\tau}{2})$). The vector orthogonal to $\langle\zeta|$ is $|\zeta\rangle^\perp = \begin{pmatrix} \bar{\theta}_3(\zeta) \\ -\bar{\theta}_4(\zeta) \end{pmatrix}$, $\langle\zeta|\zeta\rangle^\perp = 0$. More generally, we have

$$\langle\xi|\zeta\rangle^\perp = 2\theta_1(\xi + \zeta)\theta_1(\xi - \zeta). \tag{4.4}$$

Note also that


$$|\zeta + \frac{1}{2}(1 + \tau)\rangle = e^{-\frac{\pi i \tau}{2} - 2\pi i \zeta} |\zeta\rangle^\perp. \tag{4.5}$$

Let us specify the general definition of the vacuum vectors for the elliptic L-operator (2.1). Writing $L(\lambda)|\zeta\rangle$ (respectively, $\langle\zeta|L(\lambda)$) we mean that the 2×2 matrix (2.1) acts on the two-component vector from the left (respectively, on the two-component covector from the right), as usual. Similarly, we introduce right and left vacuum vectors X_R, X_L according to the relations

$$\langle\zeta|L(\lambda)X_R = \langle\xi|X'_R \quad X_L\langle\zeta|L(\lambda) = X'_L\langle\xi|. \tag{4.6}$$

In the latter formula the difference operators s_a act on X_L from the right.

The following pictorial representation is helpful:



Here we use the standard conventions of the quantum inverse scattering method.

Introducing the operator

$$\mathbf{K} = \mathbf{K}(\zeta, \xi) = \langle\zeta|L(\lambda)|\xi\rangle^\perp \tag{4.7}$$

we can rewrite (4.6) as $\mathbf{K}X_R = X_L\mathbf{K} = 0$. The explicit form of the operator \mathbf{K} is easily found from (3.2):

$$\mathbf{K} = \mathbf{K}(\zeta, \xi) = \rho(z)e^{\eta\partial_z} + \rho(-z)e^{-\eta\partial_z} \tag{4.8}$$

where

$$\rho(z) = \frac{1}{\theta_1(2z)} \prod_{\epsilon=\pm} \theta_1(z + \epsilon\zeta - \lambda_+) \theta_1(z + \epsilon\xi + \lambda_-).$$

So, the equations for the right and left vacuum vectors read

$$\rho(z)X_R(z + \eta) = -\rho(-z)X_R(z - \eta) \tag{4.9}$$

$$\rho(-z - \eta)X_L(z + \eta) = -\rho(z - \eta)X_L(z - \eta). \tag{4.10}$$

These equations themselves are not yet enough. An important additional input is the space in which to solve them. Our choice, justified by the final results, is the following. We are going to solve the first equation in the space of combs finite from the left, \mathcal{C}^+ , while the second equation is to be solved in the space of combs finite from the right, \mathcal{C}^- . (For precise definitions of these spaces see section 3.)

Let us begin by solving equations (4.9), (4.10) in the space of meromorphic functions of z . This is easy to do in terms of the elliptic gamma function $\Gamma(z|\tau, 2\eta)$ introduced by Ruijsenaars [18] and recently studied in detail by Felder and Varchenko [19]. This function was introduced just to solve equations of such a type. Its main properties are summarized in appendix A. In what follows we write $\Gamma(z|\tau, 2\eta) = \Gamma(z)$ for brevity. Let $\mathcal{X}_R(z), \mathcal{X}_L(z)$ be meromorphic solutions of (4.9), (4.10), respectively. Using (A9), we easily find

$$\mathcal{X}_R(z) = e^{-4\pi i \ell z} \omega(\lambda, \zeta - \xi) \prod_{\epsilon=\pm} \frac{\Gamma(z + \epsilon\zeta + \lambda_+ + \eta)\Gamma(z + \epsilon\xi - \lambda_- + \eta)}{\Gamma(z + \epsilon\zeta - \lambda_+ + \eta)\Gamma(z + \epsilon\xi + \lambda_- + \eta)} \tag{4.11}$$

$$\mathcal{X}_L(z) = e^{4\pi i(\ell+1)z} \Phi_\ell(2\lambda)\theta_1(2z) \prod_{\epsilon=\pm} \frac{\Gamma(z + \epsilon\zeta - \lambda_+)\Gamma(z + \epsilon\xi + \lambda_-)}{\Gamma(z + \epsilon\zeta + \lambda_+ + 2\eta)\Gamma(z + \epsilon\xi - \lambda_- + 2\eta)}. \tag{4.12}$$

The z -independent normalization factors $\omega(\lambda, \zeta - \xi)$ in (4.11) and $\Phi_\ell(2\lambda)$ in (4.12) are introduced here for later convenience. The function $\omega(\lambda, \zeta)$ is given by

$$\omega(\lambda, \zeta) = \frac{\theta_1(2\lambda - 2\ell\eta|2\eta)}{\theta_1(2\lambda + \zeta|2\eta)} \exp\left(-\frac{\pi i \zeta^2}{2\eta} - 4\pi i \ell \lambda\right) \tag{4.13}$$

(the θ functions here have the modular parameter 2η rather than τ !). The function Φ_ℓ obeys the difference equation $\theta_1(z - 2\ell\eta)\Phi_\ell(z + 2\eta) = \theta_1(z + 2(\ell+1)\eta)\Phi_\ell(z)$. For $\ell \in \frac{1}{2}\mathbb{Z}_+$ the solution is very simple:

$$\Phi_\ell(z) = \prod_{k=-\ell}^{\ell} \theta_1(z + 2k\eta) \tag{4.14}$$

where k ranges over $-\ell, -\ell + 1, \dots, \ell$. For arbitrary complex values of ℓ the function Φ_ℓ can be defined through the elliptic gamma function (see (A16)). If necessary, we write $\mathcal{X}_{R,L}(z) = \mathcal{X}_{R,L}(z; \zeta, \xi, \lambda)$. Note that $\mathcal{X}_{R,L}(z; \zeta, \xi, \lambda)$ differs from $\mathcal{X}_{R,L}(z; \xi, \zeta, -\lambda)$ only by a z -independent factor.

An important remark is in order. The solutions (4.11), (4.12) are by no means unique. Clearly, one can multiply them by an arbitrary function $f(z)$ such that $f(z + \eta) = f(z - \eta)$. In this paper, we do not take care of such 2η -periodic factors, fixing them to be constant. In other words, (4.11), (4.12) are ‘minimal’ solutions to (4.9), (4.10) in the sense of [18]. By this we break the symmetry between the three periods $1, 2\eta, \tau$ involved in the problem. In particular, the modular symmetry is broken since 2η -periodic factors may be important for good modular properties of the solutions. Indeed, the modular transformation of the elliptic gamma function has the form [19]:

$$\Gamma(z|\tau, 2\eta) = e^{i\tau P(z)} \frac{\Gamma(\frac{z}{\tau} | -\frac{1}{\tau}, \frac{2\eta}{\tau})}{\Gamma(\frac{z-\tau}{2\eta} | -\frac{1}{2\eta}, \frac{\tau}{2\eta})}$$

where $P(z)$ is a polynomial of third degree (see (A12)). The denominator is a 2η -periodic function which does not enter the minimal solution of the modular transformed equation. We note in passing that this formula is an elliptic version of equation (40) from the paper [22], where a modular double of quantum group was introduced. We hope to discuss this point elsewhere. See also the end of section 8.

Now we can proceed to build a half-infinite comb solution. Any infinite comb of the form $X(z) = \mathcal{X}_{R,L}(z) \sum_{k \in \mathbb{Z}} \delta(z - z_0 + 2k\eta)$ automatically satisfies equation (4.9) (resp. (4.10)). How to truncate the comb holding these equations valid? Let $\mathcal{X}(z)$ be any one of $\mathcal{X}_R(z), \mathcal{X}_L(z)$. The position of zeros and poles of the Γ function suggests two possibilities:

- (1) To set z_0 to be a zero of the function \mathcal{X} : $\mathcal{X}(z_0) = 0$. If this zero comes from a pole of one of the Γ functions downstairs, then $\mathcal{X}(z_0 - 2k\eta) = 0, k \geq 0$, and the comb is truncating

from the left. If z_0 is a zero of one of the Γ functions upstairs, then $\mathcal{X}(z_0 - 2k\eta) = 0$, $k < 0$, and the comb is truncating from the right.

- (2) To set z_0 to be a pole of \mathcal{X} and define $X(z) = \sum_{k \in \mathbb{Z}} \text{res}|_{z=z_0-2k\eta} \mathcal{X}(z) \delta(z - z_0 + 2k\eta)$. If z_0 is a pole of a Γ function upstairs, then the comb is truncating from the right. If the pole comes from a zero of one of the Γ functions downstairs, then the comb is truncating from the left.

In the sequel we use both possibilities: (1) for X_R and (2) for X_L . Other options are to be further analysed.

Assume that the parameters are in general position, i.e. there is no cancellation of zeros and poles in (4.11). (For $\ell \in \frac{1}{2}\mathbb{Z}_+$ the sufficient conditions are $\lambda \neq 2m\eta$, $\xi - \zeta \neq 2m\eta$, $m \in \mathbb{Z}$.) Then $\mathcal{X}_R(\xi - \lambda_- - \eta - 2k\eta) = 0$ for all integer $k \geq 0$. At the same time this function is not identically zero when k is negative. Therefore,

$$X_R^+(z; \zeta, \xi, \lambda) = \mathcal{X}_R(z; \zeta, \xi, \lambda) \sum_{k=1}^{\infty} \delta(z - \xi + \lambda_- + \eta - 2k\eta) \tag{4.15}$$

is a non-trivial comb finite from the left which solves equation (4.9). Clearly, this is not the only possibility to truncate the infinite comb from the left, even sticking to (1). There are three other series of zeros of the function (4.11) coming from poles of the gamma functions in the denominator. Any one of them can be used to generate, in the same way, a half-infinite comb solution that belongs to C^+ . However, inasmuch as the vacuum vectors are going to be kernels of difference operators, we want the δ functions in the vacuum vectors to depend on the differences $z - \zeta$ or $z - \xi$. So, along with (4.15), we have another comb obeying equation (4.9):

$$X_R^-(z; \zeta, \xi, \lambda) = \mathcal{X}_R(z; \xi, \zeta, -\lambda) \sum_{k=1}^{\infty} \delta(z - \zeta - \lambda_+ + \eta - 2k\eta). \tag{4.16}$$

For X_L we use (2): $\mathcal{X}_L(z)$ has poles at $z = \zeta + \lambda_+ - 2k\eta, k \geq 0$. In general position, these poles are all simple. The lattice of poles is finite from the right. Therefore, we can construct a solution to (4.10) in the space of combs finite from the right as follows:

$$X_L^+(\lambda, \zeta, \xi; z) = \sum_{k=0}^{\infty} \text{res}|_{z=\zeta+\lambda_+-2k\eta} \mathcal{X}_L(z; \zeta, \xi, \lambda) \delta(z - \zeta - \lambda_+ + 2k\eta). \tag{4.17}$$

Similarly to the case of the right vacuum vectors, we have a second solution:

$$X_L^-(\lambda, \zeta, \xi; z) = \sum_{k=0}^{\infty} \text{res}|_{z=\xi-\lambda_--2k\eta} \mathcal{X}_L(z; \xi, \zeta, -\lambda) \delta(z - \xi + \lambda_- + 2k\eta). \tag{4.18}$$

To make these formulae closer to (4.15), (4.16), we introduce the auxiliary function

$$\begin{aligned} \mathcal{X}_L^\varepsilon(z; \zeta, \xi, \lambda) &= e^{4\pi i(\ell+1)z} \Phi_\ell(2\lambda) \theta_1(2z) \\ &\times (\Gamma(z - \zeta - \lambda_+ + \varepsilon) \Gamma(z + \zeta - \lambda_+) \Gamma(z - \xi + \lambda_- + \varepsilon) \Gamma(z + \xi + \lambda_-)) \\ &\times \{\Gamma(z - \zeta + \lambda_+ + 2\eta) \\ &\times \Gamma(z + \zeta + \lambda_+ + 2\eta) \Gamma(z - \xi - \lambda_- + 2\eta) \Gamma(z + \xi - \lambda_- + 2\eta)\}^{-1} \end{aligned}$$

then, up to an irrelevant constant factor,

$$X_L^+(z; \zeta, \xi, \lambda) = \lim_{\varepsilon \rightarrow 0} \varepsilon \mathcal{X}_L^\varepsilon(z; \zeta, \xi, \lambda) \sum_{k=0}^{\infty} \delta(z - \zeta - \lambda_+ + 2k\eta) \tag{4.19}$$

$$X_L^-(\lambda, \zeta, \xi; z) = \lim_{\varepsilon \rightarrow 0} \varepsilon \mathcal{X}_L^\varepsilon(z; \xi, \zeta, -\lambda) \sum_{k=0}^{\infty} \delta(z - \xi + \lambda_- + 2k\eta). \tag{4.20}$$

Here one should first substitute the values of z in the δ functions and then take the limit.

Note that the function $\rho(z)$ and the meromorphic solutions $\mathcal{X}_{R,L}$ are invariant with respect to the simultaneous change $\lambda \rightarrow -\lambda, \zeta \leftrightarrow \xi$ while the half-infinite combs $X_{R,L}^\pm$ are transformed one into another:

$$\begin{aligned} X_R^-(z; \zeta, \xi, \lambda) &= X_R^+(z; \xi, \zeta, -\lambda) \\ X_L^-(\lambda, \zeta, \xi; z) &= X_L^+(-\lambda, \xi, \zeta; z). \end{aligned} \tag{4.21}$$

The construction of local vacuum vectors is completed. The vacuum vectors are defined up to an arbitrary normalization factor which may depend on ζ, ξ and λ . In our normalization, the final formulae have the most simple form.

By a straightforward computation, which uses some standard identities for theta functions, one can prove the following relations valid for $X_{R,L}^\pm = X_{R,L}$:

$$\langle \zeta | L(\lambda) X_R(z; \zeta, \xi, \lambda) = \theta_1(2\lambda_-) \langle \xi | X_R(z; \zeta, \xi, \lambda + \eta) \tag{4.22}$$

$$X_L(\zeta, \xi, \lambda; z) \langle \zeta | L(\lambda) = \theta_1(2\lambda_+) X_L(\zeta, \xi, \lambda - \eta; z) \langle \xi | \tag{4.23}$$

which are simply (4.6) in the specified parametrization. In the first equation, each operator entry of the matrix $L(\lambda)$ acts on the vacuum vector from the left while in the second one they act from the right. In the both equations, the two-component covector $\langle \zeta |$ is applied from the left. Using (3.3) and (4.5), one can represent these equations in the form where the two-component vector is applied from the right:

$$L(\lambda) |\xi\rangle^\perp X_R(z; \zeta, \xi, \lambda) = \theta_1(2\lambda_+) |\zeta\rangle^\perp X_R(z; \zeta, \xi, \lambda - \eta) \tag{4.24}$$

$$X_L(\zeta, \xi, \lambda; z) L(\lambda) |\xi\rangle^\perp = \theta_1(2\lambda_-) |\zeta\rangle^\perp X_L(\zeta, \xi, \lambda + \eta; z). \tag{4.25}$$

For the holomorphic vacuum vectors (4.11) at $\ell \in \mathbb{Z}_+$ these relations hold [12] under the additional restriction $\xi - \zeta \in 2\eta\mathbb{Z}$.

5. Global vacuum vectors

The next step, in accordance with [1], is to find the vacuum vectors for each $L^i(\lambda)$ and consider the action of the transfer matrix $T(\lambda)$ (2.3) on their direct product along the chain. It then appears that the result of this action has a very special and simple form: it is a sum of two terms, each of them being a vacuum vector that belongs to the same family. From the technical point of view, the easiest way to see this is to use the gauge transformation argument, now standard.

Let us make the ‘gauge transformation’ of the L-operator:

$$L^i(\lambda) \longrightarrow G(\zeta_i) L^i(\lambda) (G(\zeta_{i+1}))^{-1} = L'(\lambda) \tag{5.1}$$

where $G(\zeta)$ are some 2×2 c -number matrices depending on the parameter ζ . If $\zeta_{N+1} = \zeta_1$, then the trace is not changed:

$$T(\lambda) = \text{tr}(L'(\lambda)^1 L'(\lambda)^2 \dots L'(\lambda)^N).$$

We set the matrix $G(\zeta)$ to be

$$G(\zeta) = \begin{pmatrix} \bar{\theta}_4(\zeta) & \bar{\theta}_3(\zeta) \\ 0 & 1 \end{pmatrix}. \tag{5.2}$$

Then, acting on a vacuum vector X_R by each element of the gauge-transformed L-operator and using the relations (4.22), (4.24), we obtain

$$\begin{pmatrix} L'_{11} X_R & L'_{12} X_R \\ L'_{21} X_R & L'_{22} X_R \end{pmatrix} = \frac{1}{\bar{\theta}_4(\xi)} \begin{pmatrix} \bar{\theta}_4(\zeta) & \bar{\theta}_3(\zeta) \\ 0 & 1 \end{pmatrix} \begin{pmatrix} L_{11} X_R & L_{12} X_R \\ L_{21} X_R & L_{22} X_R \end{pmatrix} \begin{pmatrix} 1 & -\bar{\theta}_3(\xi) \\ 0 & \bar{\theta}_4(\xi) \end{pmatrix}$$

$$= \begin{pmatrix} X'_R & 0 \\ * & \frac{\bar{\theta}_4(\zeta)}{\bar{\theta}_4(\xi)} X''_R \end{pmatrix}. \tag{5.3}$$


Here $X'_R = \theta_1(2\lambda_-)X_R(z_i; \zeta, \xi, \lambda + \eta)$, $X''_R = \theta_1(2\lambda_+)X_R(z_i; \zeta, \xi, \lambda - \eta)$ and we imply that $\zeta_i = \zeta, \zeta_{i+1} = \xi$. The star in the lower left corner stands for something irrelevant for us.

We define the global vacuum vectors $\vec{X}_{R,L}^\pm$ to be products of the local ones over all sites of the chain:

$$\vec{X}_R^\pm(\lambda) = \vec{X}_R^\pm(\vec{z}, \vec{\zeta}, \lambda) = \prod_{i=1}^N X_R^\pm(z_i; \zeta_i, \zeta_{i+1}, \lambda) \tag{5.4}$$

$$\vec{X}_L^\pm(\lambda) = \vec{X}_L^\pm(\lambda, \vec{\zeta}, \vec{z}) = \prod_{i=1}^N X_L^\pm(\lambda, \zeta_i, \zeta_{i+1}; z_i) \tag{5.5}$$

where the notation $\vec{z} = \{z_1, \dots, z_N\}, \vec{\zeta} = \{\zeta_1, \dots, \zeta_N\}$ is used. The parameters ζ_i are naturally associated with the horizontal edges of the lattice:



Then

$$T(\lambda) \vec{X}_R^\pm(\lambda) = \text{tr} \prod_{i=1}^N (L^i(\lambda) X_R^\pm(z_i; \zeta_i, \zeta_{i+1}, \lambda))$$

and similarly for the $\vec{X}_L^\pm(\lambda)$. The triangularity of the matrix (5.3) allows one to compute the trace. Taking into account the periodic boundary condition, we arrive at the following relations:

$$T(\lambda) \vec{X}_R^\pm(\lambda) = \theta_1^N(2\lambda_-) \vec{X}_R^\pm(\lambda + \eta) + \theta_1^N(2\lambda_+) \vec{X}_R^\pm(\lambda - \eta) \tag{5.6}$$

$$\vec{X}_L^\pm(\lambda) T(\lambda) = \theta_1^N(2\lambda_-) \vec{X}_L^\pm(\lambda + \eta) + \theta_1^N(2\lambda_+) \vec{X}_L^\pm(\lambda - \eta). \tag{5.7}$$

Let us introduce left and right Q -operators, Q_L^\pm and Q_R^\pm (the *pre- Q -operators*). They are difference operators acting on functions $f(z_1, \dots, z_N)$ of N variables, respectively from the left and from the right, as follows:

$$(Q_L^\pm f)(\vec{\zeta}) = \int d\vec{z} \vec{X}_L^\pm(\lambda, \vec{\zeta}, \vec{z}) f(\vec{z}) \tag{5.8}$$

$$(f Q_R^\pm)(\vec{\zeta}) = \int d\vec{z} f(\vec{z}) \vec{X}_R^\pm(\vec{z}, \vec{\zeta}, \lambda). \tag{5.9}$$

The kernels here are the global vacuum vectors defined above. The symbol $\int d\vec{z} = \int \dots \int dz_1 \dots dz_N$ means the natural pairing in the tensor product of the local functional spaces. On each factor, this pairing coincides with (3.8).

The formulae (5.6), (5.7) can be written as the operator relations:

$$T(\lambda) Q_R^\pm(\lambda) = \theta_1^N(2\lambda_-) Q_R^\pm(\lambda + \eta) + \theta_1^N(2\lambda_+) Q_R^\pm(\lambda - \eta) \tag{5.10}$$

$$Q_L^\pm(\lambda) T(\lambda) = \theta_1^N(2\lambda_-) Q_L^\pm(\lambda + \eta) + \theta_1^N(2\lambda_+) Q_L^\pm(\lambda - \eta). \tag{5.11}$$

One might multiply the pre- Q -operators by a scalar normalization factor $f(\lambda)$ such that $f(\lambda) = f(\lambda + \eta)$. Clearly, these relations remain unchanged.

6. Basic commutation relations

In this section we prove the basic commutation relations


$$Q_L^\pm(\lambda)Q_R^\pm(\lambda') = Q_L^\pm(\lambda')Q_R^\pm(\lambda) \tag{6.1}$$

$$Q_L^+(\lambda)Q_R^-(\lambda') = Q_L^-(\lambda')Q_R^+(\lambda). \tag{6.2}$$

Let us start from the proof of (6.1) for $Q_{L,R}^+$. To this end, consider the function

$$\int d^N \vec{z} \bar{X}_L^+(\lambda, \vec{\zeta}, \vec{z}) \bar{X}_R^+(\vec{z}, \vec{\zeta}', \lambda') = \prod_{i=1}^N \int dz_i X_L^+(\lambda, \zeta_i, \zeta_{i+1}; z_i) X_R^+(z_i; \zeta'_i, \zeta'_{i+1}, \lambda') \tag{6.3}$$

which is the kernel of the operator product in the left-hand side of (6.1). Schematically, the structure of the kernel can be illustrated as follows:



As is clear from (6.3), the kernel is a product of local multipliers associated with each pair $(i, i + 1)$ of adjacent sites of the chain. For a while, we set $\zeta_i = \zeta$, $\zeta_{i+1} = \xi$. Therefore, we should investigate the function

$$F_{\zeta'_i \xi'_i}^{\zeta \xi}(\lambda, \lambda') = \int dz X_L^+(\lambda, \zeta, \xi; z) X_R^+(z; \zeta', \xi', \lambda'). \tag{6.4}$$

Let us begin with the X_L^+ . Up to an irrelevant constant factor, (4.19) reads

$$X_L^+(\lambda, \zeta, \xi; z) = \lim_{\varepsilon \rightarrow 0} [\varepsilon \mathcal{X}_L^\varepsilon(\zeta + \lambda_+; \zeta, \xi, \lambda)] \sum_{k \geq 0} x_k(\zeta, \xi, \lambda) \delta(z - \zeta - \lambda_+ + 2k\eta) \tag{6.5}$$

where

$$\begin{aligned} x_k(\zeta, \xi, \lambda) = & \frac{1}{[k]!} \frac{\theta_1(2\zeta + 2\lambda_+ - 4k\eta)}{\theta_1(2\zeta + 2\lambda_+)} \\ & \times \prod_{j=0}^{k-1} (\theta_1(-2\lambda_+ + 2j\eta)\theta_1(-2\lambda_+ - 2\zeta + 2j\eta)\theta_1(\xi - \zeta - 2\ell\eta + 2j\eta) \\ & \times \theta_1(-\xi - \zeta - 2\ell\eta + 2j\eta)) \{ \theta_1(-2\zeta + 2(j+1)\eta)\theta_1(\xi - \zeta - 2\lambda \\ & + 2(j+1)\eta)\theta_1(-\xi - \zeta - 2\lambda + 2(j+1)\eta) \}^{-1}. \end{aligned} \tag{6.6}$$

Substituting (6.5) and $X_R^+(z; \zeta', \xi', \lambda') = \mathcal{X}_R(z; \zeta', \xi', \lambda') \sum_{k' \geq 0} \delta(z - \xi' + \lambda'_- - \eta - 2k'\eta)$ (see (4.15)) into (6.4), performing the z -integral according to the pairing rule (3.9), and passing to summation over $n = k + k' \geq 0$, we get

$$\begin{aligned} F_{\zeta'_i \xi'_i}^{\zeta \xi}(\lambda, \lambda') = & \lim_{\varepsilon \rightarrow 0} [\varepsilon \mathcal{X}_L^\varepsilon(\zeta + \lambda_+; \zeta, \xi, \lambda)] \sum_{n \geq 0} \delta(\zeta - \xi' + \lambda + \lambda' - \eta - 2n\eta) \\ & \times \sum_{k=0}^n x_k(\zeta, \xi, \lambda) \mathcal{X}_R(\zeta + \lambda_+ - 2k\eta; \zeta', \xi', \lambda'). \end{aligned} \tag{6.7}$$

The next step is to identify the finite sum over k with the terminating modular hypergeometric function ${}_{10}W_9$. For the definition see appendix A. Plugging here the explicit formula for \mathcal{X}_R (4.11) and using (A13), one can represent (6.7) in the form

$$F_{\zeta'\xi'}^{\zeta\xi}(\lambda, \lambda') = F_{\text{sym}}^{\zeta\xi, \zeta'\xi'}(\lambda, \lambda')\omega(\lambda', \zeta' - \xi')F_1^{\zeta\xi}(\lambda)F_2^{\zeta'\xi'}(\lambda, \lambda')\frac{g_{\xi\xi'}(\lambda - \lambda')}{g_{\zeta\zeta'}(\lambda - \lambda')} \times \sum_{n \geq 0} {}_{10}W_9(\alpha_1; \alpha_4, \dots, \alpha_9, -n)\delta(\zeta - \xi' + \lambda + \lambda' - \eta - 2n\eta) \tag{6.8}$$

where

$$F_{\text{sym}}^{\zeta\xi, \zeta'\xi'} = \frac{e^{2\pi i \zeta} \Gamma(2\zeta)}{\Gamma(\zeta - \xi + 2(\ell + 1)\eta)\Gamma(\zeta + \xi + 2(\ell + 1)\eta)} \prod_{\epsilon = \pm} \frac{\Gamma(\zeta + \epsilon\zeta' + \lambda_+ + \lambda'_+ + \eta)}{\Gamma(\zeta + \epsilon\xi' + \lambda + \lambda' + \eta)}$$

is a symmetric function of λ, λ' for all $\zeta, \xi, \zeta', \xi', \omega(\lambda', \zeta' - \xi')$ is defined in (4.13), and

$$g_{\zeta\zeta'}(\lambda) = \Gamma(\zeta + \zeta' + \lambda + \eta)\Gamma(\zeta - \zeta' + \lambda + \eta) \tag{6.9}$$

$$F_1^{\zeta\xi}(\lambda) = e^{-4\pi i \ell \lambda} \frac{\Gamma(\zeta - \xi + 2\lambda)\Gamma(\zeta + \xi + 2\lambda)}{\Gamma(2\lambda_-)\Gamma(2\zeta + 2\lambda_+)} \tag{6.10}$$

$$F_2^{\zeta'\xi'}(\lambda, \lambda') = \frac{\Gamma(\zeta + \xi' + \lambda_+ - \lambda'_- + \eta)\Gamma(\zeta - \xi' + \lambda_+ - \lambda'_- + \eta)}{\Gamma(\xi + \xi' + \lambda - \lambda' + \eta)\Gamma(\xi - \xi' + \lambda - \lambda' + \eta)}. \tag{6.11}$$

The function ${}_{10}W_9$ is defined in (A20). It is an elliptic analogue of the very well poised hypergeometric series. The values of parameters are as follows:

$$\begin{aligned} -2\eta\alpha_1 &= 2\zeta + 2\lambda + 2\ell\eta & -2\eta\alpha_7 &= \zeta + \xi + 2\ell\eta \\ -2\eta\alpha_4 &= 2\lambda + 2\ell\eta & -2\eta\alpha_8 &= \zeta - \xi + 2\ell\eta \\ -2\eta\alpha_5 &= \zeta + \zeta' + \lambda - \lambda' - \eta & -2\eta\alpha_9 &= \zeta + \xi' + \lambda + \lambda' - \eta \\ -2\eta\alpha_6 &= \zeta - \zeta' + \lambda - \lambda' - \eta & -2\eta\alpha_{10} &= \zeta - \xi' + \lambda + \lambda' - \eta = 2n\eta. \end{aligned} \tag{6.12}$$

Note that the parameters in the right column are symmetric under the permutation $\lambda \leftrightarrow \lambda'$. The series is *balanced*, which means that the balancing condition

$$2 + 3\alpha_1 = \sum_{m=1}^7 \alpha_{m+3} \tag{6.13}$$

is satisfied. The series is terminating since $\alpha_{10} = -n$, where n is a non-negative integer number.

Let us move the factor $F_2(\lambda, \lambda')$ inside the sum $\sum_{n \geq 0}$ and set $\xi' = \zeta + \lambda + \lambda' - \eta - 2n\eta$ in the n th term (this is what the δ function suggests to do). After using (A13), (A14) and extracting the common factor

$$F_3^{\zeta\xi}(\lambda, \lambda') = \frac{\Gamma(2\zeta + 2\lambda_+)\Gamma(-2\lambda'_+ + 2\eta)}{\Gamma(\zeta + \xi + 2\lambda)\Gamma(-\zeta + \xi - 2\lambda' + 2\eta)} \tag{6.14}$$

the sum over n becomes equal to

$$F_3^{\zeta\xi}(\lambda, \lambda') \sum_{n \geq 0} \frac{[\alpha_1 - \alpha_7 + 1]_n [\beta_1 - \alpha_7 - \alpha_8 + 1]_n}{[\alpha_1 + 1]_n [\beta_1 - \alpha_8 + 1]_n} \times {}_{10}W_9(\alpha_1; \alpha_4, \dots, -n)\delta(\zeta - \xi' + \lambda + \lambda' - \eta - 2n\eta) \tag{6.15}$$

where we use the elliptic Pochhammer symbol $[x]_n = [x][x + 1] \dots [x + n - 1]$, and $\beta_1 = 2\alpha_1 + 1 - \alpha_4 - \alpha_5 - \alpha_6$. Now, we recall the elliptic analogue (A21) of the Bailey transformation formula for the terminating balanced ${}_{10}W_9$. It is easily seen from (6.12) that the Bailey transformation of the parameters $\alpha_i \rightarrow \beta_i$ (A22) is equivalent to the permutation

$\lambda \leftrightarrow \lambda'$ while all the other parameters remain unchanged. Therefore, as follows from (A21), each term inside the sum over n in (6.15) is symmetric under the permutation $\lambda \leftrightarrow \lambda'$. The common factor F_3 is not symmetric. Combining it with the remaining factors and taking into account (A10), we obtain

$$F_{\zeta'\xi'}^{\zeta\xi}(\lambda, \lambda') = \frac{g_{\xi\xi'}(\lambda - \lambda')}{g_{\zeta\zeta'}(\lambda - \lambda')} \sum_{n \geq 0} \mathcal{F}_{\text{sym}}^{\zeta\xi, \zeta'\xi'}(\lambda, \lambda'; n) \omega(\lambda, \zeta - \xi) \omega(\lambda', \zeta' - \xi') \times \delta(\zeta - \xi' + \lambda + \lambda' - \eta - 2n\eta) \tag{6.16}$$

where $\mathcal{F}_{\text{sym}}^{\zeta\xi, \zeta'\xi'}(\lambda, \lambda'; n)$ is symmetric under the permutation of λ and λ' for all $n, \zeta, \xi, \zeta', \xi'$, and the functions g, ω are given in (6.9), (4.13) respectively.

To obtain the kernel (6.3), we should set $\zeta = \zeta_i, \xi = \zeta_{i+1}$ and take the product of the local factors (6.16) over i from 1 to N provided $\zeta_{N+1} = \zeta_1$. Therefore, the ratios of the g functions cancel. The only factor which is not manifestly symmetric under the permutation $\lambda \leftrightarrow \lambda'$ comes from the product of the ω functions. This factor is equal to

$$\exp\left(-\frac{\pi i}{2\eta} \sum_{j=1}^N (\zeta'_j - \zeta'_{j+1})^2\right) \prod_{i=1}^N (\theta_1(\zeta_i - \zeta_{i+1} - 2\lambda|2\eta) \theta_1(\zeta'_i - \zeta'_{i+1} - 2\lambda'|2\eta))^{-1}. \tag{6.17}$$

However, we should take into account the constraints imposed by the δ functions: $\zeta'_i = \zeta_{i+1} - \lambda - \lambda' + \eta + 2n_{i-1}\eta$, where $n_i \in \mathbb{Z}$. In other words, (6.17) is zero unless $\zeta'_i - \zeta'_{i+1} = \zeta_{i-1} - \zeta_i + 2(n_{i-1} - n_i)\eta$. From the monodromy properties of the θ function it then follows that this function is symmetric, too. Therefore, the commutation relation (6.1) for $\mathcal{Q}_{L,R}^+$ is proved. To prove it for $\mathcal{Q}_{L,R}^-$, there is no need to repeat the calculations since $X_{L,R}^-$ is related to $X_{L,R}^+$ via (4.21).

The proof of (6.2) requires a similar but different calculation. Its scheme is outlined below. The kernel of the operator in the left-hand side is

$$\prod_{i=1}^N \int dz_i X_L^+(\lambda, \zeta_i, \zeta_{i+1}; z_i) X_R^-(z_i; \zeta'_i, \zeta'_{i+1}, \lambda') = \prod_{i=1}^N \int dz_i X_L^+(\lambda, \zeta_i, \zeta_{i+1}; z_i) X_R^+(z_i; \zeta'_{i+1}, \zeta'_i, -\lambda') \tag{6.18}$$

where (4.21) is used. Again, set $\zeta_i = \zeta, \zeta_{i+1} = \xi, \zeta'_i = \zeta', \zeta'_{i+1} = \xi'$, and examine the symmetry properties of the function

$$\tilde{F}_{\zeta'\xi'}^{\zeta\xi}(\lambda, \lambda') = \int dz X_L^+(\lambda, \zeta, \xi; z) X_R^+(z; \xi, \zeta, -\lambda')$$

under the simultaneous change $\zeta \leftrightarrow \xi, \zeta' \leftrightarrow \xi', \lambda \rightarrow -\lambda', \lambda' \rightarrow -\lambda$. It is enough to show that the product of such functions over the chain is invariant under this substitution in each factor. Proceeding as above, we obtain the representation of this function similar to (6.8):

$$\tilde{F}_{\zeta'\xi'}^{\zeta\xi}(\lambda, \lambda') = F_{\text{sym}}^{\zeta\xi, \zeta'\xi'}(\lambda, \lambda') \omega(-\lambda', \xi' - \zeta') F_1^{\zeta\xi}(\lambda) F_2^{\zeta\xi, \zeta'\xi'}(\lambda, \lambda') \frac{g_{\xi\xi'}(\lambda - \lambda')}{g_{\zeta\zeta'}(\lambda - \lambda')} \times \sum_{n \geq 0} {}_{10}W_9(\tilde{\alpha}_1; \tilde{\alpha}_4, \dots, \tilde{\alpha}_{10}) \delta(\zeta - \zeta' + \lambda - \lambda' - \eta - 2n\eta) \tag{6.19}$$

with the same functions $F_{\text{sym}}, \omega, F_1, F_2$, and g as before. The parameters of the series ${}_{10}W_9$ are also the same (see (6.12)); however, for the sake of convenience they are permuted in the following way: $\tilde{\alpha}_1 = \alpha_1, \tilde{\alpha}_4 = \alpha_8, \tilde{\alpha}_5 = \alpha_9, \tilde{\alpha}_6 = \alpha_{10}, \tilde{\alpha}_7 = \alpha_4, \tilde{\alpha}_8 = \alpha_5, \tilde{\alpha}_9 = \alpha_7, \tilde{\alpha}_{10} = \alpha_6$. (Clearly, the series is symmetric under the permutation of all parameters except α_1 .) What differs is the argument of the δ function. Note that the series is terminating since the δ function

is zero unless $\tilde{\alpha}_{10} = -n$. Next, we make the Bailey transformation (A21) passing from $\tilde{\alpha}_i$ to $\tilde{\beta}_i$ according to (A22). The result can be represented as a product of three factors:

$$\tilde{F}_{\zeta'\xi'}^{\zeta\xi}(\lambda, \lambda') = A_{\zeta'\xi'}^{\zeta\xi}(\lambda, \lambda') \times \left[\begin{array}{c} \text{invariant under } \zeta \leftrightarrow \xi, \\ \zeta' \leftrightarrow \xi', \lambda \leftrightarrow -\lambda' \end{array} \right] \times \left[\begin{array}{c} \text{factors that cancel in the} \\ \text{product over the chain} \end{array} \right]$$

where

$$A_{\zeta'\xi'}^{\zeta\xi}(\lambda, \lambda') = \frac{\exp(-\pi i(\xi' - \zeta')^2/(2\eta))}{\theta_1(\xi' - \zeta' - 2\lambda'/2\eta)\theta_1(\zeta - \xi + 2\lambda/2\eta)}.$$

The product $\prod_{i=1}^N A_{\zeta'_i \xi'_i}^{\zeta_i \xi_i}(\lambda, \lambda')$ is the only thing here which is not manifestly invariant under the simultaneous change $\zeta_i \leftrightarrow \zeta_{i+1}, \zeta'_i \leftrightarrow \zeta'_{i+1}, \lambda \leftrightarrow -\lambda'$ in each factor. However, the same argument as the one used for (6.17) establishes the invariance.

7. From pre- Q -operators to Q -operator

Suppose the operators Q_L^+ and Q_R^+ are invertible at some particular $\lambda = \lambda_*$. Following [1], we define

$$Q^\pm(\lambda) = (Q_L^+(\lambda_*))^{-1} Q_L^\pm(\lambda) = Q_R^\pm(\lambda) (Q_R^+(\lambda_*))^{-1}. \tag{7.1}$$

The second equality is due to the basic commutation relations (6.1), (6.2). It then follows, on applying (6.1), (6.2) once again, that $[Q^+(\lambda), Q^+(\mu)] = [Q^+(\lambda), Q^-(\mu)] = [Q^-(\lambda), Q^-(\mu)] = 0$ for all λ, μ :

$$\begin{aligned} Q^+(\lambda)Q^+(\mu) &= (Q_L^+(\lambda_*))^{-1} Q_L^+(\lambda) Q_R^+(\mu) (Q_R^+(\lambda_*))^{-1} \\ &= (Q_L^+(\lambda_*))^{-1} Q_L^+(\mu) Q_R^+(\lambda) (Q_R^+(\lambda_*))^{-1} \\ &= Q^+(\mu)Q^+(\lambda) \end{aligned} \tag{7.2}$$

and similarly for Q^- . Both equations (5.10), (5.11) lead to the Baxter T - Q relation

$$T(\lambda)Q(\lambda) = \theta_1^N(2\lambda - 2\ell\eta)Q(\lambda + \eta) + \theta_1^N(2\lambda + 2\ell\eta)Q(\lambda - \eta) \tag{7.3}$$

where Q is any one of Q^\pm . Besides, it follows that $Q^\pm(\lambda)$ commute with $T(\mu)$ for all λ, μ .

Suppose $\ell \notin \mathbb{Z}_+$. Then the convenient choice of the normalization point λ_* is $\lambda_* = -\ell\eta$. Indeed, at $\lambda = -\ell\eta$ the series (6.5) terminates at the $k = 0$ term; therefore, $Q^+(-\ell\eta)$ is a diagonal operator. Since we did not take care of ℓ -dependent constant factors, we continue to do so and redefine the Q -operator dividing it by the constant $\Gamma^N(-4\ell\eta)$. Note that this normalization allows us to define the Q -operator for $\ell \in \mathbb{Z}_+$ as well. We express the result through the function Φ_ℓ (A16):

$$Q_L^+(-\ell\eta) = \prod_{j=1}^N \Phi_\ell^{-1}(z_j - z_{j+1}) \Phi_\ell^{-1}(z_j + z_{j+1}). \tag{7.4}$$

Thus, by virtue of (5.8), the kernels of the Q -operators Q^\pm are

$$Q^\pm(\lambda; \vec{\zeta}, \vec{z}) = \prod_{i=1}^N q^\pm(\lambda; \zeta_i, \zeta_{i+1}, z_i) \tag{7.5}$$

where

$$\begin{aligned} q^+(\lambda; \zeta_i, \zeta_{i+1}, z_i) &= \frac{\Gamma(2\zeta_i)\Gamma(\zeta_i - \zeta_{i+1} + 2\lambda)\Gamma(\zeta_i + \zeta_{i+1} + 2\lambda)}{\Gamma(2\lambda_-)\Gamma(2\zeta_i + 2\lambda_+)\Gamma(\zeta_i - \zeta_{i+1} - 2\ell\eta)\Gamma(\zeta_i + \zeta_{i+1} - 2\ell\eta)} \\ &\times \sum_{k \geq 0} x_k(\zeta_i, \zeta_{i+1}, \lambda) \delta(z_i - \zeta_i - \lambda_+ + 2k\eta) \end{aligned} \tag{7.6}$$

(x_k is defined in (6.6)), and $q^-(\lambda, \zeta_i, \zeta_{i+1}, z_i) = q^+(-\lambda, \zeta_{i+1}, \zeta_i, z_i)$. Representing this in the operator form, according to the rule described in section 3, we obtain

$$\begin{aligned}
 Q^+(\lambda) &= \frac{1}{\Gamma^N(2\lambda - 2\ell\eta)} \prod_{i=1}^N \frac{\Gamma(2z_i)\Gamma(z_i - z_{i+1} + 2\lambda)\Gamma(z_i + z_{i+1} + 2\lambda)}{\Gamma(2z_i + 2\lambda + 2\ell\eta)\Gamma(z_i - z_{i+1} - 2\ell\eta)\Gamma(z_i + z_{i+1} - 2\ell\eta)} \\
 &\quad \times : \prod_{j=1}^N {}_6W_5 \left(-\frac{z_j + \lambda}{\eta} - \ell; -\frac{\lambda}{\eta} - \ell, -\frac{z_{j+1} + z_j}{2\eta} - \ell, \frac{z_{j+1} - z_j}{2\eta} - \ell; e^{-2\eta\partial_{z_j}} \middle| \eta, \tau \right) : \\
 &\quad \times \exp \left((\lambda + \ell\eta) \sum_{m=1}^N \partial_{z_m} \right) \tag{7.7}
 \end{aligned}$$

$$\begin{aligned}
 Q^-(\lambda) &= \frac{1}{\Gamma^N(-2\lambda - 2\ell\eta)} \prod_{i=1}^N \frac{\Gamma(2z_i)\Gamma(z_i - z_{i-1} - 2\lambda)\Gamma(z_i + z_{i-1} - 2\lambda)}{\Gamma(2z_i - 2\lambda + 2\ell\eta)\Gamma(z_i - z_{i-1} - 2\ell\eta)\Gamma(z_i + z_{i-1} - 2\ell\eta)} \\
 &\quad \times : \prod_{j=1}^N {}_6W_5 \left(-\frac{z_j - \lambda}{\eta} - \ell; \frac{\lambda}{\eta} - \ell, -\frac{z_j + z_{j-1}}{2\eta} - \ell, \frac{z_j - z_{j-1}}{2\eta} - \ell; e^{-2\eta\partial_{z_j}} \middle| \eta, \tau \right) : \\
 &\quad \times \exp \left((-\lambda + \ell\eta) \sum_{m=1}^N \partial_{z_m} \right) C \tag{7.8}
 \end{aligned}$$

where C is the cyclic shift operator,

$$(Cf)(z_1, z_2, \dots, z_{N-1}, z_N) = f(z_2, z_3, \dots, z_N, z_1) \tag{7.9}$$

with the kernel $C(\vec{\zeta}, \vec{z}) = \prod_{i=1}^N \delta(z_i - \zeta_{i+1})$. Clearly, C commutes with the $T(\lambda)$ and the $Q^\pm(\lambda)$.

In general, the series ${}_6W_5$ in (7.7), (7.8) is not terminating. However, at $\lambda = \mp\ell\eta \pm m\eta$, $m \in \mathbb{Z}_+$, the series ${}_6W_5$ in (7.7) (respectively, in (7.8)) terminates at the m th term. Note also that

$$Q^\pm(\pm\ell\eta \mp m\eta) = 0 \quad m = 0, 1, 2, \dots \tag{7.10}$$

due to zeros of the scalar common factor. At the points $\pm\ell\eta$ we have

$$Q^+(-\ell\eta) = \frac{1}{\Gamma^N(-4\ell\eta)} I \quad Q^-(\ell\eta) = \frac{1}{\Gamma^N(-4\ell\eta)} C \tag{7.11}$$

where I is the identity operator. So, at $\ell \in \frac{1}{2}\mathbb{Z}_+$ the Q -operators have poles at these points.

Let us evaluate the Q -operator at another distinguished point. Note that if $\lambda = (\ell + 1)\eta$, then ζ_i in the kernel of Q^+ decouples from ζ_{i+1} , so the operator simplifies:

$$Q^+((\ell + 1)\eta) = \left(\prod_{i=1}^N \Phi_\ell(z_i - z_{i+1})\Phi_\ell(z_i + z_{i+1}) \right) \prod_{j=1}^N \dot{w}_\ell. \tag{7.12}$$

The operators w_ℓ pairwise commute since they act on variables attached to different sites of the chain. From (7.7) one obtains the representation of the operator w_ℓ through the elliptic hypergeometric series ${}_4W_3$ with an operator argument:

$$w_\ell = \frac{e^{2\pi i(2\ell+1)z}\Gamma(2z)}{\Gamma(2z + 2(2\ell + 1)\eta)} : {}_4W_3 \left(-\frac{z}{\eta} - 2\ell - 1; -2\ell - 1; e^{-2\eta\partial_z} \middle| \eta, \tau \right) : e^{(2\ell+1)\eta\partial_z}. \tag{7.13}$$

At $\ell \in \frac{1}{2}\mathbb{Z}_+$ the series is terminating and gives the operator w_ℓ introduced in (3.4). So, at $\lambda = (\ell + 1)\eta$ we obtain a distinguished factorized integral of motion

$$A = \left(\prod_{i=1}^N \Phi_\ell(z_i + z_{i+1})\Phi_\ell(z_i - z_{i+1}) \right) \prod_{j=1}^N \dot{w}_\ell \tag{7.14}$$

(no normal ordering is required since the w_ℓ^j commute).

Let us give some remarks on the operator A . It is clear from section 3 that for $\ell \in \frac{1}{2}\mathbb{Z}_+$ the operator A annihilates the space $(\Theta_{4\ell}^+)^{\otimes N}$. At $\ell = 0$, this integral of motion is proportional to $T(0)$ and has an especially simple form: $2^N (\prod_{i=1}^N \theta_1(z_i + z_{i+1}) \theta_1(z_i - z_{i+1}) / \theta_1(2z_i)) \prod_{j=1}^N \sinh(\eta \partial_z)$. (In the limit to the XXX spin chain it coincides with the integral of motion found by Lipatov [23] in the context of high-energy QCD, see also [24].) Presumably, for $\ell \in \frac{1}{2}\mathbb{Z}_+$ A can be represented as the trace of the product of more general L-operators obtained from (2.1) by the fusion procedure, taken at a particular value of λ . For brevity, we denote $\varphi_\ell(z_i, z_{i+1}) = \Phi_\ell(z_i - z_{i+1}) \Phi_\ell(z_i + z_{i+1})$. Together with (3.6), the commutativity of A and $T(\lambda)$ implies the intertwining relation

$$T^{(\ell)}(\lambda) \prod_{i=1}^N \varphi_\ell(z_i, z_{i+1}) = \left(\prod_{i=1}^N \varphi_\ell(z_i, z_{i+1}) \right) T^{(-\ell-1)}(\lambda) \tag{7.15}$$

where the superscript of T indicates the value of spin. A direct proof of this relation is given in appendix B. We stress that the intertwining relation (3.6) holds for all the spin operators separately while (7.15) is valid for the transfer matrix only.

The operators $Q^\pm(\lambda)$ are two operator solutions of the Baxter T - Q relation (7.3) treated as a difference equation in λ . Therefore, their Wronskian,

$$W(\lambda) = Q^+(\lambda + \eta)Q^-(\lambda) - Q^+(\lambda)Q^-(\lambda + \eta)$$

obeys the functional equation $\theta_1^N(2\lambda - 2\ell\eta)W(\lambda) = \theta_1^N(2\lambda + 2\ell\eta)W(\lambda - \eta)$. Taking into account (7.10), one can write

$$W(\lambda) = \theta_1^N(2\lambda + 2\ell\eta|2\eta) \left(\frac{\Gamma(2\lambda + 2\ell\eta + 2\eta)}{\Gamma(2\lambda - 2\ell\eta + 2\eta)} \right)^N p(\lambda) A C \tag{7.16}$$

where $p(\lambda)$ is a diagonal operator such that $p(\lambda + 1) = p(\lambda)$, $p(\lambda + \eta) = e^{2\pi i N \eta} p(\lambda)$, and $p(\ell\eta) = I$. The operator A is given in (7.14). (The simplest way to find the operator factors is to calculate the Wronskian at $\lambda = \ell\eta$.) To find $p(\lambda)$ explicitly, a more careful analysis is required.

8. Miscellanea

Twisted boundary conditions. The twisted boundary condition means inserting a c -number matrix inside the trace of the chain of L-operators (2.3). To preserve integrability, this matrix must be a constant solution to the RLL = LLR relation. In the elliptic case, there are only three such solutions: the Pauli matrices σ_α , $\alpha = 1, 2, 3$. Therefore, the definition of the quantum transfer matrix with twisted boundary conditions is as follows:

$$T(\lambda) = \text{tr}[\underset{1}{L}(\lambda) \underset{2}{L}(\lambda) \dots \underset{N}{L}(\lambda) \sigma_\alpha]. \tag{8.1}$$

For definiteness, we consider the case of $\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.

Again, we employ the gauge transformation (5.1) with the same matrix (5.2). The whole argument remains the same with the only modification that $\zeta_{N+1} \neq \zeta_1$. The ζ_{N+1} should be adjusted in such a way that the matrix $\sigma'_1 = G(\zeta_{N+1})\sigma_1(G(\zeta_1))^{-1}$ be lower-triangular. Calculating it explicitly, we obtain

$$\sigma'_1 = \frac{1}{\bar{\theta}_4(\zeta_1)} \begin{pmatrix} \bar{\theta}_3(\zeta_{N+1}) & -2\theta_2(\zeta_1 + \zeta_{N+1})\theta_2(\zeta_1 - \zeta_{N+1}) \\ 1 & -\bar{\theta}_3(\zeta_1) \end{pmatrix} \tag{8.2}$$

where a simple identity for θ functions has been used. Therefore, if $\zeta_{N+1} = \zeta_1 + \frac{1}{2}$, then the right upper element is zero.

With this modification, the definitions of the vacuum vectors (5.4), (5.5) and of the pre- Q -operators (5.8), (5.9) remain the same. However, equations (5.10), (5.11) get modified:

$$T(\lambda)Q_R^\pm(\lambda) = \theta_1^N(2\lambda_-)Q_R^\pm(\lambda + \eta) - \theta_1^N(2\lambda_+)Q_R^\pm(\lambda - \eta) \tag{8.3}$$

$$Q_L^\pm(\lambda)T(\lambda) = \theta_1^N(2\lambda_-)Q_L^\pm(\lambda + \eta) - \theta_1^N(2\lambda_+)Q_L^\pm(\lambda - \eta). \tag{8.4}$$

The new constraint on ζ_{N+1} and ζ_1 does not destroy the basic commutation relation (6.1). Indeed, the g -factors in (6.16) still cancel in the product over the chain since $g_{\zeta_1\zeta'_1} = g_{\zeta_{N+1}\zeta'_{N+1}}$ due to (6.9) and (A7).

The normalization of the Q -operators goes through as before. The T - Q relation is

$$T(\lambda)Q(\lambda) = \theta_1^N(2\lambda - 2\ell\eta)Q(\lambda + \eta) - \theta_1^N(2\lambda + 2\ell\eta)Q(\lambda - \eta). \tag{8.5}$$

The Q -operators are given by the same formulae (7.7), (7.8), where it is implied that $\zeta_{N+1} = \zeta_1 + \frac{1}{2}$, and the cyclic shift operator C should be replaced by the operator C' such that $(C'f)(z_1, z_2, \dots, z_N) = f(z_2, z_3, \dots, z_N, z_1 + \frac{1}{2})$.

The case $N = 1$. For $N = 1$, there is only one L-operator in the chain (2.3). The trace of the $L(\lambda)$ is the simplest quantum transfer matrix $T(\lambda)$. It is proportional to s_0 :

$$T(\lambda) = \text{tr } L(\lambda) = \theta_1(2\lambda)s_0 \tag{8.6}$$

$$s_0 = \frac{\theta_1(2z - 2\ell\eta)}{\theta_1(2z)} e^{\eta\partial_z} + \frac{\theta_1(2z + 2\ell\eta)}{\theta_1(2z)} e^{-\eta\partial_z}. \tag{8.7}$$

This is the difference analogue of the Lamé operator proposed in [25]. The difference operator given by (8.7) for positive integer values of ℓ and arbitrary generic η has $2\ell + 1$ stable bands (and 2ℓ gaps) in the spectrum.

The construction of the previous sections gives a remarkable family of commuting difference operators in one variable. For $\ell \in \mathbb{Z}_+$ this family of operators commuting with (8.7) coincides with the one found in [26] by means of a different argument.

The case $N = 1$ is exceptional and must be considered separately. Some of the arguments for $N \geq 2$ are not directly applicable to $N = 1$ since the vacuum vectors (4.15), (4.17) are to be taken at $\zeta = \xi$ that for $\ell \in \mathbb{Z}_+$ means the *non-generic position* of the parameters. Nevertheless, it is a matter of direct verification to prove that the final formulae (7.7), (7.8) remain true, if just to change the normalization making them non-singular for $\ell \in \mathbb{Z}_+$:

$$Q_{(N=1)}^\pm(\pm\lambda) = \frac{\Gamma(2\lambda)\Gamma(2z)\Gamma(2z + 2\lambda)}{\Gamma(2\lambda - 2\ell\eta)\Gamma(2z - 2\ell\eta)\Gamma(2z + 2\lambda + 2\ell\eta)} \times :_6W_5 \left(-\frac{z+\lambda}{\eta} - \ell; -\frac{\lambda}{\eta} - \ell, -\frac{z}{\eta} - \ell, -\ell; e^{-2\eta\partial_z} \middle| \eta, \tau \right) : e^{(\lambda+\ell\eta)\partial_z}. \tag{8.8}$$

For $\ell \in \mathbb{Z}_+$ the series is terminating. In this case the above formula can be rewritten in a more explicit form:

$$Q_{(N=1)}^\pm(\lambda) = \sum_{k=0}^{\ell} A_k(z, \lambda) e^{(2k\eta - \ell\eta + \lambda)\partial_z} \tag{8.9}$$

where

$$A_k(z, \lambda) = (-1)^k \frac{[\ell]!}{[2\ell]!} \begin{bmatrix} \ell \\ k \end{bmatrix} \prod_{j=0}^{\ell-k-1} \frac{\theta_1(2z + 2(\ell - j)\eta)\theta_1(2\lambda + 2(\ell - j)\eta)}{\theta_1(2z + 2\lambda + 2(k - j)\eta)} \times \prod_{j=0}^{k-1} \frac{\theta_1(2z - 2(\ell - j)\eta)\theta_1(2\lambda - 2(\ell - j)\eta)}{\theta_1(2z + 2\lambda + 2(k + j - \ell)\eta)}. \tag{8.10}$$

If $k = 0$ or $k = \ell$, then the second (respectively, the first) product is absent.

The XXZ case. The construction of sections 5–7 admits several different trigonometric degenerations. Here we discuss the one corresponding to the *XXZ* spin chain with the standard 6-vertex *R*-matrix. The *L*-operator is (cf (2.1))

$$L^{(XXZ)}(u) = \frac{1}{2i} \begin{pmatrix} e^{2\pi i u + j_0} & -e^{-2\pi i u - j_0} & & \\ & j_+ & & \\ & & e^{2\pi i u - j_0} & j_- \\ & & & -e^{-2\pi i u + j_0} \end{pmatrix} \tag{8.11}$$

where *u* is the spectral parameter and j_0, j_{\pm} are operators acting on functions of a variable *x* as follows:

$$j_0 = \gamma \partial_x \quad j_{\pm} = e^{\pm 2\pi i x} (q^{\ell} e^{\mp \gamma \partial_x} - q^{-\ell} e^{\pm \gamma \partial_x}). \tag{8.12}$$

Here $q = e^{2\pi i \gamma}$, and these operators obey the commutation relations of the quantum algebra $U_q(sl(2))$:

$$[j_0, j_{\pm}] = \pm 2\pi i \gamma j_{\pm} \quad [j_+, j_-] = 2(q - q^{-1}) \sinh j_0. \tag{8.13}$$

The quantum transfer matrix *T*(*u*) is defined by the same formula (2.3).

The limiting transition from the elliptic formulae to the trigonometric ones consists of two steps:

- (i) The modular transformation $\tau \rightarrow -1/\tau$ with the change of variables $z_j = \tau x_j + \frac{1}{4}$, $\lambda = \tau u, \eta = \tau \gamma$.
- (ii) Tending τ to 0.

Then the *L*-operator defined as $L^{(XXZ)}(u) = \lim_{\tau \rightarrow i0} \sqrt{\frac{i\tau}{4}} e^{\frac{i\tau}{4}} L(\lambda)$ with the understanding that τ in the *L*(λ) (2.1) is replaced by $-1/\tau$, and $u = \lambda/\tau$ is kept fixed coincides with (8.11). The same limiting transition in (7.7) yields the trigonometric *Q*-operator.

Skipping the details (for the modular transformation of the θ functions and the elliptic gamma function see (A4), (A11), respectively), we present the final result. It is written in terms of the following special functions. The basic hypergeometric function ${}_2\phi_1$ [27] is defined as

$${}_2\phi_1 \left[\begin{matrix} a, b \\ c \end{matrix}; q, x \right] = \sum_{k=0}^{\infty} \frac{(a; q)_k (b; q)_k}{(q; q)_k (c; q)_k} x^k \tag{8.14}$$

where $(a; q)_k = (1 - a)(1 - aq)(1 - aq^2) \dots (1 - aq^{k-1})$. Up to simple redefinitions of parameters, this series gives the little *q*-Jacobi function (when the series is terminating, it is the little *q*-Jacobi polynomial). The *q*-gamma function is

$$\Gamma_q(x) = \frac{(q; q)_{\infty}}{(q^x; q)_{\infty}} (1 - q)^{1-x} \tag{8.15}$$

where, for $|q| < 1$, $(a; q)_{\infty} = \lim_{k \rightarrow \infty} (a; q)_k$. The *Q*-operator is given by

$$\begin{aligned} Q^+(u) &= e^{-2\pi i N \ell u} \prod_{i=1}^N \frac{\Gamma_{q^2}(\frac{x_i - x_{i+1} + 2u}{2\gamma})}{\Gamma_{q^2}(\frac{u}{\gamma} - \ell) \Gamma_{q^2}(\frac{x_i - x_{i+1}}{2\gamma} - \ell)} \\ &\times \prod_{j=1}^N {}_2\phi_1 \left[\begin{matrix} q^{-2\ell} e^{-4\pi i u}, q^{-2\ell} e^{2\pi i (x_{j+1} - x_j)} \\ q^2 e^{2\pi i (x_{j+1} - x_j - 2u)} \end{matrix}; q^2, q^{2(\ell+1)} e^{-2\gamma \partial_{x_j}} \right] : \\ &\times \exp \left((u + \ell \gamma) \sum_{m=1}^N \partial_{x_m} \right). \end{aligned} \tag{8.16}$$

As in (7.8), the formula for $Q^-(u)$ differs from (8.16) by the changes $u \rightarrow -u, x_i \leftrightarrow x_{i+1}$, and the additional cyclic shift *C*. The two operators $Q^{\pm}(u)$ obey the *T*-*Q* relation:

$$T(u)Q(u) = \sin^N 2\pi(u - \ell \gamma)Q(u + \gamma) + \sin^N 2\pi(u + \ell \gamma)Q(u - \gamma). \tag{8.17}$$

It is worth noting that the limit from (7.7) gives an additional operator factor which is omitted in (8.16) for it belongs to the centre of the algebra generated by the shifts $e^{2\gamma\partial_{x_j}}$ and the operator $\sum_{i=1}^N \partial_{x_i}$. This factor has the form

$$\prod_{i=1}^N \frac{\Gamma_{\tilde{q}^2}(-2u + 2\ell\gamma + 1)\Gamma_{\tilde{q}^2}(x_{i+1} - x_i + 2\ell\gamma + 1)}{\Gamma_{\tilde{q}^2}(x_{i+1} - x_i - 2u + 1)}$$

where $\tilde{q} = \exp(-\frac{\pi i}{2\gamma})$ is the modular partner of the $q = \exp(2\pi i\gamma)$ (cf [22]). It seems so that the non-degenerate Sklyanin algebra knows about both halves of the modular double.

9. Concluding remarks

In this work, using Baxter’s method of vacuum vectors, we have constructed a family of commuting difference operators in many variables. This family provides the Q -operator for the quantum integrable spin chain of XYZ type in the functional realization of the algebra of observables.

The form of the obtained Q -operators suggests a simple derivation of local integrals of motion for the XYZ chain in infinite-dimensional representations. For instance, the natural candidates for local Hamiltonians are $(Q^\pm(\mp\ell\eta))^{-1} d/d\lambda |_{\lambda=\mp\ell\eta} Q^\pm(\lambda)$. Their explicit form is, however, not very illuminating. The hermiticity properties of these operators are to be further investigated. We leave this for a future work.

Another important point remaining out of the scope of this paper is the limit to the XXX case ($\gamma \rightarrow 0$ in (8.11)). In this limit one should be able to link our results with the ones of the papers [6, 7, 11], where the Q -operator for the XXX spin chain and allied models with rational R -matrix was realized as an integral operator. However, it is hard to compare them directly since the limit is singular and, therefore, very delicate. Moreover, as it should be clear from the above, the rational limit is certainly not unique. Different rational limits of the trigonometric L -operator lead to L -operators for the XXX spin chains, where the $sl(2)$ algebra is realized in different ways. A detailed study of rational limits seems to be interesting since in this case, closely related to the representation theory of $SL(2, \mathbb{R})$ or $SU(1, 1)$, the result must admit a meaningful translation into the language of representation theory and harmonic analysis. This may give a hint of what an elliptic extension of the latter might look like.

The construction of the pre- Q -operators, $Q_L(\lambda)$ and $Q_R(\lambda)$, goes through also for inhomogeneous chains. The problem, however, is the normalization: in general there is no point λ_* such that $Q_L(\lambda_*)$ or $Q_R(\lambda_*)$ could be explicitly inverted. Generally speaking, the kernel of the Q -operator cannot be represented as a product of local factors, so local integrals of motion for inhomogeneous chains probably do not exist. In other respects, the pair of pre- Q -operators is a good substitute for the Q -operator. Their kernels are factorized though the operators do not commute. Presumably, they still obey the exchange commutation relations (6.1). Their proof given in section 6 should basically go through in the inhomogeneous case, too, since the key element of the proof is local in nature.

Finally, we note that the technique developed in this paper might help to construct the fundamental R -matrix in infinite-dimensional representations of the Sklyanin algebra.

Acknowledgments

I am grateful to E Antonov, S Kharchev, S Khoroshkin, I Krichever, A Levin, and T Takebe for useful discussions, and to V Spiridonov for bringing the paper [17] to my attention. This work was supported in part by RFBR grant 98-01-00344.

Appendix A

Theta functions. We use the following definition of the Jacobi θ functions:

$$\begin{aligned} \theta_1(z|\tau) &= -\sum_{k \in \mathbb{Z}} \exp(\pi i \tau (k + \frac{1}{2})^2 + 2\pi i (z + \frac{1}{2})(k + \frac{1}{2})) \\ \theta_2(z|\tau) &= \sum_{k \in \mathbb{Z}} \exp(\pi i \tau (k + \frac{1}{2})^2 + 2\pi i z (k + \frac{1}{2})) \\ \theta_3(z|\tau) &= \sum_{k \in \mathbb{Z}} \exp(\pi i \tau k^2 + 2\pi i z k) \\ \theta_4(z|\tau) &= \sum_{k \in \mathbb{Z}} \exp(\pi i \tau k^2 + 2\pi i (z + \frac{1}{2})k). \end{aligned} \tag{A1}$$

They also can be represented as infinite products. The infinite product representation for the $\theta_1(z|\tau)$ reads

$$\theta_1(z|\tau) = i \exp\left(\frac{i\pi\tau}{4} - i\pi z\right) \prod_{k=1}^{\infty} (1 - e^{2\pi i k \tau})(1 - e^{2\pi i((k-1)\tau+z)})(1 - e^{2\pi i(k\tau-z)}). \tag{A2}$$

Throughout the paper we write $\theta_a(x|\tau) = \theta_a(x)$. The transformation properties for shifts by the periods are

$$\theta_a(x \pm 1) = (-1)^{\delta_{a,1} + \delta_{a,2}} \theta_a(x) \quad \theta_a(x \pm \tau) = (-1)^{\delta_{a,1} + \delta_{a,4}} e^{-\pi i \tau \mp 2\pi i x} \theta_a(x). \tag{A3}$$

Under the modular transformation $\tau \rightarrow -1/\tau$ the θ functions behave as follows:

$$\begin{aligned} \theta_1(z|\tau) &= i\sqrt{i/\tau} e^{-\pi i z^2/\tau} \theta_1(z/\tau | -1/\tau) \\ \theta_2(z|\tau) &= \sqrt{i/\tau} e^{-\pi i z^2/\tau} \theta_4(z/\tau | -1/\tau) \\ \theta_3(z|\tau) &= \sqrt{i/\tau} e^{-\pi i z^2/\tau} \theta_3(z/\tau | -1/\tau) \\ \theta_4(z|\tau) &= \sqrt{i/\tau} e^{-\pi i z^2/\tau} \theta_2(z/\tau | -1/\tau). \end{aligned} \tag{A4}$$

By Θ_n we denote the space of θ functions of order n , i.e. entire functions $F(x)$, $x \in \mathbb{C}$, such that

$$F(x + 1) = F(x) \quad F(x + \tau) = (-1)^n e^{-\pi i n \tau - 2\pi i n x} F(x). \tag{A5}$$

It is easy to see that $\dim \Theta_n = n$. Let $F(x) \in \Theta_n$, then $F(x)$ has a multiplicative representation of the form $F(x) = c \prod_{i=1}^n \theta_1(x - x_i)$, $\sum_{i=1}^n x_i = 0$, where c is a constant. Imposing, in addition to (A5), the condition $F(-x) = F(x)$, we define the space $\Theta_n^+ \subset \Theta_n$ of even θ functions of order n , which plays the important role in representations of the Sklyanin algebra. If n is an even number, then $\dim \Theta_n^+ = \frac{1}{2}n + 1$.

Elliptic gamma function. Here we collect the main formulae on the elliptic gamma function [18, 19]. We use the (slightly modified) notation of [19]. The elliptic gamma function is defined by the double-infinite product

$$\Gamma(z|\tau, \tau') = \prod_{k, k'=0}^{\infty} \frac{1 - e^{2\pi i((k+1)\tau + (k'+1)\tau' - z)}}{1 - e^{2\pi i(k\tau + k'\tau' + z)}}. \tag{A6}$$

A sufficient condition for the product to be convergent is $\text{Im } \tau > 0$, $\text{Im } \tau' > 0$. We need the following properties of the elliptic gamma function:

$$\Gamma(z + 1|\tau, \tau') = \Gamma(z|\tau, \tau') \tag{A7}$$

$$\Gamma(z + \tau|\tau, \tau') = -ie^{-\frac{\pi i \tau'}{6}} \eta_D^{-1}(\tau') e^{\pi i z} \theta_1(z|\tau') \Gamma(z|\tau, \tau') \tag{A8}$$

$$\Gamma(z + \tau'|\tau, \tau') = -ie^{-\frac{\pi i \tau}{6}} \eta_D^{-1}(\tau) e^{\pi i z} \theta_1(z|\tau) \Gamma(z|\tau, \tau') \tag{A9}$$

where

$$\eta_D(\tau) = e^{\frac{\pi i \tau}{12}} \prod_{k=1}^{\infty} (1 - e^{2\pi i k \tau})$$

is the Dedekind function. Another useful property is

$$\Gamma(z|\tau, \tau') \Gamma(\tau' - z|\tau, \tau') = \frac{ie^{\pi i \tau' / 6} \eta_D(\tau')}{e^{\pi i z} \theta_1(z|\tau')} \tag{A10}$$

Under the modular transformation $\tau \rightarrow -1/\tau$ the elliptic gamma function behaves as follows [19]:

$$\Gamma(z|\tau, \tau') = e^{i\pi P(z)} \frac{\Gamma(z/\tau| -1/\tau, \tau'/\tau)}{\Gamma((z - \tau)/\tau' | -\tau/\tau', -1/\tau')} \tag{A11}$$

where

$$P(z) = -\frac{1}{3\tau\tau'} z^3 + \frac{\tau + \tau' - 1}{2\tau\tau'} z^2 - \frac{\tau^2 + \tau'^2 + 3\tau\tau' - 3\tau - 3\tau' + 1}{6\tau\tau'} z - \frac{(\tau + \tau' - 1)(\tau + \tau' - \tau\tau')}{12\tau\tau'}. \tag{A12}$$

Let us list the most frequently used formulae for $\Gamma(z) \equiv \Gamma(z|\tau, 2\eta)$. Using (A9) several times, we obtain

$$\frac{\Gamma(x + 2k\eta)}{\Gamma(x)} = e^{\pi i \eta k^2} R^{-k} e^{\pi i k x} \prod_{j=0}^{k-1} \theta_1(x + 2j\eta) \tag{A13}$$

$$\frac{\Gamma(x - 2k\eta)}{\Gamma(x)} = (-1)^k e^{\pi i \eta k^2} R^k e^{-\pi i k x} \prod_{j=0}^{k-1} (\theta_1(-x + 2\eta + 2j\eta))^{-1} \tag{A14}$$

where $R = ie^{\pi i(\eta + \tau/6)} \eta_D(\tau)$.

As is seen from (A6), the function $\Gamma(z|\tau, 2\eta)$ has zeros at the points $z = 2(k + 1)\eta + (m + 1)\tau + n$, and simple poles at the points $z = -2k\eta - m\tau + n$, where k, m run over non-negative integers and n over all integers. The residues of the elliptic gamma function at the poles at $z = -2k\eta, k = 0, 1, 2, \dots$ are

$$\text{res}_{|z=-2k\eta} \Gamma(z) = (-1)^k e^{\pi i \eta k^2} R^k r_0 \prod_{j=1}^k (\theta_1(2j\eta))^{-1} \tag{A15}$$

where

$$r_0 = \text{res}_{|z=0} \Gamma(z) = -\frac{e^{\pi i(\tau + 2\eta)/12}}{2\pi i \eta_D(\tau) \eta_D(2\eta)}.$$

The function $\Phi_\ell(z)$ (4.14) can be extended to arbitrary complex values of ℓ as follows:

$$\Phi_\ell(z) = e^{-\pi i \eta(2\ell + 1)} R^{2\ell + 1} e^{-\pi i(2\ell + 1)z} \frac{\Gamma(z + 2(\ell + 1)\eta)}{\Gamma(z - 2\ell\eta)}. \tag{A16}$$

Elliptic hypergeometric series. Here we follow [17]. We define the elliptic Pochhammer symbol (the shifted elliptic factorial) by

$$[x]_k \equiv [x][x + 1] \dots [x + k - 1] \tag{A17}$$

where $[x] = \theta_1(2x\eta)$ (cf (3.5)). By definition, the elliptic hypergeometric series is

$${}_{r+1}W_r(\alpha_1; \alpha_4, \alpha_5, \dots, \alpha_{r+1}; z|\eta, \tau) = \sum_{k=0}^{\infty} z^k \frac{[\alpha_1 + 2k][\alpha_1]_k}{[\alpha_1][k]!} \prod_{m=1}^{r-2} \frac{[\alpha_{m+3}]_k}{[\alpha_1 - \alpha_{m+3} + 1]_k}. \tag{A18}$$

This is an elliptic analogue of the very well poised basic hypergeometric series [27]. The series is said to be *balanced* if $z = 1$ and

$$r - 5 + (r - 3)\alpha_1 = 2 \sum_{m=1}^{r-2} \alpha_{m+3}. \tag{A19}$$

For a series $\sum_{k \geq 0} c_k$ of the form (A18), the balancing condition (A19) means that the ratio c_{k+1}/c_k of the coefficients is an elliptic function of k . For balanced series (A18), we drop the argument $z = 1$ and the parameters η, τ writing it simply as ${}_{r+1}W_r(\alpha_1; \alpha_4, \dots, \alpha_{r+1})$. For instance,

$${}_{10}W_9(\alpha_1; \alpha_4, \alpha_5, \dots, \alpha_{10}) = \sum_{k=0}^{\infty} \frac{[\alpha_1 + 2k][\alpha_1]_k}{[\alpha_1][k]!} \prod_{m=1}^7 \frac{[\alpha_{m+3}]_k}{[\alpha_1 - \alpha_{m+3} + 1]_k}. \tag{A20}$$

The series is called *terminating* if at least one of the parameters $\alpha_4, \dots, \alpha_{r+1}$ is equal to a negative integer number. In this case there is no problem of convergence. If, say $\alpha_{r+1} = -n$, then the series terminates at $k = n$. The terminating balanced hypergeometric series were shown [17] to possess nice modular properties. That is why they were called *modular hypergeometric series*.

The modular hypergeometric series obey a number of spectacular identities. One of them is the elliptic analogue of the Bailey transformation for the ${}_{10}W_9$ [27]:

$$\begin{aligned} {}_{10}W_9(\alpha_1; \alpha_4, \dots, \alpha_{10}) &= {}_{10}W_9(\beta_1; \beta_4, \dots, \beta_{10})([\alpha_1 + 1]_n [\beta_1 - \beta_7 + 1]_n [\beta_1 - \beta_8 + 1]_n \\ &\quad \times [\alpha_1 - \alpha_7 - \alpha_8 + 1]_n \\ &\quad \times \{[\beta_1 + 1]_n [\alpha_1 - \alpha_7 + 1]_n [\alpha_1 - \alpha_8 + 1]_n [\beta_1 - \beta_7 - \beta_8 + 1]_n\}^{-1} \end{aligned} \tag{A21}$$

where α_j obey the balancing condition $2 + 3\alpha_1 = \sum_{m=1}^7 \alpha_{m+3}$, $\alpha_{10} = -n$, and the parameters β_j are as follows:

$$\begin{aligned} \beta_1 &= 2\alpha_1 + 1 - \alpha_4 - \alpha_5 - \alpha_6 \\ \beta_i &= \beta_1 - \alpha_1 + \alpha_i & i = 4, 5, 6 \\ \beta_i &= \alpha_i & i = 7, \dots, 10. \end{aligned} \tag{A22}$$

This identity was proved in [17].

Appendix B

Here we present the proofs of the intertwining relations (3.6) and the commutation relation (7.15).

For general complex values of ℓ the operator w_ℓ is given by the series (7.13):

$$w_\ell = \frac{e^{2\pi i(2\ell+1)z} \Gamma(2z)}{\Gamma(2z + 2(2\ell + 1)\eta)} : {}_4W_3 \left(-\frac{z}{\eta} - 2\ell - 1; -2\ell - 1; e^{-2\eta\partial_z} | \eta, \tau \right) : e^{(2\ell+1)\eta\partial_z}. \tag{B1}$$

The proof of (3.6) is a straightforward computation. Substituting (2.2), (B1) into (3.6) and comparing the coefficients of $e^{2(\ell+1-k)\eta\partial_z}$, $k > 0$, we get, after cancelling out common factors:

$$\begin{aligned} &\theta_1(2(\ell + k + 1)\eta)\theta_{a+1}(2z + 2(\ell + 1)\eta)\theta_1(2z + 4k\eta)\theta_1(2z - 2(\ell - k + 1)\eta) \\ &\quad + \theta_1(2(\ell - k + 1)\eta)\theta_{a+1}(-2z + 2(\ell + 1)\eta)\theta_1(2z + 4k\eta)\theta_1(2z + 2(\ell + k + 1)\eta) \\ &= \theta_1(2(\ell + k + 1)\eta)\theta_{a+1}(2z + 2(2k - \ell - 1)\eta)\theta_1(2z)\theta_1(2z + 2(\ell + k + 1)\eta) \\ &\quad + \theta_1(2(\ell - k + 1)\eta)\theta_{a+1}(-2z - 2(2k + \ell + 1)\eta)\theta_1(2z)\theta_1(2z - 2(\ell - k + 1)\eta). \end{aligned}$$

All the four terms, as functions of $2z$, have the same monodromy properties. They are θ functions of 3D order. The linear space of such functions is three-dimensional, so it is enough

to verify the equality at three points, say $z = 0, z = -2k\eta$, and $z = (\ell - k + 1)\eta$. The coefficients of the highest shift $e^{2(\ell+1)\eta\partial_z}$ in both sides of (3.6) are equal, too, so the intertwining relation is proved.

To prove (7.15), we use the representation (3.2). For any two adjacent sites of the chain (say 1 and 2) we define the c -number matrix $M = M^{12, 12}(\lambda, \ell) = V^{-1}(\lambda_-, z_1)V(\lambda_+, z_2)$ explicitly given by

$$M^{12}(\lambda, \ell) = \frac{1}{\theta_1(2z_2)} \begin{pmatrix} \theta_1(z_1 + z_2 - 2\ell\eta) & \theta_1(z_1 - z_2 + 2\lambda) \\ -\theta_1(z_1 - z_2 - 2\ell\eta) & \theta_1(z_1 + z_2 + 2\lambda) \\ \theta_1(z_1 - z_2 + 2\ell\eta) & \theta_1(z_1 + z_2 - 2\lambda) \\ -\theta_1(z_1 + z_2 + 2\ell\eta) & \theta_1(z_1 - z_2 - 2\lambda) \end{pmatrix}.$$

Using the cyclic property of the trace, one can rewrite (2.3) as follows:

$$T(\lambda) = : \text{tr} \left[M^{12} \begin{pmatrix} e^{\eta\partial_2} & 0 \\ 0 & e^{-\eta\partial_2} \end{pmatrix} M^{23} \begin{pmatrix} e^{\eta\partial_3} & 0 \\ 0 & e^{-\eta\partial_3} \end{pmatrix} \cdots M^{N1} \begin{pmatrix} e^{\eta\partial_1} & 0 \\ 0 & e^{-\eta\partial_1} \end{pmatrix} \right] :$$

where $\partial_i = \partial_{z_i}$. Computing the trace, we obtain

$$T(\lambda) = \sum_{i_1, \dots, i_N=1,2} M^{12}_{i_1 i_2} M^{23}_{i_2 i_3} \cdots M^{N1}_{i_N i_1} e^{(3-2i_1)\eta\partial_1} e^{(3-2i_2)\eta\partial_2} \cdots e^{(3-2i_N)\eta\partial_N}. \quad (\text{B2})$$

Recall that $\varphi_\ell(z_i, z_{i+1}) = \Phi_\ell(z_i - z_{i+1})\Phi_\ell(z_i + z_{i+1})$. The commutation relation (7.15),

$$T^{(\ell)}(\lambda) \prod_{i=1}^N \varphi_\ell(z_i, z_{i+1}) = \left(\prod_{i=1}^N \varphi_\ell(z_i, z_{i+1}) \right) T^{(-\ell-1)}(\lambda)$$

follows from (B2) and the local commutation rule

$$e^{(3-2i_1)\eta\partial_1 + (3-2i_2)\eta\partial_2} \varphi_\ell(z_1, z_2) = \frac{\theta_1(z_1 + (-1)^{i_1+i_2}z_2 - 2(-1)^{i_1}(\ell+1)\eta)}{\theta_1(z_1 + (-1)^{i_1+i_2}z_2 + 2(-1)^{i_1}\ell\eta)} \times \varphi_\ell(z_1, z_2) e^{(3-2i_1)\eta\partial_1 + (3-2i_2)\eta\partial_2} \quad i_1, i_2 = 1, 2$$

which is easy to verify. Writing out the entries of the matrix M in the form

$$M^{12}_{ik} = \theta_1(z_1 + (-1)^{i+k}z_2 + 2(-1)^i\ell\eta)C_{ik} \quad i, k = 1, 2 \quad (\text{B3})$$

(the second factor stands for something that does not depend on ℓ), and subsequently moving the factors $\varphi_\ell(z_i, z_{i+1})$ from the right to the left through each term of (B2), we arrive at (7.15).

References

[1] Baxter R 1972 Partition function of the eight-vertex lattice model *Ann. Phys., NY* **70** 193–228
 Baxter R 1973 Eight-vertex model in lattice statistics and one-dimensional anisotropic Heisenberg chain I, II, III *Ann. Phys., NY* **76** 1–24, 25–47, 48–71

[2] Faddeev L D and Takhtadzhian L A 1979 The quantum method of the inverse problem and the Heisenberg XYZ model *Russian Math. Surveys* **34** 11–68 (Engl. transl.)

[3] Korepanov I G 1994 Vacuum curves of \mathcal{L} -operators connected with the six-vertex model and construction of \mathcal{R} -operators *Algebra i Analiz* **6** 176–94
 (Korepanov I G 1994 *Preprint Chelyabinsk Polytechn. Inst.* Deposited in the VINITI 02.04.86 N 2271-B86)
 Bazhanov V V and Yu G Stroganov 1990 Chiral Potts model as a descendant of the six-vertex model *J. Stat. Phys.* **59** 799–817

[4] Batchelor M, Baxter R, O'Rourke M and Yung C 1995 Exact solution and interfacial tension of the six-vertex model with anti-periodic boundary conditions *J. Phys. A: Math. Gen.* **28** 2759–70
 Batchelor M and Yung C 1995 Exact solution for the spin- s XXZ quantum chain with non-diagonal twists *Nucl. Phys. B* **446** 461–84

[5] Pasquier V and Gaudin M 1992 The periodic Toda chain and a matrix generalization of the Bessel function recursion relations *J. Phys. A: Math. Gen.* **28** 5243–52

- [6] Derkachov S 1999 Baxter's Q -operator for the homogeneous XXX spin chain *J. Phys. A: Math. Gen.* **32** 5299–316 (Derkachov S 1999 *Preprint* solv-int/9902015)
- [7] Pronko G 1999 On the Baxter's Q -operator for the XXX spin chain *Commun. Math. Phys.* in press (*Preprint* hep-th/9908179)
- [8] Bazhanov V, Lukyanov S and Zamolodchikov A 1997 Integrable structure of conformal field theory II. Q -operator and DDV equation *Commun. Math. Phys.* **190** 247–78
Bazhanov V, Lukyanov S and Zamolodchikov A 1999 Integrable structure of conformal field theory III. The Yang–Baxter relation *Commun. Math. Phys.* **200** 297–324
- [9] Antonov A and Feigin B 1997 Quantum group representations and Baxter equation *Phys. Lett. B* **392** 115–22
- [10] Kuznetsov V and Sklyanin E 1998 On Bäcklund transformations for many-body systems *J. Phys. A: Math. Gen.* **31** 2241–51
- [11] Kuznetsov V, Salerno M and Sklyanin E 2000 Quantum Bäcklund transformation for the integrable DST model *J. Phys. A: Math. Gen.* **33** 171–89
(Kuznetsov V, Salerno M and Sklyanin E 1999 *Preprint* solv-int/9908002)
- [12] Takebe T 1992 Generalized Bethe ansatz with the general spin representation of the Sklyanin algebra *J. Phys. A: Math. Gen.* **25** 1071–83
Takebe T 1995 Bethe ansatz for higher spin eight vertex models *J. Phys. A: Math. Gen.* **28** 6675–706
- [13] Bazhanov V and Reshetikhin N 1989 Critical RSOS models and conformal field theory *Int. J. Mod. Phys. A* **4** 115–42
- [14] Krichever I, Lipan O, Wiegmann P and Zabrodin A 1997 Quantum integrable models and discrete classical Hirota equations *Commun. Math. Phys.* **188** 267–304
- [15] Sklyanin E K 1982 On some algebraic structures related to the Yang–Baxter equation *Funk. Anal. i ego Pril.* **16** 27–34
- [16] Sklyanin E K 1983 On some algebraic structures related to the Yang–Baxter equation: representations of the quantum algebra *Funk. Anal i ego Pril.* **17** 34–48
- [17] Frenkel I and Turaeu V 1997 Elliptic solutions of the Yang–Baxter equation and modular hypergeometric functions *The Arnold–Gelfand Mathematical Seminars* (Cambridge, MA: Birkhäuser Boston) 171–204
- [18] Ruijsenaars S N M 1997 First order analytic difference equations and integrable quantum systems *J. Math. Phys.* **38** 1069–146
- [19] Felder G and Varchenko A 1999 The elliptic gamma function and $SL(3, \mathbb{Z}) \times \mathbb{Z}^3$ *Preprint* math.QA/9907061
- [20] Zabrodin A 1999 On the spectral curve of the difference Lamé operator *Int. Math. Res. Notices* vol 11 589–614
- [21] Krichever I 1981 Baxter's equations and algebraic geometry *Funk. Anal. i ego Pril.* **15** 22–35
- [22] Faddeev L D 1999 Modular double of quantum group *Preprint* math.QA/9912078
- [23] Lipatov L N 1994 High energy asymptotics of multi-colour QCD and exactly solvable lattice models *JETP Lett.* **59** 571–4
- [24] Faddeev L and Korchemsky G 1995 High energy QCD as a completely integrable model *Phys. Lett. B* **342** 311–22
- [25] Krichever I and Zabrodin A 1995 Spin generalization of the Ruijsenaars–Schneider model, non-abelian 2D Toda chain and representations of Sklyanin algebra *Usp. Mat. Nauk* **50** 3–56
(Krichever I and Zabrodin A 1995 *Preprint* hep-th/9505039)
- [26] Felder G and Varchenko A 1998 Algebraic integrability of the two-body Ruijsenaars operator *Funk. Anal. i ego Pril.* **32** 8–25
(Felder G and Varchenko A 1996 *Preprint* q-alg/9610024)
- [27] Gasper G and Rahman M 1990 Basic hypergeometric series *Encyclopedia of Mathematics and its Applications* vol 35 (Cambridge: Cambridge University Press)