

Lie point symmetries of difference equations and lattices

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2000 J. Phys. A: Math. Gen. 33 8507

(<http://iopscience.iop.org/0305-4470/33/47/313>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.124

The article was downloaded on 02/06/2010 at 08:43

Please note that [terms and conditions apply](#).

Lie point symmetries of difference equations and lattices

D Levi[†], S Tremblay[‡] and P Winternitz[§]

[†] Dipartimento di Fisica, Università Roma Tre and INFN–Sezione di Roma Tre, Via della Vasca Navale 84, 00146 Rome, Italy

[‡] Centre de Recherches Mathématiques and Département de Physique, Université de Montréal, CP 6128, succ. Centre-ville, Montréal (QC), Canada H3C 3J7

[§] Centre de Recherches Mathématiques and Département de Mathématiques et Statistiques, Université de Montréal, CP 6128, succ. Centre-ville, Montréal (QC), Canada H3C 3J7

Received 25 July 2000

Abstract. A method is presented for finding the Lie point symmetry transformations acting simultaneously on difference equations and lattices, while leaving the solution set of the corresponding difference scheme invariant. The method is applied to several examples. The found symmetry groups are used to obtain particular solutions of differential-difference equations.

1. Introduction

Lie groups have long been used to study differential equations. As a matter of fact, they originated in that context [1, 2]. They have been put to good use to solve differential equations, to classify them and to establish properties of their solution spaces [3–8].

Applications of Lie group theory to discrete equations, such as difference equations, differential-difference equations or q -difference equations, are much more recent [9–37].

Several different approaches are being pursued. One philosophy is to consider a given system of discrete equations on a given fixed lattice and to search for a group of transformations, taking solutions into solutions, while leaving the lattice invariant. Within this philosophy different approaches differ by the restrictions imposed on the transformations and by the methods used to find the symmetries. One thing that is clear is that within this philosophy it is necessary to generalize the concept of point symmetries for difference equations, if we wish to recover all point symmetries of a differential equation in the continuous limit [9–26].

A different philosophy is to consider a difference equation and a lattice as two relations involving a fixed number of points, in which we give the values of the independent and dependent variables, say x_-, x, x_+ and u_-, u, u_+ respectively. The group transformations act on the equation and on the lattice. This philosophy was mainly developed by Dorodnitsyn and collaborators [27–33]. In this approach, the given object was a Lie group and its Lie algebra. Invariants of this Lie group, depending on x and u , calculated at a predetermined number of points were obtained. They were used to obtain invariant equations and lattices. The emphasis was on discretizing differential equations while preserving all of their point symmetries, or at least most of them.

The purpose of this paper is to combine the two philosophies. More specifically, we will consider given equations on given lattices, but the lattice will also be given by some equation. We will then look for Lie point transformations, acting on both equations, and leaving the common solution sets of both equations invariant.

In section 2 we develop the formalism necessary for calculating simultaneous symmetries of difference or differential-difference equations and lattices. Section 3 is devoted to examples of symmetries of purely difference equations, both linear and nonlinear ones. In section 4 we also consider examples, this time of differential-difference equations. Some conclusions are drawn in the final section 5.

2. Symmetries of differential-difference equations

2.1. The differential-difference scheme

In this paper we shall only consider a restricted class of problems, for reasons of simplicity and clarity. However, the formalism involved can easily be extended to quite general systems of equations.

Thus we shall consider one scalar function $u(x, t)$ of two variables only. The variable t is continuous and varies in some interval $I \subset \mathbb{R}$. The variable x is also continuous and varies in some interval $\tilde{I} \subset \mathbb{R}$. However, x will be ‘sampled’ in a set of discrete points $\{\dots, x_{n-2}, x_{n-1}, x_n, x_{n+1}, \dots\}$. The points x_k are not necessarily equally spaced.

We shall study the symmetries of a pair of equations, which we postulate to have the form

$$E = E(t, \{x_k\}_{k=n-n_1}^{n+n_2}, \{u_k\}_{k=n-n_1}^{n+n_2}, u_{n,t}, u_{n,tt}) = 0 \quad (1)$$

$$\Omega = \Omega(t, \{x_k\}_{k=n-n_3}^{n+n_4}, \{u_k\}_{k=n-n_3}^{n+n_4}) = 0 \quad n_i \geq 0. \quad (2)$$

We have $k, n, n_i \in \mathbb{Z}$, all n_i are finite. Equation (1) is a differential equation in t and a difference equation in x , since we define

$$\begin{aligned} x_n &\equiv x & x_{n-1} &\equiv x_n - h_-(x_n, t) \\ x_{n+1} &\equiv x_n + h_+(x_n, t) & x_{n+2} &\equiv x_n + h_+(x_n, t) + h_+(x_{n+1}, t), \dots \\ u_n &\equiv u(x_n, t) & u_{n+k} &\equiv u(x_{n+k}, t). \end{aligned} \quad (3)$$

At this stage we are not imposing any boundary conditions, so we assume that equations (1) and (2) can be shifted arbitrarily to the left and to the right. Thus, equations (1) and (2) involve any $n_1 + n_2 + 1$ or $n_3 + n_4 + 1$ neighbouring points, respectively.

The fact that (1) involves only first and second derivatives and that there are no derivatives in (2) is also for simplicity only. The same goes for the fact that derivatives are evaluated at the reference point n only (i.e. we do not consider terms such as $\partial u(x_{n+1}, t)/\partial t$).

In order to be able to consider equations (1) and (2) as a difference scheme, we must be able to obtain x_{n+N}, u_{n+N} and also x_{n-M}, u_{n-M} ($N = \max(n_2, n_4)$, $M = \max(n_1, n_3)$). In other words, we impose two conditions:

$$\det \left(\frac{\partial(E, \Omega)}{\partial(x_{n+N}, u_{n+N})} \right) \neq 0 \quad \det \left(\frac{\partial(E, \Omega)}{\partial(x_{n-M}, u_{n-M})} \right) \neq 0. \quad (4)$$

If necessary, when calculating (4) we shift one of the equations, (1) or (2), to the left or right, so that the same values $n + N$ and $n - M$ figure in both equations.

In general, we do not require that a continuous limit should exist. If it does, then equation (1) should go into a differential equation in x and t and equation (2) should go into the identity $0 = 0$. When taking the continuous limit it is convenient to introduce ‘discrete derivatives’, e.g.

$$u_{,x} = \frac{u_{n+1} - u_n}{x_{n+1} - x_n} \quad u_{,\underline{x}} = \frac{u_n - u_{n-1}}{x_n - x_{n-1}} \quad u_{,x\bar{x}} = 2 \frac{u_{,x} - u_{,\underline{x}}}{x_{n+1} - x_{n-1}} \quad (5)$$

etc. In the continuous limit we have $h_+(x_k, t) \rightarrow 0$, $h_-(x_k, t) \rightarrow 0$, $x_{n+k} \rightarrow x$, $u_k \rightarrow u(x)$ and the discrete derivatives go to the continuous ones.

A solution of the system (1), (2) will have the form $x_n = \Phi(n, c_1, \dots, c_k)$, $u_n = f(x_n, c_1, \dots, c_k)$ where c_1, \dots, c_k are constants needed to satisfy initial conditions and the functions Φ and f are such that (1) and (2) become identities.

As a clarifying example of equations (1) and (2), let us consider a three-point purely difference scheme, namely

$$E = \frac{u_{n+1} - 2u_n + u_{n-1}}{(x_{n+1} - x_n)^2} - u_n = 0 \tag{6}$$

$$\Omega = x_{n+1} - 2x_n + x_{n-1} = 0. \tag{7}$$

The equation $\Omega = 0$ determining the lattice has constant coefficients and its solution is $x_n = hn + x_0$, where $h = h_+ = h_-$ and x_0 are constants. The equation $E = 0$ on this lattice also has constant coefficients (since we have $x_{n+1} - x_n = h$) and its general solution is

$$u(x_n) = c_1 K_+^{x_n} + c_2 K_-^{x_n} \quad K_{\pm} = \left(\frac{2 + h^2 \pm h\sqrt{4 + h^2}}{2} \right)^{1/h}. \tag{8}$$

In the continuous limit we obtain $E = 0 \rightarrow u'' - u = 0$, $\Omega = 0 \rightarrow 0 = 0$, $u(x) = c_1 e^x + c_2 e^{-x}$. Equation (7) happens to determine a regular (equally spaced) lattice. Below we shall see examples of other lattices.

2.2. Symmetries of differential-difference schemes

Let us consider a one-parameter group of local point transformations of the form

$$\tilde{x} = \Xi_{\lambda}(x, t, u) \quad \tilde{t} = \Gamma_{\lambda}(t) \quad \tilde{u}(\tilde{x}, \tilde{t}) = \Phi_{\lambda}(x, t, u). \tag{9}$$

We shall require that they leave the system of equations (1), (2) invariant on the solution set of this system. Since we are interested in continuous transformations (of discrete systems), we use an infinitesimal approach and write the transformations up to order λ as

$$\tilde{x} = x + \lambda \xi(x, t, u(x, t)) \tag{10}$$

$$\tilde{t} = t + \lambda \tau(t) \tag{11}$$

$$\tilde{u}(\tilde{x}, \tilde{t}) = u(x, t) + \lambda \phi(x, t, u(x, t)) \quad |\lambda| \ll 1. \tag{12}$$

This assumption is quite restrictive. Not only do we consider only point transformations, but we require that both t and \tilde{t} are continuous. No dependence, explicit or implicit, on the discretely sampled variable x is allowed. Indeed, once the lattice equation is solved, we obtain a discrete set of points $\{x_n\}$ and this would introduce discrete values $\tilde{t} = \tilde{t}_n$, which we do not allow. Moreover, the x -dependence of t , if allowed, remains unspecified, since the considered equations involve only time derivatives. This would lead to wrong results, i.e. infinite-dimensional transformation groups that do not take solutions into solutions.

We must now prolong the action of the transformation (10) to the prolonged space. This space includes the derivatives $u_t(x, t)$, $u_{tt}(x, t)$, the shifted points $x_{\pm} = x_{n\pm 1}, \dots$ and the function at shifted points $u_{\pm} = u(x_{\pm}, t), \dots$

It is convenient to express the invariance condition for the system (1), (2) in terms of a formalism involving vector fields and their prolongations. The vector field itself has the form

$$\hat{X} = \xi(x, t, u) \partial_x + \tau(t) \partial_t + \phi(x, t, u) \partial_u \tag{13}$$

with ξ , τ and ϕ the same as in equations (10)–(12). The prolongation of the vector field (13) acting on the system (1), (2) is

$$\text{pr}^{(M+N)} \hat{X} = \hat{X} + \sum_{k=n-M}^{n+N} \xi(x_k, t, u_k) \partial_{x_k} + \sum_{k=n-M}^{n+N} \phi^{(k)} \partial_{u_k} + \phi^t \partial_{u_t} + \phi^{tt} \partial_{u_{tt}} \tag{14}$$

with

$$\phi^{(k)} = \phi(x_k, t, u_k) \tag{15}$$

$$\phi^t = D_t \phi - (D_t \xi) u_x - (D_t \tau) u_t \tag{16}$$

$$\phi^{tt} = D_t \phi^t - (D_t \xi) u_{xt} - (D_t \tau) u_{tt}. \tag{17}$$

Thus the prolongation coefficients ϕ^t, ϕ^{tt} are the same as for differential equations; the coefficients $\phi^{(k)}$ are as in [10–27].

The requirement that the system (1), (2) be invariant under the considered one-parameter group translates into the requirement

$$\text{pr}\hat{X} E|_{E=0, \Omega=0} = 0 \quad \text{pr}\hat{X} \Omega|_{E=0, \Omega=0} = 0. \tag{18}$$

In equations (18), once the equations (1), (2) are taken into account, all involved variables are to be considered as independent. Equations (18) are thus the determining equations for the infinitesimal coefficients ξ, τ and ϕ .

For purely difference equations (u_t and u_{tt} absent from (1)) the procedure is the following.

- (1) Extract u_{n+N} and x_{n+N} (or u_{n-M} and x_{n-M}) from equations (1) and (2) and substitute into equation (18). This provides us with two functional equations for ξ, τ and ϕ .
- (2) Assuming an analytical dependence of ξ, τ and ϕ on their own variables, we convert these two equations into differential equations by differentiating them with respect to appropriately chosen variables u_{n+k} and x_{n+k} . Use the fact that the coefficients ξ, τ and ϕ depend on x and u evaluated at one point only to simplify the equations. Differentiate sufficiently many times to obtain differential equations that we can integrate.
- (3) Solve the differential equations, substitute back into the two original functional equations and solve them.

For differential-difference equations, we solve for the highest derivative (in our case u_{tt}) and for either x_{n+N} or u_{n+N} (or x_{n-M} or u_{n-M}) and substitute into equation (18). In this case, the determining equation will be a polynomial expression in the derivatives of u with respect to t (in our case u_t only) and all their coefficients must vanish. For the remaining terms, which depend on shifted variables, we proceed as in the case of purely difference equations.

3. Examples of symmetries of difference equations

We shall give several examples of the calculation of symmetries acting on difference schemes. They will involve either three or four points on a lattice. Equations (1) and (2) simplify to

$$E(x, x_-, x_+, x_{++}, u, u_-, u_+, u_{++}) = 0 \tag{19}$$

$$\Omega(x, x_-, x_+, x_{++}, u, u_-, u_+, u_{++}) = 0 \tag{20}$$

for a four-point scheme. A three-point scheme is obtained if E and Ω are independent of x_{++} and u_{++} . Here $x = x_n$ is the reference point and $x_- = x_{n-1}, x_+ = x_{n+1}$ and $x_{++} = x_{n+2}$ and similarly for u .

The prolongation (14) of the vector field simplifies to

$$\begin{aligned} \text{pr}\hat{X} = & \xi(x, u) \partial_x + \phi(x, u) \partial_u + \xi(x_-, u_-) \partial_{x_-} + \xi(x_+, u_+) \partial_{x_+} + \phi(x_-, u_-) \partial_{u_-} \\ & + \xi(x_{++}, u_{++}) \partial_{x_{++}} + \phi(x_+, u_+) \partial_{u_+} + \phi(x_{++}, u_{++}) \partial_{u_{++}} \end{aligned} \tag{21}$$

(for three-point schemes we drop the x_{++} and u_{++} terms).

A symmetry classification of three-point schemes was provided in a recent paper [35]. Here we solve a different problem. The equations and lattices are given and we determine their symmetries.

3.1. Polynomial nonlinearity on a uniform lattice

Let us consider the nonlinear ordinary differential equation

$$u_{xx} - u^N = 0 \quad N \neq 0, 1. \tag{22}$$

A straightforward calculation shows that for $N \neq -3$ equation (22) is invariant under a two-dimensional Lie group, the Lie algebra of which is spanned by

$$\hat{P} = \partial_x \quad \hat{D} = (N - 1)x\partial_x - 2u\partial_u. \tag{23}$$

For $N = -3$ the symmetry algebra is $sl(2, \mathbb{R})$ with a basis

$$\hat{P} = \partial_x \quad \hat{D} = 2x\partial_x + u\partial_u \quad \hat{C} = x^2\partial_x + xu\partial_u. \tag{24}$$

A natural way to discretize equation (22) is to use a uniform lattice and put

$$E = \frac{u_+ - 2u + u_-}{(x_+ - x_-)^2} - u^N = 0 \tag{25}$$

$$\Omega = x_+ - 2x + x_- = 0. \tag{26}$$

Let us now apply the symmetry algorithm (18). The condition $\text{pr}X\Omega = 0$ for $E = 0, \Omega = 0$ implies

$$\xi(2x - x_-, (x - x_-)^2u^N + 2u - u_-) - 2\xi(x, u) + \xi(x_-, u_-) = 0. \tag{27}$$

Differentiating first by ∂_{u_-} , then by ∂_u , we obtain

$$-\xi_{u_+}(2x - x_-, (x - x_-)^2u^N + 2u - u_-) + \xi_{u_-}(x_-, u_-) = 0 \tag{28}$$

$$[N(x - x_-)^2u^{N-1} + 2]\xi_{u_+,u_+}(2x - x_-, (x - x_-)^2u^N + 2u - u_-) = 0. \tag{29}$$

Equation (29) implies that ξ is linear in u

$$\xi(x, u) = a(x)u + b(x). \tag{30}$$

Equation (28) reduces to $a(x_+) = a(x)$, i.e. a is a constant. Substituting these results into equation (27) we obtain

$$a[u_+ - 2u + u_-] + b(x_+) - 2b(x) + b(x_-) = 0. \tag{31}$$

This implies $a = 0$ and

$$b(x_+) - 2b(x) + b(x_-) = 0. \tag{32}$$

Differentiating successively with respect to x and x_- we find $b_{x_+,x_+}(x_+) = 0$, i.e.

$$b(x) = b_1x + b_0. \tag{33}$$

Thus, the invariance of equation (26) implies $\xi = b_1x + b_0$ with b_1, b_0 constants. The function $\phi(x, u)$ is restricted by the requirement $\text{pr}XE = 0$ for $E = 0, \Omega = 0$. This invariance condition is given by

$$\begin{aligned} &\phi(2x - x_-, (x - x_-)^2u^N + 2u - u_-) - 2\phi(x, u) + \phi(x_-, u_-) \\ &- (x - x_-)^2[N\phi(x, u)u^{N-1} + 2b_1u^N] = 0. \end{aligned} \tag{34}$$

We successively differentiate this equation with respect to u_- and u and we obtain

$$-\phi_{u_+}(x_+, u_+) + \phi_{u_-}(x_-, u_-) = 0 \tag{35}$$

$$\phi_{u_+,u_+}(x_+, u_+) = 0. \tag{36}$$

These two equations require that $\phi = \phi_1u + \phi_0(x)$ with ϕ_1 a constant. Substituting back into equation (34) we obtain the remaining determining equation

$$\begin{aligned} &\phi_0(2x - x_-) - 2\phi_0(x) + \phi_0(x_-) - (x - x_-)^2[(N - 1)\phi_1 + 2b_1]u^N \\ &- N(x - x_-)^2\phi_0u^{N-1} = 0. \end{aligned} \tag{37}$$

Since we have $N \neq 0, 1$ equation (37) implies $\phi_0(x) = 0$ and $\phi_1(1 - N) = 2b_1$. Finally, we obtain the symmetry algebra of the difference system (25), (26). It is two dimensional and coincides with the algebra (23) of the differential equation (22), the continuous limit of equation (25).

Notice that the case $N = -3$ is not distinguished from the generic case. As a matter of fact, no difference equation on a uniform lattice can be invariant under the $SL(2, \mathbb{R})$ group corresponding to the algebra (24). A basis for the difference invariants of this algebra in the space $\{x, x_-, x_+, u, u_-, u_+\}$ is

$$\rho_1 = \frac{h_- u_+}{(h_+ + h_-)u} \quad \rho_2 = \frac{h_+ u_-}{(h_+ + h_-)u} \quad \rho_3 = \frac{h_+ h_-}{(h_+ + h_-)u^2} \quad (38)$$

where h_+ and h_- are defined as $h_+ = x_+ - x$, $h_- = x - x_-$, and no function of x , x_+ and x_- alone can be set equal to a constant. An $SL(2, \mathbb{R})$ invariant scheme must be constructed from these invariants. For instance, an invariant scheme approximating equation (22) for $N = -3$ is

$$\frac{h_-(u_+ - u) - h_+(u - u_-)}{h_+ h_- (h_+ + h_-)} = \frac{2h_+ h_-}{(h_+ + h_-)^2} \frac{1}{u^3} \quad h_- u_+ = h_+ u_- \quad (39)$$

3.2. Discrete versions of linear second-order equations

3.2.1. *Discretization of $u_{xx} = u$.* Consider the ordinary differential equation

$$u_{xx} = u. \quad (40)$$

Like every second-order linear ODE, it is invariant under $SL(3, \mathbb{R})$ with the Lie algebra realized in this case by the vector fields

$$\begin{aligned} \hat{X}_1 &= \partial_x & \hat{X}_2 &= u \partial_u & \hat{X}_3 &= e^x \partial_u & \hat{X}_4 &= e^{-x} \partial_u & \hat{X}_5 &= e^{2x} (\partial_x + u \partial_u) \\ \hat{X}_6 &= u e^x (\partial_x + u \partial_u) & \hat{X}_7 &= e^{-2x} (\partial_x - u \partial_u) & \hat{X}_8 &= u e^{-x} (\partial_x - u \partial_u). \end{aligned} \quad (41)$$

A very straightforward discretization of equation (40) on a uniform lattice is

$$\frac{u_+ - 2u + u_-}{(x_+ - x_-)^2} = u \quad (42)$$

$$x_+ - 2x + x_- = 0. \quad (43)$$

Applying the same procedure to the system (42), (43) that was applied to the system (25), (26) (with $N \neq 0, 1$), we again obtain a two-dimensional symmetry algebra

$$\hat{P} = \partial_x \quad \hat{D} = u \partial_u. \quad (44)$$

At first glance the absence of symmetries of the form $\phi(x) \partial_u$, representing the linear superposition principle, seems surprising. However, viewed as a system of two equations, the system (42), (43) is really nonlinear. Equation (43) defines a uniform lattice with an arbitrary step $h = x_+ - x = x - x_-$, where the step h can be scaled by a dilatation of x .

An alternative approach to the system (42), (43) is to first integrate equation (43) once, thus fixing the step on the x -axis. The system (42), (43) is then replaced by the equation

$$\frac{u_+ - 2u + u_-}{h^2} = u \quad (45)$$

where $h = x_+ - x = x - x_-$ is a fixed (non-scalable) constant. The symmetry algorithm described in section 2 and applied in section 3.1 yields a four-dimensional symmetry algebra

$$\hat{P} = \partial_x \quad \hat{D} = u \partial_u \quad \hat{S}_1 = K_+^x \partial_u \quad \hat{S}_2 = K_-^x \partial_u \quad (46)$$

with K_{\pm} as in equation (8). The symmetries \hat{S}_1, \hat{S}_2 represent the linear superposition formula for the linear system (45).

We mention that equation (40) (and any linear ODE) can be discretized in a manner that exactly preserves all of its solutions. To do this we must preserve a subalgebra of the symmetry algebra of the ODE, containing the elements corresponding to the linear superposition formula. In our case these are \hat{X}_3 and \hat{X}_4 of equation (41). Let us consider the subalgebra $\{\hat{X}_1, \dots, \hat{X}_6\}$. Its second-order discrete prolongation allows no invariants. It does, however, allow an invariant manifold, namely

$$I = ue^{-x}(e^{-2x_+} - e^{-2x_-}) + u_+e^{-x_+}(e^{-2x_-} - e^{-2x}) + u_-e^{-x_-}(e^{-2x} - e^{-2x_+}) = 0. \tag{47}$$

The expression

$$S = \frac{e^{-2x} - e^{-2x_-}}{e^{-2x_+} - e^{-2x}} \tag{48}$$

is an invariant on the manifold (47).

Indeed, we have

$$\begin{aligned} (\hat{X}_1 + 3\hat{X}_2)I = 0 & \quad \hat{X}_3I = \hat{X}_4I = \hat{X}_5I = \hat{X}_6I = 0 & \quad \hat{X}_2I = I \\ \hat{X}_iS = 0 & \quad (i = 1, \dots, 5) & \quad \hat{X}_6S = \frac{2I}{(e^{-2x_+} - e^{-2x})^2} \end{aligned} \tag{49}$$

so we have

$$\hat{X}_iI|_{I=0} = 0 \quad \hat{X}_iS|_{I=0} = 0 \quad i = 1, \dots, 6. \tag{50}$$

A uniform lattice, to first order in h and an equation with (40) as its continuous limit, is obtained by putting

$$S = 1 \quad \frac{e^{3x}I}{2h^3} = 0. \tag{51}$$

Equation (51), or $I = 0$, has $u = e^x$ and $u = e^{-x}$ as solutions and the general solution is

$$u = c_1e^x + c_2e^{-x} \tag{52}$$

just as in the continuous case (40).

To check this, let us solve the system $S = 1, I = 0$ directly, with I and S given in equations (47) and (48), respectively. We linearize $S = 1$ by a change of variables and obtain

$$z = e^{-2x} \quad z_+ - 2z + z_- = 0. \tag{53}$$

The solution is

$$z_n = c_3n + c_4 \quad x_n = -\frac{1}{2} \ln(c_3n + c_4) \tag{54}$$

so that the lattice in x is logarithmic (c_3 and c_4 are integration constants). On this lattice equation (47) reduces to

$$2u\sqrt{c_3n + c_4} - u_+\sqrt{c_3(n + 1) + c_4} - u_-\sqrt{c_3(n - 1) + c_4} = 0. \tag{55}$$

To solve this linear equation we put $u(x) = e^x f(x)$ or, on the lattice

$$u(x_n) = \frac{1}{\sqrt{c_3n + c_4}} f(x_n) \tag{56}$$

so that $f(x)$ satisfies

$$f(x_+) - 2f(x) + f(x_-) = 0. \tag{57}$$

We write the general solution of equation (57) as

$$f(x_n) = f \circ x(n) = An + B. \tag{58}$$

By rewriting A and B in terms of the new integration constants c_1 and c_2 , i.e. by putting $A = c_2 c_3$ and $B = c_2 c_4 + c_1$, we obtain the general solution of the system (51) as

$$u = \frac{c_1}{\sqrt{c_3 n + c_4}} + c_2 \sqrt{c_3 n + c_4} = c_1 e^x + c_2 e^{-x} \tag{59}$$

in full agreement with equation (52).

3.2.2. *Discrete version of $u_{xx} = 1$.* Let us consider the simplest three-point difference scheme for the ODE $u_{xx} = 1$

$$\frac{u_+ - 2u + u_-}{(x_+ - x_-)^2} = 1 \quad x_+ - 2x + x_- = 0. \tag{60}$$

Applying the prolonged vector field to these equations and eliminating x_+ and u_+ , we obtain two equations

$$\xi(2x - x_-, (x - x_-)^2 + 2u - u_-) - 2\xi(x, u) + \xi(x_-, u_-) = 0 \tag{61}$$

$$\begin{aligned} \phi(2x - x_-, (x - x_-)^2 + 2u - u_-) - 2\phi(x, u) + \phi(x_-, u_-) \\ = 2(x - x_-)[\xi(2x - x_-, (x - x_-)^2 + 2u - u_-) - \xi(x, u)]. \end{aligned} \tag{62}$$

We first concentrate on equation (61). Taking the second derivative with respect to u and u_- we find that ξ is linear in u . Substituting back into (61) and differentiating with respect to x and x_- we find

$$\xi(x, u) = \alpha \left(u - \frac{x^2}{2} \right) + \beta_1 x + \beta_0 \tag{63}$$

where α , β_1 and β_0 are constants. Substituting ξ into equation (62) and solving for ϕ in a similar manner, we obtain

$$\phi(x, u) = \alpha \left(xu - \frac{x^3}{2} \right) + c \left(u - \frac{x^2}{2} \right) + \beta_1 x^2 + \beta_2 x + \beta_3. \tag{64}$$

Finally, a basis for the symmetry algebra of the system (60) is

$$\begin{aligned} \hat{X}_1 = \partial_x \quad \hat{X}_2 = \partial_u \quad \hat{X}_3 = x\partial_u \quad \hat{X}_4 = x\partial_x + x^2\partial_u \\ \hat{X}_5 = \left(u - \frac{x^2}{2} \right) \partial_u \quad \hat{X}_6 = \left(u - \frac{x^2}{2} \right) \partial_x + \left(u - \frac{x^2}{2} \right) x\partial_u. \end{aligned} \tag{65}$$

It is easy to check that this Lie algebra is isomorphic to the general affine Lie algebra $\mathfrak{gaff}(2, \mathbb{R})$. This is the symmetry algebra of the scheme [35]

$$w_+ - 2w + w_- = 0 \quad t_+ - 2t + t_- = 0. \tag{66}$$

Indeed the system (60) is transformed into (66) by putting

$$u = w + \frac{t^2}{2} \quad x = t. \tag{67}$$

3.3. *Discrete versions of the equation $u_{xxx} = 0$*

The symmetry algebra of the ODE $u_{xxx} = 0$ is seven dimensional. A basis for this algebra is

$$\begin{aligned} \hat{X}_1 = \partial_x \quad \hat{X}_2 = \partial_u \quad \hat{X}_3 = x\partial_x \quad \hat{X}_4 = u\partial_u \quad \hat{X}_5 = x\partial_u \\ \hat{X}_6 = x^2\partial_u \quad \hat{X}_7 = x^2\partial_x + 2xu\partial_u. \end{aligned} \tag{68}$$

The generators \hat{X}_2 , \hat{X}_5 and \hat{X}_6 correspond to the linear superposition principle. We can add $u = c_2 x^2 + c_1 x + c_0$ to any solution and indeed, this itself is the general solution.

Let us now consider discretizations of this ODE.

3.3.1. *Discretization on a uniform lattice.* We consider the system

$$E = u_{++} - 3u_+ + 3u - u_- = 0 \tag{69}$$

$$\Omega_1 = x_+ - 2x + x_- = 0. \tag{70}$$

The lattice is uniform, since the general solution of (70) is $x_n = hn + x_0$ with h and x_0 constants. Equation (70) must be shifted once to the right to obtain x_{++} .

The prolonged vector fields have the form (21). We apply the same method as in section 3.2 to obtain the symmetry algebra of the system (69), (70). The result is a six-dimensional Lie algebra generated by $\{\hat{X}_1, \hat{X}_2, \hat{X}_3, \hat{X}_4, \hat{X}_5, \hat{X}_6\}$ of equation (68). The system hence has exactly the same solutions as the ODE $u_{xxx} = 0$; however, the lattice is not invariant under the projective transformations generated by \hat{X}_7 .

3.3.2. *Discretization on a four-point lattice.* We take equation (69) on the lattice

$$\Omega_2 = x_{++} - 3x_+ + 3x - x_- = 0. \tag{71}$$

The lattice given by equation (71) is not uniform but satisfies $x_n = L_2n^2 + L_1n + L_0$, where L_i are constants. We assume $L_2 \neq 0$, otherwise the lattice is the same as for $\Omega_1 = 0$.

The symmetry algebra in this case is given by

$$\{\hat{X}_1, \hat{X}_2, \hat{X}_3, \hat{X}_4, \hat{X}_5, \hat{Y} = u\partial_x\} \tag{72}$$

with $\hat{X}_1, \dots, \hat{X}_5$ as in equation (68). Thus \hat{X}_6 of (68) is absent. This reflects the fact that $u = x^2$ is not an exact solution on the lattice $\Omega_2 = 0$. Indeed, if we take $L_2 = 1$ and $L_1 = L_0 = 0$ in equation (68) we have $u = n^4$, which would solve a fourth-order equation, not, however, equation (69).

3.3.3. *Discretization preserving the entire symmetry group.* The third prolongation of the algebra (68) acts on an eight-dimensional space with coordinates $(x, x_+, x_{++}, x_-, u, u_+, u_{++}, u_-)$. If the seven prolonged fields are linearly independent, they will allow only one invariant. This invariant can be calculated directly. It lies entirely in the subspace $\{x, x_+, x_{++}, x_-\}$ and is given by the anharmonic ratio of four points, namely

$$\frac{(x_{++} - x)(x_+ - x_-)}{(x - x_-)(x_{++} - x_+)} = K. \tag{73}$$

This is the invariant of the projective action of $sl(2, \mathbb{R})$ on the real line \mathbb{R} , given by the ∂_x part of the subalgebra $\{\hat{X}_1, \hat{X}_3, \hat{X}_7\}$ of the algebra (68). Equation (73) provides us with a lattice. The invariant equation is obtained by requiring that the third prolongation of $(\hat{X}_1, \dots, \hat{X}_7)$ be linearly connected on some manifold. This manifold is given by the condition

$$I = -(u_+ - u)(x_{++} - x)(x - x_-)(x_{++} - x_-) + (u_{++} - u)(x_+ - x)(x - x_-)(x_+ - x_-) + (u - u_-)(x_+ - x)(x_{++} - x)(x_{++} - x_+) = 0. \tag{74}$$

It is easy to check that I is indeed invariant, i.e.

$$\text{pr}^{(3)}\hat{X}_i I|_{I=0} = 0 \quad i = 1, \dots, 7. \tag{75}$$

Finally, a difference scheme, invariant under the group generated by the algebra (68), having $u_{xxx} = 0$ as a continuous limit, is given by

$$u_{,x\bar{x}\bar{x}} = \frac{6I}{(x_{++} - x_-)(x_{++} - x)(x_{++} - x_+)(x_+ - x_-)(x_+ - x)(x - x_-)} = 0 \tag{76}$$

and equation (71).

We define discrete derivatives as

$$\begin{aligned} u_{,x} &= \frac{u_+ - u_-}{x_+ - x_-} & u_{,x\bar{x}} &= \frac{u_{++} - u_+}{x_{++} - x_+} & u_{,x\bar{x}} &= \frac{u - u_-}{x - x_-} \\ u_{,x\bar{x}} &= 2 \frac{u_{\bar{x}} - u_x}{x_{++} - x_-} & u_{,x\bar{x}} &= 2 \frac{u_x - u_{\bar{x}}}{x_+ - x_-} & u_{,x\bar{x}\bar{x}} &= 3 \frac{u_{x\bar{x}} - u_{x\bar{x}}}{x_{++} - x_-}. \end{aligned} \quad (77)$$

Any four solutions of a Riccati equation satisfy equation (73) and we use this fact to solve this equation. Indeed, consider, for example, the Riccati equation

$$\dot{x} = Ax^2 + Bx + C \quad B^2 - 4AC > 0 \quad (78)$$

where A , B and C are real constants and $A \neq 0$. The general solution of equation (78) is

$$x = \frac{x_1 + x_2 \omega e^{A(x_1 - x_2)t}}{1 - \omega e^{A(x_1 - x_2)t}} \quad x_{1,2} = \frac{-B \pm \sqrt{B^2 - 4AC}}{2A}. \quad (79)$$

Let us take $\omega = n$, $x_1 = \alpha$, $x_2 = \beta$ and $e^{A(x_1 - x_2)t} = \gamma$. A solution of equation (79) is

$$x \equiv x(n) = \frac{\alpha n + \beta}{\gamma n + \delta} \quad \alpha, \beta, \gamma, \delta = \text{const} \quad \alpha\delta - \beta\gamma = 1. \quad (80)$$

Substituting into equation (73) we find $K = 4$. The value $K = 4$ is also required to obtain the correct continuous limit. Indeed, putting $x_+ - x_- = \epsilon\sigma_1$, $x - x_- = \epsilon\sigma_2$, $x_{++} - x_+ = \epsilon\sigma_3$, $\sigma_i \in \mathbb{R}$ and $\epsilon \rightarrow 0$ we have

$$\frac{\epsilon^2(\sigma_1 + \sigma_3)(\sigma_1 + \sigma_2)}{\epsilon^2\sigma_2\sigma_3} = K \quad (81)$$

and for $\sigma_1 = \sigma_2 = \sigma_3$ we have $K = 4$ and also $u_{,x} \rightarrow u'$, $u_{,x\bar{x}} \rightarrow u''$, $u_{,x\bar{x}} \rightarrow u''$, $u_{,x\bar{x}} \rightarrow u''$ and $u_{,x\bar{x}\bar{x}} \rightarrow u'''$, where the primes denote (continuous) derivatives.

Plots of $x(n)$ for lattices (70), (71) and (80) are shown in figures 1, 2 and 3, respectively. The expression (80) is singular for $\gamma = \delta/n$, so such values of γ are to be avoided.

4. Examples for differential-difference equations

In this section we shall need the complete formalism of section 2, in particular the vector field prolongation (14)–(17).

4.1. Symmetries of the discrete Volterra equation

The discrete Volterra equation [17] on a uniform lattice is represented by the two equations

$$E \equiv u_t + u \frac{u_+ - u_-}{x_+ - x_-} = 0 \quad (82)$$

$$\Omega \equiv x_+ - 2x + x_- = 0 \quad (83)$$

where t is a continuous variable, $u = u(x, t)$ and $u_t = \partial u / \partial t$. The Volterra equation is integrable [17] but we make no use of that here.

The invariance condition for the lattice (83) is

$$\xi(2x - x_-, t, u_+) - 2\xi(x, t, u) + \xi(x_-, t, u_-) = 0. \quad (84)$$

In contrast to the cases in section 3, the values u_+ , u and u_- in equation (84) are independent, since the equation $E = 0$ involves u_t (in addition to u_+ , u and u_-). Differentiating equation (84) with respect to, for example, u we obtain $\xi_u = 0$. Differentiating with respect to x_- and then x , we obtain $\xi_{x_+, x_+}(x_+, t) = 0$. The function $\xi(x, t, u)$ hence reduces to

$$\xi = a(t)x + b(t) \quad (85)$$




Figure 1. Variable x as a function of n for the lattice (70), $x_n = hn + x_0$ ($h = 1, x_0 = 5$).

with $a(t)$ and $b(t)$ so far arbitrary functions of t .

Invariance of equation (82) implies

$$\phi^t + \phi \frac{u_+ - u_-}{x_+ - x_-} + \frac{u}{x_+ - x} (\phi^{(+)} - \phi^{(-)}) - \frac{u(u_+ - u_-)}{(x_+ - x_-)^2} (\xi^{(+)} - \xi^{(-)})|_{E=\Omega=0} = 0. \tag{86}$$

The coefficients in the prolongation satisfy

$$\phi^t = \phi_t + (\phi_u - \tau_t)u_t - \xi_t u_x - \xi_u u_t u_x - \tau_u u_t^2 \tag{87}$$

$$\phi^{(\pm)} = \phi(x_{\pm}, t, u(x_{\pm}, t)). \tag{88}$$

We substitute (85), (87) and (88) into equation (86) and eliminate $u_t(x, t)$ and x_+ using equations (82) and (83). The only term involving u_x is in ϕ^t . Its coefficient ξ_t must vanish and we find $\dot{a} = \dot{b} = 0$ in the expression (85).

The remaining determining equation is

$$\left\{ \phi_t + [\phi - u(\phi_u - \tau_t - au)] \frac{u_+ - u_-}{x_+ - x_-} + \frac{u}{x_+ - x_-} [\phi(x_+, t, u(x_+, t)) - \phi(x_-, t, u(x_-, t))] \right\}_{x_+ = 2x - x_-} = 0. \tag{89}$$

We differentiate twice with respect to u_+ and obtain $\phi_{u_+ u_+} = 0$, so that we have $\phi(x, t, u) = \phi_1(x, t)u + \phi_0(x, t)$. Substituting back into equation (89) we obtain the final result, namely

$$\xi = ax + b \quad \tau = c_1 t + c_2 \quad \phi = (a - c_1)u. \tag{90}$$




Figure 2. Variable x as a function of n for the lattice (71), $x_n = L_2 n^2 + L_1 n + L_0$ ($L_2 = 1/\sqrt{10}$, $L_1 = -\pi$, $L_0 = 1$).

Thus, the difference scheme (82), (83), which is the usual Volterra equation, is invariant under a four-dimensional group of Lie point transformations. The symmetry algebra is spanned by

$$\hat{P}_0 = \partial_t \quad \hat{P}_1 = \partial_x \quad \hat{D}_0 = t\partial_t - u\partial_u \quad \hat{D}_1 = x\partial_x + u\partial_u \quad (91)$$

(two translations and two dilatations).

The continuous limit of the system (82), (83) is the Euler equation in 1 + 1 dimensions

$$u_t + uu_x = 0. \quad (92)$$

Its symmetry group is infinite dimensional and can be obtained by standard techniques [3–8] (though we have not found it given explicitly in the literature). Its symmetry algebra is spanned by

$$\begin{aligned} \hat{X}(\xi) &= \xi(z, u)\partial_x & \hat{T}(\tau) &= \tau(z, t, u)(\partial_t + u\partial_x) \\ \hat{F}(\phi) &= \phi(z, u)(t\partial_x + \partial_u) & z &= x - ut \end{aligned} \quad (93)$$

where ξ , τ and ϕ are arbitrary functions of their arguments.

The Volterra equation (82) is certainly not a ‘symmetry preserving’ discretization of the Euler equation (92) on a uniform lattice. It only preserves the four-dimensional subalgebra (91) of the infinite-dimensional symmetry algebra (93). Let us mention here that equation (82) is well known to be a bad numerical scheme for equation (92).




Figure 3. Variable x as a function of n for the lattice (80), $x_n = (\alpha n + \beta)(\gamma n + \delta)^{-1}$ ($\alpha = \sqrt{2}$, $\beta = -\sqrt{3}$, $\gamma = 3$, $\delta = -\sqrt{3}\pi$).

4.2. A general nearest-neighbour interaction equation

Let us consider the difference scheme

$$E \equiv u_{tt} - F(t, x_+, x, x_-, u_+, u, u_-) = 0 \tag{94}$$

$$\Omega \equiv x_+ - 2x + x_- = 0 \tag{95}$$

where F is an arbitrary smooth function satisfying

$$(F_{u_+}, F_{u_-}) \neq (0, 0). \tag{96}$$

A symmetry analysis of a similar class of equations was recently performed for a fixed (non-transformable) regular lattice [12]. More specifically, the assumption was $x_n = n, n \in \mathbb{Z}$.

The prolongation formula for the vector field (13) is (14)–(17). Applying it to equation (95) we obtain that ξ has the form (85), just as for the Volterra equation. Applying the prolongation to equation (94) we obtain

$$\begin{aligned} \phi^{tt} - \tau F_t - (ax + b)F_x - (ax_+ + b)F_{x_+} - (ax_- + b)F_{x_-} - \phi F_u \\ - \phi^{(+)}F_{u_+} - \phi^{(-)}F_{u_-} |_{E=\Omega=0} = 0. \end{aligned} \tag{97}$$

We substitute the expression for ϕ^{tt} , $\phi^{(+)}$ and $\phi^{(-)}$ and set the coefficients of $u_t^3, u_t^2, u_t^2 u_x, u_t u_{xt}, u_{xt}$ and u_t equal to zero, after eliminating u_{tt} and x_+ , using equations (94) and (95). The result is that for any interaction F satisfying condition (96), we have

$$\tau = \tau(t) \quad \xi = ax + b \quad \phi = \left[\frac{\dot{t}}{2} + \alpha(x) \right] u + B(x, t). \tag{98}$$

The as yet unspecified functions $\tau(t)$, $\alpha(x)$ and $B(x, t)$ and constants a and b satisfy a remaining determining equation, namely

$$\begin{aligned} & \left\{ \frac{1}{2} \tau_{ttt} u + B_{tt} - \left(\frac{3}{2} \tau_t - \alpha \right) F + \tau F_t - (ax + b) F_x - (ax_+ + b) F_{x_+} \right. \\ & \quad - (ax_- + b) F_{x_-} - \left[\left(\frac{1}{2} \tau_t + \alpha(x) \right) u + B \right] F_u - \left[\left(\frac{1}{2} \tau_t + \alpha(x_+) \right) u_+ \right. \\ & \quad \left. \left. + B(x_+, t) \right] F_{u_+} - \left[\left(\frac{1}{2} \tau_t + \alpha(x_-) \right) u_- + B(x_-, t) \right] F_{u_-} \right\}_{x_+ = 2x - x_-} = 0. \end{aligned} \quad (99)$$

The results (98) and (99) agree with those of [12], but are more general. The reason for the increase in generality is that here the lattice is not fixed *a priori* and hence the vector field (13) contains a term proportional to ∂_x .

To proceed further, we restrict the interaction F to have a specific form.

4.3. Equation with $F = (x_+ - x)^6 (u_+ - 2u + u_-)^{-3}$

Let us consider a special case of the system (94), (95), namely

$$u_{tt} = \frac{(x_+ - x)^6}{(u_+ - 2u + u_-)^3} \quad (100)$$

$$x_+ - 2x + x_- = 0. \quad (101)$$

We substitute F of equation (100) into the determining equation (99) and clear the denominator. The dependence on u , u_+ and u_- is explicit and we obtain

$$\begin{aligned} \tau_{ttt} = 0 \quad B_{tt} = 0 \quad B(x_+, t) - 2B(x, t) + B(x_-, t) = 0 \\ \alpha(x)(x_+ - x) + 6(ax + b) - 6(ax_+ + b) + 3\alpha(x_+)(x_+ - x) = 0. \end{aligned} \quad (102)$$

Analysing the system (102) in the usual manner, we obtain a nine-dimensional Lie algebra with basis

$$\begin{aligned} \hat{P}_0 = \partial_t \quad \hat{P}_1 = \partial_x \quad \hat{D}_1 = 2t\partial_t + u\partial_u \quad \hat{D}_2 = 2x\partial_x + 3u\partial_u \\ \hat{C} = t^2\partial_t + tu\partial_u \quad \hat{W}_1 = \partial_u \quad \hat{W}_2 = t\partial_u \quad \hat{W}_3 = x\partial_u \quad \hat{W}_4 = tx\partial_u. \end{aligned} \quad (103)$$

A related system was studied earlier [12], namely

$$\ddot{u}_n(t) = [(\gamma_n - \gamma_{n-1})u_{n+1} + (\gamma_{n+1} - \gamma_{n-1})u_n + (\gamma_{n-1} - \gamma_n)u_{n+1}]^{-3} \quad (104)$$

where γ_n is any function of n , satisfying $\gamma_{n+1} \neq \gamma_n$. If we take $\gamma_n = n$ in equation (104) and $x = n$ in (100), (101) the two systems coincide. The symmetry algebra found in [12] is the subalgebra $\{\hat{P}_0, \hat{D}_1, \hat{C}, \hat{W}_1, \hat{W}_2, \hat{W}_3, \hat{W}_4\}$ of the algebra (103). The elements \hat{P}_1 and \hat{D}_2 are absent, since the lattice is fixed. Shifts $n' = n + N$ are allowed, but are not infinitesimal.

The system (100), (101) has a continuous limit

$$u_{tt} = \frac{1}{u_{xx}^3}. \quad (105)$$

The symmetry algebra of equation (105) coincides with (103), i.e. the system (100), (101) is a symmetry preserving discretization of equation (105). We emphasize that equation (100) was obtained as part of a classification of difference equations [12], not in any connection with the PDE (105).

4.4. Equation without a continuous limit

Let us now consider another special case of the system (94), (95), namely

$$u_{tt} = \frac{1}{(u_+ - 2u + u_-)^3} \quad x_+ - 2x + x_- = 0. \quad (106)$$

Substituting for F into equation (99) and proceeding as in section 4.3, we again obtain a nine-dimensional symmetry algebra. It differs from that given in equation (103) only in that D_2 is replaced by $\hat{D}_2 = x\partial x$. For $h = x_+ - x$ satisfying $h \rightarrow 0$ we find u_{tt} finite, but $(u_+ - 2u + u_-)^{-3} \rightarrow \infty$, so the limit $h \rightarrow 0$ does not exist.

5. Conclusions

The main questions to be addressed in a program aiming at using Lie group theory to solve difference equations are the following. (i) How does one define the symmetries? (ii) How does one calculate the symmetries? (iii) What does one do with the symmetries?

In this paper we define the symmetries as in equation (9), that is we consider only Lie point transformations that act simultaneously in a difference equation (1) and lattice equation (2). The fact that the lattice also transforms is in the spirit of Dorodnitsyn's approach to discretizing differential equations. In most symmetry studies of difference equations [9–26] the lattice is fixed and nontransformable, for example, given by the equation $x = n$, $n \in \mathbb{Z}$. For nontransforming lattices we need to go beyond point symmetries to catch transformations of interest [17].

Once the class of symmetries that we wish to consider is defined, the matter of calculating them becomes purely technical. We proposed an algorithm for calculating symmetries in section 2 (see equations (13)–(18)) and applied it in sections 3 and 4. Symmetry algorithms for fixed lattices were presented elsewhere [10–14].

Equations (100) and (104) provide good examples of different approaches. The symmetry algebra (103) of the system (100), (101) happens to coincide with the symmetry algebra of the continuous limit (105). The symmetry algebra of the related equation (104) was calculated elsewhere [12]. It is a seven-dimensional subalgebra of the algebra (103), obtained by dropping \hat{P}_1 and \hat{D}_2 . It was obtained by the 'intrinsic method' [11]. The symmetry algebra of equation (104) can also be obtained from that of the system (100), (101) by taking a specific solution $x = n$ of equation (101) and reducing the algebra (103) to the one that preserves this solution.

As far as applications of symmetries are concerned, they are the same for differential equations and difference ones, in particular, symmetry reduction.

First, consider translationally invariant solutions, i.e. solutions invariant under the subgroup generated by $\hat{X} = \hat{P}_0 - v\hat{P}_1$ with v constant and \hat{P}_0, \hat{P}_1 as in equation (103). We find that the solution, the differential-difference equations (DΔE) (100), (101) and the PDE (106) reduce to

$$u(x, t) = G(\eta) \quad \eta = x + vt \quad (107)$$

$$v^2 G_{\eta\eta} [G(\eta + h) - 2G(\eta) + G(\eta - h)]^3 = h^6 \quad (108)$$

$$v^2 G_{\eta\eta}^4 = 1 \quad (109)$$

respectively. Surprisingly, the difference equation (108) and the ODE (109) have exactly the same solution for all values of the spacing h , namely

$$G = \pm \frac{1}{2\sqrt{v}} \eta^2 + A\eta + B \quad v \neq 0 \quad (110)$$

where A and B are integration constants. Thus, the system (100), (101) is not only a symmetry preserving discretization. It also preserves translationally invariant solutions.

As a second example, consider solutions invariant under dilations generated by \hat{D}_1 of equation (103). The reduction formula, reduced DΔE and reduced PDE are

$$u(x, t) = t^{1/2} G(x) \quad (111)$$

$$G(x)[G(x+h) - 2G(x) + G(x-h)]^3 = -4h^6 \quad (112)$$

and

$$GG_{xx}^3 = -4 \quad (113)$$

respectively. A particular solution of equation (113) is $G(x) = 4(-3)^{-3/4}(x - x_0)^{3/2}$. This is not an exact solution of equation (112), but the solutions of (112) and (113) coincide to order h^2 , rather than just h .

As a final example of symmetry reduction, consider the subgroup corresponding to $\hat{D}_2 - 3\hat{D}_1$ of equation (103). The reduction formulae are

$$u(x, t) = G(\eta) \quad \eta = x^3 t \quad (114)$$

$$G_{\eta\eta} = \frac{(\eta_+^{1/3} - \eta_-^{1/3})^6}{\eta^2[G(\eta_+) - 2G(\eta) + G(\eta_-)]^3} \quad \eta_+^{1/3} - 2\eta^{1/3} + \eta_-^{1/3} = 0 \quad (115)$$

$$27\eta^3 G_{\eta\eta}[3\eta G_{\eta\eta} + 2G_\eta]^3 = 1. \quad (116)$$

While we are not able to solve the ODE (116), or the difference scheme (115), we see that in both cases we obtain a reduction of the number of independent variables. We mention that this last reduction would not be obtained on a fixed lattice.

Let us sum up the situation with this particular approach to symmetries of difference equations.

- (1) Lie point symmetries acting simultaneously on given equations and lattices can be calculated using the reasonably simple algorithm presented in this paper.
- (2) Symmetries can be used to perform symmetry reduction for $D\Delta E$.

Work is in progress on other applications of symmetries of discrete equations, in particular solving ordinary difference equations.

Acknowledgments

The authors thank V Dorodnitsyn, R Kozlov and S Lafortune for stimulating discussions. The research of PW is partially supported by grants from NSERC of Canada and FCAR du Québec. The research reported here is also partly supported by the NATO grant CRG960717 and a Cultural Agreement Università Roma Tre–Université de Montréal. ST and PW thank the Università Roma Tre for hospitality; DL similarly thanks the Centre de Recherches Mathématiques, Université de Montréal.

References

- [1] Lie S 1888 Klassifikation und integration von gewöhnlichen differentialgleichungen zwischen x, y die eine gruppe von transformationen gestatten *Math. Ann.* **32** 213
- [2] Lie S 1888, 1890, 1893 *Theorie der Transformationsgruppen* (Leipzig: Teubner)
- [3] Olver P J 1993 *Applications of Lie Groups to Differential Equations* (New York: Springer)
- [4] Ibragimov N H 1985 *Transformation Groups Applied to Mathematical Physics* (Boston: Reidel)
- [5] Ovsiannikov L V 1982 *Group Analysis of Differential Equations* (New York: Academic)
- [6] Bluman G W and Kumei S 1989 *Symmetries and Differential Equations* (Berlin: Springer)
- [7] Gaeta G 1994 *Nonlinear Symmetries and Nonlinear Equations* (Dordrecht: Kluwer)
- [8] Winternitz P 1993 Group theory and exact solutions of nonlinear partial differential equations *Integrable Systems, Quantum Groups and Quantum Field Theories* (Dordrecht: Kluwer) pp 429–95
- [9] Maeda S 1980 Canonical structure and symmetries for discrete systems *Math. Japan* **25** 405
- [10] Levi D and Winternitz P 1991 Continuous symmetries of discrete equations *Phys. Lett. A* **152** 335
- [11] Levi D and Winternitz P 1993 Symmetries and conditional symmetries of differential-difference equations *J. Math. Phys.* **34** 3713

- [12] Levi D and Winternitz P 1996 Symmetries of discrete dynamical systems *J. Math. Phys.* **37** 5551
- [13] Levi D, Vinet L and Winternitz P 1997 Lie group formalism for difference equations *J. Phys. A: Math. Gen.* **30** 663
- [14] Hernández Heredero R, Levi D and Winternitz P 1999 Symmetries of the discrete Burgers equation *J. Phys. A: Math. Gen.* **32** 2685
- [15] Gomez-Ullate D, Lafortune S and Winternitz P 1999 Symmetries of discrete dynamical systems involving two species *J. Math. Phys.* **40** 2782
- [16] Lafortune S, Martina L and Winternitz P 2000 Point symmetries of generalized Toda field theories *J. Phys. A: Math. Gen.* **33** 2419
- [17] Hernández Heredero R, Levi D, Rodriguez M A and Winternitz P 2000 Lie algebra contractions and symmetries of the Toda hierarchy *J. Phys. A: Math. Gen.* **33** 5025
- [18] Levi D and Yamilov R 1997 Conditions for the existence of higher symmetries of evolutionary equations on the lattice *J. Math. Phys.* **38** 6648
- [19] Levi D and Yamilov R 2000 Non-point integrable symmetries for equations on the lattice *J. Phys. A: Math. Gen.* **33** 4809
- [20] Levi D and Rodriguez M A 1992 Symmetry group of partial differential equations and of differential-difference equations: the Toda lattice vs the Korteweg–de Vries equations *J. Phys. A: Math. Gen.* **25** 975
- [21] Levi D and Yamilov R 1999 Dilatation symmetries and equations on the lattice *J. Phys. A: Math. Gen.* **32** 8317
- [22] Levi D and Rodriguez M A 1999 Lie symmetries for integrable equations on the lattice *J. Phys. A: Math. Gen.* **32** 8303
- [23] Floreanini R, Negro J, Nieto L M and Vinet L 1996 Symmetries of the heat equation on a lattice *Lett. Math. Phys.* **36** 351
- [24] Floreanini R and Vinet L 1995 Lie symmetries of finite-difference equations *J. Math. Phys.* **36** 7024
- [25] Quispel G R W, Capel H W and Sahadevan R 1992 Continuous symmetries of difference equations; the Kac–van Moerbeke equation and Painleve reduction *Phys. Lett. A* **170** 379
- [26] Quispel G R W and Sahadevan R 1993 Lie symmetries and integration of difference equations *Phys. Lett. A* **184** 64
- [27] Dorodnitsyn V A 1989 Transformation groups in a space of difference variables *VINITI Acad. Sci. USSR, Itogi Nauki Tech.* **34** 149–90 (Engl. Transl. 1991 *J. Sov. Math.* **55** 1490)
- [28] Ames W F, Anderson R L, Dorodnitsyn V A, Ferapontov E V, Gazizov R K, Ibragimov N H and Svirshchevskii S R 1994 *CRC Handbook of Lie Group Analysis of Differential Equations (Symmetries, Exact Solutions and Conservation Laws Vol 1)* ed N Ibragimov (Boca Raton, FL: Chemical Rubber Company)
- [29] Dorodnitsyn V A 1994 Finite-difference models entirely inheriting continuous symmetry of original differential equations *Int. J. Mod. Phys. C* **5** 723
- [30] Dorodnitsyn V 1996 Continuous symmetries of finite-difference evolution equations and grids *Symmetries and Integrability of Difference Equations (CRM Proceedings and Lecture Notes vol 9)* ed D Levi, L Vinet and P Winternitz (Providence, RI: American Mathematical Society) pp 103–12
see also Dorodnitsyn V 1994 Invariant discrete model for the Korteweg–de Vries equation *Preprint CRM-2187*, Montreal
- [31] Bakirova M, Dorodnitsyn V and Kozlov R 1997 Invariant difference schemes for heat transfer equations with a source *J. Phys. A: Math. Gen.* **30** 8139
See also Dorodnitsyn V and Kozlov R 1997 The complete set of symmetry preserving discrete versions of a heat transfer equation with a source NO.4/1997 *NTNU, Numerics Preprint Trondheim*
- [32] Dorodnitsyn V 1993 Finite-difference models entirely inheriting symmetry of original differential equations *Modern Group Analysis: Advanced Analytical and Computational Methods in Mathematical Physics* (Dordrecht: Kluwer) p 191
- [33] Dorodnitsyn V A 1993 Finite-difference analog of the Noether theorem *Dokl. Akad. Nauk* **328** 678 (in Russian)
Dorodnitsyn V 1998 Noether-type theorems for difference equation *Preprint IHES/M/98/27 Bures-sur-Yvette*
- [34] Dorodnitsyn V and Winternitz P 2000 Lie point symmetry preserving discretizations for variable coefficient Korteweg–de Vries equations *Nonlinear Dynamics* **22** 49
Dorodnitsyn V and Winternitz P 1999 *Preprint CRM-2607* Universite de Montreal
- [35] Dorodnitsyn V, Kozlov R and Winternitz P 2000 Lie group classification of second order ordinary difference equations *J. Math. Phys.* **41** 480
- [36] Levi D, Vinet L and Winternitz P (ed) 1996 *Symmetries and Integrability of Difference Equations (CRM Proceedings and Lecture Notes vol 9)* (Providence, RI: American Mathematical Society)
- [37] Clarkson P A and Nijhoff F W (ed) 1999 *Symmetries and Integrability of Difference Equations* (Cambridge: Cambridge University Press)