

Quantum Calogero-Moser models: integrability for all root systems

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2000 J. Phys. A: Math. Gen. 33 9033

(<http://iopscience.iop.org/0305-4470/33/49/303>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.124

The article was downloaded on 02/06/2010 at 08:45

Please note that [terms and conditions apply](#).

Quantum Calogero–Moser models: integrability for all root systems

S P Khastgir, A J Pocklington and R Sasaki

Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan

Received 5 June 2000, in final form 3 October 2000

Abstract. The issues related to the integrability of quantum Calogero–Moser models based on any root systems are addressed. For the models with degenerate potentials, i.e. the rational with/without the harmonic confining force, the hyperbolic and the trigonometric, we demonstrate the following for all the root systems. (i) Construction of a complete set of quantum conserved quantities in terms of a total sum of the Lax matrix L , i.e. $\sum_{\mu, \nu \in \mathcal{R}} (L^n)_{\mu\nu}$, in which \mathcal{R} is a set of \mathbb{R}^r vectors invariant under the action of the Coxeter group. They form a single Coxeter orbit. (ii) Proof of Liouville integrability. (iii) Triangularity of the quantum Hamiltonian and the entire discrete spectrum. Generalized Jack polynomials are defined for all root systems as unique eigenfunctions of the Hamiltonian. (iv) Equivalence of the Lax operator and the Dunkl operator. (v) Algebraic construction of all excited states in terms of creation operators. These are mainly generalizations of the results known for the models based on the A series, i.e. $su(N)$ -type, root systems.

1. Introduction

Calogero–Moser models are one-dimensional dynamical systems with long-range interactions having the remarkable property that they are integrable at both classical and quantum levels. In fact the integrability or more precisely the triangularity of the quantum Hamiltonian was first discovered by Calogero [1] for the model with inverse square potential plus a confining harmonic force and by Sutherland [2] for the particles on a circle with the inverse square potential. Later classical integrability of the models in terms of Lax pairs was proved by Moser [3]. Olshanetsky and Perelomov [4] showed that these models were based on A_r root systems and generalizations of the models based on other root systems including the non-crystallographic ones were introduced [5].

In this paper we discuss quantum Calogero–Moser models with degenerate potentials, that is the rational with/without harmonic force, the hyperbolic and the trigonometric potentials based on all root systems. We demonstrate, based on previous works on universal Lax pairs for classical [6–8] and quantum models [9], that various results known for the quantum A_r models can be generalized to the models based on any root systems as well. They are the following. (i) Construction of a complete set of quantum conserved quantities in terms of quantum Lax pairs and other methods. (ii) Universal proof of Liouville integrability for the rational, hyperbolic and trigonometric potential models. Namely, the quantum conserved quantities commute among themselves. (iii) Triangularity of the quantum Hamiltonian is demonstrated explicitly for all the models. In other words, the Hamiltonian is shown, in certain bases, to be decomposed into a sum of finite-dimensional triangular matrices. Thus any eigenvalue equation can be solved by finite steps of linear algebraic processes only. This also gives the entire discrete spectrum of the models. As unique eigenfunctions of the Hamiltonian,

generalization of Jack polynomials and multivariable Laguerre (Hermite) polynomials are defined for all root systems. (iv) Equivalence of the quantum Lax pair method and that of so-called differential-reflection (Dunkl) operators [10] is demonstrated. (v) For rational models with harmonic confining force, an algebraic construction of all excited states in terms of creation (annihilation) operators is achieved.

For the A_r models, the Lax pairs, conserved quantities and their involution have been discussed by many authors with varying degrees of completeness and rigour (see, e.g., [5, 11–20]). Point (iv) was shown by Wadati and collaborators [18] and point (v) was initiated by Perelomov [12] and developed by Brink and collaborators [17] and Wadati and collaborators [18]. A rather different approach by Heckman and Opdam [21, 22] to Calogero–Moser models with degenerate potentials based on any root systems should also be mentioned in this connection.

For the general background and the motivations of this series of papers and the physical applications of the Calogero–Moser models with various potentials to lower-dimensional physics, ranging from solid state to particle physics and supersymmetric Yang–Mills theory, we refer to our previous papers [6, 7] and references therein.

This paper is organized as follows. In section 2, the quantum Calogero–Moser Hamiltonian with degenerate potentials is introduced as a factorized form (2.5). Connection with root systems and the Coxeter invariance is emphasized. Some rudimentary facts of the root systems and reflections are summarized in appendix A. A universal Coxeter invariant ground state wavefunction together with its energy eigenvalue are presented. In section 3 we show that all the excited states are also Coxeter invariant and that the Hamiltonian is *triangular* in certain bases. Complete sets of quantum conserved quantities are derived from the quantum Lax operator L in section 4. Instead of the trace, the *total sum* of L^n is conserved. That is $\text{Ts}(L^n) = \sum_{\mu, \nu \in \mathcal{R}} (L^n)_{\mu\nu}$, in which \mathcal{R} is a set of \mathbb{R}^r vectors invariant under the action of the Coxeter group. They form a single Coxeter orbit. The details of the complete set for each root system are given in appendix B. In section 5 the creation and annihilation operators for the rational models with harmonic force are derived. In section 6, the equivalence of the Lax pair operator formalism and the so-called differential-reflection (Dunkl) operators is demonstrated and the quantum conserved quantities are expressed in terms of the latter. In section 7 an algebraic construction of excited states in terms of the differential-reflection (Dunkl) operators for rational models with harmonic force is presented. The complete sets of explicit eigenfunctions for the rank two models are derived in terms of separation of variables based on the Coxeter invariant variables. Section 8 gives a universal proof of the Liouville integrability for models with rational (without the confining force), hyperbolic and trigonometric potentials. For rational models with harmonic force, the involution is demonstrated for those based on classical root systems. A simple use of the quantum Lax pair with a spectral parameter is mentioned. The final section is for a summary and comments.

2. Quantum Calogero–Moser models

In this section we briefly introduce the *quantum* Calogero–Moser models along with appropriate notation and background for the main body of this paper. A Calogero–Moser model is a Hamiltonian system associated with a root system Δ of rank r , which is a set of vectors in \mathbb{R}^r with its standard inner product. A brief review of the properties of the root systems and the associated reflections will be found in appendix A.

2.1. Factorized Hamiltonian

The dynamical variables of the Calogero–Moser model are the coordinates $\{q_j\}$ and their canonically conjugate momenta $\{p_j\}$, with the canonical commutation relations

$$[q_j, p_k] = i\delta_{jk} \quad [q_j, q_k] = [p_j, p_k] = 0 \quad j, k = 1, \dots, r. \tag{2.1}$$

These will be denoted by vectors in \mathbb{R}^r

$$q = (q_1, \dots, q_r) \quad p = (p_1, \dots, p_r). \tag{2.2}$$

The momentum operator p_j acts as

$$p_j = -i \frac{\partial}{\partial q_j} \quad j = 1, \dots, r.$$

As for the interactions we consider only the degenerate potentials, that is the rational (with/without harmonic force), hyperbolic and trigonometric potentials:

$$V(\rho \cdot q) = \begin{cases} 1/(\rho \cdot q)^2 & \text{type I} \\ a^2/\sinh^2 a(\rho \cdot q) & \text{type II} \\ a^2/\sin^2 a(\rho \cdot q) & \text{type III} \end{cases} \quad \rho \in \Delta \tag{2.3}$$

in which a is an arbitrary real positive constant, determining the period of the trigonometric potentials. They imply integrability for all of the Calogero–Moser models based on the crystallographic root systems. Those models based on the non-crystallographic root systems, the dihedral group $I_2(m)$, H_3 and H_4 , are integrable only for the rational potential. The rational potential models are also integrable if a confining harmonic potential

$$\frac{1}{2}\omega^2 q^2 \quad \omega > 0 \quad \text{type V} \tag{2.4}$$

is added to the Hamiltonian. Since we will discuss the universal properties and solutions applicable to all the interaction types as well as those for specific interaction potentials, let us adopt the conventional nomenclature for them. We call the models with rational, hyperbolic, trigonometric and rational with harmonic force the type I, II, III and V models, respectively. (Type IV models have elliptic potentials, which we will not discuss in this paper.)

The Hamiltonian for the *quantum* Calogero–Moser model can be written in a ‘factorized form’:

$$\mathcal{H} = \frac{1}{2} \sum_{j=1}^r \left(p_j - i \frac{\partial W}{\partial q_j} \right) \left(p_j + i \frac{\partial W}{\partial q_j} \right) \tag{2.5}$$

$$= \frac{1}{2} \sum_{j=1}^r \left(p_j^2 + \left(\frac{\partial W}{\partial q_j} \right)^2 \right) + \frac{1}{2} \sum_{j=1}^r \frac{\partial^2 W}{\partial q_j^2} \tag{2.6}$$

$$= \frac{1}{2} p^2 + \frac{1}{2} \sum_{\rho \in \Delta_+} g_{|\rho|} (g_{|\rho|} - 1) |\rho|^2 V(\rho \cdot q) + \left(\frac{\omega^2}{2} q^2 \right) - \mathcal{E}_0. \tag{2.7}$$

It should be noted that the above factorized Hamiltonian (2.7) consists of an operator part $\hat{\mathcal{H}}$, which is the Hamiltonian in the usual definition, and a constant \mathcal{E}_0 , which is the ground state energy to be discussed later:

$$\mathcal{H} = \hat{\mathcal{H}} - \mathcal{E}_0 \tag{2.8}$$

$$\hat{\mathcal{H}} = \frac{1}{2} p^2 + \frac{1}{2} \sum_{\rho \in \Delta_+} g_{|\rho|} (g_{|\rho|} - 1) |\rho|^2 V(\rho \cdot q) + \left(\frac{\omega^2}{2} q^2 \right). \tag{2.9}$$

The real *positive* coupling constants $g_{|\rho|}$ are defined on orbits of the corresponding Coxeter group, i.e. they are identical for roots in the same orbit. That is, for the simple Lie algebra cases

Table 1. Functions appearing in the Lax pair and superpotential.

Potential	Type	$w(u)$	$x(u)$	$y(u)$
Rational	I and V	u	$1/u$	$-1/u^2$
Hyperbolic	II	$\sinh au$	$a \coth au$	$-a^2/\sinh^2 au$
Trigonometric	III	$\sin au$	$a \cot au$	$-a^2/\sin^2 au$

there is one coupling constant $g_{|\rho|} = g$ for all roots in simply laced models and two independent coupling constants, $g_{|\rho|} = g_L$ for long roots and $g_{|\rho|} = g_S$ for short roots, in non-simply laced models. For the $I_2(m)$ models, there is one coupling if m is odd, and two independent ones if m is even. Let us call them g_e for even roots and g_o for odd roots. Throughout this paper we consider the coupling constants at generic values. We parametrize the positive roots of $I_2(m)$ as

$$\rho_j = (\cos((j - 1)\pi/m), \sin((j - 1)\pi/m)) \quad j = 1, \dots, m. \tag{2.10}$$

The H_3 and H_4 models have one coupling constant $g_{|\rho|} = g$, since these root systems are simply laced. It should be noted that the operator part of the Hamiltonian $\hat{\mathcal{H}}$ is strictly positive for $g_{|\rho|} \geq 1$, which we will assume throughout this paper for simplicity of the arguments.

The simplest way to introduce the factorized form is through supersymmetry [9, 23], in which function W is called a superpotential:

$$W(q) = \sum_{\rho \in \Delta_+} g_{|\rho|} \ln |w(\rho \cdot q)| + \left(-\frac{\omega}{2} q^2\right) \quad g_{|\rho|} > 0 \quad \omega > 0. \tag{2.11}$$

The potential $V(u)$ (2.3) and the function $w(u)$ are related by

$$y(u) \equiv \frac{d}{du} x(u) \quad \frac{dw(u)}{du} / w(u) \equiv x(u) \tag{2.12}$$

$$V(u) = -y(u) = x^2(u) + a^2 \times \begin{cases} 0 & \text{rational} \\ -1 & \text{hyperbolic} \\ 1 & \text{trigonometric.} \end{cases} \tag{2.13}$$

Table 1 gives these functions for each potential.

For proofs that the factorized Hamiltonian (2.6) actually gives the quantum Hamiltonian (2.7) for *all the root systems and potentials* see [5, 9, 11]. It is easy to verify that, for any potential $V(u)$, the Hamiltonian is invariant under reflection of the phase space variables in the hyperplane perpendicular to any root

$$\mathcal{H}(s_\alpha(p), s_\alpha(q)) = \mathcal{H}(p, q) \quad \forall \alpha \in \Delta \tag{2.14}$$

with s_α defined by (A.2).

Some remarks are in order. For all of the root systems and for any choice of potential (2.3), the Calogero–Moser model has a hard repulsive potential $\sim 1/(\alpha \cdot q)^2$ near the reflection hyperplane $H_\alpha = \{q \in \mathbb{R}^r, \alpha \cdot q = 0\}$. The strength of the singularity is given by the coupling constant $g_{|\alpha|}(g_{|\alpha|} - 1)$, which is *independent* of the choice of the normalization of the roots. (Thus for rational models with/without harmonic force there is equivalence, $A_2 \cong I_2(3)$, $B_2 \cong I_2(4)$, $G_2 \cong I_2(6)$, $B_r \cong C_r \cong BC_r$.) This determines the form of the ground state wavefunction, as we will see in section 2.2. This repulsive potential is classically insurmountable. Thus the motion is always confined within one Weyl chamber. This feature allows us to constrain the configuration space to the principal Weyl chamber (Π : the set of simple roots, see appendix A)

$$PW = \{q \in \mathbb{R}^r \mid \alpha \cdot q > 0, \alpha \in \Pi\} \tag{2.15}$$

without loss of generality. In the case of the trigonometric potential, the configuration space is further limited due to the periodicity of the potential to

$$PW_T = \{q \in \mathbb{R}^r \mid \alpha \cdot q > 0, \alpha \in \Pi, \alpha_h \cdot q < \pi/a\} \tag{2.16}$$

where α_h is the highest root.

The fact that the classical motions are confined in the Weyl chamber (alcove) $PW(PW_T)$ does not necessarily mean that the corresponding quantum wavefunctions vanish identically outside the region. On the contrary, as we will see soon, the ground state wavefunction (section 2.2) and all the other excited state wavefunctions (section 3) are Coxeter invariant, reflecting the Coxeter invariance of the Hamiltonian (2.14). In the early years of Calogero–Moser models in which those based on the A_r root system were mainly discussed, these Coxeter invariant solutions were considered as totally symmetric states of bosonic systems. We will not, however, adopt this interpretation, for in the models based on the other root systems the reflection is not the same as particle interchange. The quantum theory we are discussing is the so-called first quantized theory. That is, the notions of identical particles and the associated statistics are non-existent.

2.2. Ground state wavefunction and energy

One merit of the factorized Hamiltonian (2.5) is the ease of derivation of the ground state wavefunction and of the Hamiltonian (3.3) derived by the similarity transformation in terms of the ground state wavefunction. Supersymmetric formulation of the Calogero–Moser models [9, 16, 23] provides a natural setting for the introduction of the factorized Hamiltonian. The *universal* ground state wavefunction is

$$\Phi_0(q) = e^{W(q)} = \prod_{\rho \in \Delta_+} |w(\rho \cdot q)|^{g_{|\rho|}} e^{-\frac{\omega}{2} q^2}. \tag{2.17}$$

The exponential factor $e^{-\frac{\omega}{2} q^2}$ exists only for the rational potential case with the harmonic confining force. It is easy to see that it is an eigenstate of the Hamiltonian (2.5) with *zero* eigenvalue:

$$\mathcal{H}\Phi_0(q) = \frac{1}{2} \sum_{j=1}^r \left(p_j - i \frac{\partial W}{\partial q_j} \right) \left(p_j + i \frac{\partial W}{\partial q_j} \right) \Phi_0(q) = 0 \tag{2.18}$$

since it satisfies

$$\left(p_j + i \frac{\partial W}{\partial q_j} \right) e^{W(q)} = 0 \quad j = 1, \dots, r. \tag{2.19}$$

By using the decomposition of the factorized Hamiltonian into the operator Hamiltonian (2.9) and a constant, we obtain

$$\hat{\mathcal{H}} e^W \equiv \left(\frac{1}{2} p^2 + \frac{1}{2} \sum_{\rho \in \Delta_+} g_{|\rho|} (g_{|\rho|} - 1) |\rho|^2 V(\rho \cdot q) + \left(\frac{\omega^2}{2} q^2 \right) \right) e^W = \mathcal{E}_0 e^W. \tag{2.20}$$

In other words, the above solution (2.17) provides an eigenstate of the Hamiltonian operator $\hat{\mathcal{H}}$ with energy \mathcal{E}_0 . The fact that it is a ground state (for type I, III and V) can be easily shown within the framework of the supersymmetric model [9] thanks to the positivity of the supersymmetric Hamiltonian. It should be stressed that \mathcal{E}_0 is determined purely algebraically [9], without really applying the operator on the left-hand side of (2.20) to the wavefunction. This type of ground state has been known for some time. It is derived by various methods, see for example [3, 5], and also by using supersymmetric quantum mechanics for the models based on classical root systems [16, 23].

The ground state energy for the rational potential cases is

$$\mathcal{E}_0 = \begin{cases} 0 & \text{type I} \\ \omega \left(\frac{r}{2} + \sum_{\rho \in \Delta_+} g_{|\rho|} \right) & \text{type V.} \end{cases} \tag{2.21}$$

The same for the hyperbolic and trigonometric potential cases is

$$\mathcal{E}_0 = 2a^2 \varrho^2 \times \begin{cases} -1 & \text{hyperbolic} \\ 1 & \text{trigonometric} \end{cases} \tag{2.22}$$

in which

$$\varrho = \frac{1}{2} \sum_{\rho \in \Delta_+} g_{|\rho|} \rho \tag{2.23}$$

can be considered as a ‘deformed Weyl vector’ [5, 22]. Again these formulae are universal, that is, they apply to all of the Calogero–Moser models based on any root systems. A negative \mathcal{E}_0 for the obviously positive Hamiltonian of the hyperbolic potential model indicates that the interpretation of e^W as an eigenfunction is not correct. This function diverges as $|q| \rightarrow \infty$ for the hyperbolic and the rational potential cases, destroying the hermiticity of the Hamiltonian. Obviously we have

$$\int_{\text{PW (PW}_T)} e^{2W(q)} dq = \begin{cases} \infty : & \text{type I and II} \\ \text{finite:} & \text{type III and V} \end{cases} \tag{2.24}$$

in which PW and PW_T denote that the integration is over the regions defined in (2.15) and (2.16). Naturally, most existing results in quantum Calogero–Moser models are for the models with trigonometric potential and the rational potential with harmonic force, which have normalizable states and discrete spectra.

It should be remarked that the domain of the universal ground state wavefunction Φ_0 could be considered as the entire \mathbb{R}^r space except for the points on the reflection hyperplanes, that is the disjoint union of all the Weyl chambers (alcoves), instead of the initial Weyl chamber/alcove (PW, PW_T) in which the classical motions are restricted due to the singular potential. In fact, Φ_0 and W are characterized as *Coxeter invariant*:

$$\check{s}_\rho \Phi_0 = \Phi_0 \quad \check{s}_\rho W = W \quad \forall \rho \in \Delta \tag{2.25}$$

in which \check{s}_ρ is the representation of the reflection in the function space. For an arbitrary function f of q , its action is defined by

$$(\check{s}_\rho f)(q) = f(s_\rho(q)). \tag{2.26}$$

This definition can be generalized to the entire Coxeter group G_Δ : for an arbitrary element g of G_Δ , \check{g} is defined by

$$(\check{g}f)(q) = f(g^{-1}(q)) \quad \forall g \in G_\Delta. \tag{2.27}$$

In the rest of this paper we discuss mainly the type III and V models which have normalizable states and discrete spectra.

3. Coxeter invariant excited states, triangularity and spectrum

In this section we show that all the excited state wavefunctions are Coxeter invariant, too. In other words, the Fock space consists of Coxeter invariant functions only. With the knowledge of the ground state wavefunction e^W , the other states of the Calogero–Moser models can be

easily obtained as eigenfunctions of a differential operator $\tilde{\mathcal{H}}$ obtained from \mathcal{H} by a similarity transformation:

$$\tilde{\mathcal{H}} = e^{-W} \mathcal{H} e^W = e^{-W} \left(\frac{1}{2} p^2 + \frac{1}{2} \sum_{\rho \in \Delta_+} g_{|\rho|} (g_{|\rho|} - 1) |\rho|^2 V(\rho \cdot q) + \left(\frac{\omega^2}{2} q^2 \right) - \mathcal{E}_0 \right) e^W \quad (3.1)$$

$$\tilde{\mathcal{H}} \Psi_\lambda = \lambda \Psi_\lambda \iff \mathcal{H} \Psi_\lambda e^W = \lambda \Psi_\lambda e^W. \quad (3.2)$$

Thanks to the factorized form of the Hamiltonian \mathcal{H} (2.5), (2.6), the transformed Hamiltonian $\tilde{\mathcal{H}}$ takes a simple form:

$$\tilde{\mathcal{H}} = -\frac{1}{2} \sum_{j=1}^r \left(\frac{\partial^2}{\partial q_j^2} + 2 \frac{\partial W}{\partial q_j} \frac{\partial}{\partial q_j} \right). \quad (3.3)$$

The Coxeter invariance of W implies those of \mathcal{H} and $\tilde{\mathcal{H}}$:

$$\check{s}_\rho \mathcal{H} \check{s}_\rho = \mathcal{H} \quad \check{s}_\rho \tilde{\mathcal{H}} \check{s}_\rho = \tilde{\mathcal{H}} \quad \forall \rho \in \Delta. \quad (3.4)$$

For type V and III models we introduce proper bases of Fock space consisting of Coxeter invariant functions and show that the above Hamiltonian $\tilde{\mathcal{H}}$ (3.3) is *triangular* in these bases. This establishes the integrability of the type V and III models universally[†] and also gives the entire spectrum of the Hamiltonian (see (3.12), (3.13) and (3.44)).

3.1. Rational potential with harmonic force

First, let us determine the structure of the set of eigenfunctions of the transformed Hamiltonian $\tilde{\mathcal{H}}$, for the type V models:

$$\tilde{\mathcal{H}} = \omega q \cdot \frac{\partial}{\partial q} - \frac{1}{2} \sum_{j=1}^r \frac{\partial^2}{\partial q_j^2} - \sum_{\rho \in \Delta_+} \frac{g_{|\rho|}}{\rho \cdot q} \rho \cdot \frac{\partial}{\partial q}. \quad (3.5)$$

Obviously a constant and $\omega q^2 - \mathcal{E}_0/\omega$ are its eigenfunctions with eigenvalue 0 and 2ω , respectively. Let us suppose that a polynomial $P(q)$ is an eigenfunction of $\tilde{\mathcal{H}}$:

$$\tilde{\mathcal{H}} P(q) = \lambda P(q). \quad (3.6)$$

Due to the Coxeter invariance of $\tilde{\mathcal{H}}$ (3.4), we know that $\check{s}_\rho P$ together with the difference

$$Q = (1 - \check{s}_\rho) P$$

are also eigenfunctions with the same eigenvalue:

$$\tilde{\mathcal{H}} Q(q) = \lambda Q(q) \quad (3.7)$$

if the latter is not identically zero. Since Q is a polynomial which is odd under reflection \check{s}_ρ

$$\check{s}_\rho Q(q) = -Q(q)$$

it can be factorized as

$$Q(q) = (\rho \cdot q)^{2n+1} \tilde{Q}(q) \quad \tilde{Q}|_{\rho \cdot q=0} \neq 0 \quad (3.8)$$

with a non-negative integer n and a polynomial \tilde{Q} . By substituting (3.8) into (3.7) and using the explicit form of $\tilde{\mathcal{H}}$ near the reflection hyperplane $\rho \cdot q = 0$, we obtain

$$-(\rho \cdot q)^{2n-1} (2n+1)(n+g_{|\rho|}) \rho^2 \tilde{Q} + \mathcal{O}[(\rho \cdot q)^{2n}] = \lambda (\rho \cdot q)^{2n+1} \tilde{Q} \quad (3.9)$$

[†] Triangularity of the A_r type V and III Hamiltonians was noted in the original papers of Calogero [1] and Sutherland [2]. That of rank two models in the Coxeter invariant bases was shown in [22, 24].

Table 2. The degrees f_j in which independent Coxeter invariant polynomials exist.

Δ	$f_j = 1 + e_j$	Δ	$f_j = 1 + e_j$
A_r	$2, 3, 4, \dots, r + 1$	E_8	$2, 8, 12, 14, 18, 20, 24, 30$
B_r	$2, 4, 6, \dots, 2r$	F_4	$2, 6, 8, 12$
C_r	$2, 4, 6, \dots, 2r$	G_2	$2, 6$
D_r	$2, 4, \dots, 2r - 2; r$	$I_2(m)$	$2, m$
E_6	$2, 5, 6, 8, 9, 12$	H_3	$2, 6, 10$
E_7	$2, 6, 8, 10, 12, 14, 18$	H_4	$2, 12, 20, 30$

which would imply the vanishing of \tilde{Q} on the reflection hyperplane

$$\tilde{Q}|_{\rho \cdot q=0} = 0$$

an obvious contradiction. Thus we are led to the conclusion that *the eigenfunctions are Coxeter invariant polynomials* and that the Hamiltonian \mathcal{H} (3.5) maps one Coxeter invariant polynomial to another.

An obvious basis in the space of Coxeter invariant polynomials is the homogeneous polynomials of various degrees. This basis has a natural order given by the degree. For a given degree the space of homogeneous Coxeter invariant polynomials is finite dimensional. The explicit form of $\tilde{\mathcal{H}}$ (3.5) shows that it is *lower triangular* in this basis and the diagonal elements are $\omega \times \text{degree}$ as given by the first term. Independent Coxeter invariant polynomials exist at the degrees f_j listed in table 2:

$$f_j = 1 + e_j \quad j = 1, \dots, r \tag{3.10}$$

in which $\{e_j\}$, $j = 1, \dots, r$, are the *exponents* of Δ . Let us denote them by

$$z_1(q), \dots, z_r(q) \quad z_j(\kappa q) = \kappa^{f_j} z_j(q). \tag{3.11}$$

Thus we arrive at: the quantum Calogero–Moser model with the rational potential and the harmonic confining force is algebraically solvable for any (crystallographic and non-crystallographic) root system Δ . The spectrum of the operator Hamiltonian $\hat{\mathcal{H}}$ is

$$\omega N + \mathcal{E}_0 \tag{3.12}$$

with a non-negative integer N , which can be expressed as

$$N = \sum_{j=1}^r n_j f_j \quad n_j \in \mathbb{Z}_+ \tag{3.13}$$

and the degeneracy of the above eigenvalue (3.12) is the number of different solutions of (3.13) for given N . This is a generalization of Calogero’s original argument for the A_r model [1] to the models based on arbitrary root systems. Now let us denote by \vec{N} the set of non-negative integers in (3.13):

$$\vec{N} = (n_1, n_2, \dots, n_r) \tag{3.14}$$

and by $\phi_{\vec{N}}(q)$ the homogeneous Coxeter invariant polynomial determined by \vec{N} and the above basis $\{z_j\}$ (3.11):

$$\phi_{\vec{N}}(q) = \prod_{j=1}^r z_j^{n_j}(q). \tag{3.15}$$

As shown above, there exists a unique eigenstate $\psi_{\vec{N}}(q)$ for each $\phi_{\vec{N}}(q)$:

$$\psi_{\vec{N}}(q) = \phi_{\vec{N}}(q) + \sum_{\vec{N}' < \vec{N}} d_{\vec{N}'} \phi_{\vec{N}'}(q) \quad d_{\vec{N}'}: \text{const} \tag{3.16}$$

$$\tilde{\mathcal{H}} \psi_{\vec{N}}(q) = \omega N \psi_{\vec{N}}(q). \tag{3.17}$$

This satisfies the orthogonality relation

$$(\psi_{\vec{N}}, \phi_{\vec{N}'}) = 0 \quad \vec{N}' < \vec{N} \tag{3.18}$$

with respect to the inner product in PW:

$$(\psi, \varphi) = \int_{\text{PW}} \psi^*(q)\varphi(q) e^{2W(q)} dq. \tag{3.19}$$

These polynomials $\{\psi_{\vec{N}}(q)\}$ are generalizations of the multivariable Laguerre (Hermite) polynomials [14] known for the A_r (B_r , D_r) root systems to arbitrary root systems.

Some remarks are in order.

- (1) There is no Coxeter invariant linear function in q . The quadratic invariant polynomial $q^2 = q \cdot q$ exists in all the root systems. This corresponds to the universal fact that $f_1 = 2$, ($e_1 = 1$) for all the root systems. Moreover, this is related to the fact that a special sub-series of the excited states with $N = 2n_1 = n_1 f_1$, $n_j = 0$, $j \geq 2$, can be expressed universally in terms of Laguerre polynomials in q^2 . This will be discussed at the end of section 5 and in section 7.2.
- (2) The other Coxeter invariants corresponding to the degrees f_2, \dots, f_r could be interpreted as special ‘angular’ variables of a unit sphere S^{r-1} ($q^2 = 1$), with the first Coxeter invariant $\sqrt{q^2}$ being the radial coordinate. These would provide proper variables for describing solutions. Solutions in terms of separation of variables are in general possible only for the simplest cases, that is the rank two models, which will be demonstrated in section 7.3.
- (3) For $\Delta = A_1$, the simplest root system of rank one, the Hamiltonian $\tilde{\mathcal{H}}$ can be rewritten in terms of a Coxeter invariant variable $u = \omega q^2$ as

$$\tilde{\mathcal{H}} = \omega q \frac{d}{dq} - \frac{1}{2} \frac{d^2}{dq^2} - \frac{g}{q} \frac{d}{dq} = -2\omega \left\{ u \frac{d^2}{du^2} + \left(g + \frac{1}{2} - u \right) \frac{d}{du} \right\}. \tag{3.20}$$

The Laguerre polynomial satisfying the differential equation

$$\left\{ u \frac{d^2}{du^2} + \left(g + \frac{1}{2} - u \right) \frac{d}{du} + n \right\} L_n^{(g-\frac{1}{2})}(u) = 0 \tag{3.21}$$

provides an eigenfunction with eigenvalue $2\omega n$, which corresponds to the eigenvalue $2\omega n + \mathcal{E}_0$ of $\hat{\mathcal{H}}$. This is a well known result.

3.2. Trigonometric potential

Here we consider those root systems associated with Lie algebras. In order to determine the excited states of the type III models, we have to consider the periodicity. The superpotential W and the Hamiltonian \mathcal{H} are invariant under the following translation:

$$W(q') = W(q) \quad \mathcal{H}(p, q') = \mathcal{H}(p, q) \quad q' = q + l^\vee \pi/a \tag{3.22}$$

in which l^\vee is an element of the dual weight lattice, that is

$$l^\vee = \sum_{j=1}^r l_j \frac{2}{\alpha_j^2} \lambda_j \quad l_j \in \mathbb{Z} \quad \alpha_j \in \Pi \quad \alpha_j^\vee \cdot \lambda_k = \delta_{jk}. \tag{3.23}$$

As is well known in quantum mechanics with periodic potentials, the wavefunctions diagonalizing the translation operators are expressed as

$$e^{2ia\mu \cdot q} \left(\sum_{\alpha \in L(\Delta)} b_\alpha e^{2ia\alpha \cdot q} \right) e^W \quad b_\alpha: \text{const} \quad L(\Delta): \text{root lattice} \tag{3.24}$$

in which a vector $\mu \in \mathbb{R}^r$ is as yet unspecified. In other words, up to the overall phase factor $e^{2ia\mu \cdot q}$, this is a Fourier expansion in terms of the simple roots.

Let $P_T(q)$ be a polynomial in $e^{\pm 2ia\alpha_j \cdot q}$, $\alpha_j \in \Pi$ and suppose that a function $\phi(q)$

$$\phi(q) = e^{2ia\mu \cdot q} P_T(q) \quad \mu \in \mathbb{R}^r \tag{3.25}$$

is an eigenfunction of $\tilde{\mathcal{H}}$:

$$\tilde{\mathcal{H}}\phi(q) = \lambda\phi(q) \tag{3.26}$$

in which the explicit form of $\tilde{\mathcal{H}}$ is given by

$$\tilde{\mathcal{H}} = -\frac{1}{2} \sum_{j=1}^r \frac{\partial^2}{\partial q_j^2} - a \sum_{\rho \in \Delta_+} g_{|\rho|} \cot(a\rho \cdot q) \rho \cdot \frac{\partial}{\partial q}. \tag{3.27}$$

As above, due to the Coxeter invariance of $\tilde{\mathcal{H}}$ (3.4), we know that $\check{s}_\rho\phi$ together with the difference

$$\varphi = (1 - \check{s}_\rho)\phi$$

are also eigenfunctions with the same eigenvalue:

$$\tilde{\mathcal{H}}\varphi(q) = \lambda\varphi(q). \tag{3.28}$$

Since φ is odd under reflection \check{s}_ρ

$$\check{s}_\rho\varphi(q) = -\varphi(q)$$

it can be expressed as

$$\varphi(q) = (\rho \cdot q)^{2n+1} \tilde{\varphi}(q) + \mathcal{O}[(\rho \cdot q)^{2n+3}] \quad \tilde{\varphi}|_{\rho \cdot q=0} \neq 0 \tag{3.29}$$

in a neighbourhood of the reflection hyperplane $\rho \cdot q = 0$. In this neighbourhood, the singularity structure of $\tilde{\mathcal{H}}$ for the trigonometric potential is the same as that of $\tilde{\mathcal{H}}$ for the rational potential discussed in the previous section. Thus we obtain, as before, a contradiction $\tilde{\varphi}|_{\rho \cdot q=0} = 0$. In other words, the eigenfunction ϕ (3.25) must be *Coxeter invariant*. This in turn requires the unspecified vector μ in (3.25) to be an element of the weight lattice

$$\mu \in \Lambda(\Delta). \tag{3.30}$$

Since

$$\check{s}_\rho\phi(q) = e^{2ia s_\rho(\mu) \cdot q} \check{s}_\rho P_T(q) = e^{2ia(\mu \cdot q - \rho^\vee \cdot \mu \rho \cdot q)} \check{s}_\rho P_T(q)$$

the condition

$$\rho^\vee \cdot \mu \in \mathbb{Z} \quad \forall \rho \in \Delta \tag{3.31}$$

is necessary, but not sufficient, for the Coxeter invariance of ϕ . Thus we arrive at (3.30).

Let us introduce a basis for the Coxeter invariant functions of the form (3.25). Let λ be a dominant weight

$$\lambda = \sum_{j=1}^r m_j \lambda_j \quad m_j \in \mathbb{Z}_+ \tag{3.32}$$

and W_λ be the orbit of λ by the action of the Weyl group:

$$W_\lambda = \{\mu \in \Lambda(\Delta) | \mu = g(\lambda) \quad \forall g \in G_\Delta\}. \tag{3.33}$$

We define

$$\phi_\lambda(q) \equiv \sum_{\mu \in W_\lambda} e^{2ia\mu \cdot q} \tag{3.34}$$

which is Coxeter invariant. The set of functions $\{\phi_\lambda\}$ has an order $>$:

$$|\lambda|^2 > |\lambda'|^2 \Rightarrow \phi_\lambda > \phi_{\lambda'}. \tag{3.35}$$

Next we show that $\tilde{\mathcal{H}}$ is *lower triangular* in this basis. By using (3.27) we obtain

$$\tilde{\mathcal{H}}\phi_\lambda = 2a^2\lambda^2\phi_\lambda - 2ia^2 \sum_{\rho \in \Delta_+} \sum_{\mu \in W_\lambda} g_{|\rho|} \cot(a\rho \cdot q)(\rho \cdot \mu)e^{2ia\mu \cdot q}. \tag{3.36}$$

First let us fix one positive root ρ and a weight μ in W_λ such that $\rho \cdot \mu \neq 0$. Then

$$\mu' \equiv s_\rho(\mu) = \mu - (\rho^\vee \cdot \mu)\rho \in W_\lambda \quad \rho \cdot \mu' = -\rho \cdot \mu. \tag{3.37}$$

Without loss of generality we may assume

$$\rho^\vee \cdot \mu = k > 0 \quad k \in \mathbb{Z}. \tag{3.38}$$

The contribution of the pair (μ, μ') in the summation of (3.36) reads

$$|\rho \cdot \mu|e^{2ai\mu \cdot q}(1 - e^{-2aik\rho \cdot q}) \cot(a\rho \cdot q) = i|\rho \cdot \mu| \left(e^{2ai\mu \cdot q} + e^{2ai\mu' \cdot q} + 2 \sum_{j=1}^{k-1} e^{2ai(\mu - j\rho) \cdot q} \right) \tag{3.39}$$

which is the generalization of Sutherland’s fundamental identity equation (15) in [2] to arbitrary root systems. The summation in the expression corresponds to $\phi_{\lambda'}$ with λ' being lower than λ . Thus (3.36) reads

$$\tilde{\mathcal{H}}\phi_\lambda = 2a^2\lambda^2\phi_\lambda + 2a^2 \sum_{\rho \in \Delta_+} \sum_{\mu \in W_\lambda} g_{|\rho|} |\rho \cdot \mu| e^{2ia\mu \cdot q} + \sum_{|\lambda'| < |\lambda|} c_{\lambda'} \phi_{\lambda'} \tag{3.40}$$

in which $\{c_{\lambda'}\}$ are constants. It is easy to see that $(\mu = g(\lambda), \exists g \in G_\Delta)$

$$\sum_{\rho \in \Delta_+} g_{|\rho|} |\rho \cdot \mu| = \sum_{\rho \in \Delta_+} g_{|\rho|} |g(\rho) \cdot \lambda| = \left(\sum_{\rho \in \Delta_+} g_{|\rho|} \rho \right) \cdot \lambda = 2\rho \cdot \lambda \tag{3.41}$$

which is independent of μ . Thus we have demonstrated the triangularity of $\tilde{\mathcal{H}}$:

$$\tilde{\mathcal{H}}\phi_\lambda = 2a^2(\lambda^2 + 2\rho \cdot \lambda)\phi_\lambda + \sum_{|\lambda'| < |\lambda|} c_{\lambda'} \phi_{\lambda'} \tag{3.42}$$

or that of $\hat{\mathcal{H}}$

$$\hat{\mathcal{H}}\phi_\lambda e^W = 2a^2(\lambda + \rho)^2 \phi_\lambda e^W + \sum_{|\lambda'| < |\lambda|} c_{\lambda'} \phi_{\lambda'} e^W \tag{3.43}$$

with the eigenvalue

$$2a^2(\lambda + \rho)^2. \tag{3.44}$$

In other words, for each dominant weight λ there exists an eigenstate of $\tilde{\mathcal{H}}$ with eigenvalue proportional to $\lambda(\lambda + 2\rho)$. Let us denote this eigenfunction by $\psi_\lambda(q)$:

$$\psi_\lambda(q) = \phi_\lambda(q) + \sum_{|\lambda'| < |\lambda|} d_{\lambda'} \phi_{\lambda'}(q) \quad d_{\lambda'}: \text{const} \tag{3.45}$$

$$\tilde{\mathcal{H}}\psi_\lambda(q) = 2a^2\lambda(\lambda + 2\rho)\psi_\lambda(q) \tag{3.46}$$

and call it a *generalized Jack polynomial* [25–28]. It satisfies the orthogonality relation

$$(\psi_\lambda, \phi_{\lambda'}) = 0 \quad |\lambda'| < |\lambda| \tag{3.47}$$

with respect to the inner product in PW_T :

$$(\psi, \varphi) = \int_{PW_T} \psi^*(q)\varphi(q) e^{2W(q)} dq. \tag{3.48}$$

In the A_r model, specifying a dominant weight λ is the same as giving a Young diagram which designates a Jack polynomial. It should be emphasized, however, that $\{\psi_\lambda\}$ are not identical to the Jack polynomials even for the A_r root systems, because of different treatments of the centre of mass coordinates. Detailed properties of these polynomials for various root systems will be published elsewhere.

Thus we arrive at the following.

The quantum Calogero–Moser models with the trigonometric potential are algebraically solvable for any crystallographic root system Δ . The spectrum of the Hamiltonian $\hat{\mathcal{H}}$ is given by (3.44) in which λ is an arbitrary dominant weight. This is a generalization of Sutherland’s original argument [2] to the models based on arbitrary root systems.

Some remarks are in order.

- (1) The weights μ appearing in the lower-order terms $\{\phi_{\lambda'}\}$ are those weights contained in the Lie algebra representation belonging to the highest weight λ .
- (2) As a simple corollary we find that for a minimal weight λ ,

$$\psi_\lambda(q) = \phi_\lambda(q) = \sum_{\mu \in W_\lambda} e^{2ia\mu \cdot q}$$

is an eigenfunction of $\tilde{\mathcal{H}}$. A minimal representation [6] consists of a single Weyl orbit and all of its weights μ satisfy $\rho^\vee \cdot \mu = 0, \pm 1, \forall \rho \in \Delta$.

- (3) If $\lambda = \alpha_h$, the highest root of a simply laced root system, W_λ is the set of roots itself. Then the lower-order terms are constants only. We find that

$$\psi_{\alpha_h}(q) = 2 \sum_{\rho \in \Delta_+} (\cos(2a\rho \cdot q) + g\rho^2/\alpha_h \cdot (\alpha_h + 2\rho))$$

is an eigenstate of $\tilde{\mathcal{H}}$.

- (4) If $\lambda = \alpha_{Sh}$, the highest short root of a non-simply laced root system, W_λ is the set of short roots itself. The lower-order terms are constants, too. Similarly as above, we find that

$$\psi_{\alpha_{Sh}}(q) = 2 \sum_{\rho \in \Delta_{L+}} \cos(2a\rho \cdot q) + 2 \sum_{\rho \in \Delta_{S+}} (\cos(2a\rho \cdot q) + g_S \rho_S^2/\alpha_{Sh} \cdot (\alpha_{Sh} + 2\rho))$$

is an eigenstate of $\tilde{\mathcal{H}}$. Here $\Delta_{L(S)}$ is the set of long (short) roots.

- (5) If $-\lambda \notin W_\lambda$ then there is another set of functions containing the weight $-\lambda$ which belongs to the same eigenvalue.
- (6) The Coxeter invariant trigonometric polynomials specified by the fundamental weights $\{\lambda_j\}$

$$\phi_{\lambda_j}(q) = \sum_{\mu \in W_{\lambda_j}} e^{2ia\mu \cdot q} \quad \lambda_j: \text{fundamental weight} \quad j = 1, \dots, r \tag{3.49}$$

are expected to play the role of the fundamental variables [22, 24].

- (7) Let us consider the well known case $\Delta = A_1$, the simplest root system of rank one. By rewriting the Hamiltonian $\tilde{\mathcal{H}}$ in terms of the Coxeter invariant variable $z = \cos(a\rho q)$, we obtain

$$\hat{\mathcal{H}} = -\frac{1}{2} \frac{d^2}{dq^2} - ag\rho \cot(a\rho q) \frac{d}{dq} = -\frac{1}{2} a^2 |\rho|^2 \left\{ (1 - z^2) \frac{d^2}{dz^2} - (1 + 2g)z \frac{d}{dz} \right\}. \tag{3.50}$$

The Gegenbauer polynomials [5], a special case of Jacobi polynomials $P_n^{(\alpha, \beta)}$, provide eigenfunctions

$$P_n^{(g-\frac{1}{2}, g-\frac{1}{2})}(\cos(a\rho q)) \quad \mathcal{E} = a^2 |\rho|^2 (n + g)^2 / 2 \quad n \in \mathbb{Z}_+. \tag{3.51}$$

The Jacobi polynomial $P_\ell^{(\alpha,\beta)}(z)$ satisfies differential equation

$$\left\{ (1 - z^2) \frac{d^2}{dz^2} + (\beta - \alpha - (2 + \alpha + \beta)z) \frac{d}{dz} + \ell(\ell + \alpha + \beta + 1) \right\} P_\ell^{(\alpha,\beta)}(z) = 0. \quad (3.52)$$

Here we follow the notation of [29]. They form orthogonal polynomials with weight $e^{2W} = |\sin(a\rho q)|^{2g}$ in the interval $q \in [0, \pi/a\rho]$, (2.16). The Gegenbauer polynomials have a definite parity, $(-1)^n$, reflecting the periodicity. The fundamental period is $\pi/a\rho$, the length of the interval itself. The (odd-) even-degree ones corresponding to (half-) odd integer spin representations are (anti-) periodic.

(8) Triangularity of type II models follows from the same algebraic reasoning.

4. Quantum Lax pair and quantum conserved quantities

Historically, Lax pairs for Calogero–Moser models were presented in terms of Lie algebra representations [3, 5], in particular, the vector representation of the A_r models. However, the invariance of Calogero–Moser models is that of the Coxeter group but not that of the associated Lie algebra, which does not exist for the non-crystallographic root systems. Thus the universal and Coxeter covariant Lax pairs are given in terms of the representations of the Coxeter group.

4.1. General case

Here we recapitulate the essence of the quantum Lax pair operators for the Calogero–Moser models with degenerate potentials and without spectral parameter. The case with spectral parameter will be discussed briefly in section 8.3 in connection with the proof of involution of the quantum conserved quantities for the A_r model [15]. The quantum Lax pair in this section applies to all the degenerate potential cases except for the case of the rational potential with the harmonic force, which will be treated separately in section 4.2. For details and a full exposition, see [9]. The Lax operators without spectral parameter are

$$L(p, q) = p \cdot \hat{H} + X(q) \quad X(q) = i \sum_{\rho \in \Delta_+} g_{|\rho|} (\rho \cdot \hat{H}) x(\rho \cdot q) \hat{s}_\rho \quad (4.1)$$

$$M(q) = \frac{i}{2} \sum_{\rho \in \Delta_+} g_{|\rho|} |\rho|^2 y(\rho \cdot q) \hat{s}_\rho - \frac{i}{2} \sum_{\rho \in \Delta_+} g_{|\rho|} |\rho|^2 y(\rho \cdot q) \times I \quad (4.2)$$

in which I is the identity operator and $\{\hat{s}_\alpha, \alpha \in \Delta\}$ are the reflection operators of the root system. In contrast with $\{\check{s}_\alpha\}$ operators (2.26), which act in function space, $\{\hat{s}_\alpha\}$ act on a set of \mathbb{R}^r vectors $\mathcal{R} = \{\mu^{(k)} \in \mathbb{R}^r, k = 1, \dots, d\}$, permuting them under the action of the reflection group. The vectors in \mathcal{R} form a basis for the representation space V of dimension d . The operator M satisfies the relation

$$\sum_{\mu \in \mathcal{R}} M_{\mu\nu} = \sum_{\nu \in \mathcal{R}} M_{\mu\nu} = 0 \quad (4.3)$$

which is essential for deriving quantum conserved quantities. The matrix elements of the operators $\{\hat{s}_\alpha, \alpha \in \Delta\}$ and $\{\hat{H}_j, j = 1, \dots, r\}$ are defined as follows:

$$(\hat{s}_\rho)_{\mu\nu} = \delta_{\mu, s_\rho(\nu)} = \delta_{\nu, s_\rho(\mu)} \quad (\hat{H}_j)_{\mu\nu} = \mu_j \delta_{\mu\nu} \quad \rho \in \Delta \quad \mu, \nu \in \mathcal{R}. \quad (4.4)$$

The form of the function x depends on the chosen potential, and the function y is defined by (2.12), (2.13). Note that these relations are only valid for the degenerate potentials (2.3).

The underlying idea of the Lax operator L , (4.1), is quite simple. As seen from (4.9), L is a ‘square root’ of the Hamiltonian. Thus one part of L contains p , which is not associated

with roots, and another part contains $x(\rho \cdot q)$, a ‘square root’ of the potential $V(\rho \cdot q)$, which being associated with a root ρ is therefore accompanied by the reflection operator \hat{s}_ρ . Another explanation is the factorized Hamiltonian \mathcal{H} (2.5). We obtain, roughly speaking, $L \sim \sqrt{\mathcal{H}} \sim p + i \frac{\partial W}{\partial q} \hat{s}$ and the property of reflection $\hat{s}^2 = 1$ explains the sign change in the first term in (2.5).

It is straightforward to show that the quantum Lax equation

$$\frac{d}{dt}L = i[\mathcal{H}, L] = [L, M] \tag{4.5}$$

is equivalent to the quantum equations of motion derived from the Hamiltonian (2.7). From this it follows that

$$\frac{d}{dt}(L^n)_{\mu\nu} = i[\mathcal{H}, (L^n)_{\mu\nu}] = [L^n, M]_{\mu\nu} = \sum_{\lambda \in \mathcal{R}} ((L^n)_{\mu\lambda} M_{\lambda\nu} - M_{\mu\lambda} (L^n)_{\lambda\nu}) \quad n = 1, \dots \tag{4.6}$$

Thanks to the property of the M operator (4.3)

$$\sum_{\mu \in \mathcal{R}} M_{\mu\nu} = \sum_{\nu \in \mathcal{R}} M_{\mu\nu} = 0$$

we obtain *quantum conserved quantities* as the *total sum* (Ts) of all the matrix elements of L^n †:

$$Q_n = \text{Ts}(L^n) \equiv \sum_{\mu, \nu \in \mathcal{R}} (L^n)_{\mu\nu} \quad [\mathcal{H}, Q_n] = 0 \quad n = 1, \dots \tag{4.7}$$

Independent conserved quantities appear at such a power n that

$$n = 1 + \text{exponent} \tag{4.8}$$

of each root system. These are the degrees at which independent Coxeter invariant polynomials exist. There are r exponents for each root system Δ of rank r . Thus we have r independent conserved quantities in Calogero–Moser models. We list in table 2 these powers for each root system. In particular, the power 2 is universal to all the root systems and the quantum Hamiltonian (2.7) is given by

$$\mathcal{H} = \frac{1}{2C_{\mathcal{R}}} \text{Ts}(L^2) + \text{const} \tag{4.9}$$

where the constant $C_{\mathcal{R}}$ is the quadratic Casimir invariant, which depends on the representation. It is defined by

$$\text{Tr}(\hat{H}_j \hat{H}_k) \equiv \sum_{\mu \in \mathcal{R}} (\hat{H}_j \hat{H}_k)_{\mu\mu} = \sum_{\mu \in \mathcal{R}} \mu_j \mu_k = C_{\mathcal{R}} \delta_{jk}. \tag{4.10}$$

Some remarks are in order.

- (1) The Lax pair is Coxeter covariant:

$$\begin{aligned} L(s_\rho(p), s_\rho(q))_{\mu\nu} &= L(p, q)_{\mu'\nu'} & M(s_\rho(q))_{\mu\nu} &= M(q)_{\mu'\nu'} \\ \mu' &\equiv s_\rho(\mu) & \nu' &\equiv s_\rho(\nu) \end{aligned} \tag{4.11}$$

which ensures the Coxeter invariance of the conserved quantities.

- (2) Lax pairs can be written down in various representations and the quantum conserved quantities Q_n do depend on the representations, in general. If necessary, we denote by $Q_n^{\mathcal{R}}$ the explicit representation dependence.

† This type of conserved quantity is known for A_r models [16, 18].

- (3) The availability of plural representations of the Lax pair and the conserved quantities is essential for the completeness of the set of conserved quantities as polynomials of the momentum operators. For example, let us consider the case of D_r with even r , which has two independent conserved quantities at power r (see table 2). At least two different representations of the Lax pair are necessary in order to represent them in the form of (4.7). Those based on the vector and the (anti-) spinor representations give two independent conserved quantities. For the D_4 case, we obtain

$$Q_4^v = 2 \sum_{j=1}^4 p_j^4 \quad Q_4^s - Q_4^a = 24 \prod_{j=1}^4 p_j \quad (4.12)$$

in which v, s and a stand for the vector, spinor and anti-spinor representations and we have set $g = 0$ for simplicity. If two conserved quantities are independent for zero coupling constants, surely they are so at non-vanishing couplings. Here we have used an explicit parametrization of the D_r root system:

$$D_r \text{ root system: } \Delta = \{\pm e_j \pm e_k, j < k = 1, \dots, r | e_j \in \mathbb{R}^r, e_j \cdot e_k = \delta_{jk}\}. \quad (4.13)$$

- (4) If a representation \mathcal{R} contains a vector μ and its negative $-\mu$ at the same time, then we have $\text{Ts}(L^{\text{odd}}) = 0$. In such a case the corresponding Lie algebra representations are called real. In order to construct the odd-power conserved quantities appearing in A_r for all r , D_r for odd r , E_6 and $I(m)$ for odd m , we need a Lax pair in non-real representations. For A_r all the fundamental representations corresponding to the fundamental weights λ_j , $j = 1, \dots, r$, except for the middle one $\lambda_{(r+1)/2}$ for odd r are non-real. For D_r with odd r , the spinor and anti-spinor representations are non-real. For E_6 the **27** and $\overline{\mathbf{27}}$ are non-real. $I_2(m)$ is the symmetry group of a regular m -sided polygon. The set of m vectors representing the vertices of the regular m -gon to be denoted by R_m (B.3), provides a non-real representation when m is odd.
- (5) In appendix B we list for each root system how the full set of independent conserved quantities is obtained by choosing proper representations of the Lax pair.

4.2. Rational potential with harmonic force

The quantum Lax pair for the type V models needs a separate formulation. The explicit form of the Hamiltonian is

$$\mathcal{H} = \frac{1}{2} p^2 + \frac{1}{2} \omega^2 q^2 + \frac{1}{2} \sum_{\rho \in \Delta_+} g_{|\rho|} (g_{|\rho|} - 1) \frac{|\rho|^2}{(\rho \cdot q)^2} - \mathcal{E}_0. \quad (4.14)$$

The canonical equations of motion are equivalent to the following Lax equations for L^\pm :

$$\frac{d}{dt} L^\pm = i[\mathcal{H}, L^\pm] = [L^\pm, M] \pm i\omega L^\pm \quad (4.15)$$

in which (see section 4 of [7]) M is the same as before (4.2), and L^\pm and Q are defined by

$$L^\pm = L \pm i\omega Q \quad Q = q \cdot \hat{H} \quad (4.16)$$

with L and \hat{H} as earlier (4.1), (4.4). If we define Hermitian operators \mathcal{L}_1 and \mathcal{L}_2 by

$$\mathcal{L}_1 = L^+ L^- \quad \mathcal{L}_2 = L^- L^+ \quad (4.17)$$

they satisfy Lax-like equations

$$\dot{\mathcal{L}}_k = [\mathcal{L}_k, M] \quad k = 1, 2. \quad (4.18)$$

From these we can construct conserved quantities

$$\text{Ts}(\mathcal{L}_j^n) \quad j = 1, 2 \quad n = 1, 2, \dots \tag{4.19}$$

as before. Such quantum conserved quantities have been previously reported for models based on A_r root systems [16, 18]. It should be remarked that $\text{Ts}(\mathcal{L}_2^n)$ is no longer the same as $\text{Ts}(\mathcal{L}_1^n)$ due to quantum corrections. It is elementary to check that the first conserved quantities give the Hamiltonian (4.14)

$$\mathcal{H} \propto \text{Ts}(\mathcal{L}_1) = \text{Ts}(\mathcal{L}_2) + \text{const.} \tag{4.20}$$

This then completes the presentation of the quantum Lax pairs and quantum conserved quantities for all of the quantum Calogero–Moser models with non-elliptic potentials.

5. Algebraic construction of excited states I

In this section we show that all the excited states of the type V Calogero–Moser models can be constructed algebraically. Later in section 7 we show the same results in terms of the ℓ operators to be introduced in section 6. The main result is surprisingly simple and can be stated universally, as follows.

Corresponding to each partition of an integer N which specifies the energy level (3.12) into the sum of the degrees of Coxeter invariant polynomials (3.13), we have an eigenstate of the Hamiltonian \mathcal{H} with eigenvalue $\omega N + \mathcal{E}_0$:

$$\prod_{j=1}^r (B_{f_j}^+)^{n_j} e^W \quad N = \sum_{j=1}^r n_j f_j \quad n_j \in \mathbb{Z}_+ \tag{5.1}$$

in which the integers $\{f_j\}$, $j = 1, \dots, r$, are listed in table 2. They exhaust all the excited states. In other words the above states give the complete basis of the Fock space. The creation operators $B_{f_j}^+$ and the corresponding annihilation operators[†] $B_{f_j}^-$ are defined in terms of the Lax operators L^\pm (4.16) as follows:

$$B_{f_j}^\pm = \text{Ts}(L^\pm)^{f_j} \quad j = 1, \dots, r. \tag{5.2}$$

They are Hermitian conjugate to each other

$$(B_{f_j}^\pm)^\dagger = B_{f_j}^\mp \tag{5.3}$$

with respect to the standard Hermitian inner product of the states defined in PW:

$$(\psi, \varphi) = \int_{\text{PW}} \psi^*(q) \varphi(q) dq. \tag{5.4}$$

We will show later in section 6, (6.15), that the creation (annihilation) operators commute among themselves:

$$[B_k^+, B_l^+] = [B_k^-, B_l^-] = 0 \quad k, l \in \{f_j \mid j = 1, \dots, r\} \tag{5.5}$$

so the state (5.1) does not depend on the order of the creation.

The proof is very simple. By using (4.15) we obtain

$$\frac{d}{dt}(L^\pm)^n = i[\mathcal{H}, (L^\pm)^n] = [(L^\pm)^n, M] \pm in\omega(L^\pm)^n \tag{5.6}$$

from which

$$[\mathcal{H}, B_n^\pm] = \pm n\omega B_n^\pm \tag{5.7}$$

[†] We adopt the notation by Olshanetsky and Perelomov [5, 12].

follows after taking the total sum. This simply says that B_n^\pm creates (annihilates) a state having energy $n\omega$. In other words, we have

$$\hat{\mathcal{H}} \prod_{j=1}^r (B_{f_j}^+)^{n_j} e^W = \left(\mathcal{E}_0 + \omega \sum_{k=1}^r n_k f_k \right) \prod_{j=1}^r (B_{f_j}^+)^{n_j} e^W.$$

Moreover, it is trivial to show that

$$\sum_{\nu \in \mathcal{R}} (L^-)_{\mu\nu} e^W = \left(p \cdot \mu - i\omega q \cdot \mu + i \sum_{\rho \in \Delta_+} \frac{\rho \cdot \mu}{\rho \cdot q} \right) e^W = \mu \cdot \left(p + i \frac{\partial W}{\partial q} \right) e^W = 0 \tag{5.8}$$

which implies that the ground state is annihilated by all the annihilation operators

$$B_{f_j}^- e^W = 0 \quad j = 1, \dots, r. \tag{5.9}$$

Some remarks are in order.

- (1) In most cases the energy levels are highly degenerate. The above basis is neither orthogonal nor normalized.
- (2) The independence of the creation–annihilation operators can also be shown in a way similar to that of the conserved quantities. As with the conserved quantities, plural representations are necessary to define the full set of creation–annihilation operators in some models. This aspect will be discussed in later sections in connection with the ℓ operators.
- (3) Reflecting the universality of the first exponent, $f_1 = 2$, the creation and annihilation operators of the least quanta, 2ω , exist in all the models. They form an $sl(2, \mathbb{R})$ algebra together with the Hamiltonian $\hat{\mathcal{H}}$:

$$[\hat{\mathcal{H}}, b_2^\pm] = \pm 2\omega b_2^\pm \quad [b_2^+, b_2^-] = -\omega^{-1} \hat{\mathcal{H}} \tag{5.10}$$

in which b_2^\pm are normalized forms of B_2^\pm :

$$b_2^\pm = \sum_{\mu, \nu \in \mathcal{R}} (L^\pm)_{\mu\nu}^2 / (4\omega C_{\mathcal{R}}). \tag{5.11}$$

The $sl(2, \mathbb{R})$ algebra was discussed by many authors (see, e.g., [12, 17, 21, 30] and others) in connection with the models based on classical root systems. We will show later in section 7.2 that the states created by B_2^+ (b_2^+) only can be expressed by the Laguerre polynomial:

$$(b_2^+)^n e^W = n! L_n^{(\tilde{\mathcal{E}}_0 - 1)}(\omega q^2) e^W \quad \tilde{\mathcal{E}}_0 \equiv \mathcal{E}_0 / \omega. \tag{5.12}$$

It is trivial to verify that $L_n^{(\tilde{\mathcal{E}}_0 - 1)}(\omega q^2)$ is an eigenfunction of $\tilde{\mathcal{H}}$ (3.5)

$$\tilde{\mathcal{H}} L_n^{(\tilde{\mathcal{E}}_0 - 1)}(\omega q^2) = 2n\omega L_n^{(\tilde{\mathcal{E}}_0 - 1)}(\omega q^2). \tag{5.13}$$

The normalization of the state

$$\| (b_2^+)^n e^W \|^2 = n! \mathcal{N}_0 / \Gamma(n + \tilde{\mathcal{E}}_0) \quad \mathcal{N}_0 \equiv \| e^W \|^2 \Gamma(\tilde{\mathcal{E}}_0) \tag{5.14}$$

is also dictated by the $sl(2, \mathbb{R})$ relations. The Laguerre polynomial wavefunctions appear as ‘radial’ wavefunctions in all cases [31]. This will be shown explicitly for for the rank two models given in section 7.3.

- (4) As is emphasized by Perelomov [12] and Gambardella [30] the $sl(2, \mathbb{R})$ algebra and the corresponding Laguerre wavefunctions are more universal than Calogero–Moser models. They arise when the potentials are homogeneous functions in q of degree -2 with the confining harmonic force.

(5) The operators $\{Q_n\}$ and $\{B_n^\pm\}$ do not form a Lie algebra. They satisfy interesting nonlinear relations, for example,

$$[[B_n^+, b_2^-], b_2^+] = nB_n^+ \quad [[B_n^-, b_2^+], b_2^-] = nB_n^- \tag{5.15}$$

This tells us, for example, that, although B_n^+ and b_2^+ create different units of quanta n and 2 , they are not independent:

$$[B_n^+, b_2^-] \neq 0 \neq [B_n^-, b_2^+].$$

Clarification of the algebraic structure (5.15) for each root system is desired.

6. ℓ operators

In this section we will show the equivalence of the quantum conserved quantities obtained in the Lax operator formalism of section 4 and those derived in the ‘commuting differential operators’ formalism initiated by Dunkl [10] and followed by many authors. Again the equivalence is universal, applicable to the models based on any root systems. We propose to call the operators in the latter approach simply ‘ ℓ operators’, since they are essentially the same as the L operator in the Lax pair formalism and they are not mutually commuting, as we will show presently, when the interaction potentials are trigonometric (hyperbolic) (6.14). Although these two formalisms are formally equivalent, the ℓ operator formalism has many advantages over the Lax pair one. Roughly speaking, the ‘vector-like’ objects ℓ_μ are easier to handle than the matrix $L_{\mu\nu}$.

Let us fix a representation \mathcal{R} of the Coxeter group G_Δ and define for each element $\mu \in \mathcal{R}$ the following differential-reflection operator:

$$\ell_\mu = \ell \cdot \mu = p \cdot \mu + i \sum_{\rho \in \Delta_+} g_{|\rho|}(\rho \cdot \mu) x(\rho \cdot q) \check{s}_\rho \quad \mu \in \mathcal{R}. \tag{6.1}$$

It is linear in μ and Coxeter covariant

$$\ell_{\mu+\nu} = \ell_\mu + \ell_\nu \quad \check{s}_\rho \ell_\mu \check{s}_\rho = \ell_{s_\rho(\mu)} \quad \forall \rho \in \Delta. \tag{6.2}$$

They are Hermitian operators, $\ell_\mu^\dagger = \ell_\mu$, with respect to the standard inner product for the states (5.4).

It is straightforward to show that the quantum conserved quantities Q_n derived in the previous section (4.7) can be expressed as polynomials in the ℓ operators as follows:

$$Q_n \psi = \sum_{\mu, \nu \in \mathcal{R}} (L^n)_{\mu\nu} \psi = \left(\sum_{\mu \in \mathcal{R}} \ell_\mu^n \right) \psi \tag{6.3}$$

in which ψ is an arbitrary Coxeter invariant state, $\check{s}_\rho \psi = \psi$. This also illustrates the Coxeter invariance of Q_n clearly, since $\check{s}_\rho (\sum_{\mu \in \mathcal{R}} \ell_\mu^n) \check{s}_\rho = \sum_{\mu \in \mathcal{R}} \ell_{s_\rho(\mu)}^n = \sum_{\mu \in \mathcal{R}} \ell_\mu^n$. For $n = 1$ this is trivial, since

$$\begin{aligned} \sum_{\nu \in \mathcal{R}} (L)_{\mu\nu} \psi &= \left(p \cdot \mu + i \sum_{\rho \in \Delta_+} g_{|\rho|}(\rho \cdot \mu) x(\rho \cdot q) \sum_{\nu \in \mathcal{R}} (\hat{s}_\rho)_{\mu\nu} \right) \psi \\ &= \left(p \cdot \mu + i \sum_{\rho \in \Delta_+} g_{|\rho|}(\rho \cdot \mu) x(\rho \cdot q) \check{s}_\rho \right) \psi = \ell_\mu \psi \end{aligned} \tag{6.4}$$

in which $\sum_{\nu \in \mathcal{R}} (\hat{s}_\rho)_{\mu\nu} = 1$ and $\check{s}_\rho \psi = \psi$ are used. Let us assume that

$$\sum_{\nu \in \mathcal{R}} (L^n)_{\mu\nu} \psi = \ell_\mu^n \psi \tag{6.5}$$

is correct, then we obtain

$$\begin{aligned} \sum_{\nu \in \mathcal{R}} (L^{n+1})_{\mu\nu} \psi &= \sum_{\lambda, \nu \in \mathcal{R}} L_{\mu\lambda} (L^n)_{\lambda\nu} \psi = \sum_{\lambda \in \mathcal{R}} L_{\mu\lambda} \ell_\lambda^n \psi \\ &= \sum_{\lambda \in \mathcal{R}} \left(p \cdot \mu \delta_{\mu\lambda} + i \sum_{\rho \in \Delta_+} g_{|\rho|} (\rho \cdot \mu) x(\rho \cdot q) (\hat{s}_\rho)_{\mu\lambda} \right) \ell_\lambda^n \psi. \end{aligned}$$

In the second summation only such λ as $\lambda = s_\rho(\mu)$ contribute and we find

$$\ell_{s_\rho(\mu)}^n \psi = (\check{s}_\rho \ell_\mu^n \check{s}_\rho) \psi = \check{s}_\rho \ell_\mu^n \psi.$$

Thus we arrive at

$$\sum_{\nu \in \mathcal{R}} (L^{n+1})_{\mu\nu} \psi = \ell_\mu^{n+1} \psi \tag{6.6}$$

and the equivalence of the two expressions of the conserved quantity (6.3) is proved.

Commutation relations among ℓ operators can be evaluated in a manner similar to those appearing in the Lax pair [7,9], that is, by decomposing the roots into two-dimensional sub-root systems. We obtain

$$[\ell_\mu, \ell_\nu] = -a^2 \sum_{\rho, \sigma \in \Delta_+} g_{|\rho|} g_{|\sigma|} (\rho \cdot \mu) (\sigma \cdot \nu) [\check{s}_\rho, \check{s}_\sigma] \times \begin{cases} 0 & \text{rational} \\ -1 & \text{hyperbolic} \\ 1 & \text{trigonometric.} \end{cases} \tag{6.7}$$

One important use of the ℓ operators is the proof of involution of quantum conserved quantities. For type I models Heckman [21] gave a universal proof based on the commutation relation (6.7):

$$[Q_n, Q_m] \psi = \sum_{\mu, \nu \in \mathcal{R}} [\ell_\mu^n, \ell_\nu^m] \psi = 0 \quad \text{rational model.} \tag{6.8}$$

This was the motivation for the introduction of the commuting differential-reflection operators by Dunkl [10]. In fact, the operators of Dunkl and Heckman were the similarity transformation of ℓ_μ by the ground state wavefunction e^W :

$$\tilde{\ell}_\mu = e^{-W} \ell_\mu e^W = p \cdot \mu + i \sum_{\rho \in \Delta_+} g_{|\rho|} \frac{(\rho \cdot \mu)}{(\rho \cdot q)} (\check{s}_\rho - 1). \tag{6.9}$$

As for type V models, we define ℓ^\pm corresponding to L^\pm (4.16):

$$\ell_\mu^\pm = \ell^\pm \cdot \mu = p \cdot \mu \pm i\omega(q \cdot \mu) + i \sum_{\rho \in \Delta_+} g_{|\rho|} \frac{(\rho \cdot \mu)}{(\rho \cdot q)} \check{s}_\rho \quad \mu \in \mathcal{R}. \tag{6.10}$$

They are linear in μ , Coxeter covariant and Hermitian conjugates of each other with respect to the standard inner product (5.4):

$$\check{s}_\rho \ell_\mu^\pm \check{s}_\rho = \ell_{s_\rho(\mu)}^\pm \quad (\ell_\mu^\pm)^\dagger = \ell_\mu^\mp. \tag{6.11}$$

The conserved quantities are expressed as polynomials in ℓ^\pm operators:

$$\begin{aligned} \text{Ts}(\mathcal{L}_1^n) \psi &= \sum_{\mu, \nu \in \mathcal{R}} (L^+ L^-)_{\mu\nu}^n \psi = \sum_{\mu \in \mathcal{R}} (\ell_\mu^+ \ell_\mu^-)^n \psi \\ \text{Ts}(\mathcal{L}_2^n) \psi &= \sum_{\mu, \nu \in \mathcal{R}} (L^- L^+)_{\mu\nu}^n \psi = \sum_{\mu \in \mathcal{R}} (\ell_\mu^- \ell_\mu^+)^n \psi. \end{aligned} \tag{6.12}$$

Likewise the creation and annihilation operators B_n^\pm (5.2) are expressed as

$$B_n^\pm \psi = \text{Ts}(L^\pm)^n \psi = \sum_{\mu, \nu \in \mathcal{R}} (L^\pm)_{\mu\nu} \psi = \sum_{\mu \in \mathcal{R}} (\ell_\mu^\pm)^n \psi. \tag{6.13}$$

The commutation relations among ℓ^\pm operators are easy to evaluate, since ℓ operators commute in the rational potential models (6.7):

$$[\ell_\mu^+, \ell_\nu^+] = [\ell_\mu^-, \ell_\nu^-] = 0 \quad [\ell_\mu^-, \ell_\nu^+] = 2\omega \left(\mu \cdot \nu + \sum_{\rho \in \Delta_+} g_{|\rho|} (\rho \cdot \mu) (\rho^\vee \cdot \nu) \check{s}_\rho \right). \quad (6.14)$$

From these it follows that the creation (annihilation) operators B_n^\pm do commute among themselves:

$$[B_n^+, B_m^+] \psi = [B_n^-, B_m^-] \psi = 0. \quad (6.15)$$

It is also clear that $\ell_\mu^\pm / \sqrt{2\omega}$ are the ‘deformation’ of the creation (annihilation) operators of the ordinary multicomponent harmonic oscillators. In fact we have

$$\ell_\mu^+ e^W = 2i\omega (\mu \cdot q) e^W \quad \text{and} \quad \ell_\mu^- e^W = 0. \quad (6.16)$$

In the next section we present an alternative scheme of algebraic construction of excited states of type V models by pursuing the analogy that ℓ^\pm are the creation and annihilation operators of the unit quantum. This method was applied to the A_r models by Brink *et al* and others [15, 17, 18].

7. Algebraic construction of excited states II

7.1. Operator solution of the triangular Hamiltonian

In section 3.1, we have shown that an eigenfunction of \mathcal{H} with eigenvalue $N\omega$ is given by

$$(P_N(q) + \tilde{P}_{N-2}(q)) e^W \quad (7.1)$$

in which $P_N(q)$ is a Coxeter invariant polynomial in q of homogeneous degree N and $\tilde{P}_{N-2}(q)$ is a Coxeter invariant polynomial in q of degree $N - 2$ and lower. The non-leading polynomial $\tilde{P}_{N-2}(q)$ is completely determined by the leading one $P_N(q)$ due to the triangularity. This solution can be written in an operator form as follows.

Suppose $P_N(q)$ is expressed as

$$P_N(q) = \sum_{\{\mu\}} c_{\{\mu\}} (q \cdot \mu_1) \dots (q \cdot \mu_N) \quad \mu_j \in \mathcal{R} \quad c_{\{\mu\}}: \text{const.} \quad (7.2)$$

We obtain a Coxeter invariant polynomial in the creation operators ℓ^+ by replacing $q \cdot \mu$ by $\ell_\mu^+ / (2i\omega)$:

$$P_N(q) \Rightarrow \frac{1}{(2i\omega)^N} P_N(\ell^+).$$

This creates the above eigenfunction of \mathcal{H} from the ground state:

$$\frac{1}{(2i\omega)^N} P_N(\ell^+) e^W = (P_N(q) + \tilde{P}_{N-2}(q)) e^W. \quad (7.3)$$

The proof is again elementary. By using the commutation relations among ℓ^\pm operators it is straightforward to derive the explicit expression of the Hamiltonian in terms of ℓ^\pm :

$$\mathcal{H} = \frac{1}{2\mathcal{C}_\mathcal{R}} \sum_{\mu \in \mathcal{R}} \ell_\mu^+ \ell_\mu^- + \sum_{\rho \in \Delta_+} g_{|\rho|} \left(\omega + \frac{1}{2} \frac{|\rho|^2}{(\rho \cdot q)^2} \right) (\check{s}_\rho - 1) \quad (7.4)$$

in which the second term vanishes upon acting on a Coxeter invariant state. Next we obtain

$$\frac{1}{2\mathcal{C}_\mathcal{R}} \sum_{\mu \in \mathcal{R}} [\ell_\mu^+ \ell_\mu^-, \ell_\nu^\pm] = [\ell_\nu^\pm, S] \pm \omega \ell_\nu^\pm \quad S \equiv \sum_{\rho \in \Delta_+} g_{|\rho|} \check{s}_\rho \quad (7.5)$$

which is an ℓ operator version of (4.15). Since a commutator is a derivation, we obtain

$$\frac{1}{2C_{\mathcal{R}}} \sum_{\mu \in \mathcal{R}} [\ell_{\mu}^{+} \ell_{\mu}^{-}, P_N(\ell^{+})] = [P_N(\ell^{+}), S] + N\omega P_N(\ell^{+}) \tag{7.6}$$

in which the first term in rhs vanishes due to the Coxeter invariance of P_N . Thus we arrive at the desired commutation relation

$$[\mathcal{H}, P_N(\ell^{+})] = N\omega P_N(\ell^{+}) + \sum_{\rho \in \Delta_{+}} g_{|\rho|} \left[\frac{1}{2} \frac{|\rho|^2}{(\rho \cdot q)^2}, P_N(\ell^{+}) \right] (\check{s}_{\rho} - 1) \tag{7.7}$$

and the eigenvalue equation

$$\mathcal{H} P_N(\ell^{+}) e^W = N\omega P_N(\ell^{+}) e^W. \tag{7.8}$$

Since the action of the creation operators on the ground state is

$$\ell_{\mu_1}^{+} \dots \ell_{\mu_N}^{+} e^W = [(2i\omega)^N (q \cdot \mu_1) \dots (q \cdot \mu_N) + \text{lower powers of } q] e^W \tag{7.9}$$

our assertion (7.3) is proved. It should be stressed that in this formalism the Coxeter invariance of the polynomial P is important but not how it is obtained.

Like the above Hamiltonian (7.4), the ℓ operator formulae of higher conserved quantities (6.12) contain extra terms:

$$\text{Ts}(\mathcal{L}_1^n) = \sum_{\mu, \nu \in \mathcal{R}} (L^{+} L^{-})_{\mu\nu}^n = \sum_{\mu \in \mathcal{R}} (\ell_{\mu}^{+} \ell_{\mu}^{-})^n + \text{VT}. \tag{7.10}$$

Here VT stands for vanishing terms when they act on a Coxeter invariant state. The same is true for most formulae derived in section 6.

7.2. States created by B_2^{+}

Here we derive the explicit forms of the subseries of eigenstates obtained by multiple applications of the least quanta creation operator B_2^{+} (5.2), or its normalized form b_2^{+} (5.11). It is convenient to work with the similarity transformed operator

$$\tilde{b}_2^{+} = e^{-W} b_2^{+} e^W = \frac{1}{4\omega C_{\mathcal{R}}} \sum_{\mu \in \mathcal{R}} (\tilde{\ell}_{\mu}^{+})^2 + \text{VT} \tag{7.11}$$

in which

$$\tilde{\ell}_{\mu}^{+} = p \cdot \mu + 2i\omega(q \cdot \mu) + i \sum_{\rho \in \Delta_{+}} \frac{\rho \cdot \mu}{\rho \cdot q} (\check{s}_{\rho} - 1). \tag{7.12}$$

Let $f(u)$ be an arbitrary function of $u \equiv \omega q^2$, then it is Coxeter invariant. We find

$$\tilde{\ell}_{\mu}^{+} f(u) = 2i\omega(q \cdot \mu) \left(1 - \frac{d}{du} \right) f(u) \quad u \equiv \omega q^2 \tag{7.13}$$

and

$$\tilde{b}_2^{+} f(u) = - \left[u \left(1 - \frac{d}{du} \right)^2 - \tilde{\mathcal{E}}_0 \left(1 - \frac{d}{du} \right) \right] f(u). \tag{7.14}$$

Since $\tilde{b}_2^{+} 1 = \tilde{\mathcal{E}}_0 - u = L_1^{(\tilde{\mathcal{E}}_0 - 1)}(u)$, we assume

$$(\tilde{b}_2^{+})^n 1 = n! L_n^{(\tilde{\mathcal{E}}_0 - 1)}(u). \tag{7.15}$$

By using the Laguerre differential equation (3.21) and the recurrence formulae of the Laguerre polynomial $L_n^{(\alpha)}(u)$,

$$u \frac{d}{du} L_n^{(\alpha)}(u) = n L_n^{(\alpha)}(u) - (n + \alpha) L_{n-1}^{(\alpha)}(u) \tag{7.16}$$

$$n L_n^{(\alpha)}(u) + (u - 2n - \alpha + 1) L_{n-1}^{(\alpha)}(u) + (n + \alpha - 1) L_{n-2}^{(\alpha)}(u) = 0 \tag{7.17}$$

we can show

$$- \left[u \left(1 - \frac{d}{du} \right)^2 - \tilde{\mathcal{E}}_0 \left(1 - \frac{d}{du} \right) \right] L_n^{(\tilde{\mathcal{E}}_0-1)}(u) = (n + 1) L_{n+1}^{(\tilde{\mathcal{E}}_0-1)}(u). \tag{7.18}$$

Thus the induction is proved and we arrive at (5.12). The orthogonality of the states

$$((B_2^+)^n e^W, (B_2^+)^m e^W) = 0 \quad n \neq m \tag{7.19}$$

can be easily understood as the du part of the measure

$$e^{2W} d^r q = e^{-u} u^{\tilde{\mathcal{E}}_0-1} du d\Omega \quad d\Omega: \text{angular part}$$

is the proper weight function for the Laguerre polynomial $L_n^{(\tilde{\mathcal{E}}_0-1)}(u)$.

7.3. Explicit solutions of the rank two models

For rank two models, the Liouville integrability, or the involution of conserved quantities, is automatically satisfied since the second conserved quantity is already obtained. For rank two type V models, the complete set of orthogonal wavefunctions can be written down explicitly in terms of separation of variables by using the Coxeter invariant polynomials. These are based on the dihedral root systems $I_2(m)$, with $A_2 \cong I_2(3)$ [31], $B_2 \cong I_2(4)$ and $G_2 \cong I_2(6)$ [32]. The Coxeter invariant polynomials exist at degree 2, i.e. q^2 and m which is

$$\prod_{j=1}^m (v_j \cdot q) \tag{7.20}$$

where $\{v_j\}$ is a set of vectors given in (B.4). If we introduce the two-dimensional polar coordinate system[†] for q

$$q = r(\sin \theta, \cos \theta) \tag{7.21}$$

then the principal Weyl chamber is

$$\text{PW} : 0 < r^2 < \infty \quad 0 < \theta < \pi/m. \tag{7.22}$$

The two Coxeter invariant variables read

$$q^2 = r^2 \quad \prod_{j=1}^m (v_j \cdot q) = 2 \left(\frac{r}{2} \right)^m \cos m\theta \tag{7.23}$$

and the latter variable varies over the full range, $-1 < \cos m\theta < 1$ in the PW. Thus solving the eigenvalue equation for \mathcal{H} (3.6) by separation of variables in the polar coordinate system is compatible with Coxeter invariance. We adopt as two independent variables

$$u \equiv \omega r^2 \quad z \equiv \cos m\theta. \tag{7.24}$$

The solutions consist of a Gegenbauer (Jacobi) polynomial in $\cos m\theta$ multiplied by a Laguerre polynomial in ωr^2 . The former we have encountered in the A_1 Sutherland problem, section 3.2, and the latter in the A_1 Calogero problem, sections 3.1 and 7.2.

[†] We believe no confusion arises here between the radial coordinate variable r and the rank of the root system r , which in this case is 2 of $I_2(m)$.

Let us demonstrate this for odd m with a single coupling constant and for even m with two independent coupling constants, in parallel. In terms of the Coxeter invariant variables (7.24) the $I_2(m)$ Hamiltonians take surprisingly simple forms:

$$\begin{aligned} \tilde{\mathcal{H}} &= \omega r \frac{\partial}{\partial r} - \frac{1}{2r} \frac{\partial}{\partial r} \left(r \frac{\partial}{\partial r} \right) - \frac{m}{r} \left\{ \frac{g}{\frac{1}{2}(g_o + g_e)} \right\} \frac{\partial}{\partial r} \\ &\quad - \frac{1}{2r^2} \left[\frac{\partial^2}{\partial \theta^2} + m \left\{ \frac{2g \cot m\theta}{-g_o \tan \frac{m\theta}{2} + g_e \cot \frac{m\theta}{2}} \right\} \frac{\partial}{\partial \theta} \right] \\ &= -2\omega \left[u \frac{\partial^2}{\partial u^2} + (\tilde{\mathcal{E}}_0 - u) \frac{\partial}{\partial u} \right] - \frac{\omega m^2}{2u} \left[(1 - z^2) \frac{\partial^2}{\partial z^2} \right. \\ &\quad \left. + \left(\left\{ \begin{matrix} 0 \\ g_o - g_e \end{matrix} \right\} - \left\{ \begin{matrix} 1 + 2g \\ 1 + g_e + g_o \end{matrix} \right\} z \right) \frac{\partial}{\partial z} \right]. \end{aligned} \tag{7.25}$$

The z part admits polynomial solutions

$$\begin{aligned} &\left[(1 - z^2) \frac{d^2}{dz^2} + \left(\left\{ \begin{matrix} 0 \\ g_o - g_e \end{matrix} \right\} - \left\{ \begin{matrix} 1 + 2g \\ 1 + g_e + g_o \end{matrix} \right\} z \right) \frac{d}{dz} \right] P_\ell^{\left\{ \begin{matrix} (g - \frac{1}{2}, g - \frac{1}{2}) \\ (g_o - \frac{1}{2}, g_e - \frac{1}{2}) \end{matrix} \right\}}(z) \\ &= \ell \left(\ell + \left\{ \begin{matrix} 2g \\ g_e + g_o \end{matrix} \right\} \right) P_\ell^{\left\{ \begin{matrix} (g - \frac{1}{2}, g - \frac{1}{2}) \\ (g_o - \frac{1}{2}, g_e - \frac{1}{2}) \end{matrix} \right\}}(z) \end{aligned} \tag{7.26}$$

in which ℓ is the degree of the polynomial. After substituting them, the radial part of the Hamiltonian $\tilde{\mathcal{H}}_r$ reads

$$\tilde{\mathcal{H}}_r = -2\omega \left[u \frac{d^2}{du^2} + (\tilde{\mathcal{E}}_0 - u) \frac{d}{du} - \frac{m^2}{4u} \ell \left(\ell + \left\{ \begin{matrix} 2g \\ g_e + g_o \end{matrix} \right\} \right) \right]. \tag{7.27}$$

By similarity transformation in terms of $u^{m\ell/2} \propto r^{m\ell}$, which is the radial part of the highest term of the polynomial $P_\ell^{(\alpha, \beta)}(r^m \cos m\theta)$, it reads

$$u^{-m\ell/2} \tilde{\mathcal{H}}_r u^{m\ell/2} = -2\omega \left[u \frac{d^2}{du^2} + (m\ell + \tilde{\mathcal{E}}_0 - u) \frac{d}{du} - \frac{m\ell}{2} \right]. \tag{7.28}$$

This is the main part of the differential equation for the Laguerre polynomial (3.21):

$$\left[u \frac{d^2}{du^2} + (m\ell + \tilde{\mathcal{E}}_0 - u) \frac{d}{du} - n \right] L_n^{(m\ell + \tilde{\mathcal{E}}_0 - 1)}(u) = 0.$$

Thus the eigenstates of the Hamiltonian are obtained:

$$\tilde{\mathcal{H}}_r u^{m\ell/2} L_n^{(m\ell + \tilde{\mathcal{E}}_0 - 1)}(u) = \omega(2n + m\ell) u^{m\ell/2} L_n^{(m\ell + \tilde{\mathcal{E}}_0 - 1)}(u) \tag{7.29}$$

$$\tilde{\mathcal{H}} u^{m\ell/2} L_n^{(m\ell + \tilde{\mathcal{E}}_0 - 1)}(u) P_\ell^{\left\{ \begin{matrix} (g - \frac{1}{2}, g - \frac{1}{2}) \\ (g_o - \frac{1}{2}, g_e - \frac{1}{2}) \end{matrix} \right\}}(z) = (\omega(2n + m\ell) + \mathcal{E}_0) u^{m\ell/2} L_n^{(m\ell + \tilde{\mathcal{E}}_0 - 1)}(u) P_\ell^{\left\{ \begin{matrix} (g - \frac{1}{2}, g - \frac{1}{2}) \\ (g_o - \frac{1}{2}, g_e - \frac{1}{2}) \end{matrix} \right\}}(z). \tag{7.30}$$

It is instructive to note that the Hamiltonians $\hat{\mathcal{H}}$ also look simple:

$$\hat{\mathcal{H}} = \frac{1}{2} \omega^2 r^2 - \frac{1}{2r} \frac{\partial}{\partial r} \left(r \frac{\partial}{\partial r} \right) - \frac{1}{2r^2} \frac{\partial^2}{\partial \theta^2} + \frac{m^2}{2r^2} \left\{ \frac{\frac{g(g-1)}{\sin^2 m\theta}}{4 \cos^2 \frac{m\theta}{2}} + \frac{g_e(g_e-1)}{4 \sin^2 \frac{m\theta}{2}} \right\}. \tag{7.31}$$

Olshanetsky and Perelomov [5] obtained the above solutions starting from these formulae.

8. Involution of conserved quantities

8.1. Universal proof of involution of quantum conserved quantities for type I–III models

Here we present a proof of involution of quantum conserved quantities $\{Q_n\}$ derived from the universal Lax pair in section 4.1 for type I–III models. The proof is applicable to all models based on any root systems. Though a universal proof of involution for type I models is given by Heckman [21] as recapitulated in section 6, we believe the universal proof applicable to type II and III models as well is new. It depends on a theorem by Olshanetsky and Perelomov [11]. Our own contribution is that we have provided a universal Lax pair and conserved quantities satisfying all the requirements of the theorem.

Liouville’s theorem states the complete integrability as the existence of an involutive set of conserved quantities as many as the degrees of freedom. We have already given conserved quantities $\{Q_n\}$ (4.7) independent and as many as the degrees of freedom (see appendix B). They have the following properties.

(1) Coxeter invariance

$$Q_n(s_\rho(p), s_\rho(q)) = Q_n(p, q) \quad \forall \rho \in \Delta. \tag{8.1}$$

(2) $Q_n(p, q)$ is a homogeneous polynomial of degree n in variables $(p_1, \dots, p_r, x(\rho \cdot q))$.

(3) Scaling property for those of type I models:

$${}^I Q_n(\kappa^{-1} p, \kappa q) = \kappa^{-n} {}^I Q_n(p, q) \tag{8.2}$$

as a consequence of the above point.

(4) For type II and III models, the asymptotic behaviour near the origin:

$$Q_n(p, q) = {}^I Q_n(p, q)(1 + \mathcal{O}(|q|)) \quad \text{for } |q| \rightarrow 0. \tag{8.3}$$

We need to show the vanishing of

$$J_{lm} \equiv [Q_l, Q_m] \tag{8.4}$$

which is a polynomial in $\{p\}$ of degree s

$$s < l + m. \tag{8.5}$$

Let us decompose J_{lm} into the leading part and the rest:

$$J_{lm} = J_{lm}^0 + J_{lm}^{\text{rest}} \quad J_{lm}^0 = \sum c^{j_1, \dots, j_s}(q) p_{j_1}, \dots, p_{j_s} \tag{8.6}$$

and J_{lm}^{rest} is a polynomial in $\{p\}$ of degree less than s . From the Jacobi identity and conservation $[\mathcal{H}, Q_{l(m)}] = 0$, we obtain

$$[\mathcal{H}, J_{lm}] = 0. \tag{8.7}$$

Considering the explicit form of the Hamiltonian (2.7) ($\omega = 0$), the leading (i.e. of degree $s + 1$ in $\{p\}$) part of $[\mathcal{H}, J_{lm}]$ arises only from the free part

$$[p^2, J_{lm}^0]$$

and it vanishes if the following conditions are satisfied:

$$\sum_{\sigma} \frac{\partial}{\partial q_t} c^{k_1, \dots, k_s}(q) = 0 \tag{8.8}$$

where the sum is taken over all permutations of indices $\sigma(t, k_1, \dots, k_s) = (j_1, \dots, j_{s+1})$. In [33] it is proved (lemma 2.5, p 407) that the system (8.8) has only polynomial solutions. Then Olshanetsky and Perelomov argue that for type I models the scaling property tells us that $c^{k_1, \dots, k_s}(\kappa q) = \kappa^{s-l-m} c^{k_1, \dots, k_s}(q)$. Since $s < l + m$ (8.5), it follows that the

only polynomial solution satisfying the condition is the null polynomial. Thus we obtain $c^{j_1, \dots, j_s}(q) = 0 \Rightarrow J_{lm}^0 = 0$ and $J_{lm} = 0$. The same results follow for type II and III models by considering the asymptotic behaviour for $|q| \rightarrow 0$. Thus the involution of all the conserved quantities $\{Q_n\}$ is proved. This result also implies the involution of classical conserved quantities by taking the classical limit ($\hbar \rightarrow 0$).

8.2. Rational models with the harmonic confining force

In this section we show the involution of quantum conserved quantities for the type V models based on the root systems of classical Lie algebras. The method is a straightforward generalization of that developed by Polychronakos on the A_r model. This is made possible by the availability of the universal Lax pair formalism [7, 9], in particular the root-type and minimal-type Lax pairs. We apply it to the models based on B_r and D_r root systems. For the rational potential, the B_r , C_r and BC_r models are equivalent. Let us choose $\mathcal{R} = \{\pm e_j \in \mathbb{R}^r \mid e_j \cdot e_k = \delta_{jk}\}$ as the representation space of the Coxeter group consisting of orthogonal vectors and their negatives. They are the set of short roots of B_r and the set of vector weights of D_r in the parametrization of roots given in (B.1) and (4.13), respectively.

The conserved quantities in the ℓ operator form are given by

$$Q_n = {}^\ell Q_n + \text{VT} \quad {}^\ell Q_n \equiv \sum_{j=1}^r (\ell_j^+ \ell_j^-)^n \quad n = 1, \dots, r \tag{8.9}$$

in which we abbreviate $\ell_{e_j}^\pm$ as ℓ_j^\pm . In this case the commutation relations among ℓ operators (6.14) are greatly simplified thanks to the orthogonality of $\{e_j\}$ s and the explicit forms of the roots:

$$[\ell_j^-, \ell_k^+] = -2\omega \tilde{g}(\check{s}_{jk} - \check{\bar{s}}_{jk}) \tag{8.10}$$

in which

$$\tilde{g} = \begin{cases} g_L & B_r \text{ model} \\ g & D_r \text{ model} \end{cases} \quad \check{s}_{jk} \equiv \check{s}_{e_j - e_k} \quad \check{\bar{s}}_{jk} \equiv \check{s}_{e_j + e_k}. \tag{8.11}$$

By repeating them we obtain

$$[\ell_j^+ \ell_j^-, \ell_k^+ \ell_k^-] = -2\omega \tilde{g}[\ell_j^+ \ell_j^-, m_{jk}] \quad m_{jk} \equiv \check{s}_{jk} + \check{\bar{s}}_{jk} = m_{kj} \quad j \neq k \tag{8.12}$$

$$[(\ell_j^+ \ell_j^-)^n, \ell_k^+ \ell_k^-] = -2\omega \tilde{g}[(\ell_j^+ \ell_j^-)^n, m_{jk}] = +2\omega \tilde{g}[(\ell_k^+ \ell_k^-)^n, m_{jk}] \tag{8.13}$$

$$[\ell_j^+ \ell_j^-, (\ell_k^+ \ell_k^-)^m] = +2\omega \tilde{g}[(\ell_k^+ \ell_k^-)^m, m_{jk}] = -2\omega \tilde{g}[(\ell_j^+ \ell_j^-)^m, m_{jk}]. \tag{8.14}$$

Here and later the identity $[(\ell_k^+ \ell_k^-)^t, m_{jk}] = -[(\ell_j^+ \ell_j^-)^t, m_{jk}]$ is used repeatedly. Then (8.13) leads to

$$[(\ell_j^+ \ell_j^-)^n, (\ell_k^+ \ell_k^-)^m] = +2\omega \tilde{g} \sum_{t=0}^{m-1} ((\ell_k^+ \ell_k^-)^{t+n} m_{jk} (\ell_k^+ \ell_k^-)^{m-t-1} - (\ell_k^+ \ell_k^-)^t m_{jk} (\ell_k^+ \ell_k^-)^{m+n-t-1}) \tag{8.15}$$

and (8.14) to

$$[(\ell_j^+ \ell_j^-)^n, (\ell_k^+ \ell_k^-)^m] = -2\omega \tilde{g} \sum_{t=0}^{n-1} ((\ell_j^+ \ell_j^-)^{t+m} m_{jk} (\ell_j^+ \ell_j^-)^{n-t-1} - (\ell_j^+ \ell_j^-)^t m_{jk} (\ell_j^+ \ell_j^-)^{m+n-t-1}). \tag{8.16}$$

Summing over j and k and adding (8.15) and (8.16) together with the interchange of the dummy indices $j \leftrightarrow k$ in the latter produces

$$2[\ell Q_n, \ell Q_m] = +2\omega\tilde{g} \sum_{t=0}^{n+m-1} \sum_{j,k} ((\ell_k^+ \ell_k^-)^t m_{jk} (\ell_k^+ \ell_k^-)^{n+m-t-1} - (\ell_k^+ \ell_k^-)^t m_{jk} (\ell_k^+ \ell_k^-)^{n+m-t-1}) = 0. \tag{8.17}$$

Thus we obtain

$$[Q_n, Q_m] = 0 \quad n, m = 1, \dots, r \tag{8.18}$$

on the Fock space of Coxeter invariant states.

8.3. Lax pair with spectral parameter

In the theory of classical Calogero–Moser models, the Lax pair with spectral parameter (ξ) plays an important role, in particular, in elliptic potential models [6–8, 34] for derivation of spectral curves etc. In quantum theory, however, the meaning and use of the spectral parameter are yet to be established, partly because of the underdeveloped stage of the quantum models with elliptic potentials. Here we point out a small use of the quantum Lax pair with spectral parameter in the quantum model with trigonometric potential. Namely, it accounts for the useful trick by Polychronakos [15] for the proof of the involution of conserved quantities in the trigonometric A_r model. (Now we have a universal proof of involution for type I, II and III models; see the previous section.)

From the theory of the generalized Lax pair [7] and its quantum version for degenerate potential models [9], we find that the L operator can contain one additional complex parameter ξ :

$$L^\xi = p \cdot \hat{H} + X^\xi \quad X^\xi = i \sum_{\rho \in \Delta_+} g_{|\rho|} (\rho \cdot \hat{H})(x(\rho \cdot q) - x(\rho^\vee \cdot \hat{H}\xi))\hat{s}_\rho \tag{8.19}$$

in which the function x is given in table 1 for the degenerate potentials. With the same M operator as before, the quantum equations of motion can be written in a matrix form:

$$\frac{d}{dt} L^\xi = i[\mathcal{H}, L^\xi] = [L^\xi, M]. \tag{8.20}$$

In other words, the ξ -dependent part decouples. This allows us to define a one-parameter family of conserved quantities

$${}^\xi Q_n = \text{Ts}(L^\xi)^n \quad n = 1, 2, \dots \tag{8.21}$$

which turns out to be a ξ -dependent sum of Q_n and the lower-order conserved quantities $Q_m, m < n$. A special limit $\xi \rightarrow -i\infty$ in the trigonometric models provides a convenient combination which allows easy proof of involution in the A_r model. (In the rational model, the limit reduces to the Lax pair without spectral parameter. In the hyperbolic models this limit is ill defined.) In the rest of this section we consider only the trigonometric potential models. Let us denote the limiting L^ξ operator by L^∞ , which reads

$$L^\infty = L + a \sum_{\rho \in \Delta_+} g_{|\rho|} |\rho \cdot \hat{H}| \hat{s}_\rho \tag{8.22}$$

and the corresponding ℓ operators are given by

$$\ell_\mu^\infty = \ell_\mu + t_\mu \quad t_\mu = a \sum_{\rho \in \Delta_+} g_{|\rho|} |\rho \cdot \mu| \check{s}_\rho. \tag{8.23}$$

For the A_r model in the vector representation $\mathcal{R} = \{\mu_j \in \mathbb{R}^r, j = 1, \dots, r + 1\}$, with the standard normalization of roots $\rho^2 = 2$, the above expression simplifies to

$$\ell_j^\infty = \ell_j + t_j \quad t_j = ag \sum_{k \neq j} \check{s}_{jk} \tag{8.24}$$

in which as before we abbreviate ℓ_{μ_j} as ℓ_j and $\check{s}_{jk} \equiv \check{s}_{\mu_j - \mu_k}$. These are the operators introduced in [15], $\ell_j \leftrightarrow \pi_j$ and $\ell_j^\infty \leftrightarrow \tilde{\pi}_j$. The Lax operator with the spectral parameter gives an ‘explanation’ for the rather *ad hoc* introduction of $\tilde{\pi}_j$. It is straightforward to show

$$[\ell_j, \ell_k] = [t_j, t_k] \tag{8.25}$$

which leads to

$$[\ell_j^\infty, \ell_k^\infty] = [\ell_j^\infty, t_k] + [t_j, \ell_k^\infty] = 2ag[\ell_j^\infty, \check{s}_{jk}] = -2ag[\ell_k^\infty, \check{s}_{jk}]. \tag{8.26}$$

This has the same structure as (8.12) in the previous section. By repeating the same argument we arrive at [15]

$$[{}^\infty Q_n, {}^\infty Q_m] = 0 \quad n, m = 1, \dots, r \quad {}^\infty Q_n = \sum_{j=1}^r (\ell_j^\infty)^n \tag{8.27}$$

which then imply the involution of the conserved quantities obtained from the original L operator

$$[Q_n, Q_m] = 0 \quad n, m = 1, \dots, r. \tag{8.28}$$

9. Summary and comments

We have discussed various issues related to quantum integrability of Calogero–Moser models based on all root systems. These are construction of quantum conserved quantities and a unified proof of their involution, the relationship between the Lax pair and the differential-reflection (Dunkl) operator formalisms, construction of excited states by creation operators etc. They are mainly generalizations of the results known for the models based on A_r root systems, which are shown to apply to the models based on any root systems. There are some interesting works discussing the integrability issues of the models based on other classical root systems and the exceptional ones including the non-crystallographic models [32, 35–38].

Here we list some comments on interesting issues which are not treated in this paper. The structure and properties of the eigenfunctions of the trigonometric potential models, which are generalizations of the Jack polynomials [25–28], will be discussed in future publications. A comprehensive treatment of Liouville integrability of rational models with harmonic force is required. Our starting point, the factorized Hamiltonian (2.5) for degenerate potential models, is closely related with supersymmetry and shape invariance [38, 39]. Further investigation in this direction is a future problem. It is a great challenge to formulate various aspects of quantum Calogero–Moser models with elliptic potentials; the Lax pair, the differential-reflection operators [40, 41], conserved quantities, supersymmetry and excited state wavefunctions.

Acknowledgments

We thank K Takasaki, S Otake and P Ghosh for useful discussion. This work is partially supported by a grant-in-aid from the Ministry of Education, Science and Culture, priority area (707) ‘Supersymmetry and unified theory of elementary particles’. SPK and AJP are supported by the Japanese Society for the Promotion of Science.

Appendix A. Root systems

In this appendix we recapitulate the rudimentary facts of the root systems and reflections to be used in the main text. The set of roots Δ is invariant under reflections in the hyperplane perpendicular to each vector in Δ . In other words,

$$s_\alpha(\beta) \in \Delta \quad \forall \alpha, \beta \in \Delta \quad (\text{A.1})$$

where

$$s_\alpha(\beta) = \beta - (\alpha^\vee \cdot \beta)\alpha \quad \alpha^\vee \equiv 2\alpha/|\alpha|^2. \quad (\text{A.2})$$

The set of reflections $\{s_\alpha, \alpha \in \Delta\}$ generates a group G_Δ , known as a Coxeter group, or finite reflection group. The orbit of $\beta \in \Delta$ is the set of root vectors resulting from the action of the Coxeter group on it. The set of positive roots Δ_+ may be defined in terms of a vector $U \in \mathbb{R}^r$, with $\alpha \cdot U \neq 0, \forall \alpha \in \Delta$, as those roots $\alpha \in \Delta$ such that $\alpha \cdot U > 0$. Given Δ_+ , there is a unique set of r simple roots $\Pi = \{\alpha_j, j = 1, \dots, r\}$ defined such that they span the root space and the coefficients $\{a_j\}$ in $\beta = \sum_{j=1}^r a_j \alpha_j$ for $\beta \in \Delta_+$ are all non-negative. The highest root α_h , for which $\sum_{j=1}^r a_j$ is maximal, is then also determined uniquely. The subset of reflections $\{s_\alpha, \alpha \in \Pi\}$ in fact generates the Coxeter group G_Δ . The products of s_α , with $\alpha \in \Pi$, are subject solely to the relations

$$(s_\alpha s_\beta)^{m(\alpha, \beta)} = 1 \quad \alpha, \beta \in \Pi. \quad (\text{A.3})$$

The interpretation is that $s_\alpha s_\beta$ is a rotation in some plane by $2\pi/m(\alpha, \beta)$. The set of positive integers $m(\alpha, \beta)$ (with $m(\alpha, \alpha) = 1, \forall \alpha \in \Pi$) uniquely specifies the Coxeter group. The weight lattice $\Lambda(\Delta)$ is defined as the \mathbb{Z} -span of the fundamental weights $\{\lambda_j, j = 1, \dots, r\}$, defined by

$$\alpha_j^\vee \cdot \lambda_k = \delta_{jk} \quad \alpha_j \in \Pi. \quad (\text{A.4})$$

The root systems for finite reflection groups may be divided into two types: crystallographic and non-crystallographic. Crystallographic root systems satisfy the additional condition

$$\alpha^\vee \cdot \beta \in \mathbb{Z} \quad \forall \alpha, \beta \in \Delta \quad (\text{A.5})$$

which implies that the \mathbb{Z} -span of Π is a lattice in \mathbb{R}^r and contains all roots in Δ . We call this the root lattice, which is denoted by $L(\Delta)$. These root systems are associated with simple Lie algebras: $\{A_r, r \geq 1\}$, $\{B_r, r \geq 2\}$, $\{C_r, r \geq 2\}$, $\{D_r, r \geq 4\}$, E_6, E_7, E_8, F_4 and G_2 . The Coxeter groups for these root systems are called Weyl groups. The remaining non-crystallographic root systems are H_3 and H_4 , whose Coxeter groups are the symmetry groups of the icosahedron and four-dimensional 600-cell, respectively, and the dihedral group of order $2m$, $\{I_2(m), m \geq 4\}$.

Appendix B. Conserved quantities

Here we list for each root system how the full set of independent conserved quantities is obtained by choosing proper representations of the Lax pair. We choose those of the lowest dimensionality for the convenience of practical calculation. Of course there are many other choices of representations giving equally good sets of conserved quantities. The independence of the conserved quantities can be easily verified by considering the free limit: $g_{|\rho|} \rightarrow 0$.

(1) A_r . For all powers, the vector representation ($r + 1$ dimensions) is enough.

- (2) B_r . For all powers, the representation consisting of short roots $\{\pm e_j : j = 1, \dots, r\}$ ($2r$ dimensions) is enough. Here we adopt the following explicit parametrization of the B_r root system:

$$B_r \text{ root system: } \Delta = \{\pm e_j \pm e_k, \pm e_j, j, k = 1, \dots, r | e_j \in \mathbb{R}^r, e_j \cdot e_k = \delta_{jk}\}. \quad (\text{B.1})$$

- (3) C_r . For all powers, the representation consisting of long roots $\{\pm 2e_j : j = 1, \dots, r\}$ ($2r$ dimensions) is enough. The following parametrization of the root system is used:

$$C_r \text{ root system: } \Delta = \{\pm e_j \pm e_k, \pm 2e_j, j, k = 1, \dots, r | e_j \in \mathbb{R}^r, e_j \cdot e_k = \delta_{jk}\}. \quad (\text{B.2})$$

- (4) D_r : For all even powers, the vector representation ($2r$ dimensions) is enough. For the additional one occurring at power r , the (anti-) spinor representation (2^{r-1} dimensions) would be necessary. They are minimal representations.
- (5) E_6 . For all powers, the **27**- (or $\overline{27}$ -) dimensional representation of the Lie algebra is enough. They are minimal representations.
- (6) E_7 . For all powers, the **56**-dimensional representation of the Lie algebra is enough. This is a minimal representation.
- (7) E_8 . For all powers, the 240-dimensional representation consisting of all the roots is enough. This is not the same as the adjoint representation of the Lie algebra.
- (8) F_4 . For all powers, either of the 24-dimensional representations consisting of all the long roots or the short roots is enough. These are not Lie algebra representations.
- (9) G_2 . For all powers, either of the six-dimensional representations consisting of all the long roots or the short roots is enough. These are not Lie algebra representations.
- (10) $I_2(m)$. For both powers 2 and m , the representation consisting of the vertices R_m of the regular m -gon is sufficient,

$$R_m = \{(\cos(2k/m + t_0), \sin(2k/m + t_0)) \in \mathbb{R}^2 | k = 1, \dots, m\} \quad (\text{B.3})$$

in which $t_0 = 0$ ($1/2m$) for m even (odd). Another set, V_m , is used in 7.3. Here, V_m is the set of vectors with ‘half’ angles of the roots (see (2.10)) given by

$$V_m = \{v_j = (\cos((2j - 1)\pi/2m), \sin((2j - 1)\pi/2m)) \in \mathbb{R}^2 | j = 1, \dots, m\}. \quad (\text{B.4})$$

- (11) H_3 . For all powers, the representation consisting of all 30 roots is enough.
- (12) H_4 . For all powers, the representation consisting of all 120 roots is enough.

References

- [1] Calogero F 1971 Solution of the one-dimensional N -body problem with quadratic and/or inversely quadratic pair potentials *J. Math. Phys.* **12** 419–36
- [2] Sutherland B 1972 Exact results for a quantum many-body problem in one-dimension. II *Phys. Rev. A* **5** 1372–6
- [3] Moser J 1975 Three integrable Hamiltonian systems connected with isospectral deformations *Adv. Math.* **16** 197–220
- Moser J 1975 Integrable systems of non-linear evolution equations *Dynamical Systems, Theory and Applications (Lecture Notes in Physics vol 38)* ed J Moser (Berlin: Springer)
- Calogero F, Marchioro C and Ragnisco O 1975 Exact solution of the classical and quantal one-dimensional many body problems with the two body potential $V_a(x) = g^2 a^2 / \sinh^2 ax$ *Lett. Nuovo Cimento* **13** 383–7
- Calogero F 1975 Exactly solvable one-dimensional many body problems *Lett. Nuovo Cimento* **13** 411–6
- [4] Olshanetsky M A and Perelomov A M 1976 Completely integrable Hamiltonian systems connected with semisimple Lie algebras *Invent. Math.* **37** 93–108
- [5] Olshanetsky M A and Perelomov A M 1983 Quantum integrable systems related to Lie algebras *Phys. Rep.* **94** 313–404
- [6] Bordner A J, Corrigan E and Sasaki R 1998 Calogero–Moser models I: a new formulation *Prog. Theor. Phys.* **100** 1107–29
(Bordner A J, Corrigan E and Sasaki R 1998 *Preprint* hep-th/9805106)

- Bordner A J, Sasaki R and Takasaki K 1999 Calogero–Moser models II: symmetries and foldings *Prog. Theor. Phys.* **101** 487–518
(Bordner A J, Sasaki R and Takasaki K 1998 *Preprint* hep-th/9809068)
- Bordner A J and Sasaki R 1999 Calogero–Moser models III: elliptic potentials and twisting *Prog. Theor. Phys.* **101** 799–829
(Bordner A J and Sasaki R 1998 *Preprint* hep-th/9812232)
- Khastgir S P, Sasaki R and Takasaki K 1999 Calogero–Moser models IV: limits to Toda theory *Prog. Theor. Phys.* **102** 749–76
(Khastgir S P, Sasaki R and Takasaki K 1999 *Preprint* hep-th/9907102)
- [7] Bordner A J, Corrigan E and Sasaki R 1999 Generalised Calogero–Moser models and universal Lax pair operators *Prog. Theor. Phys.* **102** 499–529
(Bordner A J, Corrigan E and Sasaki R 1999 *Preprint* hep-th/9905011)
- [8] D’Hoker E and Phong D H 1998 Calogero–Moser Lax pairs with spectral parameter for general Lie algebras *Nucl. Phys. B* **530** 537–610
(D’Hoker E and Phong D H 1998 *Preprint* hep-th/9804124)
- [9] Bordner A J, Manton N S and Sasaki R 2000 Calogero–Moser models V: supersymmetry, and quantum Lax pair *Prog. Theor. Phys.* **103** 463–87
(Bordner A J, Manton N S and Sasaki R 1999 *Preprint* hep-th/9910033)
- [10] Dunkl C F 1989 Differential-difference operators associated to reflection groups *Trans. Am. Math. Soc.* **311** 167–83
Buchstaber V M, Felder G and Veselov A P 1994 Elliptic Dunkl operators, root systems and functional equations *Duke Math. J.* **76** 885–911
- [11] Olshanetsky M A and Perelomov A M 1977 Quantum systems related to root systems, and radial parts of Laplace operators *Funct. Anal. Appl.* **12** 121–8
- [12] Perelomov A M 1971 Algebraic approach to the solution of a one-dimensional model of interacting particles *Theor. Math. Phys.* **6** 263–82
- [13] Wojciechowski S 1976 Involutive set of integrals for completely integrable many-body problems with pair interaction *Lett. Nuovo Cimento* **18** 103–7
- [14] Lassalle M 1991 Polynômes de Jacobi, généralisés *C. R. Acad. Sci., Paris (t. Sér. I Math)* **312** 425–8
Lassalle M 1991 Polynômes de Laguerre généralisés *C. R. Acad. Sci., Paris (t. Sér. I Math)* **312** 725–8
Lassalle M 1991 Polynômes de Hermite généralisés *C. R. Acad. Sci., Paris (t. Sér. I Math)* **313** 579–82
- [15] Polykronakos A P 1992 Exchange operator formalism for integrable systems of particles *Phys. Rev. Lett.* **69** 703–5
- [16] Shastry B S and Sutherland B 1993 Superlax pairs and infinite symmetries in the $1/r^2$ system *Phys. Rev. Lett.* **70** 4029–33
- [17] Brink L, Hansson T H and Vasiliev M A 1992 Explicit solution to the N body Calogero problem *Phys. Lett. B* **286** 109–11
(Brink L, Hansson T H and Vasiliev M A 1992 *Preprint* hep-th/9206049)
Brink L, Hansson T H, Konstein S and Vasiliev M A 1993 The Calogero model: anyonic representation, fermionic extension and supersymmetry *Nucl. Phys. B* **401** 591–612
(Brink L, Hansson T H, Konstein S and Vasiliev M A 1993 *Preprint* hep-th/9302023)
Brink L, Turbiner A and Wyllard N 1998 Hidden algebras of the (super) Calogero and Sutherland models *J. Math. Phys.* **39** 1285–315
(Brink L, Turbiner A and Wyllard N 1997 *Preprint* hep-th/9705219)
- [18] Ujino H, Wadati M and Hikami K 1993 The quantum Calogero–Moser model: algebraic structures *J. Phys. Soc. Japan* **62** 3035–43
Ujino H and Wadati M 1996 Rodrigues formula for Hi-Jack symmetric polynomials associated with the quantum Calogero model *J. Phys. Soc. Japan* **65** 2423–39
(Ujino H and Wadati M 1996 *Preprint* cond-mat/9609041)
Ujino H 1995 Orthogonal symmetric polynomials associated with the Calogero model *J. Phys. Soc. Japan* **64** 2703–6
(Ujino H 1997 *Preprint* cond-mat/9706133)
Nishino A, Ujino H and Wadati M 2000 Symmetric Fock space and orthogonal symmetric polynomials associated with the Calogero model *Chaos Solitons Fractals* **11** 657–74
(Nishino A, Ujino H and Wadati M 1998 *Preprint* cond-mat/9803284)
- [19] Sogo K 1996 A simple derivation of multivariable Hermite and Legendre polynomials *J. Phys. Soc. Japan* **65** 3097–9
Gurappa N and Panigrahi P K 1999 Equivalence of the Calogero–Sutherland model to free harmonic oscillators

- Phys. Rev. B* **59** R2490–3
(Gurappa N and Panigrahi P K 1997 *Preprint* cond-mat/9710035)
- [20] Ruijsenaars S N M 1999 *Systems of Calogero–Moser type (CRM Series in Math. Phys. vol 1)* (Berlin: Springer) pp 251–352
- [21] Heckman G J 1991 A remark on the Dunkl differential-difference operators *Harmonic Analysis on Reductive Groups* ed W Barker and P Sally (Basel: Birkhäuser)
- [22] Heckman G J and Opdam E M 1987 Root systems and hypergeometric functions I *Comput. Math.* **64** 329–52
Heckman G J 1987 Root systems and hypergeometric functions II *Comput. Math.* **64** 353–73
Opdam E M 1988 Root systems and hypergeometric functions III *Comput. Math.* **67** 21–49
Opdam E M 1988 Root systems and hypergeometric functions IV *Comput. Math.* **67** 191–209
- [23] Freedman D Z and Mende P F 1990 An exactly solvable N -particle system in supersymmetric quantum mechanics *Nucl. Phys.* **344** 317–43
- [24] Rühl W and Turbiner A 1995 Exact solvability of the Calogero and Sutherland models *Mod. Phys. Lett. A* **10** 2213–22
(Rühl W and Turbiner A 1995 *Preprint* hep-th/9506105)
Haschke O and Rühl W 2000 Is it possible to construct exactly solvable models *Lecture Notes in Physics* vol 539 (Berlin: Springer) pp 118–40
(Haschke O and Rühl W 1998 *Preprint* hep-th/9809152)
- [25] Stanley R 1989 Some combinatorial properties of Jack symmetric function *Adv. Math.* **77** 76–115
Macdonald I G 1995 *Symmetric Functions and Hall Polynomials* 2nd edn (Oxford: Oxford University Press)
- [26] Lapointe L and Vinet L 1997 Rodrigues formulas for the Macdonald polynomials *Adv. Math.* **130** 261–79
(Lapointe L and Vinet L 1996 *Preprint* q-alg/9607025)
Lapointe L and Vinet L 1996 Exact operator solution of the Calogero–Sutherland model *Commun. Math. Phys.* **178** 425–52
(Lapointe L and Vinet L 1995 *Preprint* q-alg/9509003)
- [27] Baker T H and Forrester P J 1997 The Calogero–Sutherland model and generalized classical polynomials *Commun. Math. Phys.* **188** 175–216
(Baker T H and Forrester P J 1996 *Preprint* solv-int/9608004)
- [28] Awata H, Matsuo Y, Odake S and Shiraishi J 1995 Excited states of Calogero–Sutherland model and singular vectors of W_N algebra *Nucl. Phys. B* **449** 347–74
(Awata H, Matsuo Y, Odake S and Shiraishi J 1995 *Preprint* hep-th/9503043)
- [29] Erdélyi A *et al* 1955 *Higher Transcendental Functions III* (New York: McGraw-Hill)
- [30] Gambardella P J 1975 Exact results in quantum many-body systems of interacting particles in many dimensions with $SU(1, 1)$ as the dynamical group *J. Math. Phys.* **16** 1172–87
- [31] Calogero F 1969 Solution of a three body problem in one dimension *J. Math. Phys.* **10** 2191–6
Calogero F 1969 Ground state of a one-dimensional N -body problem *J. Math. Phys.* **10** 2197–200
- [32] Wolfes J 1974 On the three-body linear problem with three-body interaction *J. Math. Phys.* **15** 1420–4
Calogero F and Marchioro C 1974 Exact solution of a one-dimensional three-body scattering problem with two-body and/or three-body inverse-square potential *J. Math. Phys.* **15** 1425–30
- [33] Berezin F A 1957 Laplace operators on semisimple Lie groups *Tr. Mosk. Mat. Ob-va* **6** 371–463
- [34] Krichever I M 1980 Elliptic solutions of the Kadomtsev–Petviashvili equation and integrable systems of particles *Funct. Anal. Appl.* **14** 282–9
- [35] Rosenbaum M, Turbiner A and Capella A 1998 Solvability of the g_2 integrable system *Int. J. Mod. Phys. A* **13** 3885–904
(Rosenbaum M, Turbiner A and Capella A 1997 *Preprint* solv-int/9707005)
Gurappa N, Khare A and Panigrahi P K 1998 Connection between Calogero–Marchioro–Wolfes type few-body models and free oscillators *Preprint* cond-mat/9804207
- [36] Haschke O and Rühl W 1998 An exactly solvable model of the Calogero type for the icosahedral group *Mod. Phys. Lett. A* **13** 3109–22
(Haschke O and Rühl W 1998 *Preprint* hep-th/9811011)
Haschke O and Rühl W 1999 The construction of trigonometric invariants for Weyl groups and the derivation of corresponding exactly solvable Sutherland models *Mod. Phys. Lett. A* **14** 937–49
(Haschke O and Rühl W 1999 *Preprint* math-ph/9904002)
- [37] Kakei S 1996 Common algebraic structure for the Calogero–Sutherland models *J. Phys. A: Math. Gen.* **29** L619–24
Kakei S 1997 An orthogonal basis for the B_N -type Calogero model *J. Phys. A: Math. Gen.* **30** L535–41
- [38] Ghosh P K, Khare A and Sivakumar M 1998 Supersymmetry, shape invariance and solvability of A_{N-1} and BC_N Calogero–Sutherland model *Phys. Rev. A* **58** 821–5

- (Ghosh P K, Khare A and Sivakumar M 1997 *Preprint* cond-mat/9710206)
Sukhatme U and Khare A 1999 Comment on self-isospectral periodic potentials and supersymmetric quantum mechanics *Preprint* quant-ph/9902072
- [39] Efthimiou C and Spector D 1997 Shape invariance in the Calogero and Calogero–Sutherland models *Phys. Rev. A* **56** 208–19
(Efthimiou C and Spector D 1997 *Preprint* quant-ph/9702017)
- [40] Cherednik I V 1995 Elliptic quantum many-body problem and double affine Knizhnik–Zamolodchikov equation *Commun. Math. Phys.* **169** 441–61
- [41] Oshima T and Sekiguchi H 1995 Commuting families of differential operators invariant under the action of a Weyl group *J. Math. Sci. Univ. Tokyo* **2** 1–75