

Elastic scattering and bound states in the Aharonov-Bohm potential superimposed by an attractive ρ^{-2} potential

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2001 J. Phys. A: Math. Gen. 34 235

(<http://iopscience.iop.org/0305-4470/34/2/303>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.95

The article was downloaded on 02/06/2010 at 08:58

Please note that [terms and conditions apply](#).

Elastic scattering and bound states in the Aharonov–Bohm potential superimposed by an attractive ρ^{-2} potential

Jürgen Audretsch¹, Vladimir D Skarzhinsky^{1,2} and Boris L Voronov²

¹ Fakultät für Physik der Universität Konstanz, Postfach M 673, D-78457 Konstanz, Germany

² P N Lebedev Physical Institute, Leninsky prospect 53, Moscow 117924, Russia

E-mail: Juergen.Audretsch@uni-konstanz.de, vdskarzh@sgi.lebedev.ru and voronov@td.lpi.ac.ru

Received 18 April 2000, in final form 17 October 2000

Abstract

We consider the elastic scattering and bound states of charged quantum particles moving in the Aharonov–Bohm and an attractive ρ^{-2} potential in a partial-wave approach. Radial solutions of the stationary Schrödinger equation are specified in such a way that the Hamiltonian of the problem is self-adjoint. It is shown that they are not uniquely fixed but depend on open parameters. The related physical consequences are discussed. The scattering cross section is calculated and the energy spectrum of bound states is obtained.

PACS numbers: 0365B, 0365G, 0365N, 0230T

1. Introduction

The famous Aharonov–Bohm (AB) effect [1] has been the subject of many investigations in the past (cf [13, 14]). During the last few years novel applications have been discussed which open up new areas of research. Detailed calculations of QED processes in the presence of the magnetic string have been carried out. For example, the cross sections of bremsstrahlung and pair production have been evaluated analytically [3]. Recently, the AB effect found an unexpected application in the domain of atomic interferometry with neutral atoms. Improving a suggestion in [17], the following experimental set-up has been proposed in [16]: the radial electric field \mathbf{E} of a straight charged wire with homogenous charge density κ_0 polarizes scattered neutral atoms. A uniform magnetic field \mathbf{B} is applied parallel to the wire. The atoms with mass M_0 and electric polarizability α moving in these two fields will then acquire a quantum phase. The respective Lagrangian for these atoms is

$$L = \frac{1}{2} M v^2 + \alpha [\mathbf{B} \times \mathbf{E}] \cdot \frac{\mathbf{v}}{c} + \frac{1}{2} \alpha E^2 \quad (1)$$

with $M = M_0 + \alpha B^2/c^2$ if terms of order v^2/c^2 are neglected. The corresponding stationary Schrödinger equation is in cylindrical coordinates ρ and φ (we omit the trivial z dependence and put $\hbar = 1, c = 1$)

$$\left\{ \frac{1}{2M} \left[\frac{\partial^2}{\partial \rho^2} + \frac{1}{\rho} \frac{\partial}{\partial \rho} - \frac{1}{\rho^2} \left(-i \frac{\partial}{\partial \varphi} - \beta \right)^2 \right] + \frac{\kappa^2}{\rho^2} + \mathcal{E} \right\} \psi(\rho, \varphi) = 0. \quad (2)$$

The *magnetic-field parameter* β is thereby $\beta = \alpha \kappa_0 / 2\pi c$ and the *charge parameter* κ is $\kappa = \alpha \kappa_0^2 / 8\pi^2$. $\mathcal{E} = p^2 / 2M$ is the energy of particles with momentum p . The magnetic field yields no forces on the particles, nevertheless a pure topological phase is attached to the wavefunction. Atom interferometer experiments to measure this effect are on their way. Results for the limiting case of a vanishing magnetic field ($B = 0$) are described in [12]. A computer simulation has been performed in [7].

There is a physically different set-up which leads to the same differential equation (2). It makes the relation to the usual AB situation evident. The motion of a particle with mass M and charge e in the AB potential,

$$\mathbf{A}(\mathbf{r}) = \frac{\Phi}{2\pi} \frac{[\mathbf{n} \times \mathbf{r}]}{\rho^2} \quad \rho^2 = r^2 - (\mathbf{n} \cdot \mathbf{r}) \quad (3)$$

(\mathbf{n} is a unit vector along the magnetic string and Φ is the magnetic flux) which is superimposed by the attractive scalar potential of an electrically charged string in the \mathbf{n} -direction

$$U(\rho) = -\frac{\kappa^2}{\rho^2} \quad \kappa^2 \geq 0 \quad (4)$$

is described quantum mechanically by the Hamiltonian operator

$$H = \frac{1}{2M} [-i\nabla_r - e\mathbf{A}(\mathbf{r})]^2 - \frac{\kappa^2}{\rho^2} \quad (5)$$

obtained by the canonical quantization procedure. Referring to cylindrical coordinates this again leads to the Schrödinger equation (2) where the magnetic-field parameter β now agrees with the flux $\beta = e\Phi/2\pi$. Below we will restrict β to the interval $0 \leq \beta < 1$ because addition of an integer number does not result in a physical effect.

In previous papers we had introduced absorption of atoms on the surface of a wire with finite radius [4, 5]. The same problem has been treated in [11]. In this paper we will take the two singular potentials \mathbf{A} and U (which both lead to terms proportional to ρ^{-2}) seriously for all values of ρ including the limit $\rho \rightarrow 0$. That severe physical problems may arise in this case, has already been pointed out in [9]. The reason for this is that it is not enough to consider the operator H of (5) on its own, but the domain has to be specified on which H becomes a self-adjoint and therefore quantum mechanically admissible operator. This can be obtained by a self-adjoint extension [15]. It is a well known fact that such an extension may be unique or non-unique. Our aim is to answer this for our case in performing the extension procedure. If it turns out to be unique, then the self-adjoint operator with singular potentials may be regarded as a limiting case of some well defined physical situation free of singular potentials. If, on the other hand, the extension is non-unique it may be regarded as the limiting case of many different physical set-ups. In both cases it is a task to specify the underlying non-singular situations.

To treat the problem we introduce a partial-wave decomposition in section 2. That this is feasible for the scattering problem although the potentials in question are far reaching, has been demonstrated [5]. In section 3 we find normalizable radial functions of positive energy

for the Schrödinger equation (2). They contain arbitrary coefficients and are non-orthogonal at different values of momentum p unless these coefficients are fixed appropriately. To do this we use a *pragmatic approach* [2] which is based on a direct evaluation of the normalization integral. Setting the non-orthogonal terms in this integral equal to zero we obtain the conditions under which the Hamiltonian H is a self-adjoint operator on the corresponding radial functions. Being equivalent to the standard procedure of the self-adjoint extension this approach is more transparent since it uses the fact that any self-adjoint operator possesses an orthogonal set of eigenfunctions. However, having cured the mathematical problem, a physical problem arises: it turns out that the resulting complete orthogonal set of scattering states depends on an infinite number of *open* or *free parameters*. What are the physical set-ups and the corresponding scattering states from which they can be obtained in an appropriate limit?

The cylindrically symmetric hard-core model is the simplest model to think of in which the singularity of the potentials in $\rho = 0$ is avoided. We show in section 4 that the limit of vanishing radius of the hard-core cylinder cannot be used to fix free parameters. So it remains an open question as to which realistic physical models lead in an appropriate limit to the admissible solutions found in the ‘pragmatic approach’. In section 5 we return to the Schrödinger equation (8) and its solutions with open parameters and discuss the general traits of the elastic scattering. We study the differential and the total cross sections as functions of the physically specified parameters β, κ and p . In section 6 we turn to the bound states and the related energy spectrum.

In appendix A the procedure of obtaining the self-adjoint extension is described in a mathematically more rigorous way. It is shown that the results agree with those obtained in the ‘pragmatic approach’.

2. Partial-wave approach to elastic scattering

To describe scattering we have to find a solution of equation (2) which satisfies the scattering boundary condition for $\rho \rightarrow \infty$

$$\psi(\vec{p}; \rho, \varphi) \rightarrow e^{i\beta(\varphi-\pi)} e^{ip\rho \cos \varphi} + f(\varphi) \frac{e^{ip\rho}}{\sqrt{\rho}} \quad (6)$$

where $p := \sqrt{2M\mathcal{E}}$. Note that the first term, which represents the wavefunction $\psi_{\text{in}}(\rho, \varphi)$ of the incoming flux, has the form of a modified plane wave (cf [5]). We assume that the angle φ_p under which the particles fall in, is equal to zero. For arbitrary φ_p replace $\varphi \rightarrow \varphi - \varphi_p$.

We assume rotational and translational symmetry and solve equation (2) using a partial-wave decomposition

$$\psi(\rho, \varphi) = \sum_{m=-\infty}^{\infty} R_m(\rho) e^{im\varphi} \quad m = 0, \pm 1, \pm 2, \dots \quad (7)$$

where m is the orbital angular-momentum quantum number. That this is a well defined approach in our case, although it includes Aharonov–Bohm scattering as a limiting case, has been shown in [5]. The radial functions $R_m(\rho)$ satisfy the Bessel equation

$$R_m''(\rho) + \frac{1}{\rho} R_m'(\rho) - \frac{v^2}{\rho^2} R_m(\rho) + p^2 R_m(\rho) = 0 \quad (8)$$

where we have introduced the *effective quantum number* v and the new *charge parameter* γ

$$v^2 := (m - \beta)^2 - \gamma^2 \quad \gamma^2 := 2M\kappa^2. \quad (9)$$

$\gamma \geq 0$ represents (because of (4)) the strength of the attractive potential, ν^2 depends apart from β and γ^2 on the orbital quantum number m (we omit from ν the index m). Because γ and the magnetic-field parameter β specify the physical situation, we call them together with the momentum p the *physical parameters*. If we describe with the equations above the scattering of neutral polarizable atoms by the charged wire in the uniform magnetic field [16], the parameter ν^2 acquires an additional term from the magnetic mass: $\gamma^2 \rightarrow \gamma^2 + \beta^2$.

The scattering amplitude $f(\varphi)$ can be expressed in terms of phase shifts δ_m which are defined by the asymptotic form of the radial functions

$$R_m(\rho) \rightarrow \sqrt{\frac{1}{2\pi p\rho}} \left[e^{-i(p\rho - \pi m - \frac{1}{4}\pi)} + S_m e^{i(p\rho - \frac{1}{4}\pi)} \right] \quad S_m := e^{2i\delta_m}. \quad (10)$$

We take into account the asymptotic behaviour of the radial functions R_m^{in} of the incoming wave (cf [5])

$$\psi^{\text{in}}(\rho, \varphi) = e^{i\beta(\varphi - \pi)} e^{ip\rho \cos \varphi} = \sum_{m=-\infty}^{\infty} R_m^{\text{in}}(\rho) e^{im\varphi} \quad (11)$$

$$R_m^{\text{in}}(\rho) \rightarrow \sqrt{\frac{1}{2\pi p\rho}} \left[e^{-i(p\rho - \pi m - \frac{1}{4}\pi)} + \cos \pi \beta e^{i(p\rho - \frac{1}{4}\pi)} \right]. \quad (12)$$

Then the scattering amplitude reads

$$f(\varphi) = \frac{1}{\sqrt{2\pi}} \sum_{m=-\infty}^{\infty} f_m e^{im\varphi} \quad f_m = \frac{e^{-\frac{1}{4}i\pi}}{\sqrt{p}} (S_m - \cos \beta). \quad (13)$$

3. Self-adjoint Hamiltonian and the appearance of open parameters

The Hamiltonian H of equation (2) contains two singular potentials, the AB-potential $A_\varphi(\rho)$ of equation (3) and the attractive scalar potential $U(\rho)$ of equation (4). Both lead to terms proportional to ρ^{-2} . To make H a quantum mechanical operator one has to fix the domain of solutions on which it becomes a self-adjoint operator. We describe the corresponding self-adjoint extension approach in detail in appendix A. An alternative quick and simple procedure leading to the same results is the ‘pragmatic approach’ which we have applied earlier in [2]. It is based on the demand that the radial functions $R_m(\rho)$ should be not only normalizable with regard to ρ but also orthogonal for different values of p :

$$\int_0^\infty R_m^*(p'\rho) R_m(p\rho) \rho \, d\rho = \frac{\delta(p - p')}{\sqrt{pp'}}. \quad (14)$$

This turns out to be equivalent to the restriction to solutions with specific boundary conditions for the radial functions at $\rho = 0$ which imply self-adjointness. For the pure AB potential this problem has been considered by many authors.

It is important to note that the parameter ν^2 of equation (9) can have negative values for any values of the physical parameters β and γ . This happens for non-vanishing charge parameter $\gamma \neq 0$ and modes with orbital quantum numbers m out of the interval $(\beta - \gamma < m < \beta + \gamma)$. For the pure Aharonov–Bohm case ($\gamma = 0$) there are no such modes. They are possible here because there is, in addition, the attractive potential $U(\rho)$.

For the normalizable solutions we therefore have to distinguish three domains of v^2 which correspond for given physical parameters β and γ to different intervals of the quantum number m . The conditions $v^2 > 0$ and $\mu := |v| > 1$ characterize the intervals of large positive and negative m extending to $\pm\infty$. (In [5] we used the definition $\mu = \text{Im } v$ which can differ in sign from the new one.) The corresponding normalizable solutions of equation (8) are

$$R_m(\rho) = c_m J_\mu(p\rho) \quad |m - \beta| > \sqrt{1 + \gamma^2} \quad |c_m| = 1. \quad (15)$$

The normalizable solutions with $v^2 > 0$ and $\mu < 1$ have the form

$$R_m(\rho) = a_m J_{-\mu}(p\rho) + b_m J_\mu(p\rho) \quad \gamma < |m - \beta| < \sqrt{1 + \gamma^2} \quad (16)$$

with arbitrary coefficients a_m and b_m . If $\gamma \neq 0$ the condition in equation (16) can be fulfilled for not more than two values of m , depending on the values of β and γ . For the pure AB case ($\gamma = 0$), for example, we can have only $m = 0$ and 1. This domain is an intermediate domain with respect to the values of m .

Turning to $v^2 < 0$ what is possible for $\gamma \neq 0$ we find

$$R_m(\rho) = a_m J_{-i\mu}(p\rho) + b_m J_{i\mu}(p\rho) \quad |m - \beta| < \gamma. \quad (17)$$

These solutions with the smallest values of m have to be taken into account whenever the scalar potential is large enough so that $\gamma > \min[\beta, 1 - \beta]$.

The solutions (15) are orthogonal for different p . This is in general not the case for the solutions (16) and (17) if the coefficients a_m or b_m are not specified further. Because of the singular potentials, the Hamiltonian of equation (8) is not self-adjoint if applied to the domain of all possible solutions (15)–(17). Following the ‘pragmatic approach’ to self-adjointness we will now impose the condition (14). Using the relations (see appendix B)

$$\int_0^\infty J_\nu(p'\rho) J_\nu(p\rho) \rho \, d\rho = \frac{\delta(p - p')}{\sqrt{pp'}} \quad (18)$$

and

$$\int_0^\infty J_{-\nu}(p'\rho) J_\nu(p\rho) \rho \, d\rho = \frac{\delta(p - p')}{\sqrt{pp'}} \cos \pi \nu + \frac{2 \sin \pi \nu}{\pi(p^2 - p'^2)} \left(\frac{p}{p'}\right)^\nu \quad (19)$$

we obtain from equation (16) for the intermediate interval of quantum numbers m :

$$\begin{aligned} \int_0^\infty R_m^*(p'\rho) R_m(p\rho) \rho \, d\rho &= [a_m^* a_m + b_m^* b_m + \cos \pi \mu (a_m^* b_m + b_m^* a_m)] \frac{\delta(p - p')}{\sqrt{pp'}} \\ &+ \frac{2i \sin \pi \mu}{\pi(p^2 - p'^2)} \left[\left(\frac{p}{p'}\right)^\mu a_m^*(p') b_m(p) - \left(\frac{p}{p'}\right)^{-\mu} b_m^*(p') a_m(p) \right]. \end{aligned} \quad (20)$$

It is the second term which may prevent orthogonality. To establish orthogonality we have to demand

$$\frac{b_m^*(p')}{a_m^*(p')} p'^{2\mu} = \frac{b_m(p)}{a_m(p)} p^{2\mu} \quad (21)$$

which implies that the expressions on both side of this equation are independent of p and real. Accordingly we have for $v^2 > 0$ and $\mu < 1$

$$R_m(\rho) = c_m \left[\lambda_m \left(\frac{p}{M}\right)^{2\mu} J_{-\mu}(p\rho) + J_\mu(p\rho) \right] \quad \gamma < |m - \beta| < \sqrt{1 + \gamma^2} \quad (22)$$

with dimensionless real parameters λ_m fulfilling

$$|c_m|^2 \left[\lambda_m^2 \left(\frac{p}{M} \right)^{4\mu} + 2\lambda_m \left(\frac{p}{M} \right)^{2\mu} \cos \pi \mu + 1 \right] = 1$$

which are not specified by the pragmatic approach. We will confirm this more rigorously in appendix A. The procedure to establish self-adjointness therefore does not lead to a unique solution for given m but to a whole set of solutions parametrized by the set of open parameters $\{\lambda_m\}$.

Turning to quantum numbers m with $\nu^2 < 0$ we obtain with equation (17)

$$\int_0^\infty R_m^*(p'\rho) R_m(p\rho) \rho \, d\rho = [a_m^* b_m + b_m^* a_m + \cosh \pi \mu (a_m^* a_m + b_m^* b_m)] \frac{\delta(p-p')}{\sqrt{pp'}} - \frac{2i \sinh \pi \mu}{\pi(p^2 - p'^2)} \left[\left(\frac{p}{p'} \right)^{-i\mu} a_m^*(p') a_m(p) - \left(\frac{p}{p'} \right)^{i\mu} b_m^*(p') b_m(p) \right]. \quad (23)$$

For orthogonality the second term has to vanish. This amounts to

$$\frac{b_m^*(p')}{a_m^*(p')} p'^{-2i\mu} = \frac{a_m(p)}{b_m(p)} p^{-2i\mu}. \quad (24)$$

Accordingly the expressions on either side of this equation must be independent of p and their complex conjugate must be equal to their inverse. The consequence is that the radial solutions R_m of equation (17) must be restricted to

$$R_m(\rho) = c_m [e^{i\theta_m + 2\mu \ln(p/M)} J_{-i\mu}(p\rho) + J_{i\mu}(p\rho)] \quad |m - \beta| < \gamma \quad (25)$$

with open dimensionless parameters θ_m out of the interval $0 \leq \theta_m < 2\pi$ and the normalization condition amounts to $2|c_m|^2 [\cosh \pi \mu + \cos(2\mu \ln p/M + \theta_m)] = 1$. Again self-adjointness does not lead to a unique solution for a given m (cf appendix A).

It is easily to check that the partial radial currents

$$j_m(\rho) = \frac{p}{2iM} [R_m^*(p\rho) R_m'(p\rho) - R_m(p\rho) R_m'^*(p\rho)] \quad (26)$$

vanish for the solutions (15), (22) and (25). The same is true for the total current. The reason for this is that ingoing and outgoing currents compensate each other, thereby reflecting the fact that the scattering is purely elastic.

4. The cylindrical hard-core model

We have shown that for the elastic scattering of particles in the two singular potentials the relevant Hamiltonian is self-adjoint and therefore quantum-mechanically admissible not only for one but for a total set of solutions parametrized by open parameters. Because there is not a unique solution, the differential equation (8) cannot be attributed to one specific physical situation. The reason may be that many different physical situations with non-singular potentials and certain boundary conditions lead in some physical limit to the same differential equation (8) with two singular potentials. One could imagine that as far as the wavefunctions are concerned, each of these situations leads to one particular solution $R_m(\rho)$ with well specified parameters λ_m and θ_m . In the following we want to examine whether in this way the physically simplest well defined physical situation is related to particular values of the parameters. For this we study the model with a hard core with surface ρ_0 so that no particles can penetrate

the surface. This amounts to $R_m(\rho) = 0$ for $\rho_0 \leq 0$ (cf [4]). The limit to be discussed is $\rho_0 \rightarrow 0$.

For quantum numbers m with $v^2 > 0$ the hard-core solutions

$$\begin{aligned} R_m^{\text{hc}}(\rho) &= \tilde{c}_m [J_\mu(p\rho_0)J_{-\mu}(p\rho) - J_{-\mu}(p\rho_0)J_\mu(p\rho)] \\ &\sim -[J_\mu(p\rho_0)/J_{-\mu}(p\rho_0)]J_{-\mu}(p\rho) + J_\mu(p\rho) \quad |m - \beta| > \gamma \end{aligned} \quad (27)$$

do not contain open parameters because there are no singular potentials. Because of $\lim_{\rho_0 \rightarrow 0} J_\mu(p\rho_0)/J_{-\mu}(p\rho_0) = 0$ this leads for $\rho_0 \rightarrow 0$ to the solution (15) for $\mu > 1$ and to the solution (22) with fixed parameter $\lambda_m = 0$ in the intermediate region with $\mu < 1$.

For the domain of quantum numbers m with $v^2 < 0$ the parameter-free hard-core solutions are

$$R_m^{\text{hc}}(\rho) = \tilde{c}_m [J_{-i\mu}(p\rho_0)J_{i\mu}(p\rho) - J_{i\mu}(p\rho_0)J_{-i\mu}(p\rho)] \quad |m - \beta| < \gamma. \quad (28)$$

In order to see to which values of the parameters θ_m this is related, we go to the limit $\rho_0 \rightarrow 0$ and find

$$e^{i(\theta_m + 2\mu \ln p/M)} = \lim_{\rho_0 \rightarrow 0} \left[-\frac{J_{i\mu}(p\rho_0)}{J_{-i\mu}(p\rho_0)} \right] \quad \theta_m = \lim_{\rho_0 \rightarrow 0} 2\mu \ln \frac{M\rho_0}{2} + \pi - 2\xi_m \quad (29)$$

with $\xi_m := \arg \Gamma(1 + i\mu)$. For $\rho_0 \rightarrow 0$ we have $\theta_m \rightarrow -\infty$. Therefore, this limit of the hard-core model does not fix the open parameters θ_m .

This is not surprising. The problem in question, as specified by equation (2), is explicitly scale invariant. On the other hand, by the appearance of open parameters caused by the self-adjoint extension, scales are introduced (compare with the example in section 6). To obtain a physical interpretation of the open parameters from more physical models in going to a certain limit, these models must contain scale parameters which do not disappear in this limit. The hard-core model contains the parameter ρ_0 , but it disappears in the limit $\rho_0 \rightarrow 0$.

Taken these results together we have shown that the hard-core model in the limit of vanishing ρ_0 does not lead to a solution $\psi(\rho, \varphi)$ based on the radial solutions $R_m(\rho)$ above if values of m out of the interval $|m - \beta| < \gamma$ have to be taken into account. For large values of the charge parameter γ this will be the case.

5. Scattering amplitude and cross section

To work out the scattering cross section in the general case of equations (2) and (8) we refer to the limit of the hard-core model and take $\lambda_m = 0$. In this case there is one type of solution for the whole quantum number domain $v^2 > 0$ so we have

$$R_m = \begin{cases} c_m J_\mu(p\rho) & \text{for } v^2 > 0 \\ c_m [e^{i(\theta_m + 2\mu \ln(p/M))} J_{-i\mu}(p\rho) + J_{i\mu}(p\rho)] & \text{for } v^2 < 0. \end{cases} \quad (30)$$

Based on the scattering boundary condition (6) and the asymptotic form (12) of the partial waves of the distorted ingoing plane wave, we find for the coefficients

$$c_m = \begin{cases} e^{i\pi(m - \frac{1}{2}\mu)} & \text{for } v^2 > 0 \\ e^{i\pi m} [e^{-\frac{1}{2}\pi\mu} + e^{\frac{1}{2}\pi\mu} e^{i(\theta_m + 2\mu \ln p/M)}]^{-1} & \text{for } v^2 < 0. \end{cases} \quad (31)$$

The partial scattering amplitudes of equation (13) then turn out to be

$$S_m = \begin{cases} e^{i\pi(m-\mu)} & \text{for } \nu^2 > 0 \\ e^{i\pi m - i(\theta_m + 2\mu \ln p/M)} \frac{e^{\pi\mu} + e^{i(\theta_m + 2\mu \ln p/M)}}{e^{\pi\mu} + e^{-i(\theta_m + 2\mu \ln p/M)}} & \text{for } \nu^2 < 0. \end{cases} \quad (32)$$

Equation (8) is explicitly scale invariant because the parameter γ and, consequently, the parameter ν are dimensionless. One could therefore expect that the phase functions are independent of the momentum p and the related energy [10]. This is the case if the operator of equation (8) is essentially self-adjoint. If not, the procedure of self-adjoint extension may lead to a p dependence. Equation (32) shows this explicitly for $\nu^2 < 0$. It is then convenient to replacing θ_m by dimensional open parameters $p_{m,0}$ out of the interval $Me^{-\pi/\mu} \leq p_{m,0} < M$ according to

$$\theta_m := 2\mu \ln \frac{M}{p_{m,0}} \quad (33)$$

so that S_m of equation (32) depend on $p/p_{m,0}$. We will see below that because of the necessity for a self-adjoint extension for bound states, the energy spectrum will depend on $p/p_{m,0}$ too, so that an arbitrary energy scale appears.

To work out the scattering amplitude $f(\varphi)$ it is useful to separate the contribution of the AB potential in order to obtain convergent series (cf [5]). This leads to

$$\begin{aligned} f(\varphi) &= \frac{e^{-i\pi/4}}{\sqrt{2\pi p}} \sum_{m=-\infty}^{\infty} (S_m - \cos \beta) e^{im\varphi} \\ &= f_{\text{AB}}(\varphi) + \frac{e^{-i\pi/4}}{\sqrt{2\pi p}} \sum_{m=-\infty}^{\infty} (S_m - S_m^{\text{AB}}) e^{im\varphi} := f_{\text{AB}}(\varphi) + \frac{e^{-i\pi/4}}{\sqrt{2\pi p}} \delta\Sigma(\varphi) \end{aligned} \quad (34)$$

with

$$S_m^{\text{AB}} = e^{i\pi(m-|m-\beta|)} \quad f_{\text{AB}}(\varphi) = -\frac{1}{\sqrt{2\pi p}} e^{-i\pi/4} e^{i\varphi/2} \frac{\sin \pi\beta}{\sin(\varphi/2)}. \quad (35)$$

For the subsequent numerical evaluation we split $\delta\Sigma(\varphi)$ into three parts,

$$\delta\Sigma(\varphi) = \sum_{m>\gamma+\beta}^{\infty} (S_m - S_m^{\text{AB}}) e^{im\varphi} + \sum_{m>\gamma-\beta}^{\infty} (S_{-m} - S_{-m}^{\text{AB}}) e^{-im\varphi} + \sum_{\substack{m<\beta+\gamma \\ m>\beta-\gamma}} (S_m - S_m^{\text{AB}}) e^{im\varphi}. \quad (36)$$

The differential cross section $d\sigma/d\varphi$ is a function of the scattering angle φ and depends on the physical parameters β and γ and on momentum p which specify the underlying physical situation. The open parameters θ_m in equation (32) are by definition functions of $M/p_{m,0}$ (cf equation (33)). In the following we chose θ_m to be equal for all m and correspondingly $p_{m,0} := p_0$. Equations (32) and (34) then show that the dimensionless differential scattering cross section is a function of φ and of the combination p/p_0

$$2\pi p \frac{d\sigma}{d\varphi} = \sqrt{2\pi p} |f(\varphi)|^2 := \eta(\varphi, p/p_0). \quad (37)$$

In figure 1 we show the function $\eta(\varphi, p/p_0)$ obtained by a numerical calculation for $p/p_0 = 1$ and $p/p_0 = 80$ (full curves) and values of the physical parameters $\beta = 0.5$ and $\gamma = 9.9$. The broken curve represent the AB limit $\gamma = 0$. For small scattering angles the AB contribution dominates the effect due to the factor $\sin^2(\varphi/2)$ in the denominator of $f_{\text{AB}}(\varphi)$. It can be




Figure 1. Dimensionless differential cross section $\eta = 2\pi p d\sigma/d\varphi$ of elastic scattering as a function of the scattering angle φ for two domains of φ . The AB cross section ($\gamma = 0$) is shown as broken curve.

seen that η for small angles φ is independent of p/p_0 and therefore independent of the open parameter p_0 . For larger scattering angles the AB contribution becomes negligibly small. We notice here oscillations as functions of φ as well as a weak dependence on p/p_0 .

We now turn to the total elastic scattering cross section σ . It is divergent in the presence of the AB potential which decreases too slowly at infinity. Therefore, we work out σ for the special case of a vanishing flux parameter $\beta = 0$. Accordingly it depends only on the momentum p , the charge parameter γ and on the open parameters $p_{m,0}$. We obtain

$$\sigma = \int_0^{2\pi} |f(\varphi)|^2 d\varphi = \sum_m \sigma_m = \sum_m f_m^* f_m = \frac{2}{p} \sum_m (1 - \text{Re } S_m) \quad (38)$$

with

$$1 - \text{Re } S_m = \begin{cases} 1 - \cos \pi m \cos \pi \mu & \text{for } v^2 > 0 \\ 1 - \cos \pi m \cosh \pi \mu + \frac{\sinh^2 \pi \mu \cos \pi m}{\cosh \pi \mu + \cos(\theta + 2\mu \ln p/M)} & \text{for } v^2 < 0, \end{cases} \quad (39)$$

θ_m of equation (33) and $\mu = |v|$ of equation (9). Decomposing according to

$$\sigma = \sigma_{v^2>0} + \sigma_{v^2<0} \quad (40)$$

we see that $\sigma_{v^2>0}$ has a simple p dependence of the form

$$\sigma_{v^2>0} = \frac{2\Sigma_1(\gamma)}{p} \quad (41)$$




Figure 2. The function $\Sigma_1(\gamma)$ of equation (41) (bold curve). The thin and broken curves show the functions $\pi^4\gamma^4/24$ and $\pi^2\gamma^2/2$, respectively.




Figure 3. $\Sigma_2(p)$ of equation (42) for the parameter choice $\beta = 0$, $\gamma = 0.9$ and $p_{0,0} = 10$. The dimensionless total cross section $0.5 p \sigma(p)$ is obtained in adding to Σ_2 the constant $\Sigma_1 \approx 3.48$ thus shifting up the curves.

where $\Sigma_1(\gamma)$ is an increasing function of γ . $\Sigma_1(\gamma)$ is close to $\pi^4\gamma^4/24$ for small γ , whereas it approaches asymptotically the parabola $\pi^2\gamma^2/2$ for large γ (see figure 2).

$\sigma_{v^2 < 0}$ has in contrast to $\sigma_{v^2 > 0}$ no simple dependence on the momentum p . Writing

$$\sigma_{v^2 < 0} = \frac{2}{p} \Sigma_2(\gamma, p) \quad (42)$$

we can discuss the p dependence of Σ_2 for a given value of the charge parameter γ . Restricting to the domain $0 < \gamma < 1$ of small γ , it is only the s-wave which contributes to $\sigma_{v^2 < 0}$. Fitting the remaining open parameter as $p_{0,0} = 10$ we obtain for $\gamma = 0.9$ the curves of figure 3. They show an oscillatory dependence on p which becomes infinitely fast as $p \rightarrow 0$. The contributions to $\sigma_{v^2 < 0}$ with $m \neq 0$ which appear for larger values of γ show the same behaviour.

6. Bound states

Bound states in the attractive potential $U(x) = -\kappa^2/x^2$ have been considered long ago in [6]. We generalize these results by including the AB potential ($\beta \neq 0$). Bound states can only be found for values of m with $v^2 < 0$. The energy \mathcal{E} is negative so that we have $p^2 = -2M\mathcal{E}$.

The corresponding radial equation

$$R_m''(\rho) + \frac{1}{\rho} R_m'(\rho) + \frac{\mu^2}{\rho^2} R_m(\rho) - p^2 R_m(\rho) = 0 \quad (43)$$

has the general solution

$$R_m(\rho) = a_m I_{-i\mu}(p\rho) + b_m I_{i\mu}(p\rho). \quad (44)$$

Because the wavefunctions of bound states have to decrease for $\rho \rightarrow \infty$, this leads to a real $R_m(\rho)$

$$R_m(\rho) = ic_m [I_{i\mu}(p\rho) - I_{-i\mu}(p\rho)] = \frac{2c_m}{\pi} \sinh \pi \mu K_{i\mu}(p\rho) \quad (45)$$

with $K_\nu(z)$ being the Bessel function of imaginary argument. For $\rho \rightarrow 0$ we find

$$\begin{aligned} R_m(\rho) &\sim ic_m \left[\left(\frac{p\rho}{2} \right)^{i\mu} \frac{1}{\Gamma(1+i\mu)} - \left(\frac{p\rho}{2} \right)^{-i\mu} \frac{1}{\Gamma(1-i\mu)} \right] \\ &= \frac{ic_m}{\sqrt{\Gamma(1+i\mu)\Gamma(1-i\mu)}} \left[e^{i\mu \ln(p\rho/2) - i\xi} - e^{-i\mu \ln(p\rho/2) + i\xi} \right] \\ &= -\frac{2c_m \sinh \pi \mu}{\pi \mu} \sin \left(\mu \ln \frac{p\rho}{2} - \xi \right) \end{aligned} \quad (46)$$

where $\xi := \arg \Gamma(1+i\mu)$. To obtain self-adjointness we again demand orthogonality of the wavefunctions for different values of p :

$$\begin{aligned} (p^2 - p'^2) \int_0^\infty \rho \, d\rho R_m(p'\rho) R_m(p\rho) &= \int_0^\infty \frac{d}{d\rho} [R_m(p'\rho) R_m'(p\rho) - R_m(p\rho) R_m'(p'\rho)] \\ &= \rho [R_m(p\rho) R_m'(p'\rho) - R_m(p'\rho) R_m'(p\rho)] \Big|_{\rho=0} = 0. \end{aligned} \quad (47)$$

This condition will be fulfilled if we have for given m

$$\rho [R_m(p\rho) R_m'(p'\rho) - R_m(p'\rho) R_m'(p\rho)] \Big|_{\rho=0} = \frac{4c_m^2 \sinh^2 \pi \mu}{\pi^2 \mu} \sin \left(\mu \ln \frac{p}{p'} \right) = 0$$

or

$$\frac{p'}{p} = e^{\pi n/\mu} \quad \text{i.e.} \quad p_{m,n} = p_{m,0} e^{-\pi n/\mu} \quad \mathcal{E}_{m,n} = \mathcal{E}_{m,0} e^{-2\pi n/\mu} \quad \mathcal{E}_{m,0} = -\frac{p_{m,0}^2}{2M} \quad (48)$$

with $n = 0, \pm 1, \pm 2, \dots$. Accordingly there are an infinite number of bound states related to a point spectrum of energy values. It extends from $-\infty$ to zero which is a point of accumulation. Because of the open parameters $p_{m,0}$ there appears a not specified energy scale. As for scattering we therefore have the problem of fixing the parameters $p_{m,0}$ in discussing models which lead to equation (43) in some limiting case.

7. Conclusions

We have considered quantum mechanical elastic scattering in the presence of two cylindrically symmetric electric and magnetic potentials which are singular for $\rho = 0$. The approach was based on a decomposition with regard to partial waves with angular quantum number m . The underlying Hamiltonian has been made self-adjoint by a self-adjoint extension which fixed the form of the Schrödinger solutions for given m . It thereby turned out that for fixed m and

given particle momentum p the wavefunctions belonging to the self-adjoint extension contain open parameters. This is the case for quantum numbers m with $v^2 := (m - \beta)^2 - \gamma^2 < 1$, where β and γ are the magnetic-field and charge parameters. To specify the quantum states it is therefore not enough to fix the strength of the singular potentials by these physical parameters. This goes back to the fact that many different physical situations with non-singular electric and magnetic potentials (for example, $\rho = 0$ may be excluded for some reason) can lead in some limit to the radial differential equation treated above. Dependent on the situation, the open parameters may thereby be fixed. It has not been answered above what type of physical situations these are. We showed that a cylinder with hard a core is not a representative.

Independent of the answer to this question one can study the elastic scattering in these singular potentials. We have shown how the AB contribution dominates the behaviour for small scattering angles. The dependence on the charge parameter γ and the momentum p is discussed.

There is a lesson to be learned: the open parameters appear mode by mode. The angular momentum quantum number m is essentially an impact parameter. One could think of avoiding the singularities by modifying the physical situation for certain modes (for example, by introducing absorption) and keeping other modes unaltered. It may then be that for these unaltered modes one is again confronted with the problem of open parameters which are to be fixed by some physical argument.

Appendix A. Self-adjoint extension approach

In this appendix we present very briefly the ideas and results concerning the self-adjoint extension approach to the problem under consideration. We do not dwell on the rigorous formulations and details. A more extended version will be presented elsewhere.

The H of equation (3) is only a formal differential expression. It must be supplemented by specifying the dense domain $\mathcal{D}_H \subset \mathcal{H} = \mathcal{L}_2(\mathbb{R}_3)$, $\overline{\mathcal{D}_H} = \mathcal{H}$ on which H of equation (3) defines a self-adjoint operator: $H = H^\dagger$. This is the problem to be solved. In contrast to the case of regular potentials, in this case the differential expression (3) does not define an operator at all (even a self-adjoint one) on the natural set of sufficiently smooth functions. The reason is the strong singularity of the potentials (1) and (2) on the z -axis.

The differential expression (3) naturally defines an operator H_0 on the domain \mathcal{D}_{H_0} of functions ψ_0 which are sufficiently smooth and sufficiently rapidly vanishing when approaching the z -axis (and infinity). The operator H_0 is evidently *symmetric* (Hermitian)¹ but it is *not self-adjoint*: $H_0 \subset H_0^\dagger$. The adjoint operator H_0^\dagger is defined by the same differential expression (3) but on the different domain $\mathcal{D}_{H_0^\dagger} \subset \mathcal{H}$, which is the extension of \mathcal{D}_{H_0} with $\mathcal{D}_{H_0} \subset \mathcal{D}_{H_0^\dagger}$. The functions ψ of $\mathcal{D}_{H_0^\dagger}$ may have singularities on the z -axis².

The physically meaningful self-adjoint Hamiltonian $H = H^\dagger$ can be defined as a self-adjoint extension of the symmetric operator H_0 (and consequently a restriction of H_0^\dagger): $H_0 \subset H \subset H_0^\dagger$ and therefore $\mathcal{D}_{H_0} \subset \mathcal{D}_H \subset \mathcal{D}_{H_0^\dagger}$.

Now we recall the basic points of the self-adjoint extension theory [15]. For an arbitrary symmetric operator $H_0 \subseteq H_0^\dagger$ the possibility to have a self-adjoint extension depends on the

¹ In fact, the equality $(\psi_0^{(1)}, H_0 \psi_0^{(2)}) = (H_0 \psi_0^{(1)}, \psi_0^{(2)})$, $\psi_0^{(1)}, \psi_0^{(2)} \in \mathcal{D}_{H_0}$ holds true because integration by parts is possible without any boundary terms at the z -axis (and at infinity).

² The integration by parts in $(\psi, H_0 \psi_0)$ is possible without boundary terms because of the vanishing of ψ_0 on the z -axis to give $(H_0^\dagger \psi, \psi_0)$.

so-called deficiency subspaces \mathcal{H}_+ and \mathcal{H}_- which are the eigensubspaces of the operator H_0^\dagger corresponding to the respective eigenvalues $+i$ and $-i$ ³: $\mathcal{H}_\pm = \{\psi_\pm : \psi_\pm \in \mathcal{D}_{H_0^\dagger}, H_0^\dagger \psi_\pm = \pm i \psi_\pm\}$. The extension does exist if the dimensions of these subspaces, the so-called deficiency indices $n_\pm = \dim \mathcal{H}_\pm$, coincide: $n_+ = n_-$.

If \mathcal{H}_\pm is trivial, i.e. for $n_+ = n_- = 0$, the self-adjoint extension is unique. It is the closure of $H_0 : H = \overline{H_0} = H_0^\dagger$. In this case the symmetric H_0 is called essentially self-adjoint. If \mathcal{H}_\pm is non-trivial, i.e. for $n_+ = n_- = n > 0$, the self-adjoint extension is non-unique. The concrete self-adjoint extension H_U is specified by a unitary mapping $U : \mathcal{H}_+ \rightarrow \mathcal{H}_-$ with $\mathcal{D}_{H_U} = \{\psi_U : \psi_U = \psi_0 + \psi_+ + U\psi_+, \forall \psi_0 \in \mathcal{D}_{H_0}, \forall \psi_+ \in \mathcal{H}_+; U\psi_+ \in \mathcal{H}_-\}$. H_U coincides with H_0^\dagger on $\mathcal{D}_{H_U} : H_U \psi_U = H_0 \psi_0 + i\psi_+ - iU\psi_+$ ⁴. Therefore, if $n_+ = n_- = n > 0$ there is a manifold of self-adjoint extensions of the given symmetric operator H_0 which is the group $U(n)$. In this case additional considerations are necessary to select a particular extension.

In our case we have $n_+ = n_- = \infty$. It is easy to verify that the functions

$$\psi_\pm = e^{im\varphi} \int_{-\infty}^{\infty} dk e^{ikz} \tilde{\psi}_\pm(k) H_v^{(1)}(\sqrt{\pm 2iM - k^2} \rho)$$

are normalizable solutions of the respective differential equations $H_0^\dagger \psi_\pm = \pm i \psi_\pm$ where $\tilde{\psi}_\pm(k)$ are arbitrary functions with a compact support. $v^2 \equiv v^2(m) = (m - \beta)^2 - \gamma^2 < 1$ and may be negative, $m = 0, \pm 1, \dots$, $H_v^{(1)}$ is the Hankel function and $\text{Im} \sqrt{\pm 2iM - k^2} > 0$ ⁵. Therefore, there exists an infinite-dimensional $U(\infty)$ manifold of self-adjoint extensions H_U of the symmetric operator H_0 in question.

In order to specify further concrete extensions we impose as an additional demand that H_U be invariant under translations along and rotations around the z -axis as has been done in the sections above. H_0 and H_0^\dagger show this invariance but H_U does not, in general. The reason is that although H_0 and H_0^\dagger are invariant under these transformations, the mapping U may not respect this invariance. Such a possibility corresponds to the case where the potential has additional singularities of the $\delta(z)$ -type that break the translation and rotation invariance. The operator H_0 remains the same.

The requirement of this invariance greatly restricts the arbitrariness in the choice of U but, in fact, does still not eliminate it. This is in agreement with the results of the ‘pragmatic approach’ of section 3. This can be demonstrated as follows: if we require the rotation and translation invariance with respect to the z -axis, then the problem of a self-adjoint extension of H_0 reduces to the problem of self-adjoint extensions of the radial symmetric operators

$$H_{m0} = -\frac{d^2}{d\rho^2} - \frac{1}{\rho} \frac{d}{d\rho} + \frac{v^2}{\rho^2} \quad (\text{A1})$$

in the radial Hilbert space $\mathcal{L}_2(R_+, \rho d\rho)$ for each given quantum number $m = 0, \pm 1, \dots$. These operators are defined on functions $R_{m0}(\rho)$ which vanish sufficiently rapidly for $\rho \rightarrow 0$. The deficiency indices of H_{m0} are fitted by $v^2 = v^2(m)$. It is easy to see that if $v^2 \geq 1$ we have $n_+ = n_- = 0$. Consequently, H_{m0} is essentially self-adjoint and there is no arbitrariness in defining the self-adjoint extension H_m . However, if $v^2 < 1$, we have $n_+ = n_- = 1$, and for

³ $\pm i$ can be replaced by any complex numbers $\zeta, \bar{\zeta}$, $\text{Im} \zeta > 0$

⁴ If $\{\psi_{\pm k}\}$ with $k = 1, 2, \dots, n$ are respective orthonormal bases in \mathcal{H}_\pm , then U and \mathcal{H}_U are specified by a unitary matrix $U_{kl} : \mathcal{D}_{H_U} = \{\psi_U : \psi_U = \psi_0 + \sum_{k=1}^n c_k \psi_{+,k} + \sum_{k,l=1}^n c_k U_{kl} \psi_{-,l}\}$, $H_U \psi_U = H_0 \psi_0 + i \sum_{k=1}^n c_k \psi_{+,k} - i \sum_{k,l=1}^n c_k U_{kl} \psi_{-,l}$.

⁵ It is also evident that if ψ_+ satisfies the equation $H_0^\dagger \psi_+ = +i\psi_+$, then $\psi_- = \overline{\psi_+}$ satisfies the equation $H_0^\dagger \psi_- = -i\psi_-$.

each m we have a one-parameter $U(1)$ manifold of self-adjoint extensions H_m . We can specify the domains \mathcal{D}_{H_m} of these extensions in terms of the boundary behaviour of the functions $R_m(\rho)$ for $\rho \rightarrow 0$. For $v^2 \geq 1$ we have $R_m(0) = 0$. If $0 < v^2 < 1$ ⁶ a concrete self-adjoint extension H_{ml_m} is specified by the (asymptotic) boundary condition of the form

$$R_m(\rho) := R_{ml_m}(\rho) \sim \rho^\mu + l_m \rho^{-\mu} \quad \rho \rightarrow 0 \quad (\text{A2})$$

with particular parameters $l_m \in \mathbb{R}_1$. If $v^2 < 0$, the concrete self-adjoint extension $H_{m\vartheta_m}$ is defined by the (asymptotic) boundary condition of the form

$$R_m(\rho) := R_{m\vartheta_m}(\rho) \sim \rho^{i\mu} + e^{i\vartheta_m} \rho^{-i\mu} \quad \rho \rightarrow 0 \quad (\text{A3})$$

with the particular parameters $\vartheta_m, 0 \leq \vartheta_m < 2\pi$. The same behaviour has been obtained in section 3.

To sum up, we can determine the eigenfunctions of the self-adjoint Hamiltonian for the problem under consideration by separating variables in cylindrical coordinates and imposing the boundary conditions stated above on the radial wavefunctions at $\rho = 0$ in fixing λ_m and θ_m . This specifies the Hamiltonian as a self-adjoint operator.

The arbitrariness in defining the Hamiltonian is finite-parametric. The number of parameters agrees with the number of the orbital quantum numbers $m = 0, \pm 1, \dots$, for which $v^2 = (m - \beta)^2 - \gamma^2 < 1$. The further reduction of this finite-parametric arbitrariness requires physical arguments in addition (to the rotation and translation invariance).

Appendix B. Some integrals with the Bessel functions

Using formula (5.53) of [8] we obtain

$$\begin{aligned} \int_0^\infty J_\nu(p'\rho) J_\nu(p\rho) \rho \, d\rho &= \lim_{R \rightarrow \infty} \frac{1}{p^2 - p'^2} \left[p' \rho J_\nu(p\rho) J_{\nu-1}(p'\rho) - p \rho J_{\nu-1}(p\rho) J_\nu(p'\rho) \right] \Big|_0^R \\ &= \lim_{R \rightarrow \infty} \frac{2}{\pi(p^2 - p'^2)} \left[\sqrt{\frac{p'}{p}} \cos\left(pR - \frac{\pi}{2}\nu - \frac{\pi}{4}\right) \cos\left(p'R - \frac{\pi}{2}(\nu-1) - \frac{\pi}{4}\right) \right. \\ &\quad \left. - \sqrt{\frac{p}{p'}} \cos\left(pR - \frac{\pi}{2}(\nu-1) - \frac{\pi}{4}\right) \cos\left(p'R - \frac{\pi}{2}\nu - \frac{\pi}{4}\right) \right] \\ &= \frac{1}{\pi\sqrt{pp'}} \lim_{R \rightarrow \infty} \left[\frac{\sin(p-p')R}{p-p'} - \frac{\cos(p+p')R}{p+p'} \cos \pi p \right. \\ &\quad \left. - \frac{\sin(p+p')R}{p+p'} \sin \pi p \right] \\ &= \frac{\delta(p-p')}{\sqrt{pp'}} \end{aligned} \quad (\text{B1})$$

⁶ The case of $\nu = 0$ requires special consideration.

and

$$\begin{aligned}
& \int_0^\infty J_{-\nu}(p'\rho) J_\nu(p\rho) \rho \, d\rho \\
&= \lim_{R \rightarrow \infty} \frac{1}{p^2 - p'^2} [p'\rho J_\nu(p\rho) J_{-\nu-1}(p'\rho) - p\rho J_{-\nu-1}(p\rho) J_{-\nu}(p'\rho) \\
&\quad + 2\nu J_\nu(p\rho) J_{-\nu}(p'\rho)] \Big|_0^R \\
&= \lim_{R \rightarrow \infty} \frac{2}{\pi(p^2 - p'^2)} \left[\sqrt{\frac{p'}{p}} \cos\left(pR - \frac{\pi}{2}\nu - \frac{\pi}{4}\right) \cos\left(p'R + \frac{\pi}{2}(\nu+1) - \frac{\pi}{4}\right) \right. \\
&\quad \left. - \sqrt{\frac{p}{p'}} \cos\left(pR - \frac{\pi}{2}(\nu-1) - \frac{\pi}{4}\right) \cos\left(p'R + \frac{\pi}{2}\nu - \frac{\pi}{4}\right) \right] \\
&\quad - \frac{2}{p^2 - p'^2} \left(\frac{p}{p'}\right)^\nu \left[\frac{1}{\Gamma(\nu+1)\Gamma(-\nu)} - \frac{1}{\Gamma(\nu)\Gamma(-\nu+1)} \right. \\
&\quad \left. + \frac{2\nu}{\Gamma(\nu+1)\Gamma(-\nu+1)} \right] \\
&= \frac{1}{\pi\sqrt{pp'}} \lim_{R \rightarrow \infty} \left[\frac{\sin[(p-p')R - \pi\nu]}{p-p'} - \frac{\cos(p+p')}{p+p'} \right] \\
&\quad - \frac{2}{(p^2 - p'^2)} \left(\frac{p}{p'}\right)^\nu \frac{1}{\Gamma(\nu+1)\Gamma(-\nu)} \\
&= \frac{\delta(p-p')}{\sqrt{pp'}} \cos\pi\nu + \frac{2 \sin\pi\nu}{\pi(p^2 - p'^2)} \left(\frac{p}{p'}\right)^\nu. \tag{B2}
\end{aligned}$$

We have thereby made use of

$$\begin{aligned}
\int_0^\infty e^{ixt} dt &= \pi \delta_+(x) = \pi \delta(x) + iP \left(\frac{1}{x}\right) \\
&= \lim_{R \rightarrow \infty} \int_0^R e^{ixt} dt = \lim_{R \rightarrow \infty} \left(\frac{\sin xR}{x} + i \frac{1 - \cos xR}{x} \right) \tag{B3}
\end{aligned}$$

so that we may write

$$\delta(x) = \lim_{R \rightarrow \infty} \frac{\sin xR}{\pi x} \quad P \left(\frac{1}{x}\right) = \lim_{R \rightarrow \infty} \frac{1 - \cos xR}{x}. \tag{B4}$$

Acknowledgments

VS thanks Professor J Audretsch and the members of his group for the friendly atmosphere and the hospitality at the University of Konstanz. This work was supported by the Deutsche Forschungsgemeinschaft. BV is grateful for the support from RFBR (grant no 99-01-00376).

References

- [1] Aharonov Y and Bohm D 1959 *Phys. Rev.* **119** 485
- [2] Audretsch J, Jasper U and Skarzhinsky V D 1995 *J. Phys. A: Math. Gen.* **28** 2359

- [3] Audretsch J, Jasper U and Skarzhinsky V D 1996 *Phys. Rev. D* **53** 2178–89
Skarzhinsky V D, Audretsch J and Jasper U 1996 *Phys. Rev. D* **53** 2190–200
- [4] Audretsch J and Skarzhinsky V D 1998 *Phys. Lett. A* **241** 7
- [5] Audretsch J and Skarzhinsky V D 1999 *Phys. Rev. A* **60** 1854
- [6] Case K M 1950 *Phys. Rev.* **80** 797
- [7] Denschlag J and Schmiedmayer J 1997 *Europhys. Lett.* **38** (6) 405
- [8] Gradshteyn I S and Ryzhik I M 1980 *Table of Integrals, Series and Products* (New York: Academic) p 634
- [9] Hagen C R 1996 *Phys. Rev. Lett.* **77** 1656
- [10] Jackiw R 1972 *Phys. Today* **25** January, 23
- [11] Leonhardt U and Wilkens M 1998 *Europhys. Lett.* **42** 365
- [12] Novak S *et al* 1998 *Phys. Rev. Lett.* **81** 5792
- [13] Olariu S and Popescu I I 1985 *Rev. Mod. Phys.* **47** 339
- [14] Peshkin M and Tonomura A 1989 *The Aharonov–Bohm Effect* (Berlin: Springer)
- [15] Reed M and Simon B 1975 *Methods of Modern Mathematical Physics vol II, Fourier Analysis, Self-Adjointness* (New York: Academic)
Akhiezer N I and Glazman I M 1993 *Theory of Linear Operators in Hilbert Space* (New York: Dover)
- [16] Wei H, Han R and Wei X 1995 *Phys. Rev. Lett.* **75** 2071
- [17] Wilkens M 1994 *Phys. Rev. Lett.* **72** 5