

Tomograms and other transforms: a unified view

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2001 J. Phys. A: Math. Gen. 34 8321

(<http://iopscience.iop.org/0305-4470/34/40/309>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.98

The article was downloaded on 02/06/2010 at 09:20

Please note that [terms and conditions apply](#).

Tomograms and other transforms: a unified view

M A Man'ko^{1,2}, V I Man'ko^{1,2} and R Vilela Mendes^{2,3}

¹ P N Lebedev Physical Institute, Leninskii Prospect 53, Moscow 119991, Russia

² Zentrum für interdisziplinäre Forschung, Universität Bielefeld, Wellenberg 1, 33615 Bielefeld, Germany

³ Grupo de Física Matemática, Complexo Interdisciplinar, Universidade de Lisboa, Av. Gama Pinto, 2-P1699 Lisboa Codex, Portugal

E-mail: mmanko@sci.lebedev.ru, manko@sci.lebedev.ru and vilela@cii.fc.ul.pt

Received 30 April 2001

Published 28 September 2001

Online at stacks.iop.org/JPhysA/34/8321

Abstract

A general framework is presented which unifies the treatment of wavelet-like, quasidistribution and tomographic transforms. Explicit formulae relating the three types of transforms are obtained. The case of transforms associated with the symplectic and affine groups is treated in some detail. Special emphasis is given to the properties of the scale–time and scale–frequency tomograms. Tomograms are interpreted as a tool to sample the signal space by a family of curves or as the matrix element of a projector.

PACS numbers: 03.65.Wj, 03.65.Ta, 42.65.Re

1. Introduction

Several types of integral transforms [1, 2] are used for signal processing in physics, engineering, medicine, etc. In addition to the traditional Fourier analysis [3], wavelet analysis has been extensively developed in the last two decades [4–6]. These two types of transforms are linear transforms. In addition, the Wigner–Ville quasidistribution [7, 8], a bilinear transform, provides optimal energy resolution in the joint time–frequency domain. A joint time–frequency description of signals is important, because in many applications (biomedical, seismic, radar, etc) the signals are of finite (sometimes very short) duration. However, the oscillating cross-terms in the Wigner–Ville quasidistribution make the interpretation of this transform a difficult matter. Even if the average of the cross-terms is very small, their amplitude may be greater than the signal in time–frequency domains that carry no physical information. To profit from the time–frequency energy resolution of the bilinear transforms while controlling the cross-terms problem, modifications to the Wigner–Ville transform have been proposed. Transforms in the Cohen class [9] make a two-dimensional filtering of the Wigner–Ville quasidistribution and the Gabor spectrogram [10] is a truncated version of this quasidistribution.

Recently, a new type of strictly positive bilinear transforms has been proposed, namely the Radon–Wigner transform [11, 12] or, more generally, the noncommutative tomography [13] which, in addition to the time–frequency domain, also applies to other noncommutative pairs like time–scale, frequency–scale, etc. It is this last class of transforms that will be called *tomograms* in this paper. The tomograms are strictly positive probability densities, provide a full characterization of the signal and are robust in the presence of noise.

Developing a general operator scheme, we show how linear transforms, like Fourier transforms or wavelets, are related to the quasidistributions and the tomograms. Explicit general formulae are derived relating the three types of transforms. The time–frequency plane is then briefly discussed because most of the material has been treated before [13]. Special emphasis is, however, given to the transforms associated with the affine group, namely to the time–scale and frequency–scale tomograms. To clarify the physical meaning of the tomograms, we also propose an interpretation as a sampling of the signal space by families of phase-space curves or as the action of a projection operator.

2. Wavelet-like transforms, quasidistributions and tomograms

We present a unified general construction of three types of transforms used in signal analysis. The first class consists of wavelet-type transforms, the second of quasidistributions, and the third class of tomographic transforms. Quasidistributions are transforms like the Wigner–Ville one [7, 8] or the P -quasidistributions of Glauber and Sudarshan [14, 15]. Husimi–Kano positive quasidistributions [16, 17] will be discussed as well. These types of quasidistributions are unified in the class of s -ordered quasidistributions [18].

In quantum mechanics, quasidistributions describe a quantum state in terms of phase-space quasiprobability densities. In signal analysis, quasidistributions describe the structure of analytic signals in the time–frequency plane. There also exist quasidistributions characterizing the signal structure in the time–scale plane [19–22]. We refer to quasiprobability densities because the corresponding functions are not conventional probabilities, being either complex or nonpositive. In the case of positive quasiprobabilities like the Husimi–Kano function, an interpretation as a joint probability distribution is also not possible because the two arguments of the function are not simultaneously measurable random variables. The corresponding observables do not commute and the uncertainty relation prevents the existence of a joint probability distribution function for noncommuting observables. Time \hat{t} and frequency $\hat{\omega}$, time and scale $\frac{1}{2}(\hat{\omega}\hat{t} + \hat{t}\hat{\omega})$ or frequency and scale are common examples of such pairs of noncommuting observables.

The general setting for our construction is as follows.

Signals $f(t)$ are considered to be vectors $|f\rangle$ belonging to a dense nuclear subspace \mathcal{N} of a Hilbert space \mathcal{H} with dual space \mathcal{N}^* (and the canonical identification $\mathcal{N} \subset \mathcal{N}^*$). $\{U(\alpha) : \alpha \in I\}$ is a family of operators defined on \mathcal{N}^* , and *a fortiori* on \mathcal{N} by the canonical identification $\mathcal{N} \subset \mathcal{N}^*$. In many cases, the family of operators $U(\alpha)$ generates a unitary group. However, this is not a necessary condition for the consistency of the formalism, provided the completeness conditions discussed below are satisfied.

In this setting, three types of transforms are defined. Consider a reference vector $h \in \mathcal{N}^*$ chosen in such a way that the linear span of $\{U(\alpha)h \in \mathcal{N}^* : \alpha \in I\}$ is dense in \mathcal{N}^* . This means, in particular, that, out of the set $\{U(\alpha)h\}$, a complete set of vectors can be chosen to serve as a basis. Two of the transforms considered are

$$W_f^{(h)}(\alpha) = \langle U(\alpha)h | f \rangle \quad (1)$$

$$Q_f(\alpha) = \langle U(\alpha)f | f \rangle. \quad (2)$$

If $U(\alpha)$ is a unitary operator generated by $B(\vec{\alpha}) = \alpha_1 t + i\alpha_2 d/dt$ and h is a (generalized) eigenvector of the time-translation operator, $W_f^{(h)}(\alpha)$ becomes a Fourier transform. With the same $B(\vec{\alpha})$ plus the parity operator, $Q_f(\alpha)$ would be the Wigner–Ville transform. Similarly, for $B(\vec{\alpha}) = \alpha_1 D + i\alpha_2 d/dt$, where D is the dilation operator

$$D = -\frac{1}{2} \left(it \frac{d}{dt} + i \frac{d}{dt} t \right)$$

$W_f^{(h)}(\alpha)$ is a wavelet transform and $Q_f(\alpha)$ the Bertrand transform.

We will denote the transforms of the $W_f^{(h)}$ -type as *wavelet-type* transforms and those of the Q_f -type as *quasidistribution* transforms.

In general, if $U(\alpha)$ are unitary operators, by Stone's theorem, there are self-adjoint operators $B(\alpha)$ such that

$$W_f^{(h)}(\alpha) = \langle h | e^{iB(\alpha)} | f \rangle \quad (3)$$

$$Q_f^{(B)}(\alpha) = \langle f | e^{iB(\alpha)} | f \rangle. \quad (4)$$

In this case, because $B(\alpha)$ has a real valued spectrum, another transform may be defined, namely

$$M_f^{(B)}(X) = \langle f | \delta(B(\alpha) - X) | f \rangle. \quad (5)$$

The delta operator $\delta(B(\alpha) - X)$ is to be interpreted in terms of the spectral theorem for unbounded operators [23]:

$$\delta(B(\alpha) - X) = \int_{-\infty}^{\infty} \delta(\lambda - X) dP_\lambda^{(\alpha)} \quad (6)$$

$dP_\lambda^{(\alpha)}$ being the projection-valued measure associated with $B(\alpha)$:

$$B(\alpha) = \int_{-\infty}^{\infty} \lambda dP_\lambda^{(\alpha)}. \quad (7)$$

Equation (5) defines what we call the tomographic transform or *tomogram*. In contrast to the quasiprobabilities, the transform $M_f^{(B)}(\alpha)$ is positive and, as we will see below, it can be correctly interpreted as a probability distribution. Therefore, it benefits from the properties of the bilinear transforms, without being plagued by the interpretation ambiguities associated with the quasidistribution transforms.

For a normalized vector $|f\rangle$,

$$\langle f | f \rangle = 1 \quad (8)$$

the tomogram is a normalized function:

$$\int M_f^{(B)}(X) dX = 1 \quad (9)$$

and, therefore, it may be interpreted as a probability distribution for the random variable X corresponding to the observable defined by the operator $B(\alpha)$. The tomogram is a homogeneous function:

$$M_f^{(B/p)}(X) = |p| M_f^{(B)}(pX). \quad (10)$$

The three classes of transforms are mutually related:

$$M_f^{(B)}(X) = \frac{1}{2\pi} \int Q_f^{(kB)}(\alpha) e^{-ikX} dk \quad (11)$$

and

$$Q_f^{(B)}(\alpha) = \int M_f^{(B/p)}(X) e^{ipX} dX. \quad (12)$$

Wavelet-type transforms, quasidistributions, and tomograms are related by the formulae

$$Q_f^{(B)}(\alpha) = W_f^{(f)}(\alpha) \quad (13)$$

$$W_f^{(h)}(\alpha) = \frac{1}{4} \int e^{iX} [M_{f_1}^{(B)}(X) - iM_{f_2}^{(B)}(X) - M_{f_3}^{(B)}(X) + iM_{f_4}^{(B)}(X)] dX \quad (14)$$

where

$$\begin{aligned} |f_1\rangle &= |h\rangle + |f\rangle & |f_3\rangle &= |h\rangle - |f\rangle \\ |f_2\rangle &= |h\rangle + i|f\rangle & |f_4\rangle &= |h\rangle - i|f\rangle. \end{aligned}$$

Another important case concerns operators $U(\alpha)$, which can be represented in the form

$$U(\alpha) = e^{ib(\alpha)} P_h e^{-ib(\alpha)} \quad (15)$$

P_h being a projector on a reference vector $|h\rangle$. This creates a quasidistribution of the Husimi–Kano type

$$H_f^{(b)}(\alpha) = \langle f|U(\alpha)|f\rangle.$$

In the following sections, we show how known examples of wavelet-like and quasidistribution transforms are described within the framework presented above. We will consider quasidistributions such as Wigner–Ville [7, 8], Bertrand and Bertrand [19] and Husimi–Kano [16, 17]. We reformulate the standard wavelet analysis in terms of this general scheme using an operator $U(\vec{\alpha})$ belonging to the two-dimensional affine group. Tomographic transform schemes for time–frequency, time–scale and frequency–scale pairs [13] will be studied within the framework of the general approach. Inversion formulae are obtained for the tomograms as well as the explicit connection of the wavelet transform to the time–scale and frequency–scale tomograms.

3. Time–frequency transforms

Here we discuss the case where the operator $B(\vec{\alpha})$, $\vec{\alpha} = (\mu, \nu)$ is

$$B^{(S)}(\alpha) = \mu\hat{t} + \nu\hat{\omega}$$

(with $\hat{\omega} = -i\partial/\partial t$) or is equal to this one plus a parity operator. The wavelet-like transform in this case is just the Fourier transform and we discuss only the tomograms and quasidistributions.

For the tomogram, that is,

$$M_f^{(S)}(X, \mu, \nu) = \langle f|\delta(\mu\hat{t} + \nu\hat{\omega} - X)|f\rangle \quad (16)$$

using equation (6) or, equivalently, a Fourier transform representation of the delta-operator, one obtains [13]

$$M_f^{(S)}(X, \mu, \nu) = \frac{1}{2\pi|\nu|} \left| \int \exp\left[\frac{i\mu t^2}{2\nu} - \frac{itX}{\nu}\right] f(t) dt \right|^2. \quad (17)$$

The tomogram (17) is normalized if $\langle f|f\rangle = 1$

$$\int M_f^{(S)}(X, \mu, \nu) dX = 1.$$

From the tomogram $M_f^{(S)}(X, \mu, \nu)$, the signal $f(t)$ may be recovered up to a phase

$$f(t) f^*(0) = \frac{1}{2\pi} \int M_f^{(S)}(X, \mu, t) \exp \left[i \left(X - \mu \frac{t}{2} \right) \right] dX d\mu. \quad (18)$$

According to the general scheme, the corresponding quasidistribution is

$$Q_f^{(S)}(\mu, \nu) = \langle f | e^{iB^{(S)}(\mu, \nu)} | f \rangle = \int M_f^{(S)}(X, \mu, \nu) e^{iX} dX \quad (19)$$

or

$$Q_f^{(S)}(\mu, \nu) = \int f^* \left(t - \frac{\nu}{2} \right) f \left(t + \frac{\nu}{2} \right) e^{i\mu t} dt. \quad (20)$$

This quasidistribution is called the ambiguity function in the signal processing literature [24].

The tomogram (17) and this quasidistribution are related by equations (11) and (12). The tomogram (17) is also related to another quasidistribution, namely to the Wigner–Ville quasidistribution $WV(\tau, \omega)$ [7, 8] by

$$M_f^{(S)}(X, \mu, \nu) = \int \exp \left[-ik(X - \mu\omega - \nu\tau) \right] WV(\tau, \omega) \frac{dk d\omega d\tau}{(2\pi)^2}. \quad (21)$$

The Wigner–Ville quasidistribution is given by the formula

$$WV(\tau, \Omega) = \int f \left(\tau + \frac{u}{2} \right) f^* \left(\tau - \frac{u}{2} \right) e^{-i\Omega u} du. \quad (22)$$

The unitary operator $U(\tau, \Omega)$, which determines the Wigner–Ville quasidistribution by

$$WV(\tau, \Omega) = \langle f | U^{(WV)}(\tau, \Omega) | f \rangle \quad (23)$$

is

$$U^{(WV)}(\tau, \Omega) = e^{2i(\Omega\hat{t} - \tau\hat{\omega})} e^{i\pi(\hat{t}^2 + \hat{\omega}^2 - 1)/2} \quad (24)$$

the generator being

$$B^{(WV)}(\tau, \Omega) = 2\tau\hat{\omega} - 2\Omega\hat{t} + \frac{\pi(\hat{t}^2 + \hat{\omega}^2 - 1)}{2}. \quad (25)$$

4. Wavelets and quasidistributions in the affine group

4.1. Wavelets

The wavelet transform of a signal $f(t)$ is a linear integral transform decomposing the signal into a set of basis functions:

$$W_f^{(A)}(s, \tau) = \int f(t) h_{s, \tau}^*(t) dt. \quad (26)$$

The wavelets $h_{s, \tau}(t)$ are kernel functions generated from a basic wavelet $h(\tau)$ by means of a translation and a rescaling ($-\infty < \tau < \infty, s > 0$):

$$h_{s, \tau}(t) = \frac{1}{\sqrt{s}} h \left(\frac{t - \tau}{s} \right). \quad (27)$$

Using the operator

$$U^{(A)}(\tau, s) = \exp(i\tau\hat{\omega}) \exp(i \log s D) \quad (28)$$

where

$$D = \frac{1}{2}(\hat{t}\hat{\omega} + \hat{\omega}\hat{t}) = \hat{\omega}\hat{t} + \frac{i}{2} \quad \text{with} \quad \hat{\omega} = -i \frac{\partial}{\partial t}$$

equation (27) can be represented in the form

$$h_{s,\tau}(t) = U^{(A)\dagger}(\tau, s)h(t). \quad (29)$$

For normalized $h(t)$, the wavelets $h_{s,\tau}(t)$ satisfy the normalization condition

$$\int |h_{s,\tau}(t)|^2 dt = 1. \quad (30)$$

The basic wavelet (reference vector) may have different forms, for example,

$$h(t) = \frac{1}{\sqrt{\pi}} e^{i\omega_0 t} e^{-t^2/2} \quad (31)$$

or

$$h(t) = (1 - t^2) e^{-t^2/2} \quad (32)$$

called the Mexican hat wavelet.

The inverse of the wavelet transform $W_f^{(A)}(s, \tau)$ is

$$f(t) = N_h^{-1} \int W_f^{(A)}(s, \tau) \frac{1}{\sqrt{s}} h\left(\frac{t - \tau}{s}\right) \frac{d\tau ds}{s^2} \quad (33)$$

with

$$N_h = \int \frac{|H(\omega)|^2}{|\omega|} d\omega \quad H(\omega) = \int h(t) e^{-i\omega t} dt. \quad (34)$$

One has the property

$$\int |W_f^{(A)}(s, \tau)|^2 \frac{d\tau ds}{s^2} = N_h \int |f(t)|^2 dt. \quad (35)$$

Let us consider the operator (28), with the parameters μ and ν being

$$\nu = \log s \quad \mu = \frac{\tau \log s}{s - 1}. \quad (36)$$

We obtain for the unitary operator (28)

$$U^{(A)\dagger}(\tau, s) \equiv U^{(A)}(\mu, \nu) = \exp(i\mu\hat{\omega} + i\nu D) \quad (37)$$

and the operator $B(\vec{\alpha})$ becomes

$$B_1^{(A)} = \mu\hat{\omega} + \nu D. \quad (38)$$

According to the general scheme with the operator (38), the wavelet transform (26) can be rewritten in the form (1)

$$W_f^{(A)}(s, \tau) = \langle h | U^{(A)}(\mu, \nu) | f \rangle. \quad (39)$$

The commutation relation for the operators $\hat{\omega}$ and D is

$$[\hat{\omega}, D] = -i\hat{\omega}. \quad (40)$$

The commutation relations (40) define the Lie algebra of the affine group. Therefore, the wavelet transform is the nondiagonal matrix element of a unitary irreducible representation of the affine group. The parameters μ and ν are the group parameters and the Hermitian operator $B_1^{(A)}$ belongs to the Lie algebra of the affine group.

4.2. Quasidistributions

The diagonal matrix elements of the irreducible representation determine a quasidistribution for the signal $f(t)$

$$Q_f^{(A)}(s, \tau) = \langle f | e^{i(\mu\hat{\omega} + \nu D)} | f \rangle \quad (41)$$

with μ and ν expressed in terms of shift and scaling parameters by equation (36).

Defining the action on the vector $|f\rangle$ as

$$e^{i(\mu\hat{\omega} + \nu D)/2} f(t) = F(\mu, \nu, t) \quad (42)$$

the quasidistribution (41) may be rewritten

$$Q_f^{(A)}(s, \tau) = \int F^*(-\mu, -\nu, t) F(\mu, \nu, t) dt. \quad (43)$$

Using the known kernel (Green function) of the operator (37) [25], one obtains for the quasidistribution (43) the following expression in terms of the parameters τ and s :

$$Q_f^{(A)}(s, \tau) = \int f^* \left(\frac{t - \tau/2}{\sqrt{s}} \right) f(\sqrt{s}(t + \tau/2)) dt. \quad (44)$$

The wavelet transform (26) may also be written in a similar form:

$$W_f^{(A)}(s, \tau) = \int h^* \left(\frac{t - \tau/2}{\sqrt{s}} \right) f(\sqrt{s}(t + \tau/2)) dt \quad (45)$$

as follows from equation (13).

The relation between wavelets and the corresponding affine group quasidistributions is also treated in [26], where the approach of [27, 28] is extended for general Lie groups.

4.3. Relation of wavelets to time–frequency tomograms

In view of (33) and (17), one relates the wavelet transform to the time–frequency tomogram:

$$M_f^{(S)}(X, \mu, \nu) = \frac{1}{2\pi |v| |N_h|^2} \left| \int W_f^{(A)}(s, \tau) \frac{h(t)}{|s|\sqrt{s}} \right. \\ \left. \times \exp \left\{ \frac{i\mu}{2v} (s^2 t^2 + 2st\tau + \tau^2) - \frac{iX}{v} (st + \tau) \right\} ds dt d\tau \right|^2. \quad (46)$$

The inverse transform is

$$W_f^{(A)}(s, \tau) = \frac{\sqrt{s}}{2\pi \mathcal{D}} \int h^*(t) M_f^{(S)}(X, \mu, st + \tau) \exp \left\{ iX - \frac{i\mu(st + \tau)}{2} \right\} dX d\mu dt \quad (47)$$

where

$$\mathcal{D} = \left[\frac{1}{2\pi} \int M_f^{(S)}(X, \mu, 0) e^{iX} dX d\mu \right]^{1/2}.$$

For the Mexican hat wavelet (32), with the admissibility condition

$$\int h(t) dt = 0$$

one has the explicit form

$$M_f^{(S)}(X, \mu, \nu) = \frac{1}{2\pi |v| |N_h|^2} \left| \int W_f^{(A)}(s, \tau) K_M(s, \tau, \mu, \nu, X) ds d\tau \right|^2 \quad (48)$$

where

$$K_M(s, \tau, \mu, \nu, X) = \sqrt{2\pi} \left(s - \frac{i\mu s^3}{\nu} \right)^{-3/2} \left[\frac{(s\mu\tau - sX)^2}{\nu^2 - i\mu\nu s^2} - \frac{i\mu s^2}{\nu} - 1 \right] \\ \times \exp \left[-\frac{(s\mu\tau - sX)^2}{2(\nu^2 - i\mu\nu s^2)} + \frac{i\mu\tau^2}{2\nu} - \frac{iX\tau}{\nu} \right]$$

and

$$N_h = \int |\omega|^3 e^{-\omega^2} d\omega = 1.$$

The inverse transform for the Mexican hat wavelet is

$$W_f^{(A)}(s, \tau) = \int R(s, \tau, X, \mu, \nu) M_f^{(S)}(X, \mu, \nu) dX d\mu d\nu \quad (49)$$

the kernel being

$$R(s, \tau, X, \mu, \nu) = \frac{1}{2\pi\mathcal{D}\sqrt{s}} \left[1 - \left(\frac{\tau - \nu}{s} \right)^2 \right] \exp \left[-\frac{i}{2} \left(\frac{\nu - \tau}{s} \right)^2 - \frac{i\mu\nu}{2} + iX \right].$$

4.4. Tomograms. Frequency–scale and time–scale

The tomogram associated with the operator $B_1^{(A)} = \mu\hat{\omega} + \nu D$ has been computed in [13]. It is

$$M_f^{(A\omega)}(s, \mu, \nu) = \frac{1}{2\pi|\nu|} \left| \int_{\omega>0} d\omega \frac{f(\omega)}{\sqrt{\omega}} \exp \left[-i \left(\frac{\mu}{\nu} \omega - \frac{s}{\nu} \log \omega \right) \right] \right|^2 \\ + \frac{1}{2\pi|\nu|} \left| \int_{\omega<0} d\omega \frac{f(\omega)}{\sqrt{|\omega|}} \exp \left[-i \left(\frac{\mu}{\nu} \omega - \frac{s}{\nu} \log |\omega| \right) \right] \right|^2 \quad (50)$$

$f(\omega)$ being the Fourier transform of the signal $f(t)$.

The tomogram corresponding to the operator

$$B_2^{(A)} = \mu\hat{t} + \nu D$$

was also computed, namely

$$M_f^{(A_t)}(s, \mu, \nu) = \frac{1}{2\pi|\nu|} \left| \int_{t>0} dt \frac{f(t)}{\sqrt{t}} \exp \left[i \left(\frac{\mu}{\nu} t - \frac{s}{\nu} \log t \right) \right] \right|^2 \\ + \frac{1}{2\pi|\nu|} \left| \int_{t<0} dt \frac{f(t)}{\sqrt{|t|}} \exp \left[i \left(\frac{\mu}{\nu} t - \frac{s}{\nu} \log |t| \right) \right] \right|^2. \quad (51)$$

The quasidistribution $Q_f^{(B)}(\mu, \nu)$ related to the above tomogram is constructed from the affine group. It was discussed in [19]. To compare signal analysis based on the time–scale tomograms and based on wavelets, it is useful to write the tomogram $M_f^{(A_t)}(s, \mu, \nu)$ in terms of the wavelet transform $W_f^{(A)}(s, \tau)$:

$$M_f^{(A_t)}(s_1, \mu, \nu) = \frac{1}{2\pi|\nu N_h^2|} \\ \times \left\{ \left| \int_{t>0, s>0} \frac{dt d\tau ds}{\sqrt{ts} s^2} h \left(\frac{t - \tau}{s} \right) W_f(s, \tau) \exp \left[i \left(\frac{\mu}{\nu} t - \frac{s_1}{\nu} \log t \right) \right] \right|^2 \right. \\ \left. + \left| \int_{t<0, s>0} \frac{dt d\tau ds}{\sqrt{|t|s} s^2} h \left(\frac{t - \tau}{s} \right) W_f(s, \tau) \exp \left[i \left(\frac{\mu}{\nu} t - \frac{s_1}{\nu} \log |t| \right) \right] \right|^2 \right\}. \quad (52)$$

The tomographic transform is invertible, that is, the signal may be recovered from the tomogram, namely

$$f(t) = \left[\int M_f^{(A_t)}(s, \mu, 1) e^{2is} d\mu ds \right]^{-1/2} \int M_f^{(A_t)}(s, \mu, 1) e^{2is - i\mu t} d\mu ds \quad (53)$$

and

$$f(t)f^*(t') = \frac{1}{4\pi^2} \int M_f^{(A_\omega)}(s, \mu, \nu) \times \exp \left\{ i \left[s - \mu\omega - \nu\omega \frac{t+t'}{2} + \omega(t-t') \right] \right\} |\omega| d\mu d\nu d\omega ds. \quad (54)$$

5. Meaning of the tomograms

5.1. Sampling the phase space

In the time–frequency space, the tomogram

$$M_f^{(S)}(X, \mu, \nu) = \langle f | \delta(\mu\hat{t} + \nu\hat{\omega} - X) | f \rangle \quad (55)$$

is the expectation value of an operator delta function in the state $|f\rangle$. The support of the delta function in (55) is a line in the time–frequency plane:

$$X = \mu t + \nu \omega. \quad (56)$$

Therefore, $M_f^{(S)}(X, \mu, \nu)$ is the marginal distribution of the variable X along this line in the time–frequency space. The line is rotated and rescaled when one changes the parameters μ and ν . In this way, the whole time–frequency space is sampled and the tomographic transform contains all information on the signal.

It is clear that, instead of marginals collected along straight lines on the time–frequency plane, one may use other curves to sample this space. For the tomograms associated with the affine group, one has

$$M_f^{(A_t)}(S_1, \mu, \nu) = \langle f | \delta \left(\mu\hat{t} + \nu \frac{\hat{t}\hat{\omega} + \hat{\omega}\hat{t}}{2} - S_1 \right) | f \rangle \quad (57)$$

and

$$M_f^{(A_\omega)}(S_2, \mu, \nu) = \langle f | \delta \left(\mu\hat{\omega} + \nu \frac{\hat{t}\hat{\omega} + \hat{\omega}\hat{t}}{2} - S_2 \right) | f \rangle. \quad (58)$$

The curves in the time–frequency space, defined by

$$S_1 = \mu t + \nu t \omega \quad S_2 = \mu \omega + \nu t \omega \quad (59)$$

are hyperbolas. This becomes clear using the system of coordinates

$$q = \frac{1}{\sqrt{2}}(t - \omega) \quad p = \frac{1}{\sqrt{2}}(t + \omega). \quad (60)$$

In the new coordinates, the curves are

$$\begin{aligned} \left(p + \frac{\mu}{\sqrt{2}\nu} \right)^2 - \left(q - \frac{\mu}{\sqrt{2}\nu} \right)^2 &= \frac{2S_1}{\nu} \\ \left(p + \frac{\mu}{\sqrt{2}\nu} \right)^2 - \left(q + \frac{\mu}{\sqrt{2}\nu} \right)^2 &= \frac{2S_2}{\nu} \end{aligned} \quad (61)$$

which are equations for a parametric family of hyperbolas. This means that for the tomograms $M_f^{(A_t)}(S_1, \mu, \nu)$ and $M_f^{(A_\omega)}(S_2, \mu, \nu)$ the probability distribution is collected not on straight lines but on hyperbolas. Other generalizations are obvious. One might use marginals on ellipses, parabolas or any other algebraic curves.

5.2. Operator delta function as a projector density

While constructing tomograms, the non-negative operator

$$\delta(B(\alpha) - X) \quad (62)$$

plays an essential role, $B(\alpha)$ being an Hermitian operator with nondegenerate continuous spectrum. The random variable X takes values on the spectrum of $B(\alpha)$. Considering a set of generalized eigenstates (in \mathcal{N}^*) of $B(\alpha)$, one obtains for the kernel

$$\langle Y | \delta(B(\alpha) - X) | Y' \rangle = \delta(Y' - X) \delta(Y - X) = \langle Y | X \rangle \langle X | Y' \rangle. \quad (63)$$

Therefore, we may identify $\delta(B(\alpha) - X)$ with the projector $|X\rangle\langle X|$

$$\delta(B(\alpha) - X) = |X\rangle\langle X| = P_X. \quad (64)$$

From this, it follows

$$M_f^{(B)} = \langle f | \delta(B(\alpha) - X) | f \rangle = \langle f | X \rangle \langle X | f \rangle = |\langle X | f \rangle|^2 \quad (65)$$

displaying the positivity of the tomogram. This means that there is always a basis in \mathcal{N}^* such that, by projecting on this basis, the tomogram is the product of two complex conjugate functions. By a unitary transformation S , $B(\alpha)$ may be transformed to

$$SB(\alpha)S^\dagger = B'(\alpha). \quad (66)$$

Then if $\{|Z\rangle\}$ is the set of (generalized) eigenvectors of $B'(\alpha)$, $\{S^\dagger|Z\rangle\}$ is a set of eigenvectors for B . Therefore

$$M_f^{(B)}(Z) = \langle f | \delta(B(\alpha) - Z) | f \rangle = |\langle Z | S | f \rangle|^2 = \langle f | S^\dagger | Z \rangle \langle Z | S | f \rangle. \quad (67)$$

In this case, the operator $U(\alpha)$ in the general scheme described in section 2 would be

$$U(\alpha) = S^\dagger |Z\rangle\langle Z| S \quad (68)$$

which, in this case, is not represented as an exponent of an operator. The form (68), with the presence of a projector operator $P_Z = |Z\rangle\langle Z|$, also shows the relation of the tomograms to transforms of the Husimi–Kano type. Notice, however, that, for example, the time–frequency Husimi–Kano transform

$$Q_f(t, \omega) = |\langle \beta | f \rangle|^2 \quad \beta = \frac{t + i\omega}{\sqrt{2}} \quad (69)$$

where $|\beta\rangle$ is a coherent state, does not describe a joint probability distribution in the time–frequency plane, because time and frequency do not commute and a joint probability distribution of two noncommuting observables cannot exist due to the uncertainty relation. Therefore, the correct way to interpret the Husimi–Kano quasidistribution is not as a joint time–frequency probability but as a unitarily transformed tomogram.

6. Discrete spectrum

In the case where the operator $B(\alpha)$ has a discrete spectrum, one uses a Kronecker delta function $\delta_K(B(\alpha) - n)$ and associates to the operator the Fourier integral on the circle. For example, for the number operator

$$\delta_K(a^\dagger a - n) = \frac{1}{2\pi} \int_0^{2\pi} e^{i\varphi(a^\dagger a - n)} d\varphi \quad n = 0, 1, 2, 3 \dots \quad (70)$$

and the matrix elements of this operator in the Fock basis $|m\rangle$ are

$$\langle m' | \delta_K(a^\dagger a - n) | m \rangle = \delta_{m'm} \delta_{mn} \quad m, m' = 0, 1, 2, 3 \dots \quad (71)$$

which equal the matrix elements of the projector

$$P_n = |n\rangle\langle n|. \quad (72)$$

This means that, also in the discrete-spectrum case, the Kronecker delta function of the operator $B(\alpha)$ is reduced to a projector. The tomogram associated with the Wigner–Ville function by this method is given by the so-called photon-number tomography [29–31], i.e.

$$\begin{aligned} w(n, \beta) &= \langle f | D^\dagger(\beta) | n \rangle \langle n | D(\beta) | f \rangle \\ &= \frac{1}{\sqrt{\pi} 2^n n!} \left| \int dt e^{-(t^2/2) + i\sqrt{2}t \operatorname{Im} \beta} H_n(t) f(t + \sqrt{2} \operatorname{Re} \beta) \right|^2 \end{aligned} \quad (73)$$

where the complex number β is a linear combination of the parameters μ and ν .

One may also construct a tomogram using a Dirac delta function of the same operator $B(\alpha)$:

$$\begin{aligned} M(X, \Omega, \tau) &= \int dk dt' dt'' dt f(t' + k\tau) f^*(t'' - k\tau) e^{-ik(X + \pi/2)} \frac{i}{2\pi \sin(k\pi/2)} \\ &\quad \times e^{-ik\Omega(t' + t'')} \exp \left[-\frac{i}{2} \cot \frac{k\pi}{2} (2t^2 + t'^2 + t''^2) + \frac{it}{\sin(k\pi/2)} (t' + t'') \right]. \end{aligned} \quad (74)$$

The inverse of the transform is

$$f(t) f^*(t') = \frac{1}{\pi} \int M \left(X - \frac{\pi}{2}, \Omega, \frac{t' - t}{2} \right) \exp \left[-i\Omega(t + t') + i \left(X + \frac{\pi}{2} \right) \right] d\Omega dX. \quad (75)$$

In this case, the Dirac delta function of $B(\alpha) - X$ is not reduced to a projector density. The tomogram (74) corresponds to marginals collected from shifted circles in the classical phase space.

7. Remarks and conclusions

- (1) The main result in this work is the formulation of a unified view for some linear and nonlinear transforms through the operator formulation developed in section 2. The formulation emphasizes the basic unity of these transforms, which are related by explicit formulae. Nevertheless, for each particular application, one type of transform may be more convenient than the others. In particular, when non-ambiguous joint information on noncommutative observable planes is desired, tomograms seem to be the most competitive type of transforms.

The formulation applies both to unitary operators or nonunitary ones of the form (15). It is also an appropriate framework to construct new transforms once a particular aspect of the signal is defined and this is expressed through the corresponding operator set.

- (2) The operators $U(\vec{\alpha})$ may belong to group representations or be operators of a deformed group. In the cases we have considered in detail, the operators belong to the Lie algebra of the Weyl-symplectic group, which has the following six generators:

$$L_1 = \hat{t} \quad L_2 = \hat{\omega} \quad L_3 = 1 \quad L_4 = \hat{t}^2 \quad L_5 = \hat{\omega}^2 \quad L_6 = \frac{1}{2} (\hat{t}\hat{\omega} + \hat{\omega}\hat{t}).$$

When it is unitary, the operator $U(\vec{\alpha})$ may be considered to be an evolution operator associated with a Hamiltonian operator formed from the generators of the group. Quasidistributions evolve the state $|f\rangle$ and project the evolved state on the initial condition. On the other hand, the tomograms collect the probability density on a family of lines in phase space. In the cases that were studied, hyperbolas, straight lines and circles were considered. Other types of tomograms might be considered, for example, those corresponding to parabolas, that is, $X = \mu t + \nu \omega^2$. A similar construction might be done for other groups and other algebraic structures like quantum groups.

Acknowledgment

MAM thanks the Russian Foundation for Basic Research for partial support under project no 01-02-17745.

References

- [1] Pouarikas A D (ed) 1996 *The Transforms and Applications Handbook* (Boca Raton, FL: CRC Press)
- [2] Wolf K-B 1979 *Integral Transforms in Science and Engineering* (New York: Plenum)
- [3] Fourier J B J 1988 *Théorie Analytique de la Chaleur* ed G Darbous *Oeuvres de Fourier Tome Premier* (Paris: Gauthiers-Villars)
- [4] Combes J M, Grossmann A and Tchamitchian Ph (ed) 1990 *Wavelets* 2nd edn (Berlin: Springer)
- [5] Daubechies I 1990 The wavelet transform: time–frequency localization and signal analysis *IEEE Trans. Inf. Theory* **36** 961–1005
- [6] Chui C K (ed) 1992 *Wavelets: A Tutorial. Theory and Applications* vol 2 (Boston: Academic)
- [7] Wigner E 1932 On the quantum correction for thermodynamic equilibrium *Phys. Rev.* **40** 749–59
- [8] Ville J 1948 Théorie et applications de la notion de signal analytique *Cables Transm. A* **2** 61–74
- [9] Cohen L 1966 Generalized phase-space distribution functions *J. Math. Phys.* **7** 781–806
- [10] Qian S and Chen D 1965 *Joint Time–Frequency Analysis* (Englewood Cliffs, NJ: Prentice-Hall)
- [11] Woods J C and Barry D T 1994 Linear signal synthesis using the Radon–Wigner transform *IEEE Trans. Signal Process.* **42** 2105–11
- [12] Granieri S, Furlan W D, Saavedra G and Andrés P 1997 Radon–Wigner display: a compact optical implementation with a single varifocal lens *Appl. Opt.* **36** 8363–9
- [13] Man'ko V I and Mendes R V 1999 Noncommutative time–frequency tomography *Phys. Lett. A* **263** 53–9
- [14] Glauber R J 1963 Coherent and incoherent states of the radiation fields *Phys. Rev.* **131** 2766–88
- [15] Glauber R J 1963 Photon correlations *Phys. Rev. Lett.* **10** 84–6
- [15] Sudarshan E C G 1963 Equivalence of semiclassical and quantum-mechanical descriptions of statistical light beams *Phys. Rev. Lett.* **10** 277–9
- [16] Husimi K 1940 Some formal properties of the density matrix *Proc. Phys. Math. Soc. Japan* **22** 264–314
- [17] Kano Y 1965 A new phase-space distribution function in the statistical theory of the electromagnetic field *J. Math. Phys.* **6** 1913–5
- [18] Cahill K E and Glauber R J 1969 Density operators and quasiprobability distributions *Phys. Rev.* **177** 1882–902
- [19] Bertrand J and Bertrand P 1992 A class of affine Wigner functions with extended covariance properties *J. Math. Phys.* **33** 2515–27
- [20] Goncalvés P and Baraniuk R J 1996 A pseudo-Bertrand distribution for time–scale analysis *IEEE Signal Process. Lett.* **3** 82–4
- [21] Flandrin P and Goncalvés P 1994 From wavelets to time–scale energy distributions *Recent Advances in Wavelet Analysis* ed L L Schumaker and J Webb (New York: Academic) pp 309–34
- [22] Cohen L 1993 The scale representation *IEEE Trans. Signal Process.* **41** 3275–92
- [23] Reed M and Simon B 1980 *Functional Analysis* (San Diego: Academic) ch 8
- [24] Cohen L 1989 Time–frequency distributions—a review *Proc. IEEE* **77** 941–81
- [25] Dodonov V V and Man'ko V I 1989 Invariants and evolution of nonstationary quantum systems *Proc. P N Lebedev Physical Institute* vol 183 (Commack, NY: Nova Science)
- [26] Ali T, Atakishiyev N M, Chumakov S M and Wolf K-B 2000 The Wigner function for general Lie groups and the wavelet transform *Ann. Inst. Henri Poincaré* **1** 685–714
- [27] Atakishiyev N M, Chumakov S M and Wolf K-B 1998 Wigner distribution function for finite systems *J. Math. Phys.* **139** 6244–61
- [28] Nieto L M, Atakishiyev N M, Chumakov S M and Wolf K-B 1998 Wigner distribution function for Euclidean systems *J. Phys. A: Math. Gen.* **31** 3875–95
- [29] Mancini S, Tombesi P and Man'ko V I 1997 Density matrix from photon number tomography *Europhys. Lett.* **37** 79–83
- [30] Wallentowitz S and Vogel W 1996 Unbalanced homodyning for quantum state measurements *Phys. Rev. A* **53** 4528–33
- [31] Banaszek K and Wodkiewicz K 1996 Direct probing of quantum phase space by photon counting *Phys. Rev. Lett.* **76** 4344–7