

## Strongly coupled quantum discrete Liouville theory: II. Geometric interpretation of the evolution operator

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2002 J. Phys. A: Math. Gen. 35 4043

(<http://iopscience.iop.org/0305-4470/35/18/304>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.106

The article was downloaded on 02/06/2010 at 10:03

Please note that [terms and conditions apply](#).

# Strongly coupled quantum discrete Liouville theory: II. Geometric interpretation of the evolution operator

L D Faddeev and R M Kashaev

Steklov Mathematical Institute at St Petersburg, Fontanka 27, St Petersburg 191011, Russia

E-mail: faddeev@pdmi.ras.ru and kashaev@pdmi.ras.ru

Received 6 February 2002

Published 26 April 2002

Online at stacks.iop.org/JPhysA/35/4043

## Abstract

It is shown that the  $N$ th power of the light-cone evolution operator of the  $2N$ -periodic quantum discrete Liouville model can be identified with the Dehn twist operator in quantum Teichmüller theory.

PACS numbers: 02.60.Lj, 02.10.Hh, 02.40.Re, 03.65.Fd, 03.70.+k, 11.25.Hf

## 1. Introduction

Integrable lattice regularization of quantum Liouville theory has been developed in [2, 4, 5]. According to the recent development in [3], the model is expected to describe the quantum Liouville equation with Virasoro central charge  $c_L > 1$ , including the ‘strongly coupled regime’  $1 < c_L < 25$ .

This paper can be considered as a second part of our previous work [3] dedicated to the discrete Liouville model. Here we show that the evolution operator of the model can be interpreted in pure geometrical terms within quantum Teichmüller theory [1, 6–9]. Namely, we identify the  $N$ th power of the light-cone evolution operator of the quantum discrete Liouville model of spatial length  $2N$  (which is the number of sites in a chain) with the Dehn twist operator in quantum Teichmüller theory.

The paper is organized as follows. The quantum discrete Liouville system is briefly described in section 2. The relation to quantum Teichmüller theory is explained in section 3.

## 2. Quantum discrete Liouville system

### 2.1. Algebra of observables

Following [3], algebra of observables  $\mathcal{A}_N$ ,  $N > 1$ , is generated by self-adjoint elements  $f_j$ ,  $j \in \mathbb{Z}$ , with periodicity condition  $f_{j+2N} = f_j$  and commutation relations

$$[f_m, f_n] = \begin{cases} (-1)^m (2\pi i)^{-1}, & \text{if } n = m \pm 1 \pmod{2N} \\ 0, & \text{otherwise.} \end{cases} \quad (2.1)$$

## 2.2. Equations of motion

The field variables

$$\chi_{j,t} \equiv \mathbf{U}_{\text{lc}}^t e^{2\pi b f_{j+t}} \mathbf{U}_{\text{lc}}^{-t}, \quad j+t = 1 \pmod{2}$$

are defined so that

$$\mathbf{U}_{\text{lc}} \chi_{j,t} \mathbf{U}_{\text{lc}}^{-1} = \chi_{j-1,t+1}.$$

Here the ‘light-cone’ evolution operator  $\mathbf{U}_{\text{lc}}$  is defined explicitly

$$\mathbf{U}_{\text{lc}} = \mathbf{G} \prod_{j=1}^N \varphi_b(f_{2j}) = \prod_{j=1}^N \varphi_b(f_{2j-1}) \mathbf{G} \quad (2.2)$$

where

$$\varphi_b(z) \equiv \exp\left(\frac{1}{4} \int_{i0-\infty}^{i0+\infty} \frac{e^{-i2zw} dw}{\sinh(wb) \sinh(w/b)}\right) = (e^{2\pi(z+c_b)b}; q^2)_{\infty} / (e^{2\pi(z-c_b)b^{-1}}; \bar{q}^2)_{\infty},$$

$$q \equiv e^{i\pi b^2}, \quad \bar{q} \equiv e^{-i\pi b^{-2}}, \quad c_b \equiv i(b+b^{-1})/2$$

while operator  $\mathbf{G}$  is defined through the equations

$$\mathbf{G} f_j = (-1)^j f_{j-1} \mathbf{G}. \quad (2.3)$$

The field variables solve the quantum discrete Liouville equation<sup>1</sup>

$$\chi_{j,t+1} \chi_{j,t-1} = (1+q\chi_{j+1,t})(1+q\chi_{j-1,t})$$

with spatial periodic boundary condition

$$\chi_{j+2N,t} = \chi_{j,t}$$

and the initial data given by exponentiated generators

$$\chi_{2j+1,0} = e^{2\pi b f_{2j+1}}, \quad \chi_{2j,-1} = e^{2\pi b f_{2j}}.$$

## 3. Interpretation within quantum Teichmüller theory


In this section we interpret the evolution operator  $\mathbf{U}_{\text{lc}}$  in geometrical terms by using the formalism of decorated ideal triangulations and their transformations within quantum Teichmüller theory described in [9].

### 3.1. Geometric realization


We consider an annulus with  $N$  marked points on each of its boundary components ( $2N$  points in total) and choose decorated ideal triangulation  $\tau_N$  of it, shown in figure 1. Equivalently, we can think of  $\tau_N$  as an infinite triangulated strip where triangles are numerated by integers in accordance with figure 1 with periodicity condition  $\bar{\tau}_N(n+2N) = \bar{\tau}_N(n)$ ,  $\forall n \in \mathbb{Z}$ . In this way we come to identification of the integers from 1 to  $2N$ , numbering triangles in  $\tau_N$ , with elements of the ring of residues  $\mathbb{Z}_{2N} \equiv \mathbb{Z}/2N\mathbb{Z}$ . Such identification will be assumed in algebraic expressions, when necessary.

Denote  $D^{1/N}$  the isotopy class of a homeomorphism of the annulus which rotates the top boundary component wrt the bottom through angle  $2\pi/N$  so that the marked points of the top

<sup>1</sup> Using invariance of  $\mathbf{U}_{\text{lc}}$  with respect to symmetry  $b \leftrightarrow b^{-1}$ , one can also define the dual fields satisfying the dual equation, see [3].



**Figure 1.** Decorated ideal triangulation  $\tau_N$  of an annulus with  $N$  marked points on each boundary component. The leftmost and the rightmost vertical edges are identified.



**Figure 2.** Continuous transformation  $D^{1/N}$  of triangulated annulus  $\tau_N$  is a cyclic shift of the top boundary wrt the bottom boundary to the right by one spacing.


boundary are cyclically shifted by one period. The reason for using fractional power notation comes from the fact that

$$\underbrace{D^{1/N} \circ \dots \circ D^{1/N}}_{N \text{ times}} = D$$


is nothing else but the Dehn twist. From figure 2 it follows that the following composition of geometric transformations is identity,

$$(\dots, j, j-1, \dots) \circ \prod_k \rho_k^{-1} \circ \prod_l \omega_{2l+1, 2l}^{-1} \circ D^{1/N} = \text{id}$$

where elementary geometric transformations  $\rho_i$  and  $\omega_{ij}$  have the form



and



with

$$\omega_{k, \check{l}} \equiv \rho_l \circ \omega_{k, l} \circ \rho_l^{-1}$$

while  $(\dots, j, j-1, \dots)$  denotes the index shift transformation  $j \mapsto j-1$ . Equivalently we can write

$$D^{1/N} = \prod_l \omega_{2l+1, 2l} \circ \prod_k \rho_k \circ (\dots, j, j+1, \dots)$$

so, quantum realization of  $D^{1/N}$  in  $L^2(\mathbb{R}^{2N})$  has the form

$$F(\tau_N, D^{1/N}(\tau_N)) \simeq D^{1/N} \equiv \zeta^{-N-6/N} P_{(\dots j, j+1 \dots)} \prod_{k=1}^{2N} A_k \prod_{l=1}^N T_{2l+1, \check{l}} \tag{3.1}$$

with the normalization factor chosen in accordance with the convention for Dehn twists used in [8]. Here  $P_{(\dots j, j+1 \dots)}$  is the natural realization of the cyclic permutation,

$$\begin{aligned} \zeta &= e^{-i\pi(b+b^{-1})^2/12} \\ A_k &\equiv e^{-i\pi/3} e^{i3\pi q_k^2} e^{i\pi(p_k+q_k)^2} \\ T_{k, \check{l}} &= e^{-i2\pi p_k p_l} \bar{\varphi}_b(q_k + q_l), \quad \bar{\varphi}_b(z) \equiv (\varphi_b(z))^{-1} \end{aligned}$$

where self-adjoint operators  $p_j, q_j$  satisfy Heisenberg commutation relations

$$[p_j, p_k] = [q_j, q_k] = 0, \quad [p_j, q_k] = \delta_{j,k} (2\pi i)^{-1}.$$

We can rewrite equation (3.1) in the form

$$D^{-1/N} = \zeta^{N+6/N} \prod_{m=1}^N \varphi_b(q_{2m} + q_{2m+1}) e^{i2\pi \sum_{j=1}^N p_{2j} p_{2j+1}} \prod_{k=1}^{2N} A_k^{-1} P_{(\dots l, l-1, \dots)}. \tag{3.2}$$

**Proposition 1. Operators**

$$\kappa(f_j) = \begin{cases} p_j + p_{j-1}, & \text{if } j = 0 \pmod{2} \\ q_j + q_{j-1}, & \text{otherwise} \end{cases} \tag{3.3}$$

$$\kappa(G) = \zeta^{N+6/N} e^{i2\pi \sum_{j=1}^N p_{2j} p_{2j+1}} \prod_{k=1}^{2N} A_k^{-1} P_{(\dots l, l-1, \dots)} \tag{3.4}$$

define faithful (reducible) realization of the observable algebra  $\mathcal{A}_N$  in  $L^2(\mathbb{R}^{2N})$ .

**Proof.** Let us check that these definitions are consistent with relations (2.1) and (2.3). First, evidently,

$$[\kappa(f_{2j}), \kappa(f_{2k})] = [\kappa(f_{2j+1}), \kappa(f_{2k+1})] = 0$$

while

$$[\kappa(f_{2j}), \kappa(f_{2k+1})] = [p_{2j} + p_{2j-1}, q_{2k+1} + q_{2k}] = (2\pi i)^{-1} (\delta_{j,k} + \delta_{j,k+1})$$

thus reproducing relations (2.1). Next,

$$\begin{aligned} \text{Ad}(\kappa(G))\kappa(f_{2j}) &= \text{Ad}(\kappa(G))(p_{2j} + p_{2j-1}) = \text{Ad}(e^{i2\pi \sum_{k=1}^N p_{2k} p_{2k+1}})(p_{2j-1} + p_{2j-2}) \\ &= \text{Ad}(e^{i2\pi \sum_{k=1}^N p_{2k} p_{2k+1}})(q_{2j-1} - p_{2j-1} + q_{2j-2} - p_{2j-2}) = q_{2j-1} + q_{2j-2} \\ &= \kappa(f_{2j-1}) \end{aligned}$$

and similarly

$$\begin{aligned} \text{Ad}(\kappa(G))\kappa(f_{2j+1}) &= \text{Ad}(\kappa(G))(q_{2j+1} + q_{2j}) \\ &= \text{Ad}(e^{i2\pi \sum_{k=1}^N p_{2k} p_{2k+1}})(q_{2j} + q_{2j-1}) = -p_{2j} - p_{2j-1} = -\kappa(f_{2j}) \end{aligned}$$

in agreement with equation (2.3). □

Now, comparing equations (2.2) and (3.2), we come to our main result

$$\kappa(U_C) = D^{-1/N}. \tag{3.5}$$

### 3.2. A similarity transformation

Here, we give (without proof) the result of similarity transformation which simplifies the  $N$ th power of the evolution operator.

Define

$$W \equiv \prod_{N \geq j > 1} (\bar{\varphi}_b(f_{2j})\varphi_b(f_{2j-1})\varphi_b(\mathfrak{g}_{2j,2N+1})), \quad \bar{\varphi}_b(x) \equiv (\varphi_b(x))^{-1}$$

where  $\mathfrak{g}_{j,k} \equiv \sum_{l=j+1}^k f_l$ , and the product of noncommuting operators is in decreasing order from left to right.

**Proposition 2.** *One has the following explicit expression for the transformed evolution operator*

$$\begin{aligned} \tilde{U}_{lc} \equiv \text{Ad}(W^{-1})U_{lc} &= (\zeta e^{i\pi/6})^{-N} \prod_{N > k > 1} \bar{\varphi}_b(\mathfrak{g}_{2k-1,2N-1}) \\ &\times \bar{\varphi}_b(\mathfrak{g}_{2,2N-1})\varphi_b(\mathfrak{g}_{2N-1,2N+1})\bar{\varphi}_b(f_2)e^{i\pi \sum_{l=1}^N f_{2l}} \mathbf{G} \end{aligned}$$

where the product is again in decreasing order from left to right, while the  $N$ th power has the form

$$\tilde{U}_{lc}^N = (\zeta e^{i\pi/6})^{1-N^2} \varphi_b(\mathfrak{g}_{2,2N+1})e^{-i\pi f_2^2} (e^{i\pi \sum_{l=1}^N f_{2l}} \mathbf{G})^N.$$

## 4. Conclusion

The main result of this paper is formula (3.5) which, on one side, identifies ‘zero-modes’ of the  $2N$ -periodic quantum discrete Liouville equation to be given by the  $N$ th power of the light-cone evolution operator  $U_{lc}$ , and equates these zero-modes to the (inverse of) Dehn twist operator in quantum Teichmüller theory, on the other side. The immediate consequence of this result is that now, based on the known spectrum of operator  $D$ , we know the spectrum of the model. Indeed, the complete spectrum of  $D$  is given by the formula [9]:

$$\text{Spec}(D) = \{e^{i2\pi \Delta_s} | s \in \mathbb{R}_{>0}\}$$

where

$$\Delta_s = \frac{c_L - 1}{24} + s^2, \quad c_L = 1 + 6(b + b^{-1})^2 > 1$$

are conformal weights and the Virasoro central charge in (continuous) quantum Liouville theory, see [10] for a recent review. This is consistent with interpretation of the Dehn twist spectrum as Liouville conformal weights through the formula  $\text{Spec}(D) = \text{Spec}(e^{i2\pi L_0})$ , where  $L_0$  is the Virasoro generator in continuous quantum Liouville theory with the known spectrum

$$\text{Spec}(L_0) = \{\Delta_s + m | s \in \mathbb{R}_{>0}, m \in \mathbb{Z}_{\geq 0}\}.$$

Our result implies the following spectrum of  $U_{lc}$

$$\text{Spec}(U_{lc}) = \{e^{-i2\pi(\Delta_s+m)/N} | s \in \mathbb{R}_{>0}, m \in \mathbb{Z}/N\mathbb{Z}\}$$

which coincides with the spectrum of the exponential operator  $e^{-2\pi i L_0/N}$  in quantum Liouville theory. Thus, the discrete version of quantum Liouville theory is in complete agreement with the continuous one and there is no modification in the spectrum of conformal weights.

## Acknowledgments

The authors would like to thank A Fring, M Karowski, R Schrader, S Sergeev, Yu Suris and J Teschner for useful discussions. The work is supported in part by grants CRDF RM1-2244, INTAS 99-01705 and RFBR 99-01-00101.

## References

- [1] Chekhov L and Fock V V 1999 Quantum Teichmüller space *Theor. Math. Phys.* **120** 1245–59 (math.QA/9908165)
- [2] Faddeev LD 1992 Quantum symmetry in conformal field theory by Hamiltonian methods. Cargèse 1991 *New Symmetry Principles in Quantum Field Theory* ed J Frölich *et al* (New York: Plenum) pp 159–75
- [3] Faddeev L D, Kashaev R M and Volkov A Yu 2000 Strongly coupled quantum discrete Liouville theory. I: Algebraic approach and duality *Preprint* hep-th/0006156
- [4] Faddeev L D and Takhtajan L A 1986 Liouville model on the lattice. *Field Theory, Quantum Gravity and Strings: Proc. Meudon and Paris VI, France 1984/85 (Lecture Notes in Physics vol 246)* ed H J de Vega and N Sanchez (Berlin: Springer)
- [5] Faddeev L D and Volkov A Yu 1999 Algebraic quantization of integrable models in discrete space-time *Discrete Integrable Geometry and Physics (Oxford Lecture Series in Mathematics and its Applications vol 16)* (Oxford: Clarendon) pp 301–20 (hep-th/97010039)
- [6] Fock V V 1997 Dual Teichmüller spaces *Preprint* dg-ga/9702018
- [7] Kashaev R M 1998 Quantization of Teichmüller spaces and the quantum dilogarithm *Lett. Math. Phys.* **43** 105–15 (q-alg/9705021)
- [8] Kashaev R M 1999 Liouville central charge in quantum Teichmüller theory *Proc. Steklov Inst. Math.* vol 226 pp 63–71 (hep-th/9811203)
- [9] Kashaev R M 2000 On the spectrum of Dehn twists in quantum Teichmüller theory *Preprint* HIP-2000-44/TH, math.QA/0008148
- [10] Teschner J 2001 Liouville theory revisited *Preprint* hep-th/0104158