

A comparison between star products on regular orbits of compact Lie groups*

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2002 J. Phys. A: Math. Gen. 35 5687

(<http://iopscience.iop.org/0305-4470/35/27/310>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.107

The article was downloaded on 02/06/2010 at 10:14

Please note that [terms and conditions apply](#).

A comparison between star products on regular orbits of compact Lie groups*

R Fioresi¹ and M A Lledo^{2,3}

¹ Dipartimento di Matematica, Università di Bologna, Piazza di Porta S Donato, 5, 40126 Bologna, Italy

² Dipartimento di Fisica, Politecnico di Torino, Corso Duca degli Abruzzi 24, I-10129 Torino, Italy

³ INFN, Sezione di Torino, Italy

E-mail: fioresi@dm.UniBo.it and lledo@athena.polito.it

Received 31 December 2001, in final form 1 May 2002

Published 28 June 2002

Online at stacks.iop.org/JPhysA/35/5687

Abstract

In this paper, an algebraic and a differential star product defined on a regular coadjoint orbit of a compact semisimple group are compared. It has been proved that there is an injective algebra homomorphism between the algebra of polynomials with the algebraic star product and the algebra of differential functions with the differential star product structure.

PACS numbers: 02.20.Sv, 02.20.Tw, 02.40.-k

1. Introduction

The problem of classification of differential star products on a general Poisson manifold was solved in [1]. The existence of star products on symplectic manifolds was already proved in [2, 3] and, using a different technique, a construction of a star product and a classification of all star products on a symplectic manifold were given in [4, 5]. For other special cases, as for regular manifolds, a proof of existence of tangential star products was known (see [6]).

To motivate our discussion, let us consider the Heisenberg group $H = \mathbb{R}^3$ with multiplication

$$(a_1, b_1, c_1) \cdot (a_2, b_2, c_2) = (a_1 + a_2, b_1 + b_2, c_1 + c_2 + a_1 b_2).$$

Its Lie algebra is $\mathfrak{h} = \mathbb{R}^3 = \text{span}\{Q, P, E' = -iE\}$ with commutation rules

$$[Q, P] = -iE \quad (\text{the rest trivial}).$$

* Investigation supported by the University of Bologna, funds for selected research topics.

The coadjoint orbits of H are the planes $c = \text{constant} \neq 0$ (regular orbits) and the points $(a, b, 0)$. One way of obtaining the Moyal–Weyl product on \mathbb{R}^2 is considering the Weyl map or symmetrizer in the enveloping algebra of \mathfrak{h} ,

$$\begin{aligned} \text{Sym} : \text{Pol}[\mathfrak{h}^*] &\rightarrow U(\mathfrak{h}) \\ x_1 x_2 \cdots x_k &\mapsto \frac{1}{k!} \sum_{\sigma \in S_k} X_{\sigma(1)} \cdots X_{\sigma(k)} \end{aligned} \quad (1)$$

where x_i are the coordinates on \mathfrak{h}^* on the basis dual to $\{X_i\}$ in \mathfrak{h} . By multiplying the commutation rules by a formal parameter \hbar we obtain the following star product on $\text{Pol}[\mathfrak{h}^*][[\hbar]]$:

$$f \star g = \text{Sym}^{-1}(\text{Sym}(f)\text{Sym}(g)). \quad (2)$$

This star product is *differential*, so it can be extended to $C^\infty(\mathfrak{h}^*)$, it is *tangential*, so it can be restricted to the orbits and it is *algebraic*, that is, it is closed (and convergent) on polynomials.

If instead of the Heisenberg group we take another group, say $SU(2)$, we can define a star product using the symmetrizer as in (1). The resulting star product is algebraic and differential, but it is not tangent to the coadjoint orbits, so it does not define a star product on them, in this case the spheres.

Two different approaches can be taken at this point. One is to look for a differential star product on the sphere in the spirit of [1–4]. The resulting star product is neither algebraic nor appears related to the product on the enveloping algebra. The other approach insists on using the product in the enveloping algebra. The consequence is that differentiability is lost. These kinds of star products have been considered in [7–10] and, in particular, in [8, 9] it was proved that a non-differential star product on coadjoint orbits of $SU(2)$ corresponds to the standard quantization of angular momentum. It then seems unavoidable to look for a wider class of star products than the differentiable ones. In particular, one cannot immediately assume that the canonical quantization given by Kontsevich’s theorem [1] is the one relevant for physics in all cases.

The problem of existence and classification of algebraic star products on algebraic Poisson varieties appears as a separate problem, mathematically interesting in itself, which has been recently studied in [11]. From the physical point of view it is of interest since the algebra of a physical quantum system may have a non-differential star product, as in the case of the angular momentum and its standard quantization.

Our purpose here is to compare the deformations obtained by algebraic [8, 9] and differential methods on regular coadjoint orbits of compact semisimple Lie groups. We want to establish if there is some kind of equivalence among these different star products. We work with a family of algebraic star products, not all isomorphic, and we relate them to the differential star product given by Kontsevich’s theorem or Fedosov’s construction [4]. Our result is that one of the algebraic star products can be injected homomorphically into the differential one.

The organization of the paper is as follows. In section 2 we recall known facts concerning coadjoint orbits of a semisimple compact group G and its complexification $G_{\mathbb{C}}$. In section 3 we introduce different star products on a fixed regular coadjoint orbit Θ and on a tubular neighbourhood of the orbit \mathcal{N}_Θ , and we prove that two different star products on \mathcal{N}_Θ , one tangential \star_T and one nontangential $\star_{S\mathcal{N}_\Theta}$, are equivalent. In section 4 we show our main result, that there is an injective homomorphism between an algebraic star product \star_{P_Θ} and a differential one \star_{T_Θ} on the orbit Θ . The algebraic star product belongs to the family constructed in [8], while the differential one is obtained by gluing tangential star products defined on open sets of \mathcal{N}_Θ , computed with Kontsevich’s formula [1]. In appendix A we give

for completeness some standard definitions and results on star products and deformations. In appendix B we give an explicit formula for the gluing of star products given in open sets and satisfying a compatibility condition, in terms of a partition of unity.

2. Coadjoint orbits of semisimple Lie groups

Let G be a compact semisimple group of dimension n and rank m and \mathfrak{g} its Lie algebra. Let \mathfrak{g}^* be the dual of \mathfrak{g} . On $C^\infty(\mathfrak{g}^*)$ we have the Kirillov Poisson structure:

$$\{f_1, f_2\}(\lambda) = \langle [(df_1)_\lambda, (df_2)_\lambda], \lambda \rangle \quad f_1, f_2 \in C^\infty(\mathfrak{g}^*) \quad \lambda \in \mathfrak{g}^*.$$

$(df)_\lambda : \mathfrak{g}^* \rightarrow \mathbb{R}$ can be considered as an element of \mathfrak{g} , and $[\cdot, \cdot]$ is the Lie bracket on \mathfrak{g} . Let $\{X_1 \dots X_n\}$ be a basis of \mathfrak{g} and $\{x^1, \dots, x^n\}$ the coordinates on \mathfrak{g}^* in the dual basis. We have that

$$\{f_1, f_2\}(x^1, \dots, x^n) = \sum_{ijk} c_{ij}^k x^k \frac{\partial f_1}{\partial x^i} \frac{\partial f_2}{\partial x^j}$$

where c_{ij}^k are the structure constants of \mathfrak{g} , that is $[X_i, X_j] = \sum_k c_{ij}^k X_k$.

\mathfrak{g}^* is an algebraic Poisson manifold since the ring of polynomials $\mathbb{R}[\mathfrak{g}^*]$ is closed under the Poisson bracket.

The Kirillov–Poisson structure is neither symplectic nor regular. The symplectic leaves are the orbits of the coadjoint action of G on \mathfrak{g}^* ,

$$\langle \text{Ad}^*(g)\lambda, Y \rangle = \langle \lambda, \text{Ad}(g^{-1})Y \rangle \quad \forall g \in G \quad \lambda \in \mathfrak{g}^* \quad Y \in \mathfrak{g}.$$

We denote by Θ_λ the orbit of an element $\lambda \in \mathfrak{g}^*$ under the coadjoint action.

Let $G_{\mathbb{C}}$ be the complexification of G and $\mathfrak{g}_{\mathbb{C}}$ its Lie algebra. Let $\Theta_{\lambda_{\mathbb{C}}}$ be the coadjoint orbit of $\lambda \in \mathfrak{g}^*$ in $\mathfrak{g}_{\mathbb{C}}^*$ under the action of $G_{\mathbb{C}}$. $\Theta_{\lambda_{\mathbb{C}}}$ is an algebraic variety defined over \mathbb{R} and $\Theta_\lambda = \Theta_{\lambda_{\mathbb{C}}} \cap \mathfrak{g}^*$.

Let $\mathbb{C}[\mathfrak{g}^*]$ be the ring of polynomials on $\mathfrak{g}_{\mathbb{C}}^*$. We denote by $\text{Inv}(\mathfrak{g}_{\mathbb{C}}^*)$ the subalgebra of polynomials invariant under the coadjoint action. It is generated by homogeneous polynomials, $p_i, i = 1, \dots, m$ (Chevalley generators). We have that

$$\text{Inv}(\mathfrak{g}^*) = \text{Inv}(\mathfrak{g}_{\mathbb{C}}^*) \cap \mathbb{R}[\mathfrak{g}^*].$$

If λ is regular, the ideal of $\Theta_{\lambda_{\mathbb{C}}}$ is given by [12], $\mathcal{I}_{0\mathbb{C}} = (p_i - c_i, i = 1, \dots, m), c_i \in \mathbb{R}$, and the polynomials on $\Theta_{\lambda_{\mathbb{C}}}$ by $\mathbb{C}[\Theta_{\lambda_{\mathbb{C}}}] = \mathbb{C}[\mathfrak{g}^*]/\mathcal{I}_{0\mathbb{C}}$. For the real forms the ideal of Θ_λ is $\mathcal{I}_0 = \mathcal{I}_{0\mathbb{C}} \cap \mathbb{R}[\mathfrak{g}^*]$, with the same generators as the complex one and $\mathbb{R}[\Theta_\lambda] = \mathbb{R}[\mathfrak{g}^*]/\mathcal{I}_0 = \mathbb{C}[\Theta_{\lambda_{\mathbb{C}}}] \cap \mathbb{R}[\mathfrak{g}^*]$.

3. Star products on a regular coadjoint orbit

In this section we will consider complex star products which are deformations of the complexification of a real Poisson algebra. We want to describe different star products [9] that will be compared later.

From now on we fix a regular coadjoint orbit Θ in \mathfrak{g}^* . We will consider \mathfrak{g}_h the Lie algebra over $\mathbb{C}[[h]]$ obtained by multiplying the structure constants of $\mathfrak{g}_{\mathbb{C}}$ by a formal parameter h . U_h is its enveloping algebra.

3.1. The star products \star_S and $\star_{S\mathcal{N}_\Theta}$

It is well known that U_h is a formal deformation of $\mathbb{C}[\mathfrak{g}^*]$. In [1] it was shown that this deformation is isomorphic to the star product canonically associated with the Kirillov–Poisson

structure. Moreover, since the linear coordinates on \mathfrak{g}^* are global, one can compute a star product using Kontsevich’s universal formula.

The symmetrizer Sym (1) (that can be defined in the same way for any Lie algebra) defines, through (2), a differential and algebraic star product on \mathfrak{g}^* that we denote by \star_S . Any other isomorphism that is the identity modulo \hbar could be chosen in the place of Sym . All the star products constructed in this way are equivalent to the one obtained with Kontsevich’s explicit formula. All of them are algebraic and differential, but none of them is tangential to all the orbits [13].

Since a differential star product tangential to all orbits cannot exist in the whole \mathfrak{g}^* (see appendix B), we have to look for a smaller space. We consider a regular orbit Θ and a regularly foliated neighbourhood of the orbit, a tubular neighbourhood $\mathcal{N}_\Theta \simeq \Theta \times \mathbb{R}^m$, where the global coordinates in \mathbb{R}^m are the invariant polynomials p_i , $i = 1, \dots, m$. Since \star_S is differential, it can be restricted to the open set \mathcal{N}_Θ . We will denote that restriction by $\star_{S\mathcal{N}_\Theta}$. $\star_{S\mathcal{N}_\Theta}$ is a differential star product belonging to the canonical equivalence class associated with the Kirillov Poisson structure restricted to \mathcal{N}_Θ .

Since \mathcal{N}_Θ is a regular Poisson manifold, we know that a tangential star product (with respect to the symplectic leaves) exists [6]. We want to prove that there exists a tangential star product on \mathcal{N}_Θ equivalent to $\star_{S\mathcal{N}_\Theta}$.

3.2. The star products \star_T and $\star_{T\Theta}$

We want to define a tangential star product \star_T on \mathcal{N}_Θ and its restriction $\star_{T\Theta}$ to the regular orbit Θ . We will use the gluing of star products computed in appendix B in terms of a partition of unity.

Let $\mathcal{U} = \{U_r, r \in J\}$, where J is a set of indices, be a good covering of \mathcal{N}_Θ with Darboux charts. The coordinates in an open set U_r are

$$\begin{aligned} \varphi_r : U_r &\longrightarrow \mathbb{R}^n \quad \text{with} \\ \varphi_r &= (\theta_r, \pi_r, p) = (\theta_r^1, \dots, \theta_r^{(n-m)/2}, \pi_r^1, \dots, \pi_r^{(n-m)/2}, p_1, \dots, p_m) \\ \{\theta_r^\alpha, \pi_r^\beta\} &= \delta^{\alpha\beta} \quad \{\theta_r^\alpha, p_i\} = 0 \quad \{p_i, \pi_r^\beta\} = 0. \end{aligned}$$

The invariant polynomials p_i are global coordinates, so $U_r \simeq \hat{U}_r \times \mathbb{R}^m$ and $\{(\hat{U}_r, (\theta_r, \pi_r))\}_{r \in J}$ is an atlas of Θ , with $\{\hat{U}_r, (\theta_r, \pi_r), r \in J\}$ being the symplectic charts.

We can now apply Kontsevich’s formula in a coordinate patch U_r , using the Darboux coordinates φ_r . We denote this star product by \star_r^K . It is a tangential star product. If \star_r denotes the restriction of $\star_{S\mathcal{N}_\Theta}$ to U_r , then \star_r and \star_r^K are equivalent. We will denote by

$$R_r : (C^\infty(U_r)[[\hbar]], \star_r) \longrightarrow (C^\infty(U_r)[[\hbar]], \star_r^K)$$

the isomorphism

$$R_r(f \star_r g) = R_r(f) \star_r^K R_r(g) \quad R_r = \text{Id} + \sum_{i=1}^{\infty} \hbar^i R_r^i.$$

In the intersection $U_{rs} = U_r \cap U_s$, one has that \star_r^K and \star_s^K are equivalent as in (13) of appendix B with

$$T_{rs} = R_r \circ R_s^{-1}. \tag{3}$$

We have the following.

Proposition 3.1. *Let \mathcal{N}_Θ and \mathcal{U} be the tubular neighbourhood of the orbit Θ and the covering of \mathcal{N}_Θ defined above. Let \mathcal{F}_S be the sheaf of star products defined by $\star_{S\mathcal{N}_\Theta}$ and \star_r^K the star product obtained via the Kontsevich formula in $U_r \in \mathcal{U}$.*

The assignment

$$U_r \mapsto (C^\infty(U_r), \star_r^K) \quad \forall U_r \in \mathcal{U}$$

is a sheaf of star products isomorphic to \mathcal{F}_S . There is a star product \star_T on \mathcal{N}_Θ that is tangential and gauge equivalent to $\star_{S\mathcal{N}_\Theta}$.

Proof. It is necessary that the transition functions (3) satisfy the conditions (14) of appendix B, so we have a sheaf of star products that we will denote by \mathcal{F}_T . The isomorphisms R_r give the isomorphism of sheaves among \mathcal{F}_S and \mathcal{F}_T .

Given a partition of unity subordinated to \mathcal{U} one can use the method of appendix B to construct a global star product. From the explicit formula (15), one can see that it is a tangential star product. \square

The restriction of \star_T to the orbit will be denoted by $\star_{T\Theta}$.

3.3. The star products \star_P and $\star_{P\Theta}$

We want to define an algebraic star product \star_P on \mathfrak{g}^* and its restriction to the orbit Θ , the algebraic star product $\star_{P\Theta}$.

We consider the ideal in U_h

$$\mathcal{I}_h = (P_i - c_i(h), i = 1, \dots, m)$$

where $P_i = \text{Sym}(p_i)$ and $c_i(h) \in \mathbb{C}[[h]]$ with $c_i(0) = c_i^0$. It has been proved in [8] that U_h/\mathcal{I}_h is a deformation quantization of $\mathbb{C}[\Theta] = \mathbb{C}[\mathfrak{g}^*]/\mathcal{I}_0$ where

$$\mathcal{I}_0 = (p_i - c_i^0, i = 1, \dots, m)$$

is the ideal of a regular orbit Θ . Further properties of this deformation were studied in [9]. The generalization of this construction to nonregular orbits was done in [10].

A star product associated with this deformation can be constructed by giving a $\mathbb{C}[[h]]$ -module isomorphism:

$$\psi : \mathbb{C}[\mathfrak{g}^*][[h]] \longrightarrow U_h$$

that maps the ideal \mathcal{I}_0 isomorphically onto \mathcal{I}_h . One way of choosing this map (but not the only one) is by using the decomposition of $\mathbb{C}[\mathfrak{g}^*]$ in terms of invariant and harmonic polynomials [12]

$$\mathbb{C}[\mathfrak{g}^*] \cong \text{Inv}(\mathfrak{g}_\mathbb{C}^*) \otimes \mathcal{H}.$$

The harmonic polynomials \mathcal{H} are in one to one correspondence with $\mathbb{C}[\Theta]$ and we have a monomial basis $\mathcal{B} = \{x_{i_1} \dots x_{i_k}, (i_1, \dots, i_k) \in I\}$, where I is some subset of indices such that \mathcal{B} is a basis of $\mathbb{C}[\Theta]$ (see [8] for more details). We consider the following $\mathbb{C}[[h]]$ -module isomorphism:

$$\begin{aligned} \psi : \mathbb{C}[\mathfrak{g}^*][[h]] &\longrightarrow U_h \\ (p_{i_1} - c_{i_1}^0) \cdots (p_{i_k} - c_{i_k}^0) \otimes x_{j_1} \cdots x_{j_l} &\mapsto (P_{i_1} - c_{i_1}(h)) \cdots (P_{i_k} - c_{i_k}(h)) \otimes (X_{j_1} \cdots X_{j_l}), \end{aligned} \tag{4}$$

with $x_{j_1} \cdots x_{j_l} \in \mathcal{B}$. ψ defines an algebraic star product on $\mathbb{C}[\mathfrak{g}^*][[h]]$, that we will denote by \star_P . Since ψ descends to the quotient, it also defines an algebraic star product on $\mathbb{C}[\Theta][[h]]$ and we will denote it by $\star_{P\Theta}$. The case with $c_i(h) = c_i^0$ was considered first in [7], where it was shown that the star product is not differential.

4. Comparison between $\star_{T\Theta}$ and $\star_{P\Theta}$

In this section, we want to compare the differential star product $\star_{T\Theta}$ and the algebraic star product $\star_{P\Theta}$ defined on a fixed regular coadjoint orbit Θ . We want to show that there is an injective algebra homomorphism

$$\tilde{H} : (\mathbb{C}[\Theta][[h]], \star_{P\Theta}) \longrightarrow (C^\infty(\Theta)[[h]]_{\mathbb{C}}, \star_{T\Theta}).$$

We will first show that there exists an injective algebra homomorphism

$$H : (\mathbb{C}[\mathfrak{g}^*][[h]], \star_P) \longrightarrow (C^\infty(\mathcal{N}_\Theta)[[h]]_{\mathbb{C}}, \star_T) \quad (5)$$

and then we will show that it descends appropriately to the quotients as an injective homomorphism.

In order to compare the tangential star products \star_P on \mathfrak{g}^* (algebraic, not differential) and \star_T on \mathcal{N}_Θ (differential, not algebraic) we will use the nontangential star product \star_S on \mathfrak{g}^* (algebraic and differential).

The algebraic star products \star_P and \star_S on \mathfrak{g}^* are equivalent, since they define algebra structures that are isomorphic to U_h . The equivalence is realized by the $\mathbb{C}[[h]]$ -module isomorphism:

$$\begin{aligned} \eta : (\mathbb{C}[\mathfrak{g}^*][[h]], \star_P) &\rightarrow (\mathbb{C}[\mathfrak{g}^*][[h]], \star_S) \\ \eta &= \text{Sym}^{-1} \circ \psi \quad \eta(f \star_P g) = \eta(f) \star_S \eta(g). \end{aligned}$$

By the very definition (4)

$$f \star_P p_i = f \cdot p_i$$

so, since the p_i are central, the ideal $\mathcal{I}_0 = (p_i - c_i^0)$ in $\mathbb{C}[\mathfrak{g}^*][[h]]$ with respect to the commutative product is equal to the ideal with respect to the product \star_P , $\mathcal{I}_0^{\star_P} = (p_i - c_i^0)_{\star_P}$.

The generators of the ideal are mapped as

$$\eta(p_i - c_i^0) = (\text{Sym}^{-1} \circ \psi)(p_i - c_i^0) = \text{Sym}^{-1}(P_i - c_i(h)) = p_i - c_i(h)$$

so the ideal $\mathcal{I}_0^{\star_P}$ is mapped isomorphically by η onto the ideal with respect to the product \star_S , $\mathcal{I}_{c(h)}^{\star_S} = (p_i - c_i(h))_{\star_S}$. We note that in the case of \star_S , $\mathcal{I}_{c(h)}$, the ideal generated by $p_i - c_i(h)$ with respect to the commutative product does not coincide with $\mathcal{I}_{c(h)}^{\star_S}$. Note also that one can choose the $c_i(h)$ arbitrarily, provided that $c_i(0) = c_i^0$.

Since \star_S is differential, it is well defined on the whole $C^\infty(\mathfrak{g}^*[[h]])_{\mathbb{C}}$. The commutative ideal generated by $p_i - c_i^0$ on $C^\infty(\mathfrak{g}^*[[h]])_{\mathbb{C}}$ will be denoted by $\hat{\mathcal{I}}_0$. More generally, we can define $\mathcal{I}_{c(h)} = (p_i - c_i(h)) \subset \mathbb{C}[\mathfrak{g}^*][[h]]$, $\hat{\mathcal{I}}_{c(h)} = (p_i - c_i(h)) \subset C^\infty(\mathfrak{g}^*[[h]])_{\mathbb{C}}$. We have that $\mathcal{I}_0 \subset \hat{\mathcal{I}}_0$ and $\mathcal{I}_{c(h)} \subset \hat{\mathcal{I}}_{c(h)}$.

Let us consider the restriction map

$$r : C^\infty(\mathfrak{g}^*[[h]])_{\mathbb{C}} \longrightarrow C^\infty(\mathcal{N}_\Theta)[[h]]_{\mathbb{C}}.$$

Since the commutative product and \star_S are both local, the restriction r is an algebra homomorphism, between $C^\infty(\mathfrak{g}^*[[h]])_{\mathbb{C}}$ and $C^\infty(\mathcal{N}_\Theta)[[h]]_{\mathbb{C}}$ as commutative algebras, and also between $(C^\infty(\mathfrak{g}^*[[h]])_{\mathbb{C}}, \star_S)$ and $(C^\infty(\mathcal{N}_\Theta)[[h]]_{\mathbb{C}}, \star_{S\mathcal{N}_\Theta})$.

We consider the restriction of polynomials $r(\mathbb{C}[\mathfrak{g}^*][[h]])$. Since a polynomial is determined by its values on any open set, we can identify, via r , $(\mathbb{C}[\mathfrak{g}^*][[h]], \star_S)$ with a subalgebra of $(C^\infty(\mathcal{N}_\Theta)[[h]]_{\mathbb{C}}, \star_{S\mathcal{N}_\Theta})$.

On $C^\infty(\mathcal{N}_\Theta)[[h]]_{\mathbb{C}}$ there is an equivalence among $\star_{S\mathcal{N}_\Theta}$ and \star_T (proposition 3.1). We denote it by

$$\begin{aligned} \rho : (C^\infty(\mathcal{N}_\Theta)[[h]]_{\mathbb{C}}, \star_{S\mathcal{N}_\Theta}) &\longrightarrow (C^\infty(\mathcal{N}_\Theta)[[h]]_{\mathbb{C}}, \star_T) \\ \rho(f \star_S g) &= \rho(f) \star_T \rho(g) \quad \rho = \text{Id} + \sum_{n=1}^{\infty} h^n \rho_n \end{aligned}$$

where ρ_n are bidifferential operators. We have given the injective homomorphism (5) by $H = \rho \circ r \circ \eta$.

We now want to show that $H(\mathcal{I}_0) = \hat{\mathcal{J}}_0$, where $\hat{\mathcal{J}}_0$ is the ideal with respect to \star_T in $C^\infty(\mathcal{N}_\Theta)[[h]]_{\mathbb{C}}$ generated by $p_i - c_i^0$.

We want to find out how the generators $p_i - c_i(h)$ are mapped under ρ . The scalars are mapped into scalars, since the bidifferential operators involved in the star products \star_T and $\star_{S\mathcal{N}_\Theta}$ are null on the constants, and so are the operators ρ_n . We need to know $\rho(p_i)$.

Remark 4.1. Since ρ is an isomorphism of algebras and p_i belongs to the centre of $(C^\infty(\mathcal{N}_\Theta)[[h]], \star_S)$, $\rho(p_i)$ must also be in the centre of $(C^\infty(\mathcal{N}_\Theta)[[h]], \star_T)$. A function f in the centre of $(C^\infty(\mathcal{N}_\Theta)[[h]], \star_T)$ is a function depending only on the global coordinates $f(p_1, \dots, p_m)$, since the condition

$$f \star_T g - g \star_T f = 0 \quad \forall g \in C^\infty(\mathcal{N}_\Theta)$$

implies for the Poisson bracket

$$\{f, g\} = 0 \quad \forall g \in C^\infty(\mathcal{N}_\Theta)$$

so $\{f, \cdot\}$ is a null Hamiltonian vector field and in particular does not have components tangent to the symplectic leaves.

Remark 4.2. The algebra homomorphism condition determines the form of ρ on the centre in terms of $\rho(p_i) = p_i + a_i(p, h)$, where $a_i(p, h) = h z_i(p, h)$. In fact, on the centre we have

$$\rho = \sum_{j_1 \dots j_m} a_{1 \dots m}^{j_1 \dots j_m}(p, h) \frac{\partial}{\partial p_1^{j_1}} \dots \frac{\partial}{\partial p_m^{j_m}}.$$

$a_{1 \dots m}^{0 \dots 0} = 1$ and the rest of coefficients are multiples of h . In particular, the images of p_i are $\rho(p_i) = p_i + a_{1 \dots m}^{0 \dots 1 \dots 0}(p, h) = p_i + a_i(p, h)$. Using the fact that \star_T is tangential we have

$$f \star_T p_i = f \cdot p_i \quad \forall f \in C^\infty(\mathcal{N}_\Theta)$$

the homomorphism condition reads

$$\rho \left(p_1^{i_1} \dots p_m^{i_m} \right) = (p_1 + a_1)^{i_1} \dots (p_m + a_m)^{i_m}.$$

The solution of this equation is

$$a_{1 \dots m}^{j_1 \dots j_m} = \frac{1}{j_1! \dots j_m!} a_1^{j_1} \dots a_m^{j_m}. \tag{6}$$

In particular, ρ is trivial on the centre if and only if $a_1 = \dots = a_m = 0$.

By remarks 4.1 and 4.2 we have that

$$H(p_i - c_i^0) = \rho(p_i - c_i(h)) = p_i + a_i(p, h) - c_i(h) = p_i - c_i^0 + h(z_i(p, h) - \Delta_i(h))$$

where we have denoted $a_i(p, h) = h z_i(p, h)$ and $c_i(h) = c_i^0 + h \Delta_i(h)$. Since $\Delta_i(h)$ is arbitrary, we can choose it as

$$\Delta_i(h) = z_i(c_i^0, h). \tag{7}$$

It is not difficult to see that

$$z_i(p, h) - \Delta_i(h) = \sum_{j=1}^m b_{ij} (p_j - c_j^0) \in r(\mathcal{I}_0)$$

and we have

$$p_i - c_i^0 + h(z_i(p, h) - \Delta_i(h)) = \sum_{j=1}^m (\delta_{ij} + h b_{ij}) (p_j - c_j^0).$$

The matrix $(\delta_{ij} + hb_{ij})$ is invertible, so the ideal generated by $H(p_i - c_i^0)$ in $H(\mathbb{C}[\mathfrak{g}^*][[h]])$ coincides with the ideal generated by $(p_i - c_i^0)$ in $H(\mathbb{C}[\mathfrak{g}^*][[h]])$. (For \star_T , the star ideal coincides with the commutative ideal.)

In order to state the main result we need a lemma.

Lemma 4.1. *Let \mathcal{J}_0 be the ideal in $(H(\mathbb{C}[\mathfrak{g}^*][[h]]), \star_T)$ generated by $(p_i - c_i^0)$ and let $\hat{\mathcal{J}}_0$ be the ideal in $(C^\infty(\mathcal{N}_\Theta)[[h]], \star_T)$ generated by the same generators. Then*

$$\hat{\mathcal{J}}_0 \cap H(\mathbb{C}[\mathfrak{g}^*][[h]]) = \mathcal{J}_0.$$

Proof. Since the product \star_T is tangential to the orbits the star ideals \mathcal{J}_0 and $\hat{\mathcal{J}}_0$ coincide with the ideals with respect to the commutative product, so we will limit ourselves to those.

One inclusion is obvious. For the other, let $b = \sum_{r=0}^{\infty} b_r h^r \in H(\mathbb{C}[\mathfrak{g}^*][[h]])$. Assume that

$$H(b) = \sum_{i=1}^m f^i H(p_i - c_i^0) \quad (8)$$

where $f^i = \sum_{r=0}^{\infty} f_r^i h^r \in C^\infty(\mathcal{N}_\Theta)[[h]]_{\mathbb{C}}$ are not unique. We need to prove that f^i can be chosen in $H(\mathbb{C}[\mathfrak{g}^*][[h]])$. We will show that there exist $q^i = \sum_{r=0}^{\infty} q_r^i h^r \in \mathbb{C}[\mathfrak{g}^*][[h]]$ such that $b_r = \sum_{i=1}^m q_r^i (p_i - c_i^0)$. This will clearly be enough.

By induction on r . For $r = 0$, we look at the order 0 in h of the equation (8) (we recall that $H = \text{Id mod}(h)$),

$$b_0 = \sum_i f_0^i (p_i - c_i^0) \in \mathbb{C}[\mathfrak{g}^*] \subset C^\infty(\mathcal{N}_\Theta)_{\mathbb{C}}.$$

It is not difficult to see that f_0^i can be chosen in $\mathbb{C}[\mathfrak{g}^*]$, so we set $q_0^i = f_0^i$.

We go to the general case. By the induction hypothesis, we assume that we have found q_0^i, \dots, q_r^i , with

$$b_0 + b_1 h + \dots + b_r h^r = \sum_{i=1}^m (q_0^i + q_1^i h + \dots + q_r^i h^r) (p_i - c_i^0).$$

Then

$$H(b) - H(b_0 + b_1 h + \dots + b_r h^r) = \sum_{i=1}^m (f^i - H(q_0^i + \dots + q_r^i h^r)) H(p_i - c_i^0)$$

so

$$h^{r+1} H(b_{r+1} + b_{r+2} h + \dots) = h^{r+1} \sum_{i=1}^m \left(f_{r+1}^i - \sum_{s+t=r+1} H_s(q_t^i) \right) H(p_i - c_i^0) \quad \text{mod}(h^{r+2}).$$

Since the ring $C^\infty(\mathcal{N}_\Theta)[[h]]_{\mathbb{C}}$ is torsion free we have

$$H(b_{r+1} + b_{r+2} h + \dots) = \sum_{i=1}^m \left(f_{r+1}^i - \sum_{s+t=r+1} H_s(q_t^i) \right) H(p_i - c_i^0) \quad \text{mod}(h).$$

Now if we look at the order 0 in h

$$b_{r+1} = \sum_{i=1}^m \left(f_{r+1}^i - \sum_{s+t=r+1} H_s(q_t^i) \right) (p_i - c_i^0),$$

as in the $r = 0$ case, if we set

$$q_{r+1}^i = f_{r+1}^i - \sum_{s+t=r+1} H_s(q_t^i)$$

it is not difficult to see that it can be chosen as a polynomial, which gives us the result. \square

Proposition 4.1. *Let Θ be a regular coadjoint orbit of a compact Lie group defined by the constraints*

$$p_i - c_i^0 \quad i = 1, \dots, m.$$

There is an injective homomorphism between the algebraic deformation of $\mathbb{C}[\Theta]$ defined by U_h/\mathcal{I}_h with \mathcal{I}_h generated by

$$P_i - c_i^0 + h\Delta_i(h)$$

and the differential deformation of $C^\infty(\Theta)_\mathbb{C}$ ($C^\infty(\Theta)[[h]]_\mathbb{C}, \star_{T\Theta}$), which is obtained via the Kontsevich formula (see section 3.2 for more details), provided the constants $\Delta_i(h)$ are chosen as in (7).

Proof. H is an algebra isomorphism onto its image. We have the commutative diagram

$$\begin{CD} \mathbb{C}[\mathfrak{g}^*][[h]] @>H>> H(\mathbb{C}[\mathfrak{g}^*][[h]]) \subset C^\infty(\mathcal{N}_\Theta)[[h]]_\mathbb{C} \\ @VV\pi V @VV\pi_H V \\ \mathbb{C}[\Theta][[h]] @>\tilde{H}>> H(\mathbb{C}[\mathfrak{g}^*][[h]])/H(\mathcal{I}_0) \subset C^\infty(\Theta)[[h]]_\mathbb{C} \end{CD}$$

The last inclusion follows from lemma 4.1. □

Remark 4.3. We want to note that the ideal \mathcal{I}_h used in the previous proposition is not, in general, the ideal used in geometric quantization. In fact, for $SU(2)$ it was shown in [8] that the latter is generated by

$$P - l(l + \hbar) \quad \hbar = \frac{h}{2\pi}.$$

(P is the Casimir of $\mathfrak{su}(2)$). But according to remark 4.2, (6) the ideal has either $c_i(h) = c_i^0$ or $\Delta^i(h)$ is an infinite series in h (an exponential). Then we have a contradiction.

Acknowledgments

We wish to thank A Levrero for helpful discussions.

Appendix A

In this appendix we want to give some standard definitions on deformations and star products that have been used throughout the text.

Definition A1. *Let $(\mathcal{A}, \{, \})$ be a Poisson algebra over \mathbb{R} . We say that the associative algebra $\mathcal{A}_{[[h]]}$ over $\mathbb{R}[[h]]$ is a formal deformation of \mathcal{A} if*

1. *there exists an isomorphism of $\mathbb{R}[[h]]$ -modules $\psi : \mathcal{A}[[h]] \longrightarrow \mathcal{A}_{[[h]]}$;*
2. *$\psi(f_1 f_2) = \psi(f_1)\psi(f_2) \text{ mod }(\hbar), \quad \forall f_1, f_2 \in \mathcal{A}[[h]]$;*
3. *$\psi(f_1)\psi(f_2) - \psi(f_2)\psi(f_1) = h\psi(\{f_1^0, f_2^0\}) \text{ mod }(\hbar^2), \quad \forall f_1, f_2 \in \mathcal{A}[[h]], f_i \equiv f_i^0 \text{ mod }(\hbar), i = 1, 2.$*

If $\mathcal{A}_\mathbb{C}$ is the complexification of a real Poisson algebra \mathcal{A} we can give the definition of formal deformation of $\mathcal{A}_\mathbb{C}$ by replacing \mathbb{R} with \mathbb{C} in the above definition.

The associative product in $\mathcal{A}[[h]]$ defined by

$$f \star g = \psi^{-1}(\psi(f) \cdot \psi(g)) \quad f, g \in \mathcal{A}[[h]] \tag{9}$$

is called the star product on $\mathcal{A}[[h]]$ induced by ψ .

A star product on $\mathcal{A}[[\hbar]]$ can also be defined as an associative $\mathbb{R}[[\hbar]]$ -linear product given by the formula

$$f \star g = fg + B_1(f, g)\hbar + B_2(f, g)\hbar^2 + \cdots \in \mathcal{A}[[\hbar]] \quad f, g \in \mathcal{A}$$

where the B_i are bilinear operators. By associativity of \star one has that $\{f, g\} = B_1(f, g) - B_1(g, f)$. So this definition is a special case of the previous one where $\mathcal{A}_\hbar = \mathcal{A}[[\hbar]]$ and \star is induced by $\psi = \text{Id}$.

Two star products on $\mathcal{A}[[\hbar]]$, \star and \star' are said to be *equivalent* (or *gauge equivalent*) if there exists $T = \sum_{n \geq 0} \hbar^n T_n$, with T_n linear operators on $\mathcal{A}[[\hbar]]$, $T_0 = \text{Id}$ such that

$$f \star g = T^{-1}(Tf \star' Tg).$$

If $\mathcal{A} \subset C^\infty(M)$ and the operators B_i are bidifferential operators we say that the star product is *differential*. If in addition $\mathcal{A} = C^\infty(M)$ and M is a real Poisson manifold, we will say that \star is a *differential star product on M* .

In [1] Kontsevich classifies differential star products on a manifold M up to gauge equivalence.

Theorem A1 (Kontsevich [1]). *The set of gauge equivalence classes of differential star products on a smooth manifold M can be naturally identified with the set of equivalence classes of Poisson structures depending formally on \hbar ,*

$$\alpha = \hbar\alpha_1 + \hbar^2\alpha_2 + \cdots$$

modulo the action of the group of formal paths in the diffeomorphism group of M , starting at the identity isomorphism.

In particular, for a given Poisson structure α_1 , we have the equivalence class of differential star products associated with $\hbar\alpha_1$. We will say that this is the equivalence class of star products canonically associated with the Poisson structure α_1 .

Also, an explicit universal formula to compute the bidifferential operators of the star product associated with any formal Poisson structure is given in [1] in the case of an arbitrary Poisson structure on flat space \mathbb{R}^n . The formula depends on the coordinates chosen, but it was also proved in [1] that the star products constructed with different choices of coordinates are gauge equivalent.

Let $\mathcal{A}_{\mathbb{C}} = \mathbb{C}[M_{\mathbb{C}}]$ be the coordinate ring of the complex algebraic affine variety $M_{\mathbb{C}}$ defined over \mathbb{R} whose real points are a real algebraic Poisson variety M . If the B_i are bilinear algebraic operators we will say that \star is an *algebraic star product on M* .

An example of great interest to us, of such M , is given by the dual \mathfrak{g}^* of the Lie algebra of a compact semisimple Lie group (see section 2).

The classification of algebraic star products is still an open problem [11].

Definition A2. *Let N be a submanifold of the Poisson manifold M and let \star_M be a star product on M . We say that \star_M is tangential to N if for $f, g \in C^\infty(N)$*

$$f \star_M g =_{\text{def}} (F \star_M G)|_N \quad \text{with} \quad f = F|_N \quad g = G|_N$$

is a well-defined star product on N , that is, if

$$(F - F')|_N = (G - G')|_N = 0 \quad \text{then} \quad F \star_M G|_N = F' \star_M G'|_N \quad (10)$$

for $F, F', G, G' \in C^\infty(M)$.

The same definition works for algebraic Poisson varieties, replacing the algebra of C^∞ functions with the algebra of polynomials on the varieties.

Given a Poisson manifold M , one can ask if there exists a differential star product on M , that is tangential to all the leaves of the symplectic foliation. For regular manifolds a positive answer was found in [6]. For $M = \mathfrak{g}^*$ foliated in coadjoint orbits, it was found in [13] that there is an obstruction to the existence. In particular, for a semisimple Lie algebra \mathfrak{g}^* it is not possible to find a differential star product on \mathfrak{g}^* which is tangential to all coadjoint orbits.

Appendix B

In this appendix, we want to give an explicit formula on how to construct a global star product starting from the star products defined on open sets of a manifold and satisfying certain conditions (see below). We will refer to this procedure as *gluing of star products*, and it will be used in section 3.

Let \star be a differential star product on a manifold M . Since the operators B_i that define \star are local, there are well defined star products \star_U on every open set U of M . We have a sheaf of algebras \mathcal{S} :

$$\mathcal{S}(U) = (C^\infty(U)[[h]], \star_U) \tag{11}$$

which we will call a *sheaf of star products*.

Let M be a Poisson manifold and fix an open cover $\mathcal{U} = \{U_r\}_{r \in J}$ where J is some set of indices. Assume that in each U_r there is a differential star product

$$\star_r : C^\infty(U_r)[[h]] \otimes C^\infty(U_r)[[h]] \longrightarrow C^\infty(U_r)[[h]].$$

This defines a collection of sheaves of star products

$$\mathcal{F}_r(V_r) = (C^\infty(V_r)[[h]], \star_r) \quad V_r \subset U_r. \tag{12}$$

It is a general fact in theory of sheaves that if there are isomorphisms of sheaves in the intersections

$$\begin{aligned} T_{sr} : \mathcal{F}_r(U_{rs}) &\longrightarrow \mathcal{F}_s(U_{sr}) & U_{r_1 \dots r_k} &= U_{r_1} \cap \dots \cap U_{r_k} \\ T_{sr}(f) \star_s T_{sr}(g) &= T_{sr}(f \star_r g) \end{aligned} \tag{13}$$

such that the following conditions are satisfied

$$\begin{aligned} 1. \quad T_{rs} &= T_{sr}^{-1} && \text{on } U_{sr}, \\ 2. \quad T_{ts} \circ T_{sr} &= T_{tr} && \text{on } U_{rst} \end{aligned} \tag{14}$$

then there exists a global sheaf \mathcal{F} on M isomorphic to the local sheaves \mathcal{F}_r on each U_r .

If the sheaves of star products (12) satisfy the conditions (14) with

$$T_{sr} = \text{Id} \pmod{h}$$

then we have a global sheaf of star products on M . The algebra of the global sections is $C^\infty(M)[[h]]$ together with a star product that we will call the *gluing* of local star products. We want to write an explicit formula for the star product of global sections.

We denote $U^{r_1 \dots r_k} = U_{r_1} \cup \dots \cup U_{r_k}$. Let us first consider the gluing on two open sets, say U_1 and U_2 , with non-trivial intersection. Let $\phi_1 : U_1 \rightarrow \mathbb{R}$, $\phi_2 : U_2 \rightarrow \mathbb{R}$ be a partition of unity of U^{12} ,

$$\phi_1(x) + \phi_2(x) = 1 \quad \forall x \in U^{12} \quad \text{supp}(\phi_r) \subset U_r.$$

Let $f_r \in C^\infty(U_r)[[h]]$ such that $f_s = T_{sr} f_r$ in U_{rs} . One can define an element $f \in C^\infty(U^{rs})[[h]]$ by $f = \phi_1 f_1 + \phi_2 f_2$. On the intersection U_{12} one has

$$f = \phi_1 f_1 + \phi_2 T_{21} f_1 = (\phi_1 \text{Id} + \phi_2 T_{21}) f_1 = A_{21} f_1.$$

Note that the operator $A_{21} = \text{Id} + \mathcal{O}(\hbar)$ is invertible. On U^{12} we can define the star product

$$f \star g = \begin{cases} (f_1 \star_1 g_1)(x) & \text{if } x \in U_1 - U_{12} \\ A_{21} (A_{21}^{-1}(f) \star_1 A_{21}^{-1}(g))(x) & \text{if } x \in U_{12} \\ (f_2 \star_2 g_2)(x) & \text{if } x \in U_2 - U_{12}. \end{cases} \quad (15)$$

It is easy to check that the star product is smooth.

One can do the gluing interchanging U_1 and U_2 . One has that on U_{12}

$$f = \phi_1 f_1 + \phi_2 f_2 = (\phi_1 T_{12} + \phi_2 \text{Id}) f_j = A_{12} f_2.$$

A_{12} is also invertible and

$$A_{21} = A_{12} T_{21}$$

provided $T_{12} = T_{21}^{-1}$. One can construct a star product on U^{12} using the same procedure as in (15). It is easy to check that both star products are identical.

The procedure in (15) can be generalized to an arbitrary number of open sets. Let $\phi_i : U_i \rightarrow \mathbb{R}$ be a partition of unity of M subordinate to the covering \mathcal{U} . We define $f \in C^\infty(M)$

$$f = \sum_{r \in J} \phi_r f_r \quad \text{where } f_r = T_{rs} f_s.$$

On U_r , f becomes

$$f = \left(\phi_r \text{Id} + \sum_s \phi_s T_{sr} \right) f_r = A_r f_r.$$

The star product on U_r is defined as

$$f \star g = A_r (A_r^{-1}(f) \star_r A_r^{-1}(g)). \quad (16)$$

Using conditions (14) one has

$$A_r T_{rt} = A_t.$$

Then, the star products (16) on each U_r coincide in the intersections, so they define a unique star product on M . The restriction of this star product to U_r is equivalent to \star_r . Also, using different partitions of unity one obtains equivalent star products.

References

- [1] Kontsevich M 1997 Deformation quantization of Poisson manifolds *Preprint math.QA/9709040*
- [2] Omori H, Maeda Y and Yoshioka A 1991 Weyl manifolds and deformation quantization *Adv. Math.* **85** 224–55
- [3] De Wilde M and Lecomte P B A 1983 Existence of star products and of formal deformations of the Poisson Lie algebra of arbitrary symplectic manifolds *Lett. Math. Phys.* **7** 487–96
- [4] Fedosov B 1994 A simple geometric construction of deformation quantization *J. Differential Geom.* **40** 213–38
- [5] Deligne P 1995 Déformations de l’algèbre des fonctions d’une variété symplectique: comparaison entre Fedosov et De Wilde, Lecomte *Selecta Math. (N.S.)* **1** 667–97
- [6] Masmoudi M 1992 Tangential deformation of a Poisson bracket and tangential star-products on a regular Poisson manifold *J. Geom. Phys.* **9** 155–71
- [7] Cahen M and Gutt S 1983 Produits $*$ sur les orbites des groupes semi-simples de Rang 1, *C.R. Acad. Sci. Paris* **296** 821–23
Cahen M and Gutt S 1987 An algebraic construction of $*$ product on the regular orbits of semisimple Lie groups *Gravitation and Cosmology (Monographs and Textbooks in Physical Sciences)* ed W Rundler and A Trautman A volume in honor of Ivor Robinson (Naples: Bibliopolis)
- Cahen M and Gutt S 1987 Non localité d’une déformation symplectique sur la sphère S^2 *Bull. Soc. Math. Belg.* **B 36** 207–21
- [8] Fioresi R and Lledó M A 1999 On the deformation quantization of coadjoint orbits of semisimple Lie groups 1999 *Pacific J. Math.* **198** 411–36 (*Preprint math.QA/9906104*)

- [9] Fioresi R, Levrero A and Lledó M A 2000 Algebraic and differential star products on regular orbits of compact Lie groups *Preprint* math.QA/0011172 (2000 *Pacific J. Math.* at press)
- [10] Lledó M A 2001 Deformation quantization of non regular orbits of compact Lie groups *Preprint* QA/0105191
- [11] Kontsevich M 2001 Deformation quantization of algebraic varieties *Preprint* math.AG/0106006
- [12] Kostant B 1978 Lie group representations on polynomial rings *Am. J. Math.* **85** 327
- [13] Cahen M, Gutt S and Rawnsley J 1996 On tangential star products for the coadjoint Poisson structure *Commun. Math. Phys.* **180** 99–108