

## Replica bounds for diluted non-Poissonian spin systems

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2003 J. Phys. A: Math. Gen. 36 10967

(<http://iopscience.iop.org/0305-4470/36/43/021>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.89

The article was downloaded on 02/06/2010 at 17:12

Please note that [terms and conditions apply](#).

# Replica bounds for diluted non-Poissonian spin systems

Silvio Franz<sup>1</sup>, Michele Leone<sup>2</sup> and Fabio Lucio Toninelli<sup>3</sup>

<sup>1</sup> The Abdus Salam International Center for Theoretical Physics, Condensed Matter Group, Strada Costiera 11, PO Box 586, I-34100 Trieste, Italy

<sup>2</sup> Institute for Scientific Interchange, viale Settimio Severo 65, Torino, Italy

<sup>3</sup> EURANDOM, Technische Universiteit Eindhoven, PO Box 513, 5600 MB Eindhoven, The Netherlands

E-mail: franz@ictp.trieste.it, leone@isiosf.isi.it and ftoninell@euridice.tue.nl

Received 5 June 2003, in final form 22 July 2003

Published 15 October 2003

Online at [stacks.iop.org/JPhysA/36/10967](http://stacks.iop.org/JPhysA/36/10967)

## Abstract

In this paper we extend replica bounds and free energy subadditivity arguments to diluted spin-glass models on graphs with arbitrary, non-Poissonian degree distribution. The new difficulties specific of this case are overcome introducing an interpolation procedure that stresses the relation between interpolation methods and the cavity method. As a byproduct we obtain self-averaging identities that generalize the Ghirlanda–Guerra ones to the multi-overlap case.

PACS number: 75.50.Lk

## 1. Introduction

The replica and the cavity methods play a fundamental role in the analysis of mean-field disordered systems, where they suggest a low-temperature glassy phase with a great extent of universality [1].

With the technique of ‘replica symmetry breaking’ (RSB) physicists have been since a long time able to describe in great detail the statistical properties of the configurational space of ‘fully connected’ disordered models, where each degree of freedom interacts with an extensive number of neighbours.

The last times have seen important progresses in extending the analysis to ‘diluted models’, where each degree of freedom interacts with a finite number of randomly chosen neighbours. Although the equations describing RSB in these models were well known already in the past, it is only with the recent introduction of a population dynamics algorithm [2, 3] that it has been possible to tackle the subtleties of the glassy low-temperature phases. Thanks to this algorithm and its specification to zero temperature it has been possible to analyse at the level of ‘one step replica symmetry breaking’ (1RSB) a great variety of problems relevant for physics of glassy systems or computer science or both. These range from the Viana–Bray model for

spin glasses [3], where the 1RSB is expected to give just an approximate result, to the diluted  $p$ -spin model or XOR-SAT problem [4], the Biroli–Mézard lattice glass model [5], models for efficient error correcting codes [6], the random  $K$ -SAT problem and graph colouring [7–9], where at least at zero temperature 1RSB is thought to be exact.

The population dynamic algorithm, originally thought for computing averages over random graph disorder, has been suitably generalized to deal with single samples, and in conjunction with decimation algorithms has been proved to be effective to solve random  $K$ -SAT or graph colouring in essentially linear time up to very close to the SAT/UNSAT transition threshold [8].

Despite these successes, together with a large wealth of important rigorous results on both mean-field and short-range models [10–12], the theoretical foundations of replica and cavity methods remain unsatisfactory. On one hand replica methods rely on a non-unique analytic continuation of the integer moments of the partition function which is difficult to control mathematically; on the other, the cavity method is based on physically sound, but unproven assumptions on the nature of the low-temperature pure states.

A great deal of mathematical work has been devoted to the analysis of disordered models with rigorous—i.e., conventional probabilistic—methods, which in many cases allow us to control the RS high-temperature case, and in the remarkable analysis of Talagrand [11] for the fully connected  $p$ -spin for large enough  $p$ , even the low-temperature one. A different approach has been put forward by Guerra [13], who was able to prove that in a large class of long-range models the free energy can be written as the sum of the replica expression plus a remainder which, by simple inspection, is proved to be positive. This was the first proof that the replica/cavity Ansatz is in some cases a variational Ansatz, therefore justifying the replica rule that the free energy should be maximized with respect to the replica parameters<sup>4</sup>. The technique of Guerra is based on the idea of interpolating the original model with a pure paramagnet with suitably chosen local external fields. In the original formulation the interpolation was performed tuning the relative strengths of the couplings and the external fields, and relayed heavily on the Gaussian and long-range nature of the interactions among spins. This technique has been employed in several papers dealing with long- and short-range models [14].

In a previous paper [15], hereafter referred to as I, two of us have shown how to generalize the technique of the interpolating model to diluted models on graphs with Poissonian degree distribution. In that case, the interpolation is done progressively removing addends in the Hamiltonian, and compensating in average by the introduction of appropriate external fields on random sites of the lattice. The method was then interpreted as a variational version of the cavity method. The result was that the free energy could be written as a sum of a replica/cavity like contribution plus a remainder. For the even  $p$  diluted  $p$ -spin model and the even  $K$   $K$ -SAT problems, simple considerations not relying on the actual solutions to the replica/cavity equations, allowed us to conclude that the remainder was positive. This means that the replicas give free energy lower bounds to these models. Specified to zero temperature, this proves that replicas give an upper bound for the satisfiability threshold, that is the value of the degree of connectivity below which all the terms of the Hamiltonian can be minimized at the same time. The cases of odd  $p$  or  $K$  led to an expression for the remainder whose sign cannot be obviously decided, thus leaving as an open question if the recently obtained 1RSB solution for the random 3-SAT problems or the odd  $p$ -spin case actually gives free energy lower bounds and SAT-threshold upper bounds.

<sup>4</sup> Cases are known in replica theory where the right free energy extremum is a saddle point, but this does not happen in the mentioned cases.

As we said, the results were confined to models on Erdős–Rényi (hyper)-graphs which have Poissonian degree distribution. However, the replica and the cavity methods have been successfully applied for spin models on more general graphs, where the degree distribution is basically arbitrary.

Motivated by that consideration, in this paper we generalize the analysis to random graphs with arbitrary connectivity distribution. Here, differently from the Poissonian case, the compensation in average is not possible and the erasure of a term in the Hamiltonian requires detailed compensation through the introduction of proper fields on the sites belonging to the erased clause. The new difficulties arising in this case can be overcome employing the property of self-averaging of appropriate observables, leading to identities on multi-overlap distributions that generalize the Ghirlanda–Guerra identities [16] for the usual overlaps. This new feature is common both to the RS and the RSB case. For exposition simplicity we will limit ourselves in this paper to the discussion of the RS case.

This paper is organized as follows: in section 2 we define the model and introduce the notation, in section 3 we introduce the interpolating models, which we use in sections 4 and 5 to find representations of the free energy as main terms plus remainders, suitable respectively to prove replica bounds and subadditivity. In section 6 we use the self-averaging to estimate some terms in the remainders and generalize the Ghirlanda–Guerra identities. Section 7 is devoted to showing the positivity of the remainders, and finally in section 8 we discuss some conclusions.

## 2. Definition of the models and summary of mathematical notation

### 2.1. The models

In this paper we consider diluted spin models on random graphs with arbitrary degree distribution, consisting in a collection of  $N$  Ising  $\pm 1$  spins  $\mathbf{S} = \{S_1, \dots, S_N\}$ , interacting through Hamiltonians of the kind

$$\mathcal{H}^{(M)}(\mathbf{S}, \mathbf{J}) = \sum_{\mu=1}^M H_{J^{(\mu)}}(S_{i_1^\mu}, \dots, S_{i_p^\mu}) \quad (1)$$

where the indices  $i_l^\mu$  are i.i.d. quenched random variables chosen in the following way: one first extracts a set of site degrees  $k_i$ , representing the number of clauses where  $S_i$  appears ( $i = 1, \dots, N$ ), as i.i.d. from a distribution  $p_k$ . Then the configuration of indices  $i_l^\mu$  is chosen uniformly among all the possible ways respecting the prescribed degrees. In other words, the joint probability of all the indices  $\{i_l^\mu\}$  will be proportional to

$$\prod_{i=1}^N \delta \left( \sum_{\mu=1}^M \sum_{l=1}^p \delta_{i_l^\mu, i} - k_i \right) \quad (2)$$

where both  $\delta$  appearing in (2) denote the Kronecker symbol. We will concentrate on regular degree distributions where all the moments  $\langle k^l \rangle = \sum_k p_k k^l$  are finite. The number of clauses  $M$  is therefore itself in principle a random variable given by  $M = \frac{1}{p} \sum_{i=1}^N k_i$ , its average  $\langle M \rangle = \alpha N$  will be proportional to  $N$ , with  $\alpha = \langle k \rangle / p$  and it will have small  $O(\sqrt{N})$  fluctuations around its average. For this reason, we will often treat  $M$  as a constant, making only a  $O(1/N)$  error in the free energy. As will become soon clear, the subscript  $J^{(\mu)}$  in the clauses indicates dependence on a single or on a set of quenched random variables.

We will treat in a unitary way the case of the  $p$ -spin model [17] or random XOR-SAT problem of computer science, and the random  $K$ -SAT model. In the  $p$ -spin model the clauses have the form

$$H_{J^{(\omega)}}(S_{i_1^\mu}, \dots, S_{i_p^\mu}) = J^\mu S_{i_1^\mu} \cdots S_{i_p^\mu} \quad (3)$$

where  $J^\mu$  will be taken as i.i.d. random variable with regular symmetric distribution  $\mu(J) = \mu(-J)$ . Particular attention will be given to Viana–Bray model corresponding to  $p = 2$ . In random  $K$ -SAT model the clauses have the form [18]

$$H_{J^{(\omega)}}(S_{i_1^\mu}, \dots, S_{i_p^\mu}) = \prod_{l=1}^p \frac{1 + J_{i_l^\mu}^\mu S_{i_l^\mu}}{2} \quad (4)$$

where the quenched variables  $J_{i_l^\mu}^\mu = \pm 1$  are i.i.d. with symmetric probability<sup>5</sup>. The number  $p$  of spins appearing in a clause is usually called  $K$  in the  $K$ -SAT problem, but for uniformity of notation we will deviate from this convention.

In I the indices  $i_l^\mu$  of the spins appearing in the clauses were chosen with uniform probability, giving rise to random graphs with Poissonian degree statistics, and the treatment was based on the peculiar property of the Poisson distribution. Here we show that thanks to self-averaging of certain quantities, the validity of the results does not depend on the specific form of the graph degree distribution.

## 2.2. Notation

Let us establish some notation. We will need several kinds of averages:

- The Boltzmann–Gibbs average for fixed quenched disorder: given an observable  $A(\mathbf{S})$

$$\omega(A) = \frac{\sum_{\mathbf{S}} A(\mathbf{S}) \exp(-\beta \mathcal{H}(\mathbf{S}, \mathbf{J}))}{Z} \quad (5)$$

where  $Z = \sum_{\mathbf{S}} \exp(-\beta \mathcal{H}(\mathbf{S}, \mathbf{J}))$  and  $\beta$  is the inverse temperature.

Obviously,  $\omega(A)$ , as well as  $Z$  will be functions of the quenched variables, the size of the system and the temperature. This dependence will be made explicit only when needed.


- The disorder average: given an observable quantity  $B$  dependent on the quenched variables appearing in the Hamiltonian, we will denote as  $E(B)$  its average. This will include the average with respect to the  $J$  variables and the choice of the random indices in the clauses as well as with respect to other quenched variables to be introduced later.
- We will need in several occasions the ‘replica measure’

$$\Omega(A_1, \dots, A_n) = \omega(A_1) \cdots \omega(A_n). \quad (6)$$

- Moreover, throughout the work we will consider modified versions of the original Hamiltonians that will depend on a discrete dilution parameter  $t$ . Both Boltzmann and disorder averages will depend accordingly on  $t$ . The original averages will correspond to  $t = M$ .
- Another notation we will have the occasion to use is the one for the overlaps among  $l$  spin configurations  $\{S_i^{a_1}, \dots, S_i^{a_l}\}$ , out of a population of  $n\{S_i^1, \dots, S_i^n\}$ :

$$q^{(a_1, \dots, a_l)} = \frac{1}{N} \sum_{i=1}^N S_i^{a_1} \cdots S_i^{a_l} \quad (1 \leq a_r \leq n \quad \forall r) \quad (7)$$

<sup>5</sup> While the assumption of a symmetric distribution  $\mu(J) = \mu(-J)$  will play an important role in establishing the free energy bounds, the precise form of the distribution will not be essential.



**Figure 1.** Clause erasure and balancing fields addition for a particular random clause  $\mu$ . The hyper-edge interaction is drawn in the factor-graph notation with square nodes representing clauses and circles representing spins. Dotted lines represent the added balancing fields. Only clause  $\mu$  and corresponding variables and fields are explicitly named.

and in particular

$$q^{(n)} = q^{(1, \dots, n)} = \frac{1}{N} \sum_{i=1}^N S_i^1 \cdots S_i^n. \quad (8)$$

- Finally, we will denote as  $T_i$  the set of clause indices where the  $i$ 'th spin appears and  $V_\mu$  the set of spin indices belonging to the clause  $\mu$ .

In the following we will need to consider averages where some of the variables are excluded, e.g., the averages when a variable  $u_i^{k_i}$  is erased. These averages will be denoted with a subscript  $-u_i^{k_i}$ , e.g., if an  $\omega$  average is concerned the notation will be  $\omega_{-u_i^{k_i}}(\cdot)$ . Other notation will be defined later in the text whenever needed.

Our interest will be confined to bounds to the free energy density  $F_N = -\frac{1}{\beta N} E \log Z$  and the ground-state energy density  $U_{GS} = \lim_{N \rightarrow \infty} 1/N E[\min(U_N)]$  valid in the thermodynamic limit, so that  $O(1/N)$  will be often implicitly neglected in our calculations.

### 3. Introducing the interpolating models

We will use the technique of interpolating models for the purposes of showing the existence of the infinite volume free energy, and of proving replica bounds. For exposition reasons, we will explain the method first for the replica bounds and later for the existence of the free energy.

#### 3.1. The replica/cavity bounds: clause deletion versus fields compensation

In order to prove the free energy bounds, we will use an iterative discrete graph pruning procedure where at each time step  $t$  decreasing from  $M$  to  $0$ , we erase the clause labelled by  $t$  and compensate this reduction by the introduction of some auxiliary fields  $u_{i_l}^t$  on the sites  $i_l^t$  ( $l = 1, \dots, p$ ) belonging to the clause  $t$  (see figure 1). At each time step  $t$  we have a

different model with Hamiltonian

$$\mathcal{H}^{(t)}[\mathbf{S}] = \sum_{\mu=1}^t H_{J^{(\mu)}}(S_{i_1^\mu}, \dots, S_{i_p^\mu}) - \sum_{\mu=t+1}^M \sum_{l=1}^p u_{i_l^\mu}^\mu S_{i_l^\mu} \quad (9)$$

which interpolates between a simple paramagnet made of non-interacting spins at time 0 and the original model at time  $M$ . The clause removal procedure is very similar to the analogous operation of spin and clause addition usually performed in the cavity method, the difference being that here the deletion is explicitly compensated by the introduction of external fields. A key point in obtaining the free energy replica/cavity bounds is to assume that the *external* fields are random variables obeying the statistics of the *cavity* fields in the cavity approach.

In order to explain this point, and to motivate the introduction of the definitions below, let us remind some formulae of the cavity approach. In that context, one singles out the contribution of the clauses and the sites to the free energy and defines cavity fields  $h_i^\mu$  and  $u_i^\mu$  respectively as the local field acting on the spin  $i = i_1^\mu$  in the absence of the clause  $\mu$  and the local field acting on  $i$  due to the presence of the clause  $\mu$  only. If we define  $Z[S_i]$  as the partition function of a given sample with  $N$  spins where all but the spin  $S_i$  are integrated, and  $F_{N,-i}$  as the free energy of the corresponding systems where the spin  $S_i$  and all the clauses which belong to it are removed, in the cavity approach one assumes that

$$\begin{aligned} Z[S_i] &\approx e^{-\beta F_{N,-i}} \prod_{\mu \in T_i} \sum_{S_2^\mu, \dots, S_p^\mu} e^{-\beta H_{J^{(\mu)}}(S_i, S_2^\mu, \dots, S_p^\mu) + \beta \sum_{l=2}^p h_{i_l^\mu}^\mu S_{i_l^\mu}} \\ &= e^{-\beta F_{N,-i}} \prod_{\mu \in T_i} B_\mu^{(i)} e^{\beta u_i^\mu S_i}. \end{aligned} \quad (10)$$

Note that in general, if (10) had to represent the exact integration of  $N - 1$  spins in a finite  $N$  systems, effective  $l$ -spin interactions ( $l = 2, \dots, p$ ) should also be present among the spins explicitly appearing on the rhs of (10). Given the peculiar topology of random graphs, where loops of interacting variables have generically length  $O(\log(N))$ , these coupling can be expected to be small and are neglected in the approach. This is the reason why we wrote  $\approx$  instead of  $=$ . The constant  $B_\mu^{(i)} = e^{-\beta \Delta F_\mu^{(i)}}$  is interpreted as a suitable shift in the free energy due to the contribution of the clause  $\mu$  for fixed value of the spin  $i$ . We note that denoting  $J^\mu$  as  $J$ , and renaming the fields in (10) into  $h_1, \dots, h_{p-1}$ , equation (10) defines functions

$$u_J(h_1, \dots, h_{p-1}) \quad \text{and} \quad B_J(h_1, \dots, h_{p-1}) \quad (11)$$

that one can compute explicitly using the form of the clauses of the different models. For instance, in the case of the Viana–Bray model, where  $p = 2$ , one has

$$u_J(h) = \frac{1}{\beta} \tanh^{-1}[\tanh(\beta J) \tanh(\beta h)] \quad (12)$$

$$B_J(h) = \frac{2 \cosh(\beta J) \cosh(\beta h)}{\cosh(\beta u_J(h))}. \quad (13)$$

Always in the cavity approach one closes the set of equations imposing the self-consistent equations

$$h_i^{(\mu)} = \sum_{v \in \{T_i - \mu\}} u_v^{(i)}. \quad (14)$$

Following the intuition based on equation (10) we will define the external fields  $u_{i_l^\mu}^\mu$  in the interpolating model as verifying the relation

$$u_{i_l^\mu}^\mu = u_{J^{(\mu)}}(g_{i_1^\mu}^\mu, \dots, g_{i_{l-1}^\mu}^\mu, g_{i_{l+1}^\mu}^\mu, \dots, g_{i_p^\mu}^\mu) \quad (15)$$

where the  $p - 1$  arguments  $g$  will be independent variables with suitable statistics. We do not write at this point a relation analogous to the self-consistency relation (14). This equation will be obeyed in average when the statistics of the fields  $g$  is chosen in such a way to optimize the replica bounds. The key point of the procedure consists in the choice of the distribution of the primary fields  $g_{i_t^\mu}^\mu$ . Different free energy bounds can be obtained assuming the type of statistics implied by the different replica solutions.

In this paper we concentrate on the replica symmetric bound, that will be obtained considering the primary fields  $g_{i_t^\mu}^\mu$  as i.i.d. variables drawn from a distribution  $G(g)$ . Correspondingly the  $u$  will be distributed according to  $Q(u)$  verifying

$$Q(u) = \int dg_1 G(g_1) \cdots dg_{p-1} G(g_{p-1}) \langle \delta(u - u_J(g_1, \dots, g_{p-1})) \rangle_J. \quad (16)$$

Writing the free energy of the interpolating model as

$$F_N(t) = -\frac{1}{\beta N} E \log Z(t) \quad (17)$$

we observe that the average free energy  $F_N = F_N(M)$  of the original model can be written as

$$F_N = \sum_{t=1}^M \Delta_N(t) + F(0) \quad (18)$$

where we defined the discrete time derivative of the free energy as

$$\Delta_N(t) = F_N(t) - F_N(t - 1). \quad (19)$$

The bound will consist in comparing  $\Delta_N(t)$  with the corresponding replica symmetric expression.

The extension to 1RSB bounds, although technically more involved, does not present additional conceptual difficulties. In I, it has been shown how to provide 1RSB bounds in the Poissonian case. One needs to assume that the fields  $g_{i_t^\mu}^\mu$  are still independent variables, but subject to site and clause distributions  $G_{i_t^\mu}^\mu$  which are themselves random and independent from one another and subject to the common functional distribution  $\mathcal{G}(G)$ . The appropriate definition of interpolating free energy will depend on a parameter  $m \in [0, 1]$ :


$$F_N(t) = -\frac{1}{m\beta N} E_1 \log E_2 Z(t)^m. \quad (20)$$

We denoted by  $E_2$  the average over the fields  $g_{i_t^\mu}^\mu$  for fixed distributions  $G_{i_t^\mu}^\mu$  and by  $E_1$  the average over the distributions  $G_{i_t^\mu}^\mu$  as well as over all the other quenched variables. For  $t = M$  the  $E_2$  average is immaterial and formula (18) holds. For general degree distribution one could follow the same procedure, but in order to keep this paper within a reasonable length we will not present here this case.

### 3.2. The thermodynamic limit: a cut and paste procedure

The aim of this section is to interpolate between the original system of  $N$  interacting spins and two separate models, respectively with  $N_1$  and  $N_2$  spins ( $N_1 + N_2 = N$ ) and to show that the free energy is subadditive [19]. For notational simplicity we consider here explicitly just the case of the Viana–Bray model, with Hamiltonian  $H = -\sum_{\mu=1}^M J^\mu S_{i_1^\mu} S_{i_2^\mu}$ . Inspired by the construction of the previous section we start from the model with  $N$  spins and consider the sets of the first  $N_1$  spins and that of the remaining  $N_2$  spins. Each of the  $M$  clauses will either belong to the first sub-system, if  $i_1^\mu, i_2^\mu \in \{1, \dots, N_1\}$ , or to the second if  $i_1^\mu, i_2^\mu \in \{N_1 + 1, \dots, N\}$





**Figure 2.** Cut and paste procedure in the Viana-Bray model. The groups of constraints not belonging to  $M_b$  and of graph sites not participating to the constraints in  $M_b$  are generically represented as dashed line sets.

or they will be ‘bridge’ clauses if one of the indices is less than or equal to  $N_1$  and the other is greater than  $N_1$ . Let us denote as  $M_1(0)$ ,  $M_2(0)$  and  $M_b(0)$  the number of clauses of the different types, respectively. We define our interpolating model via an iterative ‘cut and paste’ procedure where at each time step we select at random two bridge clauses, we cut them, and we reconnect the spins belonging to the first sub-system between themselves with a new random coupling, and similarly for the two spins belonging to the second sub-system (see figure 2).

In such a way at each time step  $t = 1, \dots, M_b(0)/2$  (we suppose for simplicity to choose the ordering of the spins in such a way that  $M_b(0)$  is even) each spin conserves its original connectivity, and the number of clauses of the different kinds are modified as  $M_1(t) = M_1(t - 1) + 1$ ,  $M_2(t) = M_2(t - 1) + 1$  and  $M_b(t) = M_b(t - 1) - 2$ . At the end of the procedure we have two separate models with, respectively,  $N_1$  spins and  $M_1(0) + M_b(0)/2$  clauses and  $N_2$  spins and  $M_2(0) + M_b(0)/2$  clauses. Note that  $M_1$ ,  $M_2$  and  $M_b$  are random variables with average proportional to  $N$  and small  $O(\sqrt{N})$  fluctuations. As in the case of  $M$ , we will neglect these harmless fluctuations in our analysis. It is easy to realize that the graphs associated with the resulting non-interacting models are chosen with uniform probability among the ones having the correct connectivities for each spin. This can be explicitly checked by an elementary induction calculation. Each step of the cut and paste procedure transforms the uniform distribution on the graphs with prescribed connectivities  $k_i$  of the sites and number of clauses of the three types  $M_1(t)$ ,  $M_2(t)$  and  $M_b(t)$ , into the uniform distribution with the same connectivities but with clause numbers  $M_1(t) + 1$ ,  $M_2(t) + 1$  and  $M_b(t) - 2$ .

Therefore, in analogy with (18) we have

$$F_N = \frac{N_1}{N} F_{N_1} + \frac{N_2}{N} F_{N_2} - \sum_{t=1}^{M_b(0)/2} \Delta'_N(t) \tag{21}$$

where  $\Delta'_N(t)$  is the discrete time derivative along the cut and paste procedure. As will be shown in section 5,  $\Delta'_N(t)$  is non-negative for the models we are considering, as long as  $p$  is even, thus implying subadditivity of the free energy, and therefore the existence of its infinite volume limit.

#### 4. The RS bound

Let us start with a few observations independent of the chosen statistics of the primary fields, and therefore valid both for the RS and for the RSB estimate.

In order to compare the free energies  $F(t)$  and  $F(t-1)$ , let us consider  $Z(t-1)$  and isolating the terms containing the fields  $u_{i_l}^t$  observe that this can be written as

$$Z(t-1) = Z_{-u^t}(t-1) \omega_{-u^t}^{(t-1)} \left( e^{\beta \sum_{l=1}^p u_{i_l}^t S_{i_l}^t} \right) \quad (22)$$

where  $Z_{-u^t}(t-1)$  and  $\omega_{-u^t}^{(t-1)}(\cdot)$  are respectively the partition function and the Boltzmann averages corresponding to the Hamiltonian at time  $t-1$  in the absence of the fields  $u_{i_l}^t$ . In the same way, isolating the  $t$ th clause term in  $Z(t)$  and noting that  $Z_{-J^{(t)}}(t) = Z_{-u^t}(t-1)$  and  $\omega_{-J^{(t)}}^{(t)}(\cdot) = \omega_{-u^t}^{(t-1)}(\cdot)$  we can write

$$Z(t) = Z_{-u^t}(t-1) \omega_{-u^t}^{(t-1)} \left( e^{-\beta H_{J^{(t)}}(S_{i_1}^t, \dots, S_{i_p}^t)} \right). \quad (23)$$

Using these relations in the definition of the RS free energy time derivative  $\Delta_N(t)$  we find

$$N \Delta_N(t) = -T \left[ E \left( \ln \omega_{-u^t}^{(t-1)} \left( e^{-\beta H_{J^{(t)}}(S_{i_1}^t, \dots, S_{i_p}^t)} \right) \right) - E \left( \ln \omega_{-u^t}^{(t-1)} \left( e^{\beta \sum_{l=1}^p u_{i_l}^t S_{i_l}^t} \right) \right) \right]. \quad (24)$$

Now we compare this term with that obtained in the cavity approach (supposing that the statistics of the cavity fields  $h$  coincide with that of our external primary fields  $g$ ):

$$\omega_{-u^t}^{(t-1)} \left( e^{-\beta H_{J^{(t)}}(S_{i_1}^t, \dots, S_{i_p}^t)} \right) \approx \frac{1}{\prod_{l=1}^p 2 \cosh(\beta h_{i_l}^t)} \sum_{S_1, \dots, S_p} e^{-\beta H_{J^{(t)}}(S_{i_1}^t, \dots, S_{i_p}^t) + \beta \sum_{l=1}^p h_{i_l}^t S_l} \quad (25)$$

and in the same way

$$\omega_{-u^t}^{(t-1)} \left( e^{\beta \sum_{l=1}^p u_{i_l}^t S_{i_l}^t} \right) \approx \frac{1}{\prod_{l=1}^p 2 \cosh(\beta h_{i_l}^t)} \sum_{S_1, \dots, S_p} e^{\beta \sum_{l=1}^p u_{i_l}^t S_l + \beta \sum_{l=1}^p h_{i_l}^t S_l}. \quad (26)$$

This leads to the expression

$$N \Delta_N(t) \approx -T \left[ E \log \left( 2^{-p} \sum_{S_1, \dots, S_p} e^{-\beta H_{J^{(t)}}(S_1, \dots, S_p)} \prod_{l=1}^p (1 + S_l \tanh(\beta h_{i_l}^t)) \right) - E \sum_{l=1}^p \left( \log(\cosh(\beta u_{i_l}^t)) + \log(1 + \tanh(\beta h_{i_l}^t) \tanh(\beta u_{i_l}^t)) \right) \right]. \quad (27)$$

The replica symmetric approximation to this expression consists in assuming that the statistics of the cavity fields  $h_{i_l}^t$  coincide with that of the external fields  $g_{i_l}^t$ , and we call  $\Delta^*(t)$  the expression corresponding to (27), once these substitutions have been made.

Now, we go back to our approach where, it is important to emphasize, we are not assuming the validity of approximations such as (25), (26). In order to get a control of the free energy we add and subtract  $\Delta^*(t)$  from the expression (24) of  $\Delta_N(t)$ . It is also useful to add and subtract the term

$$T \sum_{l=1}^p E \log \left( 1 + \omega_{-u_{i_l}^t}^{(t-1)}(S_{i_l}^t) \tanh(\beta u_{i_l}^t) \right). \quad (28)$$

Rearranging terms in  $\Delta_N$  and taking into account that

$$F(0) = -\frac{1}{\beta N} E \sum_{i=1}^N \log \left( 2 \cosh \left( \beta \sum_{\mu \in T_i} u_i^\mu \right) \right) \quad (29)$$

we rewrite the free energy as

$$F_N = F_{\text{var}}[G] + \frac{1}{N} \sum_{t=1}^M R[G, t] + \frac{1}{N} \sum_{t=1}^M \tilde{R}[G, t] + O(1/N). \quad (30)$$

Here, we have isolated the ‘variational term’  $F_{\text{var}}[G]$

$$\begin{aligned}
 F_{\text{var}}[G] = & -\frac{1}{\beta N} E \sum_{i=1}^N \log \left( 2 \cosh \left( \beta \sum_{\mu \in T_i} u_i^\mu \right) \right) \\
 & - \frac{1}{\beta N} \sum_{t=1}^M \left[ E \log \left( 2^{-p} \sum_{S_1, \dots, S_p} e^{-\beta H_{J^{(t)}}(S_1, \dots, S_p)} \prod_{l=1}^p (1 + S_l \tanh(\beta g_{i_l}^t)) \right) \right. \\
 & \left. - E \sum_{l=1}^p (\log(\cosh(\beta u_{i_l}^t)) + \log(1 + \tanh(\beta g_{i_l}^t) \tanh(\beta u_{i_l}^t))) \right] \quad (31)
 \end{aligned}$$

from the remainders

$$\begin{aligned}
 R[G, t] = & -T \left[ E(\log \omega_{-u^t}^{(t-1)}(e^{-\beta H_{J^{(t)}}(S_{i_1}^t, \dots, S_{i_p}^t)})) - E \left( \sum_{l=1}^p \log \left( 1 + \omega_{-u_{i_l}^t}^{(t-1)}(S_{i_l}^t) \tanh(\beta u_{i_l}^t) \right) \right) \right. \\
 & - E \log \left( 2^{-p} \sum_{S_1, \dots, S_p} e^{-\beta H_{J^{(t)}}(S_1, \dots, S_p)} \prod_{l=1}^p (1 + S_l \tanh(\beta g_{i_l}^t)) \right) \\
 & \left. + E \sum_{l=1}^p (\log(1 + \tanh(\beta g_{i_l}^t) \tanh(\beta u_{i_l}^t))) \right] \quad (32)
 \end{aligned}$$

and

$$\begin{aligned}
 \tilde{R}[G, t] = & T \left[ E \left( \log \omega_{-u^t}^{(t-1)} \left( \prod_{l=1}^p (1 + S_{i_l}^t \tanh(\beta u_{i_l}^t)) \right) \right) \right. \\
 & \left. - E \left( \sum_{l=1}^p \log \left( 1 + \omega_{-u_{i_l}^t}^{(t-1)}(S_{i_l}^t) \tanh(\beta u_{i_l}^t) \right) \right) \right]. \quad (33)
 \end{aligned}$$

Expressions (31)–(33) are suggestive on how to use the interpolating model method to analyse single samples [20]. Moreover, we can simplify them observing that

- All the  $u$  and  $g$  fields appearing are statistically independent and one can therefore assign them arbitrary indices.
- The marginal of the index probability (2) with respect to the indices appearing in the  $t$ th clause is the uniform distribution on  $\{1, \dots, N\}^p$ . The indices appearing in the averages in (31)–(33) are therefore immaterial.

We finally find

$$\begin{aligned}
 \beta F_{\text{var}}[G] = & -E \log \left( 2 \cosh \left( \beta \sum_{l=1}^k u_l \right) \right) \\
 & - \alpha \left[ E \log \left( 2^{-p} \sum_{S_1, \dots, S_p} e^{-\beta H_J(S_1, \dots, S_p)} \prod_{l=1}^p (1 + S_l \tanh(\beta g_l)) \right) \right. \\
 & \left. - E \sum_{l=1}^p (\log(\cosh(\beta u_l)) + \log(1 + \tanh(\beta g_l) \tanh(\beta u_l))) \right] \quad (34)
 \end{aligned}$$

$$\begin{aligned}
R[G, t] = -T & \left[ E \left( \log \omega_{-u'}^{(t-1)} \left( e^{-\beta H_J(S_1, \dots, S_p)} \right) \right) - pE \left( \log \left( 1 + \omega_{-u'}^{(t-1)}(S_{i'_1}) \tanh(\beta u) \right) \right) \right. \\
& - E \log \left( 2^{-p} \sum_{S_1, \dots, S_p} e^{-\beta H_J(S_1, \dots, S_p)} \prod_{l=1}^p (1 + S_l \tanh(\beta g_l)) \right) \\
& \left. + pE \left( \log(1 + \tanh(\beta g) \tanh(\beta u)) \right) \right] \quad (35)
\end{aligned}$$

$$\begin{aligned}
\tilde{R}[G, t] = T & \left[ E \left( \log \omega_{-u'}^{(t-1)} \left( \prod_{l=1}^p (1 + S_{i'_l} \tanh(\beta u_l)) \right) \right) \right. \\
& \left. - pE \left( \log \left( 1 + \omega_{-u'}^{(t-1)}(S_{i'_1}) \tanh(\beta u) \right) \right) \right]. \quad (36)
\end{aligned}$$

Direct inspection upon optimizing over the  $G$  function shows that  $F_{\text{var}}[G]$  coincides with the free energy found with the replica/cavity method in the replica symmetric approximation. The first remainder term  $R[G, t]$  is analogous to that one finds on Poissonian degree graphs and will be dealt with in section 7. Conversely,  $\tilde{R}$  was absent in the Poissonian case thanks to the possibility of compensating in average the clause removal procedure and represents a new difficulty for graphs of arbitrary connectivity. At a first sight it would seem difficult to say anything in general about its behaviour. However, we will see in the next section that, thanks to the self-averaging property of extensive quantities, this term becomes vanishingly small in the thermodynamic limit. This is the only point, besides trivial  $O(1/N)$  terms neglectation where the large- $N$  limit enters in our estimates.

## 5. Thermodynamic limit

In order to analyse the cut and paste algorithm let us introduce  $Z(M_1, M_2, M_b)$  as the partition function of the interpolating model when the number of various kinds of clauses are given by  $M_1, M_2, M_b$ , and  $\omega_{(M_1, M_2, M_b)}(\cdot)$  as the corresponding Gibbs average. A step of the algorithm, at generic time  $t$ , sends  $Z(M_1, M_2, M_b)$  to  $Z(M_1 + 1, M_2 + 1, M_b - 2)$ . Let us consider  $s$  consecutive steps. If we denote as  $i'_1, i'_2$  and  $j'_1, j'_2$ ,  $r = 1, \dots, s$  the indices of the bridge clauses we cut and as  $J_r^1$  and  $J_r^2$  the corresponding couplings, we can write

$$\begin{aligned}
Z(M_1, M_2, M_b) &= Z(M_1, M_2, M_b - 2s) \omega_{(M_1, M_2, M_b - 2s)} \left( e^{\beta \sum_r (J_r^1 S_{i'_1} S_{i'_2} + J_r^2 S_{j'_1} S_{j'_2})} \right) \\
Z(M_1 + s, M_2 + s, M_b - 2s) &= Z(M_1, M_2, M_b - 2s) \omega_{(M_1, M_2, M_b - 2s)} \\
&\quad \times \left( e^{\beta \sum_r (J_r^1 S_{i'_1} S_{j'_1} + J_r^2 S_{i'_2} S_{j'_2})} \right). \quad (37)
\end{aligned}$$

The total average free energy change  $\Delta_N^{(s)}$  after the  $s$  steps is given therefore by

$$\begin{aligned}
\beta N \Delta_N^{(s)} &= \beta N \sum_{\tau=t}^{t+s-1} \Delta'_N = E \left( \log \omega_{(M_1, M_2, M_b - 2s)} \right. \\
&\quad \times \left( \prod_r (1 + \tanh(\beta J_r^1) S_{i'_1} S_{i'_2}) (1 + \tanh(\beta J_r^2) S_{j'_1} S_{j'_2}) \right) - \log \omega_{(M_1, M_2, M_b - 2s)} \\
&\quad \left. \times \left( \prod_r (1 + \tanh(\beta J_r^1) S_{i'_1} S_{j'_1}) (1 + \tanh(\beta J_r^2) S_{i'_2} S_{j'_2}) \right) \right). \quad (38)
\end{aligned}$$

Similarly to section 4 we add and subtract suitable terms to  $\Delta_N^{(s)}$ , to write

$$\beta N \Delta_N^{(s)} = 2s E \left\{ \log \omega_{(M_1, M_2, M_b - 2s)} (1 + \tanh(\beta J^1) S_{i_1} S_{i_2}) \right\} - s E \left\{ \log \omega_{(M_1, M_2, M_b - 2s)} (1 + \tanh(\beta J^1) S_{i_1} S_{j_1}) \right\} - s E \left\{ \log \omega_{(M_1, M_2, M_b - 2s)} (1 + \tanh(\beta J^2) S_{i_2} S_{j_2}) \right\} + \tilde{T} \tag{39}$$

where, for symmetry reasons, we have omitted the index  $r$ . Since we have factored the Gibbs averages, the remainder  $\tilde{T}$  will contain terms of the type

$$E \left[ \log \omega_{(M_1, M_2, M_b - 2s)} \left( \prod_r (1 + \tanh(\beta J_r^1) S_{i_r^1} S_{j_r^1}) (1 + \tanh(\beta J_r^2) S_{i_r^2} S_{j_r^2}) \right) - \sum_r \log \omega_{(M_1, M_2, M_b - 2s)} (1 + \tanh(\beta J_r^1) S_{i_r^1} S_{j_r^1}) - \sum_r \log \omega_{(M_1, M_2, M_b - 2s)} (1 + \tanh(\beta J_r^2) S_{i_r^2} S_{j_r^2}) \right]. \tag{40}$$

As we will briefly discuss in section 6, the same kind of self-averaging properties that will ensure the vanishing of  $\tilde{R}$  in the thermodynamic limit, will also guarantee that  $\tilde{T} \rightarrow 0$ . At this point, as in I, one expands the logarithms in absolutely convergent Taylor series of the variables  $\tanh(\dots)$ . Recalling the fact that the distribution of the  $J$  variables is even, one obtains

$$\beta N \Delta_N^{(s)} = s \sum_{n=1}^{\infty} \frac{\langle \tanh^{2n} \beta J \rangle}{2n} E \left[ (\omega_{(M_1, M_2, M_b - 2s)} (S_{i_1} S_{j_1}))^{2n} - 2 (\omega_{(M_1, M_2, M_b - 2s)} (S_{i_1} S_{i_2}))^{2n} + (\omega_{(M_1, M_2, M_b - 2s)} (S_{i_2} S_{j_2}))^{2n} \right]. \tag{41}$$

Define  $\mathcal{I}_1, \mathcal{I}_2$  as the sets of  $2s$  indices in the first and second sub-systems, respectively, corresponding to the deleted bridge clauses. Then it is easy to realize that, conditionally only on all the non-deleted clauses, the random indices  $i_1, j_1$  and  $i_2, j_2$  are independently and uniformly distributed on  $\mathcal{I}_1$  and  $\mathcal{I}_2$ , respectively, apart from an error term of order  $O(1/s)$ . This error term arises because of the constraints  $i_1 \neq i_2, j_1 \neq j_2$ , which becomes weak for  $s$  large, since the number of non-diagonal configurations is  $O(s^2)$ , while that of the diagonal ones is  $O(s)$ . Introducing, in analogy with section 2,  $\Omega_{(M_1, M_2, M_b - 2s)}$  as the replicated version of the measure  $\omega_{(M_1, M_2, M_b - 2s)}$  and  $\tilde{q}_1^{(2n)}, \tilde{q}_2^{(2n)}$  as

$$\tilde{q}_1^{(2n)} = \frac{1}{2s} \sum_{i \in \mathcal{I}_1} S_i^1 \dots S_i^{2n} \tag{42}$$

$$\tilde{q}_2^{(2n)} = \frac{1}{2s} \sum_{i \in \mathcal{I}_2} S_i^1 \dots S_i^{2n} \tag{43}$$

$\Delta_N^{(s)}$  can be rewritten as

$$\beta N \Delta_N^{(s)} = s \sum_{n=1}^{\infty} \frac{\langle \tanh^{2n}(\beta J) \rangle}{2n} E \Omega_{(M_1, M_2, M_b - 2s)} [(\tilde{q}_1^{(2n)} - \tilde{q}_2^{(2n)})^2] + O(1). \tag{44}$$

The first term is clearly positive. As for the term  $O(1)$ , it becomes negligible in the limit of large  $s$ , since the sum in (21) contains  $O(N/s)$  terms of this kind and has a pre-factor  $1/N$ .

Note that this expression is different from that found in the procedures used in the Sherrington–Kirkpatrick [19] and in the Poissonian [15] cases where, in the analogous expansions, one finds polynomials in the usual multi-overlaps  $q_1^{(2n)}, q_2^{(2n)}$  defined in section 2, depending explicitly on the ratio  $N_1/N$ .

## 6. Consequences of a self-averaging property

Formulae (34)–(36) provide an exact representation of the free energy. We will see in the next section that at least for even  $p$ , the remainder  $R[G, t]$  is non-negative for all distributions  $G$  for which it makes sense. This is not enough to prove that  $F$  is limited from above by the replica free energy; a control of the term  $\tilde{R}[G, t]$  is also required. As we show below, this control is guaranteed by the self-averaging property of suitable extensive quantities, which follows from general thermodynamical convexity arguments, first employed in the context of mean-field spin glasses by Guerra in [21].

As a preliminary fact, note that, as it is clear from (30), in order to establish the lower bound for the free energy it is sufficient to show that  $\tilde{R}[G, t]$  vanishes for  $t \leq M - \epsilon N$ , for arbitrary  $\epsilon > 0$  which will be let tend to zero in the end.

Let us consider the interpolating model at time  $t - 1$ , and rewrite its Hamiltonian (9) as

$$\mathcal{H} = \mathcal{H}_0 - \sum_{\mu=t}^M \sum_{l=1}^p u_{i_l^\mu}^\mu S_{i_l^\mu} \quad (45)$$

where  $\mathcal{H}_0$  contains the clauses indexed  $1, \dots, t - 1$ . Let us now consider the quantities

$$\phi_k = \frac{1}{N} \sum_{\mu=t}^M \sum_{l=1}^p (u_{i_l^\mu}^\mu)^k S_{i_l^\mu} \quad (46)$$

which can be expected to be self-averaging with respect to the Boltzmann and the quenched averages, for any integer  $k$ . Indeed, one has that *generically*

$$L(k, l) = \lim_{N \rightarrow \infty} L_N(k, l) = \lim_{N \rightarrow \infty} E(\omega(\phi_k \phi_l) - \omega(\phi_k)\omega(\phi_l)) = 0 \quad (47)$$

where what we mean by ‘generically’ will be clarified below. For the moment, let us explore the consequences of this. Note that we can write

$$L_N(k, l) = \frac{1}{N^2} \sum_{\mu, \nu=t}^M \sum_{r, s=1}^p E \left[ (u_{i_r^\mu}^\mu)^k (u_{i_s^\nu}^\nu)^l \frac{\partial}{\partial \beta u_{i_r^\mu}^\mu} \frac{\partial}{\partial \beta u_{i_s^\nu}^\nu} \log \omega_{-u_{i_r^\mu}^\mu - u_{i_s^\nu}^\nu} (e^{\beta(u_{i_r^\mu}^\mu S_{i_r^\mu} + u_{i_s^\nu}^\nu S_{i_s^\nu})}) \right]. \quad (48)$$

Thanks to the average  $E$  the sum is immaterial and we can write that

$$\frac{(M-t)^2}{N^2} E \left[ (u_{i_r^\mu}^\mu)^k (u_{i_s^\nu}^\nu)^l \frac{\partial}{\partial \beta u_{i_r^\mu}^\mu} \frac{\partial}{\partial \beta u_{i_s^\nu}^\nu} \log \omega_{-u_{i_r^\mu}^\mu - u_{i_s^\nu}^\nu} (e^{\beta(u_{i_r^\mu}^\mu S_{i_r^\mu} + u_{i_s^\nu}^\nu S_{i_s^\nu})}) \right] \rightarrow 0 \quad (49)$$

where  $\mu \neq \nu$  or  $r \neq s$ . Note that the pre-factor is of order 1. Next we note that, conditionally only on the clauses  $1, \dots, t - 1$  which have not been removed, the random variables  $i_r^\mu, i_s^\nu$  are independent and identically distributed as

$$P(i_1^\mu = i) \propto (k_i - k_i(t)) \quad (50)$$

where  $k_i$  is the degree of the site  $i$  in the original system, and  $k_i(t)$  is its degree at time  $t - 1$ . Of course,  $k_i - k_i(t) \geq 0$  is just the number of deleted clauses which involved the  $i$ th spin. In particular, choosing  $\mu = \nu = t, r = 1, s = 2$ , one can write

$$E \left[ (u_{i_1^t}^t)^k (u_{i_2^t}^t)^l \frac{\partial}{\partial \beta u_{i_1^t}^t} \frac{\partial}{\partial \beta u_{i_2^t}^t} \log \omega_{-u_{i_1^t}^t - u_{i_2^t}^t} (e^{\beta(u_{i_1^t}^t S_{i_1^t} + u_{i_2^t}^t S_{i_2^t})}) \right] \rightarrow 0. \quad (51)$$

Observing that this identity has to be valid for all  $k$  and  $l$ , we find that with probability 1 with respect to the distribution of  $u_{i_1^t}^t$  and  $u_{i_2^t}^t$ ,

$$\frac{\partial}{\partial \beta u_{i_1^t}^t} \frac{\partial}{\partial \beta u_{i_2^t}^t} E_{-u_{i_1^t}^t - u_{i_2^t}^t} \left( \log \omega_{-u_{i_1^t}^t - u_{i_2^t}^t} (e^{\beta(u_{i_1^t}^t S_{i_1^t} + u_{i_2^t}^t S_{i_2^t})}) \right) \rightarrow 0. \quad (52)$$

This implies that

$$E_{-u'_{i_1}-u'_{i_2}} \left( \log \omega_{-u'_{i_1}-u'_{i_2}} \left( e^{\beta(u'_{i_1} S_{i_1} + u'_{i_2} S_{i_2})} \right) \right) \approx E_{-u'_{i_1}-u'_{i_2}} \left( \log \omega_{-u'_{i_1}-u'_{i_2}} \left( e^{\beta u'_{i_1} S_{i_1}} \right) \right) + E_{-u'_{i_1}-u'_{i_2}} \left( \log \omega_{-u'_{i_1}-u'_{i_2}} \left( e^{\beta u'_{i_2} S_{i_2}} \right) \right) \quad (53)$$

and that

$$E_{-u'_{i_1}-u'_{i_2}} \left( \log \omega_{-u'_{i_1}} \left( e^{\beta u'_{i_1} S_{i_1}} \right) \right) = E_{-u'_{i_1}-u'_{i_2}} \left( \log \omega_{-u'_{i_1}-u'_{i_2}} \left( e^{\beta u'_{i_1} S_{i_1} + u'_{i_2} S_{i_2}} \right) \right) - E_{-u'_{i_1}-u'_{i_2}} \left( \log \omega_{-u'_{i_1}-u'_{i_2}} \left( e^{\beta u'_{i_2} S_{i_2}} \right) \right) \approx E_{-u'_{i_1}-u'_{i_2}} \left( \log \omega_{-u'_{i_1}-u'_{i_2}} \left( e^{\beta u'_{i_1} S_{i_1}} \right) \right) \quad (54)$$

where the equivalence sign  $\approx$  means in the previous formulae and in the rest of the section that both quantities on the two sides of an equation tend to the same value in the thermodynamic limit. It is easy to realize that these identities can be generalized to ones involving arbitrary number  $j$  of fields:

$$E_{-u'_{i_1}, \dots, -u'_{i_j}} \left( \log \omega_{-u'_{i_1}, \dots, -u'_{i_j}} \left( e^{\beta \sum_{r=1}^j u'_{i_r} S_{i_r}} \right) \right) \approx \sum_{r=1}^j E_{-u'_{i_1}, \dots, -u'_{i_j}} \left( \log \omega_{-u'_{i_r}} \left( e^{\beta u'_{i_r} S_{i_r}} \right) \right). \quad (55)$$

If we now take  $j = p$  and recall (33), we conclude that self-averaging of the  $\phi_k$  implies that  $\tilde{R}[G, t]$  tends to zero in the thermodynamic limit.

Similarly, to prove that the remainder  $\tilde{T}$  of section 5 vanishes, one has to consider a Hamiltonian of the form

$$\mathcal{H} = \mathcal{H}_0 - \sum_{\mu=1}^M w_{\mu} S_{i_1}^{\mu} S_{i_2}^{\mu}$$

and to exploit self-averaging of the quantities

$$\psi_k = \frac{1}{N} \sum_{\mu} w_{\mu}^k S_{i_1}^{\mu} S_{i_2}^{\mu}.$$

Repeating the steps which led from (47) to (55), one finally finds that  $\tilde{T}$  vanishes for  $N \rightarrow \infty$ .

We are now left with the task of showing that relations such as (47) generically hold. The strategy we use has been already employed in [21], and more recently in [22]. The idea is as follows: while one is not able in general to prove that equations (47) hold for the original model (45), one can perturb it by adding a term  $N \sum_{k=1}^{\infty} \lambda_k \phi_k$  to the Hamiltonian  $\mathcal{H}$ ,  $\lambda_k$  being real numbers decreasing sufficiently fast with  $k$ , so that the infinite volume free energy remains bounded<sup>6</sup>. In the end, we will be interested in the case where all the  $\lambda$  vanish. Then, as in [21], convexity of the free energy with respect to parameters  $\lambda_k$  implies that, for almost every choice for their values, the second derivative  $\partial_{\lambda_k}^2 F_N$  is finite, also in the thermodynamic limit. Since this derivative can be written as

$$\partial_{\lambda_k}^2 F_N = N E(\omega_{\lambda}(\phi_k^2)) - \omega_{\lambda}(\phi_k)^2$$

it follows that equations (47) hold, for almost every choice of the  $\lambda$ , if the Gibbs averages are understood to correspond to the modified model. In order to conclude the argument and show

<sup>6</sup> Results similar to those described below could be obtained considering perturbations of the kind  $N^{1-\alpha} \sum_{k=1}^{\infty} \lambda_k \phi_k$  with  $0 < \alpha < 1/2$ , which even for finite  $\lambda$  modify the free energy or equations (48) only by  $O(N^{-\alpha})$  terms.

that  $\tilde{R} \rightarrow 0$ , we have to remark two important points. First, equations (48) will be modified by a harmless remainder of order  $O(\lambda)$ , which will disappear in the end since we can take the  $\lambda_k$  arbitrarily small. Next, the remainder term  $R$  of section 4 remains positive, so that equation (30) gives

$$F_N \geq F_{\text{var}}[G] + O(1/N) + O(\lambda) \tag{56}$$

where both the replica-symmetric and the true free energy are those corresponding to the modified system. Finally, one employs the fact that the free energy is always a continuous function, as a consequence of its convexity, to deduce that (56) holds also for the original system with all  $\lambda$  set to 0. It is important to emphasize that we have not used anywhere assumptions of continuity of Gibbs averages with respect to the parameters  $\lambda$ , which we cannot prove in general, but just simple positivity properties and the continuity of the free energy, which holds as a general fact.

6.1. Generalization to multi-overlaps of the replica equivalence identities

It is interesting to investigate the consequences of equation (49) on the distribution of multi-overlaps. To that scope, one can perform an expansion in powers of  $\tanh(\beta v_i)$  and  $\tanh(\beta v_j)$  in (49). Omitting tedious but conceptually simple calculations one finds that in the thermodynamic limit, for all  $r$  and  $s$ ,

$$\sum_{l=0}^{\min[2r, 2s]} \frac{(-1)^{(l+1)} (2r + 2s - l - 1)!}{(2r - l)! (2s - l)! l!} \langle (q^{(2r)} \cdot q^{(2s)})_l \rangle = 0 \tag{57}$$

where the  $q^{(2r)}$  and  $q^{(2s)}$  are multi-overlaps involving respectively  $2r$  and  $2s$  replicas, and with the notation  $(q^{(2r)} \cdot q^{(2s)})_l$  we mean that among the two groups of replica  $l$  are in common. Note that relations (57) generalize to the case of multi-overlaps the Ghirlanda–Guerra identity [16]

$$-\frac{3}{2} \langle q_{12} q_{34} \rangle + 2 \langle q_{12} q_{23} \rangle - \frac{1}{2} \langle q_{12}^2 \rangle = 0 \tag{58}$$

and reduce to it for  $r = s = 1$ . More general relations can be obtained from (55) or, in analogy with [16], considering self-averaging properties of multi-spin perturbations in the Hamiltonian. While we will not pursue this route in this paper in full generality, it is clear that  $p$ -spin perturbations to the Hamiltonian will give rise to equations similar to (57) involving multi-overlaps raised to the  $p$ th power.

We note that—as it happens in the Ghirlanda–Guerra case—within the replica method, identities (57) can be derived from the requirement of ‘replica equivalence’. Within replica method one introduces multi-overlap order parameters  $Q_{a_1, \dots, a_r}$ , and self-consistently finds that the multi-overlap averages are given by

$$\langle q^{(r)} \rangle = \lim_{n \rightarrow 0} \frac{1}{n(n-1) \cdots (n-r+1)} \sum_{a_1, \dots, a_r}^{\prime, 1, n} Q_{a_1, \dots, a_r} \tag{59}$$

where all the indices in  $\sum'$  are different. Similarly, the averages  $\langle (q^{(r)} \cdot q^{(s)})_l \rangle$  are given by replica sums of  $Q_{a_1, \dots, a_r} Q_{b_1, \dots, b_s}$  with  $l$   $a$ -indices coinciding with  $l$   $b$ -indices, normalized to the number of terms in the sum. For instance,  $\langle (q^{(2)} \cdot q^{(2)})_1 \rangle = \frac{1}{n(n-1)(n-2)} \sum_{a, b, c}^{\prime} Q_{ab} Q_{bc}$ . Replica equivalence states that for all values of  $a_1$  the sum

$$\sum_{a_2, \dots, a_r}^{1, n} Q_{a_1, \dots, a_r} \tag{60}$$



takes the same value. As a consequence, for  $n \rightarrow 0$

$$\frac{1}{n} \sum_{a_1, \dots, a_r}^{\prime, 1, n} Q_{a_1, \dots, a_r} \sum_{b_1, \dots, b_s}^{\prime, 1, n} Q_{b_1, \dots, b_s} = O(n) \rightarrow 0. \quad (61)$$

Singling out in the sum terms with coinciding indices in the overlaps and making use of (59), after some algebra one finds equation (57). Equation (49) can be seen as the generating function of these identities.

## 7. The remainder $R[G, T]$

While the control of the remainder  $\tilde{R}$  was obtained, in the previous section, without making reference to a specific form for the Hamiltonian (1), in the present section we restrict ourselves for simplicity to the  $p$ -spin case, where  $H_J(S_{i_1}, \dots, S_{i_p}) = JS_{i_1} \cdots S_{i_p}$ . Substituting in equation (30), using the independence among the external fields and the relation (16) one finds for the first remainder term:

$$\begin{aligned} R^{p\text{-spin}}[G, t] = & -\frac{1}{\beta} \left[ E \left( \log \left( 1 + \tanh(\beta J) \omega_{-u^t}^{(t-1)}(S_{i_1} \cdots S_{i_p}) \right) \right)_J \right. \\ & - p E \left\langle \log \left( 1 + \tanh(\beta J) \prod_{l=1}^{p-1} \tanh(\beta g_l) \omega_{-u_{i_1}^l}^{(t-1)}(S_{i_1}^l) \right) \right\rangle_J \\ & \left. + (p-1) E \left\langle \log \left( 1 + \tanh(\beta J) \prod_{l=1}^p \tanh(\beta g_l) \right) \right\rangle_J \right]. \quad (62) \end{aligned}$$

Through this expression we can establish that the remainder is positive for even  $p$ .

Now, we expand the logarithm of the three terms into (absolutely convergent) series of  $\tanh(\beta J)$ , and note that thanks to the parity of the  $J$  and the  $g$  distributions, they will just involve negative terms. We can then take the expected value of each term and write

$$\begin{aligned} R^{p\text{-spin}}[G, t] = & \frac{1}{\beta} \sum_{n=1}^{\infty} \frac{\langle \tanh^{2n} \beta J \rangle_J}{2n} E \left[ \left( \omega_{-u^t}^{(t-1)}(S_{i_1} \cdots S_{i_p}) \right)^{2n} \right. \\ & \left. - p \langle \tanh^{2n} \beta g \rangle_g^{p-1} \left( \omega_{-u_{i_1}^1}^{(t-1)}(S_{i_1}^1) \right)^{2n} + (p-1) \langle \tanh^{2n} \beta g \rangle_g^p \right]. \quad (63) \end{aligned}$$

In analogy with section 2, we introduce  $\Omega^{(t-1)}$  as the replicated version of the measure  $\omega^{(t-1)}$ . Next we note that, as in section 6, conditionally only on the clauses  $1, \dots, t-1$  which have not been removed, the random variable  $i_1^t$  is distributed as

$$P(i_1^t = i) \propto (k_i - k_i(t)). \quad (64)$$

Then, defining

$$\hat{\psi}_i^a = e^{-\beta u_i^1 S_i^a} \frac{Z(t-1)}{Z_{-u_i^1}(t-1)} \quad (65)$$

and

$$r^{(2n)} = \frac{\sum_i (k_i - k_i(t)) S_i^1 \hat{\psi}_i^1 \cdots S_i^{2n} \hat{\psi}_i^{2n}}{\sum_i (k_i - k_i(t))} = \frac{\sum_i (k_i - k_i(t)) S_i^1 \hat{\psi}_i^1 \cdots S_i^{2n} \hat{\psi}_i^{2n}}{p(M-t+1)} \quad (66)$$

it is easy to realize that

$$R^{p\text{-spin}}[G, t] = \frac{1}{\beta} \sum_{n=1}^{\infty} \frac{\langle \tanh^{2n} \beta J \rangle_J}{2n} E \Omega^{(t-1)} \\ \times \left[ (r^{(2n)})^p - p r^{(2n)} \langle \tanh^{2n} \beta g \rangle_g^{p-1} + (p-1) \langle \tanh^{2n} \beta g \rangle_g^p \right] + O(1/N). \quad (67)$$

The observable  $\hat{\psi}_i$  can be seen as the operator which annihilates one of the external fields  $u_i$  from site  $i$ . Indeed, for any observable  $A$  one has

$$\omega(A \hat{\psi}_i) = \omega_{-u_i^1}(A). \quad (68)$$

The harmless error term  $O(1/N)$  arises because, in order to reconstruct  $(r^{(2n)})^p$ , we added ‘diagonal’ terms where at least two indices  $i_l^t$  and  $i_{l'}^t$  are equal. Since  $M-t > \epsilon N$  with  $\epsilon > 0$ , as in section 6, these terms give altogether a vanishing contribution in the infinite volume limit. It is interesting to notice that, as found in I, for models with Poissonian connectivity degree the expansion of  $R[G, t]$  can be expressed in terms of the usual multi-overlap  $q^{(2n)}$  defined in section 2.

In the case of the  $K$ -SAT, using definition (4) for the clause  $H_J$ , we find relation:

$$u_J(g_1, \dots, g_{p-1}) \equiv u_J(\{J_l\}, \{g_l\}) = \frac{J}{\beta} \tanh^{-1} \left[ \frac{\frac{\xi}{2} \prod_{l=1}^{p-1} \left( \frac{1+J_l \tanh(\beta g_l)}{2} \right)}{1 + \frac{\xi}{2} \prod_{l=1}^{p-1} \left( \frac{1+J_l \tanh(\beta g_l)}{2} \right)} \right] \quad (69)$$

where  $\xi \equiv e^{-\beta} - 1 < 0$ . Therefore, one has

$$R^{K\text{-SAT}}[G, t] = -\frac{1}{\beta} E \left[ \left\langle \log \left( 1 + (e^{-\beta} - 1) \omega \left( \prod_{l=1}^p \frac{1 + J_l S_{i_l^t}}{2} \right) \right) \right\rangle_{\{J_l\}} \right. \\ \left. - p \left\langle \log \left( 1 + \xi \omega \left( \frac{1 + J S_{i_1^t}}{2} \prod_{l=1}^{p-1} \frac{1 + J_l \tanh(\beta g_l)}{2} \right) \right) \right\rangle_{\{g_l\}, J, \{J_l\}} \right. \\ \left. + (p-1) \left\langle \log \left( 1 + \xi \prod_{l=1}^p \frac{1 + J_l \tanh(\beta g_l)}{2} \right) \right\rangle_{\{g_l\}, \{J_l\}} \right]. \quad (70)$$

Expanding in series the logarithms, exploiting the symmetry of the probability distribution functions and taking the expectation of each term of the absolutely convergent series we obtain

$$R^{K\text{-SAT}}[G, t] = \frac{1}{\beta} \sum_{n \geq 1} \frac{(-\xi^*)^n}{n} \Omega \left[ (1 + R_n)^p - p(1 + R_n) \langle (1 + J \tanh(\beta g))^n \rangle_{J, g}^{p-1} \right. \\ \left. + (p-1) \langle (1 + J \tanh(\beta g))^n \rangle_{J, g}^p \right] \quad (71)$$

where we have defined  $\xi^* \equiv \xi/(2^p) < 0$  and  $R_n \equiv \sum_{l=1}^n \langle J^l \rangle_J \sum_{a_1 < \dots < a_l}^{1, n} r^{a_1 \dots a_l}$ .

We note that both in the  $p$ -spin and in the  $K$ -SAT problem the remainder can be written as a series of the type

$$R[G, t] = \sum_{n=1}^{\infty} C_n E \Omega[f_p(X_n, Y_n)] \quad (72)$$

where

$$f_p(x, y) = x^p - pxy^{p-1} + (p-1)y^p \quad (73)$$

$C_n \geq 0$  for any temperature,  $X_n$  are suitable combinations of overlaps and  $Y_n$  averages of  $g$ -fields hyperbolic tangents moments that correctly calculate the overlaps in the corresponding replica symmetric approximation. More specifically, for the  $p$ -spin

$$C_n = \frac{\langle \tanh^{2n}(\beta J) \rangle}{2\beta n} \quad X_n = r^{(2n)} \quad Y_n = \langle \tanh^{2n}(\beta g) \rangle \quad (74)$$

while for the  $K$ -SAT

$$C_n = \frac{(-\xi^*)^n}{\beta n} \quad X_n = 1 + R_n \quad Y_n = \langle (1 + J \tanh(\beta g))^n \rangle. \quad (75)$$

As noted in I, for even  $p$  the function  $f_p(x, y)$  is positive for all real  $x$  and  $y$  thus ensuring the positivity of the remainder. For odd  $p$  this is not the case as  $f_p(x, y)$  at fixed  $y$  becomes negative for  $x$  negative and large enough. Although physically one expects that for odd  $p$  negative values of  $X_n$  appear with probability exponentially small in  $N$ , we have not been able to prove this property in full generality.

The odd  $p$  case is however interesting. In particular, one would like to be able to control the remainder in the case of the 3-SAT problem, recently solved within 1RSB scheme [8]. When explicit solutions of the replica equations exist, one can try to plug them into the expression of the remainder, to check positivity. Results in this direction, in the RS case at zero temperature for  $K$ -SAT and  $p$ -spin models, will be reported in [23].

## 8. Summary and conclusions

In this paper we have shown that the free energy of diluted spin glass models with arbitrary random connectivity can be written as the sum of a term identical to the ones got in the cavity/replica plus an error term. The expression has been obtained through the introduction of an auxiliary model interpolating between the original model and a pure paramagnet. The interpolation can be thought of as a discrete time dynamical process in which the terms of the Hamiltonian are progressively removed, while the removal effect is compensated by the introduction of some external fields. The procedure generalizes the previous work [15] on Poissonian graphs, where the compensation could be performed in average: at each step one adds there a random number of fields on random sites. In the present case, on the other hand, a detailed compensation where one puts a field on each site involved in the erased clause is necessary. As a consequence, a new term in the remainder appears. We have shown that, thanks to self-averaging of suitable extensive quantities, this new term gives a vanishing contribution in the thermodynamic limit. The rest of the remainder is manifestly positive for even  $p$ .

It is also possible to show, as was done in [24] for models with Poissonian random connectivity, that the free energy and the ground-state energy are self-averaging quantities, and one can obtain upper bounds, exponentially small in the system size, for the probability of large fluctuations.

## Acknowledgment

We are very grateful to Andrea Montanari for pointing out a weak point in the first version of this work.

## References

- [1] Mézard M, Parisi G and Virasoro M A 1987 *Spin Glass Theory and Beyond* (Singapore: World Scientific)
- [2] Mézard M and Parisi G 2001 *Eur. Phys. J. B* **20** 217
- [3] Mézard M and Parisi G 2003 The cavity method at zero temperature *J. Stat. Phys.* **111** 1
- [4] Ricci-Tersenghi F, Mézard M and Zecchina R 2003 *J. Stat. Phys.* **111** 505–33
- [5] Biroli G and Mézard M 2002 *Phys. Rev. Lett.* **88** 025501
- [6] Franz S, Leone M, Montanari A and Ricci-Tersenghi F 2002 *Phys. Rev. E* **66** 046120
- [7] Mézard M, Parisi G and Zecchina R 2002 *Science* **297** 812–5
- [8] Mézard M and Zecchina R 2002 *Phys. Rev. E* **66** 056126
- [9] Mulet R, Pagnani A, Weigt M and Zecchina R 2002 *Phys. Rev. Lett.* **89** 268701
- [10] For a review on the work of mathematicians on spin glass systems see: Bovier A and Picco P (ed) 1997 *Mathematical aspects of spin glasses and neural networks* *Progress in Probability* vol 41 (Boston: Birkhauser)  
Talagrand M 1998 *Proc. Berlin Int. Congress of Mathematicians. Doc. Math.* extra volume p 1, and references therein (We refer to the webpage of M Talagrand for the comprehensive list of references on the work of this author: <http://www.math.ohio-state.edu/~talagrand/>)
- [11] Talagrand M 2003 *Rev. Math. Phys.* **15** 1–78
- [12] Talagrand M 2000 Mean field models for spin glasses, a first course *Proc. 2000 Saint Flour Summer School in Probability*
- [13] Guerra F 2003 *Commun. Math. Phys.* **233** 1–12
- [14] Contucci P and Graffi S 2003 Monotonicity and thermodynamic limit for short range disordered models *Preprint math-ph/0302013 (J. Stat. Phys. at press)*
- [15] Franz S and Leone M 2003 *J. Stat. Phys.* **111** 535–64
- [16] Ghirlanda S and Guerra F 1998 *J. Phys. A: Math. Gen.* **31** 9149–55
- [17] Ricci-Tersenghi F, Weigt M and Zecchina R 2001 *Phys. Rev. E* **63** 026702
- [18] Monasson R and Zecchina R 1997 *Phys. Rev. E* **56** 1357
- [19] Guerra F and Toninelli F L 2002 *Commun. Math. Phys.* **230** 71–9
- [20] Franz S, Leone M and Zecchina R in preparation
- [21] Guerra F 1996 *Int. J. Mod. Phys. B* **10** 1675–84
- [22] Talagrand M 2003 *Spin Glasses, A Challenge for Mathematicians* (Berlin: Springer)
- [23] Franz S and Leone M in preparation
- [24] Guerra F and Toninelli F L 2003 The high temperature region of the Viana–Bray diluted spin glass model *Preprint cond-mat/0302401 (J. Stat. Phys. at press)*