

Unicyclic components in random graphs

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2004 J. Phys. A: Math. Gen. 37 L189

(<http://iopscience.iop.org/0305-4470/37/18/L01>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.90

The article was downloaded on 02/06/2010 at 17:57

Please note that [terms and conditions apply](#).

LETTER TO THE EDITOR

Unicyclic components in random graphs

E Ben-Naim¹ and P L Krapivsky²¹ Theoretical Division and Center for Nonlinear Studies, Los Alamos National Laboratory, Los Alamos, NM 87545, USA² Center for Polymer Studies and Department of Physics, Boston University, Boston, MA 02215, USA

Received 18 March 2004

Published 20 April 2004

Online at stacks.iop.org/JPhysA/37/L189 (DOI: 10.1088/0305-4470/37/18/L01)**Abstract**

The distribution of unicyclic components in a random graph is obtained analytically. The number of unicyclic components of a given size approaches a self-similar form in the vicinity of the gelation transition. At the gelation point, this distribution decays algebraically, $U_k \simeq (4k)^{-1}$ for $k \gg 1$. As a result, the total number of unicyclic components grows logarithmically with the system size.

PACS numbers: 02.10.Ox, 64.60.-i, 02.50.-r

Random graphs underly processes such as polymerization [1], percolation [2] and the formation of social networks [3, 4]. Random graphs have been extensively studied, especially in theoretical computer science [5, 6]. Special families of random graphs have also been examined, e.g., planar random graphs appear in combinatorics [7, 8] and in physics [9]. The basic framework for generic random graphs naturally emerged in two different contexts [10]. Flory [1, 11] and Stockmayer [12] modelled a polymerization process in which monomers polymerize via binary chemical reactions until a giant polymer network, namely a gel, emerges. Erdős and Rényi studied an equivalent process in which connected components emerge from ensembles of nodes that are linked sequentially and randomly in pairs [13].

Different methodologies have been employed to characterize random graphs. Kinetic theory, specifically, the rate equation approach, was used to obtain the size distribution of components [14]. Using probability theory, a number of additional characteristics including in particular the complexity of random graphs have been addressed [5, 6].

In this study, we show that the rate equation approach is useful for studying the complexity of random graphs. Our main result asserts that U_k , the average number of unicyclic components of size k in a random evolving graph, is given by

$$U_k(t) = \frac{1}{2} t^k e^{-kt} \sum_{n=0}^{k-1} \frac{k^{n-1}}{n!}. \quad (1)$$

The unicyclic components size distribution becomes self-similar as the gelation transition is approached. At the gelation point ($t_g = 1$), the size distribution develops an algebraic




Figure 1. A random graph with $N = 10$ nodes and eight links. The two components on the left are trees, the two components on the right are unicyclic.

large-size tail: $U_k(1) \simeq (4k)^{-1}$. This implies that at the gelation point, the total number of unicyclic components depends logarithmically on the system size.

The random graph evolves from N disconnected nodes as follows. At each step, two nodes are selected at random and a link is drawn between them. This linking process is repeated *ad infinitum*, leading to an ensemble of components, defined as maximally connected sets of nodes. We consider multi-graphs where the two selected nodes need not be different so that self-connections are allowed.

Each component has a certain number of nodes and links. The number of nodes is the component size. The number of links minus the number of nodes is the component complexity [15]. Up to a sign, the complexity is the Euler characteristic of the component—a topological invariant. Trees have complexity -1 , unicyclic components have complexity 0 , bi-cyclic components have complexity 1 , etc (figure 1).

This linking process is treated dynamically. Initially, there are N components of size 1 and complexity -1 . Links are drawn between any two nodes with a uniform rate, set equal to $1/(2N)$ without loss of generality. It is useful to consider linking between different components and linking within the same component separately. Two different components of size and complexity (i, l) and (j, m) , respectively merge with the rate $ij/(2N)$, symbolically represented by the reaction scheme



When linking involves two nodes in the same (k, n) component, the process is



and it occurs with the rate $k^2/(2N)$. The average number of links at time t is $Nt/2$; the average number of self-links is smaller by a factor N , i.e. it is equal to $t/2$.

Let the number of components of size k and complexity n be $N_{k,n}$. Following the dynamical rules (2), (3), the quantities $N_{k,n}$ change with time according to the bi-variate Smoluchowsky equation [16, 17]

$$\frac{dN_{k,n}}{dt} = \frac{1}{2N} k^2 (N_{k,n-1} - N_{k,n}) + \frac{1}{2N} \sum_{i+j=k} \sum_{l+m=n-1} ij N_{i,l} N_{j,m} - k N_{k,n}. \quad (4)$$

The initial condition is $N_{k,n} = N \delta_{k,0} \delta_{n,-1}$. This rate equation implies that the number of trees, unicyclic components and bi-cyclic components are proportional to N , N^0 and N^{-1} , respectively. The random graph consists primarily of trees and unicyclic components, with more complex components being rare. Our goal is to derive the distribution of unicyclic components. Given the recursive nature of the rate equations, it requires the size distribution of trees, that to leading order in N equals the total size distribution.

We first recapitulate the computation of the total size distribution. Let $c_k = N_k/N$ be the size distribution of all components (formally, $N_k = \sum_l N_{k,l}$). Since linking within a component affects its complexity but does not affect its size, the corresponding (first two) terms in the rate equation (4) are irrelevant and the total size distribution satisfies the *nonlinear* equation

$$\frac{dc_k}{dt} = \frac{1}{2} \sum_{i+j=k} ij c_i c_j - kc_k. \quad (5)$$

The initial condition is $c_k(0) = \delta_{k,1}$. The size distribution is obtained via the generating function

$$F(z, t) = \sum_{k \geq 1} kc_k(t) e^{kz}. \quad (6)$$

This generating function evolves according to

$$\frac{\partial F}{\partial t} = (F - 1) \frac{\partial F}{\partial z} \quad (7)$$

subject to the initial condition $F(z, 0) = e^z$. Let us re-write the derivatives through Jacobians: $\frac{\partial F}{\partial t} = \frac{\partial(F,z)}{\partial(t,z)}$ and $\frac{\partial F}{\partial z} = \frac{\partial(F,t)}{\partial(z,t)}$. Using the relation $\frac{\partial z}{\partial t} = \frac{\partial(z,F)}{\partial(t,F)}$, we recast the nonlinear equation (7) for $F(z, t)$ into a linear equation $\frac{\partial z}{\partial t} = 1 - F$ for $z(F, t)$. Integration over time yields $z(F, t) = (1 - F)t + \ln F$ (the integration constant $\ln F$ follows from the initial condition $F(z, 0) = e^z$). Exponentiating this equality yields an implicit relation satisfied by the generating function

$$F e^{-tF} = e^{-t} \zeta \quad \zeta = e^z. \quad (8)$$

We can expand ζ in terms of F , yet we are seeking the opposite: $F = \sum kc_k(t) \zeta^k$. The size distribution can be obtained either using the Lagrange inversion formula [18] or alternatively, by performing the direct calculation

$$\begin{aligned} kc_k &= \frac{1}{2\pi i} \oint d\zeta \frac{F}{\zeta^{k+1}} \\ &= \frac{1}{2\pi i} \oint dF \frac{\zeta'(F)F}{[\zeta(F)]^{k+1}} \\ &= \frac{1}{2\pi i} \oint dF \frac{(1-tF)e^{ktF}}{F^k e^{kt}} \\ &= e^{-kt} \left[\frac{(kt)^{k-1}}{(k-1)!} - t \frac{(kt)^{k-2}}{(k-2)!} \right]. \end{aligned}$$

The size distribution is therefore [14, 19]

$$c_k(t) = \frac{(kt)^{k-1}}{k \cdot k!} e^{-kt}. \quad (9)$$

At time $t_g = 1$, the system undergoes a gelation transition: a giant component that eventually engulfs the entire mass in the system emerges. Close to the gelation time, the size distribution attains the scaling behaviour $c_k(t) \simeq k_*^{-5/2} \Phi(k/k_*)$ with the typical size scale $k_* = (1-t)^{-2}$. This size scale diverges as the gelation point is approached. The underlying scaling function $\Phi(z) = (2\pi)^{-1/2} z^{-5/2} e^{-z/2}$ exhibits an exponential large-size decay in the pre-gel regime while at the gelation time it develops an algebraic tail: $c_k \sim k^{-5/2}$. This power-law behaviour allows us to estimate the size of the giant component, $N_g \sim N^{2/3}$ [5], using the extremal statistics criterion $N \sum_{k \geq N_g} c_k \sim 1$ [20]. The time when the giant component emerges in a finite system is estimated from $N_g \sim (1-t_g)^{-2}$, i.e., $1-t_g \sim N^{-1/3}$.




Figure 2. The average number of unicyclic components at the gelation point versus the system size N .

The average size distribution of unicyclic components $U_k \equiv \langle N_{k,0} \rangle$ is coupled to the total size distribution c_k . From (4) we find that U_k satisfies the *linear* inhomogeneous equation

$$\frac{dU_k}{dt} = \frac{1}{2}k^2c_k + \sum_{i+j=k} iU_i j c_j - kU_k. \quad (10)$$

The initial condition is $U_k(0) = 0$. The first term on the right-hand side of (10) plays the role of a source—it represents the formation of unicyclic components from trees via the linking of two nodes within the same component. Such linking occurs with the rate $k^2/2N$ (recall that multigraphs where two nodes can be connected by more than one link are considered). The next two terms account for changes in the size of a unicyclic component due to mergers with different components.

Consider the average number of unicyclic components $U = \sum_k U_k$ prior to the gelation time. Summing equations (10) we find that U satisfies

$$\frac{dU}{dt} = \frac{1}{2}M_2 \quad (11)$$

where $M_2(t) = \sum_k k^2 c_k(t)$ is the second moment of the size distribution. Using $M_2(t) = F'(z, t)|_{z=0}$ and equation (8) we obtain $M_2 = (1-t)^{-1}$. The total number of unicyclic components is therefore

$$U(t) = \frac{1}{2} \ln \frac{1}{1-t}. \quad (12)$$

The number of unicyclic components diverges as the gelation point is approached. The total number of unicyclic components at the gelation point is obtained from the estimate $1-t_g \sim N^{-1/3}$; it diverges logarithmically with the system size N [21, 22]

$$U(t_g) \simeq \frac{1}{6} \ln N. \quad (13)$$

This number is much larger compared with the average number of self-connections that equals $1/2$ at the gelation point. The approach to the asymptotic behaviour (13) is shown using Monte Carlo simulations (figure 2). The data represent an average over 10^6 independent realizations.

To determine the entire distribution U_k , it is useful to manually solve for small k . The expressions

$$U_1 = \frac{1}{2}t e^{-t} \quad U_2 = \frac{3}{4}t^2 e^{-2t} \quad U_3 = \frac{17}{12}t^3 e^{-3t}$$

suggest the following structure $kU_k(t) = A_k t^k e^{-kt}$. Substituting this form and the size distribution (9) into (10) we obtain a recurrence for the coefficients

$$A_k = \frac{1}{2} \frac{k^k}{k!} + \sum_{i+j=k} A_i \frac{j^{j-1}}{j!}. \tag{14}$$

To solve this recursion, we introduce the generating functions $A(z) = \sum_{k \geq 1} A_k e^{kz}$ and $H(z) = \sum_{k \geq 1} \frac{k^{k-1}}{k!} e^{kz}$, thereby recasting (14) into

$$A = \frac{1}{2} (1 - H)^{-1} \frac{dH}{dz}. \tag{15}$$

The generating function $H(z) = F(z + 1, 1)$ is expressed via the known generating function $F(z, t)$ and therefore,

$$H e^{-H} = \zeta \quad \zeta = e^z. \tag{16}$$

Re-writing this equality as $\ln H - H = z$ and differentiating with respect to z we obtain $\frac{dH}{dz} = H/(1 - H)$, which is then inserted into (15) to give $A = \frac{1}{2} H(1 - H)^{-2}$. Combining this equation together with equation (16), the coefficients are evaluated similar to the derivation of the size distribution c_k :

$$\begin{aligned} A_k &= \frac{1}{2\pi i} \oint d\zeta \frac{A}{\zeta^{k+1}} \\ &= \frac{1}{2\pi i} \oint dH \frac{\zeta'(H)A}{[\zeta(H)]^{k+1}} \\ &= \frac{1}{2\pi i} \oint dH \frac{e^{kH}}{H^k 2(1 - H)} \\ &= \frac{1}{2} \sum_{n=0}^{k-1} \frac{k^n}{n!}. \end{aligned}$$

Thus, the size distribution of the unicyclic components (1) is obtained analytically.

Given the scaling behaviour of the size distribution, we investigate the unicyclic size distribution in the vicinity of the gelation point. To obtain the large- k behaviour, the size distribution is re-written as

$$U_k(t) = \frac{1}{2} t^k e^{-kt} \frac{k^{k-2}}{(k-1)!} \sum_{j=0}^{k-1} \prod_{i=0}^j \left(1 - \frac{i}{k}\right). \tag{17}$$

When $k \gg 1$, the product approaches an exponential, $e^{-j(j+1)/2k}$, and the summation can be replaced by integration

$$\sum_{j=0}^{k-1} \prod_{i=0}^j \left(1 - \frac{i}{k}\right) \simeq \int_0^k dj \exp\left[-\frac{j(j+1)}{2k}\right] \simeq \int_0^\infty dj \exp\left[-\frac{j^2}{2k}\right] = \sqrt{\pi k/2}.$$

The leading large- k asymptotics of the unicyclic components size distribution is therefore

$$U_k(t) \simeq \sqrt{\frac{\pi}{8k}} \frac{(kt)^k}{k!} e^{-kt}. \tag{18}$$

In the vicinity of the gelation point, this distribution approaches the scaling form $U_k(t) \simeq k_*^{-1} \Psi(k/k_*)$ with the same scaling variable as that underlying the total size distribution, $k_* = (1 - t)^{-2}$. The scaling function is also similar in form

$$\Psi(z) = (4z)^{-1} e^{-z/2}. \tag{19}$$

The unicyclic size distribution has exponential tails both above and below the gelation transition and it becomes algebraic at the gelation point

$$U_k(t_g) \simeq (4k)^{-1} \quad \text{when} \quad k \gg 1. \quad (20)$$

This result is of course consistent with the total number of unicyclic components: $U(t_g) \simeq \sum_{k \leq N^{2/3}} U_k(t_g)$ leads to equation (13).

We now probe the complexity of the giant component. In the post-gelation region the gel mass (mass not contained in finite components) $g(t)$ is given by $g = 1 - \sum_k kc_k(t) = 1 - F(z, t)|_{z=0}$. Using equation (8), the gel mass obeys $g = 1 - e^{-gt}$ for $t > 1$. The complexity of the giant component increases due to linking processes involving its internal nodes. The total number of nodes in the giant component is Ng . The linking rate is therefore $(2N)^{-1} \times (Ng)^2$, so the complexity of the giant component is

$$C(t) = \frac{N}{2} \int_1^t dt' g^2(t'). \quad (21)$$

Just above the gelation transition, the gel mass increases linearly, $g \simeq 2(t - 1)$, and therefore, $C(t) \simeq \frac{2}{3}N(t - 1)^3$ as $t \downarrow 1$. In the large time limit there are a few monomers apart from the giant component. The total number of links is $Nt/2$ and the total number of nodes is $N - N_1 = N(1 - c_1)$, so the complexity of the giant component grows according to $N(t/2 - 1 + e^{-t})$ when $t \rightarrow \infty$. This result also follows from the general formula (21) and the asymptotic behaviour of the gel mass $g(t)$.

In this study, we considered size and complexity characteristics obtained via an average over infinitely many realizations of the linking process. For some quantities, averaging is irrelevant in the thermodynamic limit $N \rightarrow \infty$. For instance, the tree distribution is an extensive random quantity ($\langle N_{k,-1} \rangle = Nc_k$) with fluctuations of the order of $N^{1/2}$ [23, 24], so relative fluctuations decrease with the systems size according to $N^{-1/2}$. Thus, for trees the average distribution Nc_k well represents the outcome of a single realization of the random evolving graph. For unicyclic components, fluctuations are of the same order as the average and an analytical computation of the correlation function $\langle N_{i,0}(t)N_{j,0}(t) \rangle$ is a challenging open problem. This correlation function is required for the determination of the average distribution of bi-cyclic components, so the naive form of the rate equation (4) is inadequate for describing bi-cyclic (and more complex) components.

In conclusion, we have obtained the size distribution of unicyclic components in a random graph. Overall, the unicyclic size distribution has similar properties as the total size distribution; it becomes self-similar near the gelation point, and generally, it has an exponential tail. Precisely at the gelation point the distribution has an algebraic tail. The main difference with the size distribution is in the value of the power-law exponent itself.

Acknowledgment

This research was supported in part by DOE (W-7405-ENG-36).

References

- [1] Flory P J 1953 *Principles of Polymer Chemistry* (Ithaca, NJ: Cornell University Press)
- [2] Stauffer D 1985 *Introduction to Percolation Theory* (London: Taylor and Francis)
- [3] Skyrms B and Pemantle R 2000 *Proc. Natl Acad. Sci.* **97** 9340
- [4] Girvan M and Newman M E J 2002 *Proc. Natl Acad. Sci.* **99** 7821
- [5] Bollobás B 1985 *Random Graphs* (London: Academic)
- [6] Janson S, Łuczak T and Rucinski A 2000 *Random Graphs* (New York: Wiley)

-
- [7] Tutte W T 1962 *Can. J. Math.* **14** 21
Tutte W T 1962 *Can. J. Math.* **14** 402
Tutte W T 1962 *Can. J. Math.* **14** 708
Tutte W T 1963 *Can. J. Math.* **15** 249
 - [8] Chassaing Ph and Schaeffer G 2004 *Probab. Theory Rel.* **128** 161
 - [9] Ambjørn J, Durhuus B and Jonsson T 1997 *Quantum Geometry: A Statistical Field Theory Approach* (Cambridge: Cambridge University Press)
 - [10] Aldous D J 1999 *Bernoulli* **5** 3
 - [11] Flory P J 1941 *J. Am. Chem. Soc.* **63** 3083
 - [12] Stockmayer W H 1943 *J. Chem. Phys.* **11** 45
 - [13] Erdős P and Rényi A 1960 *Publ. Math. Inst. Hung. Acad. Sci.* **5** 17
 - [14] McLeod J B 1962 *Q. J. Math. Oxford* **13** 119
McLeod J B 1962 *Q. J. Math. Oxford* **13** 193
McLeod J B 1962 *Q. J. Math. Oxford* **13** 283
 - [15] Janson S 2000 *Rand. Struct. Algorithms* **17** 343
 - [16] Smoluchowski M V 1916 *Physik. Z.* **17** 585
Smoluchowski M V 1917 *Z. Phys. Chem.* **92** 129
 - [17] Krapivsky P L and Ben-Naim E 1996 *Phys. Rev. E* **53** 291
 - [18] Wilf H S 1990 *Generatingfunctionology* (Boston: Academic)
 - [19] Hendriks E M, Ernst M H and Ziff R M 1983 *J. Stat. Phys.* **31** 519
 - [20] Ben-Naim E and Krapivsky P L 2004 *Europhys. Lett.* **65** 151
 - [21] Janson S, Knuth D E, Łuczak T and Pittel B 1993 *Rand. Struct. Algorithms* **3** 233
 - [22] Janson S 2003 *Comb. Probab. Comput.* **12** 27
 - [23] Lushnikov A A 1977 *J. Colloid. Inter. Sci.* **65** 276
 - [24] van Dongen P G J and Ernst M H 1987 *J. Stat. Phys.* **49** 879