

An approximation to δ' couplings on graphs

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2004 J. Phys. A: Math. Gen. 37 L329

(<http://iopscience.iop.org/0305-4470/37/29/L01>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.91

The article was downloaded on 02/06/2010 at 18:23

Please note that [terms and conditions apply](#).

LETTER TO THE EDITOR

An approximation to δ' couplings on graphs**T Cheon¹ and P Exner^{2,3}**¹ Laboratory of Physics, Kochi University of Technology, Tosa Yamada, Kochi 782-8502, Japan² Department of Theoretical Physics, Nuclear Physics Institute, Academy of Sciences, 25068, Řež, Czech Republic³ Doppler Institute, Czech Technical University, Břehová 7, 11519 Prague, Czech Republic

E-mail: taksu.cheon@kochi-tex.ac.jp and exner@ujf.cas.cz

Received 26 April 2004

Published 7 July 2004

Online at stacks.iop.org/JPhysA/37/L329

doi:10.1088/0305-4470/37/29/L01

Abstract

We discuss a general parametrization for vertices of quantum graphs and show, in particular, how the δ'_s and δ' couplings at an n -edge vertex can be approximated by means of $n + 1$ couplings of the δ type provided the latter are properly scaled.

PACS number: 02.30.Tb

Quantum graphs became in the last decade a useful and versatile tool to describe several classes of physical systems, in particular, various combinations of quantum wires. There are numerous papers devoted to the subject and we restrict ourselves to mentioning the bibliography given in [KS99, Ku04], where basic concepts of theory are also discussed.

The purpose of this letter is twofold. First, we want to draw attention to useful parametrization of a general coupling at graph vertices whose advantages in the present context have remained so far unnoticed. Second and more important, we address the question of physical meaning of such a coupling and suggest an answer illustrating it on a pair of simple nontrivial examples of the so-called δ'_s and δ' couplings [Ex95, Ex96a].

We consider a free spinless particle on a graph, with the Hamiltonian which acts as $H\psi_j = -\psi''_j$, where ψ_j denotes the wavefunction at the j th edge. Since early times, it has been known that a vertex joining n -graph edges can be characterized by n^2 real parameters [EŠ89] characterizing the boundary condition at the vertex. We use the symbol $\Psi(0)$ for the column vector of the boundary values at the vertex (identified conventionally with the origin of the coordinates), and analogously $\Psi'(0)$ for the vector of the derivatives, all taken in the outgoing direction.

The boundary conditions have to be chosen to make the Hamiltonian self-adjoint, or in physical terms, to ensure conservation of the probability current at the vertex. A general form

of such a coupling was found in [KS99]. It is described by a pair of $n \times n$ matrices A, B such that $\text{rank}(A, B) = n$ and AB^* is self-adjoint; the boundary values have to satisfy the conditions

$$A\Psi(0) + B\Psi'(0) = 0. \quad (1)$$

They have an advantage in comparison to earlier parametrizations relating $\Psi(0)$ and $\Psi'(0)$ by a single matrix, because the latter is typically singular for a subset of parameters, albeit a zero-measure one.

On the other hand, the matrix pair in (1) is non-unique; one would prefer to have a condition analogous to $\psi(0) \cos \theta + \psi'(0) \sin \theta = 0$, which is the case of a single-edge end. Such conditions exist; they were obtained independently in [FT00, CFT01] for a generalized point interaction, $n = 2$, and in [Ha00] for any $n \geq 1$. It is easy to derive them: the self-adjointness requires vanishing of the boundary form, $\sum_{j=1}^n (\bar{\psi}_j \psi'_j - \bar{\psi}'_j \psi_j)(0) = 0$, which occurs iff the norms $\|\Psi(0) \pm i\ell\Psi'(0)\|_{\mathbb{C}^n}$ with a fixed nonzero ℓ coincide, so the two vectors must be related by an $n \times n$ unitary matrix. The length parameter is not important because matrices corresponding to two different values are related by

$$U' = \frac{(\ell + \ell')U + \ell - \ell'}{(\ell - \ell')U + \ell + \ell'}. \quad (2)$$

Thus, we set $\ell = 1$, which means a choice of the length scale, and put

$$A = U - I \quad B = i(U + I). \quad (3)$$

The edges are obviously fully decoupled at the vertex iff U is diagonal. It is easy to check that any such pair satisfies the above quoted requirements from [KS99]. Conversely, for any A, B with these properties there is a $U \in U(n)$ and an invertible C such that $U = C(A - iB)$. Indeed, such a U must satisfy $UU^* = C(BB^* + AA^*)C^*$ since $AB^* = BA^*$ by assumption. The matrix $BB^* + AA^*$ is strictly positive because its null space is

$$\ker A^* \cap \ker B^* = (\text{ran } A)^\perp \cap (\text{ran } B)^\perp = (\text{ran } A \cup \text{ran } B)^\perp = \{0\}. \quad (4)$$

In particular, $AA^* + BB^*$ is Hermitian so $C = (AA^* + BB^*)^{-1/2}$ makes sense, being also Hermitian and invertible.

The parametrization (3) simplifies various previous results. For instance, the eigenspace of U with eigenvalue -1 gives the projection $P = P_1$ in [Ku04] which makes lemma 4 and the following claims of this letter rather transparent. Likewise, the on-shell scattering matrix for a star graph of n halflines with the considered coupling equals

$$S_U(k) = \frac{(k-1)I + (k+1)U}{(k+1)I + (k-1)U} \quad (5)$$

which makes a discussion of its properties simpler than in section 2 of [KS99].

To give an example of the parametrization (3), denote by \mathcal{J} the $n \times n$ matrix whose entries are all equal to 1. It is a straightforward exercise to check that $U = \frac{2}{n+i\alpha}\mathcal{J} - I$ describes the standard δ coupling,

$$\psi_j(0) = \psi_k(0) =: \psi(0) \quad j, k = 1, \dots, n \quad \sum_{j=1}^n \psi'_j(0) = \alpha\psi(0) \quad (6)$$

with $\alpha \in \mathbb{R}$; the case $\alpha = 0$ corresponds to the ‘free motion’ at the vertex, the so-called Kirchhoff boundary conditions, while $\alpha = \infty$ gives $U = -I$, the full Dirichlet decoupling. In a similar way, $U = I - \frac{2}{n-i\beta}\mathcal{J}$ describes the singular counterpart, the so-called δ'_s coupling [Ex95, Ex96a],

$$\psi'_j(0) = \psi'_k(0) =: \psi'(0) \quad j, k = 1, \dots, n \quad \sum_{j=1}^n \psi_j(0) = \beta\psi'(0) \quad (7)$$

with $\beta \in \mathbb{R}$; for $\beta = \infty$ we get $U = I$, the full Neumann decoupling.

Let us mention another ‘dual’ pair of vertex couplings in which the wavefunctions exhibit permutation symmetry. The more regular one of these is the ‘permuted’ δ , or δ_p coupling, given by the boundary conditions

$$\sum_{j=1}^n \psi_j(0) = 0 \quad \psi'_j(0) - \psi'_k(0) = \frac{\alpha}{n}(\psi_j(0) - \psi_k(0)) \quad j, k = 1, \dots, n \quad (8)$$

with $\alpha \in \mathbb{R}$. It generalizes the δ_s interaction of [TFC01] and one can check easily that the corresponding matrix equals $U = \frac{n-i\alpha}{n+i\alpha}I - \frac{2}{n+i\alpha}\mathcal{J}$. Its counterpart is the so-called δ' coupling [Ex95, Ex96a],

$$\sum_{j=1}^n \psi'_j(0) = 0 \quad \psi_j(0) - \psi_k(0) = \frac{\beta}{n}(\psi'_j(0) - \psi'_k(0)) \quad j, k = 1, \dots, n \quad (9)$$

with $\beta \in \mathbb{R}$, which corresponds to $U = -\frac{n+i\beta}{n-i\beta}I + \frac{2}{n-i\beta}\mathcal{J}$. The infinite values of the parameters refer again to the Dirichlet and Neumann decouplings of the graph edges, respectively.

Note that in these four examples, the connection condition at the origin is totally symmetric with respect to the interchange of edge indices. Consequently, their U are constructed from symmetric matrices I and \mathcal{J} .

If one wants to continue analysis of such graphs, the first question to be answered is about the physical meaning and possible use of the whole family of such general couplings. Concerning the second part, a recent inspiration comes from the domain of quantum computing, where the generalized point interactions parametrized by elements of the group $U(2)$ have been proposed as an alternative realization of a qubit [CFT04]; an extension to higher degree vertices opens, of course, interesting possibilities. To make use of them, however, one has to understand whether there is a meaningful way to ‘construct’ vertices with different couplings.

The currently available results suggest that this goal cannot be achieved in a purely geometrical way by squeezing a system of branching tubes with the same topology as the graph. Several such approximations were analysed recently [KZ01, RS01, Sa01, EP03]; they all lead either to trivial (Kirchhoff) boundary conditions, or to graphs having an extended state Hilbert space with extra dimensions due to the vertices. Their common feature was that the transverse ground state was a constant function. Hence, a nontrivial result might be obtained through tubes with Dirichlet boundaries, but this problem is open for a long time and notoriously difficult.

Approximations using potentials scaled in the usual way, i.e. preserving their integrals, do yield nontrivial results [Ex96b] but only for couplings with wavefunctions continuous at the junction, which is just the family (6). This is not sufficient and more singular couplings need other means. Our main aim here is to explore a natural alternative with approximating interactions scaled in a *nonlinear* way as a generalization of the procedure proposed in [CS98a, CS98b] and analysed from the viewpoint of the convergence topology in [AN00, ENZ01]. To keep things simple, we will analyse here the δ'_s and δ' couplings leaving the general case to a subsequent paper.

Consider first the Hamiltonian H_β on the graph Γ consisting of n halflines coupled at a single vertex by conditions (7). Consider further the same graph with additional vertices of degree 2 at each arm, all at the same distance $a > 0$ from the common junction. The approximating family will be constructed as follows. The operators act, of course, as $\psi_j \mapsto -\psi'_j$ at each arm; the wavefunctions satisfy the δ conditions (6) at the central vertex with a coupling parameter $\alpha = b$, to be specified later, and another δ coupling (6), this time with the parameter c , at each of the additional vertices (see figure 1). We call such an operator $H^{b,c}(a)$.




Figure 1. Scheme of the approximation. For simplicity the graph is featured as planar; the vertical bars denote the δ coupling strength.

The crucial feature that allows us to simplify the treatment in the present situation is a symmetry. Each of the Hamiltonians H_{β} and $H^{b,c}(a)$ decomposes into a nontrivial part which acts on the one-dimensional subspace of $\mathcal{H} = \bigoplus_{j=1}^n L^2(\mathbb{R}_+)$ consisting of functions symmetric with respect to permutations, $\psi_j(x) = \psi_k(x)$ for all j, k , and the $(n-1)$ -dimensional part corresponding to Dirichlet and Neumann conditions at the central vertex for the δ and δ'_s couplings, respectively. Note that the matrices U corresponding to these couplings each have one simple eigenvalue and another one equal to ∓ 1 , respectively, of multiplicity $n-1$.

To see what the choice of the effective coupling constants b, c should be, let us first modify to our problem the argument of [CS98a]. As we have said, in the nontrivial sector all the functions are the same, so we may drop the arm index. The boundary values at $x=0$ and $x=a$ are related by

$$\psi(a) = \psi(0) + a\psi'(0) + \mathcal{O}(a^2) \quad \psi'(a-) = \psi'(0+) + \mathcal{O}(a) \quad (10)$$

$$\psi'(a+) = \psi'(a-) + c\psi(a) \quad \psi'(0+) = b\psi(0). \quad (11)$$

Eliminating $\psi(0)$ and $\psi'(0+)$ from here, we get in the leading order the relation $B(a)\psi(a) = \psi'(a+)$, where

$$B(a) := c + \frac{b}{1+ab}. \quad (12)$$

Hence the needed limit, $\beta\psi'(0+) = n\psi(0)$, is achieved as $a \rightarrow 0+$ if we choose

$$b(a) := -\frac{\beta}{na^2} \quad c(a) := -\frac{1}{a}. \quad (13)$$

In the orthogonal complement to the permutation-symmetric subspace one we can again drop the index, because the operators act in the same way on all the linear combinations of $\sum_{j=1}^n d_j \psi_j(x)$ which we can choose as the basis here, i.e. those satisfying $\sum_{j=1}^n d_j = 0$. The second of the conditions (11) is now replaced by $\psi(0) = 0$. Eliminating then the boundary values at $x=0$, we get in the leading order the relation $\psi'(a+) = (c + a^{-1})\psi(a) + \mathcal{O}(a)$. The right-hand side vanishes with the parameter choice (13), giving Neumann condition, $\psi'(0+) = 0$, in the limit.

Now we can state and prove our main result.

Theorem 1. $H^{b,c}(a) \rightarrow H_{\beta}$ as $a \rightarrow 0+$ in the norm-resolvent sense provided the coupling constants b, c are chosen in correspondence with (13).

Proof. By the same symmetry argument as above we can again reduce the problem to investigation of a pair of halfline problems. Let us start with the one having Dirichlet condition at the origin, so the free Green's function at energy k^2 is

$$G_k(x, y) = \frac{\sin kx_{<}}{k} e^{ikx_{>}} \quad (14)$$

where as usual $x_<$ is the smaller of the values x, y and vice versa. The Green's function of the operator with the δ interaction at $x = a$ is obtained easily by Krein's formula [AGHH, appendix A]

$$G_k^c(x, y) = G_k(x, y) + \frac{G_k(x, a)G_k(a, y)}{-c^{-1} - G_k(a, a)}. \tag{15}$$

On the other hand, the Green function referring to the Neumann boundary is

$$G_k^N(x, y) = \frac{\cos kx_<}{k} e^{ikx_>}. \tag{16}$$

Our aim is to show that the last two converge to each other for some $k^2 \in \mathbb{C}$. It is convenient to choose $k = i\kappa$ with $\kappa > 0$; we will see below that the denominator of the last term on the right-hand side of (15) is then nonzero for a small enough. Since the functions involved are uniformly bounded around zero, it is sufficient to compute the difference in the case when neither of the arguments is smaller than a . For the sake of definiteness suppose that $a \leq x \leq y$; then (15) and (16) give

$$G_{ik}^c(x, y) - G_{ik}^N(x, y) = \frac{e^{-\kappa x} e^{-\kappa y}}{\kappa} \left[-1 + \frac{\sinh^2 \kappa a}{-\kappa c^{-1} - e^{-\kappa x} \sinh^2 \kappa a} \right]. \tag{17}$$

If $c = -a^{-1}$, the last term behaves as $1 + \mathcal{O}(a)$ for $a \rightarrow 0+$, so

$$\lim_{a \rightarrow 0+} G_{ik}^c(x, y) = G_{ik}^N(x, y) \tag{18}$$

holds for all $x, y > 0$.

Consider next the case with the δ coupling at the origin using the same parameter values, namely $k = i\kappa$ and $a \leq x \leq y$. We are interested in the following two Green's functions,

$$G_{ik}^b(x, y) = \frac{e^{-\kappa y}}{\kappa(b + \kappa)} (b \sinh \kappa x + \kappa \cosh \kappa x) \tag{19}$$

$$G_{ik}^\beta(x, y) = \frac{e^{-\kappa y}}{\kappa(n + \beta\kappa)} (n \sinh \kappa x + \beta\kappa \cosh \kappa x) \tag{20}$$

which replace (14) and (16), respectively, in the present case. The first of them determines the full approximating Green's function by Krein's formula,

$$G_k^{b,c}(x, y) = G_k^b(x, y) + \frac{G_k^b(x, a)G_k^b(a, y)}{-c^{-1} - G_k^b(a, a)}. \tag{21}$$

Using relations (13), we express the difference

$$G_{ik}^{b,c}(x, y) - G_{ik}^\beta(x, y) = \frac{e^{-\kappa y}}{\kappa} \left[\frac{b \sinh \kappa x + \kappa \cosh \kappa x}{b + \kappa} + \frac{\frac{e^{-\kappa x}}{(b+\kappa)^2} (b \sinh \kappa x + \kappa \cosh \kappa x)^2}{\kappa a - \frac{e^{-\kappa a}}{b+\kappa} (b \sinh \kappa x + \kappa \cosh \kappa x)} - \frac{n \sinh \kappa x + \beta\kappa \cosh \kappa x}{n + \beta\kappa} \right]. \tag{22}$$

The first term in the bracket tends to $\sinh \kappa x$ as $a \rightarrow 0+$, while the third one is independent of a , so their sum in the limit gives

$$-\frac{\beta\kappa e^{-\kappa x}}{n + \beta\kappa}. \tag{23}$$

Next we take the middle term without the factor $e^{-\kappa x}$ and expand the numerator and denominator to the second power in a ; this gives its limit which differs just by the sign from (23). In other words

$$\lim_{a \rightarrow 0+} G_{ik}^{b,c}(x, y) = G_{ik}^\beta(x, y) \tag{24}$$

holds again for all $x, y > 0$. To conclude the proof, we have just to realize that as functions of x, y the differences (17) and (22) decay exponentially, so the corresponding resolvent differences converge to zero even in the Hilbert–Schmidt norm. \square

Let us add that the proven result opens the way to approximation of H_β by Hamiltonians with more regular interactions. We have mentioned that the central δ in $H^{b,c}(a)$ can be approximated by a family of potentials scaled in the usual way; the same is true for the δ interactions at the points $x_j = a$. As in the related problem discussed in [ENZ01], the question then is how fast have these approximating potentials to shrink with respect to a .

Consider finally the case of δ' coupling, i.e. the Hamiltonian \tilde{H}_β on our star graph with the boundary conditions (9) at the vertex. The approximating family will be constructed in a similar way as above; the difference is that now the wavefunctions will satisfy the δ_p conditions (8) at the central vertex with a coupling parameter $\alpha = b$, to be specified. The rest is the same, and there is another δ coupling (6) with the parameter called again c at each of the additional vertices; we denote such an operator by $\tilde{H}^{b,c}(a)$.

To realize that this problem can be again reduced to a one-dimensional analysis, denote $\varepsilon := e^{2\pi i/n}$ and introduce

$$\mathcal{G}_r := \{(\psi(x), \varepsilon^r \psi(x), \dots, \varepsilon^{r(n-1)} \psi(x)) : \psi \in L^2(\mathbb{R}_+)\}. \quad (25)$$

The graph state Hilbert space can be written as $\mathcal{H} = \bigoplus_{r=0}^{n-1} \mathcal{G}_r$. Indeed, any vector of \mathcal{H} is a unique linear combination of the vectors from \mathcal{G}_r , because the determinant of the corresponding linear system is the Vandermond determinant of $1, \varepsilon, \dots, \varepsilon^{n-1}$, which is nonzero because the latter are mutually different. It is straightforward to see that the subspaces \mathcal{G}_r are invariant under the Hamiltonians in question; the δ_p and δ' couplings act ‘trivially’ at the origin corresponding to Dirichlet and Neumann conditions, respectively, while on each of the subspaces $\mathcal{G}_r, r = 1, \dots, n$, these boundary conditions are replaced by $\psi'(0) = \frac{\alpha}{n} \psi(0)$ and $\psi(0) = \frac{\beta}{n} \psi'(0)$. Thus, we can choose

$$b(a) := -\frac{\beta}{a^2} \quad c(a) := -\frac{1}{a} \quad (26)$$

and repeat the above considerations, arriving at the following conclusion.

Theorem 2. $\tilde{H}^{b,c}(a) \rightarrow \tilde{H}_\beta$ holds in the norm-resolvent sense as $a \rightarrow 0+$ if the coupling constant families b, c are given by (26).

Acknowledgments

PE appreciates the hospitality extended to him at the Kochi University of Technology where a part of this work was done. The research has been partially supported by the Czech Ministry of Education and ASCR under the projects ME482 and K1010104.

References

- [AGHH] Albeverio S, Gesztesy F, Høegh-Krohn R and Holden H 1988 *Solvable Models in Quantum Mechanics* (Berlin: Springer)
- [AN00] Albeverio S and Nizhnik L 2000 Approximation of general zero-range potentials *Ukrainian Math. J.* **52** 582–9
- [CFT01] Cheon T, Fülöp T and Tsutsui I 2001 Symmetry, duality, and anholonomy of point interaction in one dimension *Ann. Phys.* **294** 1–23
- [CFT04] Cheon T, Fülöp T and Tsutsui I 2004 Quantum abacus *Preprint* quant-ph/0404039

- [CS98a] Cheon T and Shigehara T 1998 Realizing discontinuous wavefunctions with renormalized short-range potentials *Phys. Lett. A* **243** 111–6
- [CS98b] Cheon T and Shigehara T 1998 Some aspects of generalized contact interaction in one-dimensional quantum mechanics *Mathematical Results in Quantum Mechanics, Proc. QMath7 Conf.* ed J Dittrich *et al* (Basel: Birkhäuser) pp 203–8
- [Ex95] Exner P 1995 Lattice Kronig–Penney models *Phys. Rev. Lett.* **74** 3503–6
- [Ex96a] Exner P 1996 Contact interactions on graph superlattices *J. Phys. A: Math. Gen.* **29** 87–102
- [Ex96b] Exner P 1996 Weakly coupled states on branching graphs *Lett. Math. Phys.* **38** 313–20
- [ENZ01] Exner P, Neidhardt H and Zagrebnov V A 2001 Potential approximations to δ' : an inverse Klauuder phenomenon with norm-resolvent convergence *Commun. Math. Phys.* **224** 593–612
- [EP03] Exner P and Post O 2003 Convergence of spectra of graph-like thin manifolds *Preprint* math-ph/0312028
- [EŠ89] Exner P and Šeba P 1989 Free quantum motion on a branching graph *Rep. Math. Phys.* **28** 7–26
- [FT00] Fülöp T and Tsutsui I 2000 A free particle on a circle with point interaction *Phys. Lett. A* **264** 366–74
- [Ha00] Harmer M 2000 Hermitian symplectic geometry and extension theory *J. Phys. A: Math. Gen.* **33** 9193–203
- [KS99] Kostykin V and Schrader R 1999 Kirchhoff's rule for quantum wires *J. Phys. A: Math. Gen.* **32** 595–630
- [Ku04] Kuchment P 2004 Quantum graphs: I. Some basic structures *Waves Random Media* **14** S107–28
- [KZ01] Kuchment P and Zeng H 2001 Convergence of spectra of mesoscopic systems collapsing onto a graph *J. Math. Anal. Appl.* **258** 671–700
- [KS01] Rubinstein J and Schatzmann M 2001 Variational problems on multiply connected thin strips: I. Basic estimates and convergence of the Laplacian spectrum *Arch. Ration. Mech. Anal.* **160** 271–308
- [Sa01] Saito T 2001 Convergence of the Neumann–Laplacian on shrinking domains *Analysis* **21** 171–204
- [TFC01] Tsutsui I, Fülöp T and Cheon T 2001 Möbius structure of the spectral space of Schrödinger operators with point interaction *J. Math. Phys.* **42** 5687–97