

Centre of quantum group in roots of unity and the restriction of integrable models

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2004 J. Phys. A: Math. Gen. 37 317

(<http://iopscience.iop.org/0305-4470/37/2/003>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.90

The article was downloaded on 02/06/2010 at 17:59

Please note that [terms and conditions apply](#).

Centre of quantum group in roots of unity and the restriction of integrable models*

A Belavin

Landau Institute for Theoretical Physics, Chernogolovka 142432, Russia

E-mail: belavin@itp.ac.ru

Received 4 June 2003, in final form 11 July 2003

Published 15 December 2003

Online at stacks.iop.org/JPhysA/37/317 (DOI: 10.1088/0305-4470/37/2/003)

Abstract

We show the connection between the extended centre of the quantum group in roots of unity and the restriction of the XXZ model. We also give explicit expressions for operators that respect the restriction and act on the state space of the restricted models. The formulae for these operators are verified by explicit calculation for third-degree roots; they are conjectured to hold in the general case.

PACS numbers: 02.20.Uw, 02.30.Ik, 11.25.Hf, 75.10.Jm

1. Introduction

Alcaraz *et al* [1] discovered a remarkable fact: the XXZ model with the special open boundary conditions (OBC) and a rational value of the anisotropy parameter admits a restriction. The model arising as a result of the restriction coincides in the thermodynamic limit with one of the minimal models of CFT. The algebraic reason for the restriction was explained in [2] and [3]. It was shown in [2] that the XXZ model with the OBC considered in [1] has not only integrability but also $U_q(sl(2))$ symmetry. In roots of unity, the state space of the model decomposes to the sum of ‘good’ and ‘bad’ representations of $U_q(sl(2))$. The restriction of Alcaraz *et al* is equivalent to throwing out ‘bad’ parts and keeping only the highest vectors of ‘good’ representations. In [3], a new monodromy matrix was constructed that is bilinear in terms of generators of the quantum group $A(u)$, $B(u)$, $C(u)$, $D(u)$ connected in the usual way with the Hamiltonian of the model in [1] and compatible with integrability. The twisted trace of this monodromy matrix (Sklyanin transfer matrix) also has $U_q(sl(2))$ symmetry and admits the restriction.

In this paper, we generalize the Pasquier–Saleur construction. We show that not only $U_q(sl(2))$ -invariant but also a much wider class of OBC indeed admits the restriction (the Sklyanin construction of the transfer matrix also works for this wider class of OBC).

* Talk given at the RAQIS03 conference, Aneccy, 25-28/03/03.

The decisive condition for compatibility of the Hamiltonian, the transfer matrix and other operators with the restriction is their ‘weak’ commutativity with a special element of the quantum group. The notion of ‘weak’ commutativity and its connection with the extension of the centre of the quantum group in roots of unity is explained below.

In section 2, we describe the conditions that the operators must have in order to admit the restriction. In section 3, some such operators are found. We discuss some possible generalizations of the construction in the last section.

2. Centre of the quantum group and the restriction in roots of unity

As usual, let the R -matrix $R(u)$ denote the solution of the Yang–Baxter equation. We consider the simplest and well-known $R(u)$ matrix corresponding to the six-vertex model, whose elements can be written as

$$R_{\alpha\alpha}^{\alpha\alpha}(u) = \rho \sin(u + \eta) \quad R_{\alpha\beta}^{\alpha\beta}(u) = \rho \sin u \quad R_{\beta\alpha}^{\alpha\beta}(u) = \rho \sin \eta \quad (1)$$

where $\alpha, \beta = 1, 2, \alpha \neq \beta$ and η is the so-called anisotropy parameter.

The quantum group \mathbf{A} connected with $R(u)$ is generated by $A(u), B(u), C(u), D(u)$, entries of the monodromy matrix $L(u)$, which satisfies

$$R_{12}(u-v)L_1(u)L_2(v) = L_2(v)L_1(u)R_{12}(u-v). \quad (2)$$

As shown by Tarasov [4], the centre of \mathbf{A} in roots of unity, i.e. $\eta = \pi m/N$, is generated by the following elements of \mathbf{A} :

$$\begin{aligned} \langle A(u) \rangle &= A(u)A(u+\eta) \cdots A(u+(N-1)\eta) \\ \langle B(u) \rangle &= B(u)B(u+\eta) \cdots B(u+(N-1)\eta) \\ \langle C(u) \rangle &= C(u)C(u+\eta) \cdots C(u+(N-1)\eta) \\ \langle D(u) \rangle &= D(u)D(u+\eta) \cdots D(u+(N-1)\eta). \end{aligned}$$

For convenience, we let $\langle B(u) \rangle$ denote the central element,

$$\langle B(u) \rangle = B(u)B_1(u)$$

where

$$B_1(u) = B(u+\eta) \cdots B(u+(N-1)\eta).$$

We now fix $V = C^2 \otimes \cdots \otimes C^2$ as the representation space of our quantum group. It is easy to see that for arbitrary v ,

$$\langle B(v) \rangle = B(v)B_1(v) = 0$$

on this space. We can then define the state space of a restricted model as

$$W(v) = \text{Ker } B(v) / \text{Im } B_1(v).$$

In the limit $v \rightarrow \infty$, $B(v)$ coincides up to a scalar factor with X , one of the generators of $U_q(\mathfrak{sl}(2))$. As a result, $W(\infty)$ coincides with the space of the ‘good’ highest vectors of Pasquier–Saleur. It was shown in [2] that the Hamiltonian of the XXZ chain with OBC of special type

$$H_{XXZ} = \sum_{n=1}^{L-1} \left[\sigma_n^+ \sigma_{n+1}^- + \sigma_n^- \sigma_{n+1}^+ + \frac{\cos \eta}{2} \sigma_n^z \sigma_{n+1}^z + i \frac{\sin \eta}{2} (\sigma_n^z - \sigma_{n+1}^z) \right]$$

is invariant under the quantum algebra $U_q(\mathfrak{sl}(2))$. Here, $q = e^\eta$. Because of this, H_{XXZ} is properly defined on $W(\infty) = \text{Ker } X / \text{Im } X^{(N-1)}$.

In the thermodynamic limit, where $L \rightarrow \infty$, the spectrum of low-lying states coincides (in Cardy’s sense) with $M(N - 1/N)$, one of the minimal models of CFT. In the next section, we show that the construction in [2] can be generalized to arbitrary values of the parameter v . The corresponding Hamiltonian is [3, 5]

$$H_{XXZ} = \sum_{n=1}^{L-1} \left[\sigma_n^+ \sigma_{n+1}^- + \sigma_n^- \sigma_{n+1}^+ + \frac{\cos \eta}{2} \sigma_n^z \sigma_{n+1}^z \right] + i \frac{\sin \eta}{2} (\cot v \sigma_1^z - \cot(v + \eta) \sigma_L^z). \tag{3}$$

We now find the sufficient conditions for any operator Q to be projected on $W(v)$. It is easy to see that sufficient conditions are that there exist some operators \hat{Q} and \hat{Q}_1 for a given Q such that

$$B(v)Q = \hat{Q}B(v) \tag{4}$$

and

$$QB_1(v) = B_1(v)\hat{Q}_1. \tag{5}$$

Indeed, equation (4) guarantees that if a vector $\psi \in \text{Ker } B(v)$, then the vector $Q\psi \in \text{Ker } B(v)$. It follows from equation (5) that if the difference of two vectors ψ_1 and ψ_2 belong to $\text{Im } B_1(v)$, i.e. if $\psi_1 - \psi_2 = B_1(v)\chi$, then the difference of $Q\psi_1$ and $Q\psi_2$ also belongs to $\text{Im } B_1(v)$.

3. Sklyanin transfer matrix and other operators respecting the restriction in roots of unity

In [3], Sklyanin explained the integrability of the XXZ model with OBC of the form

$$H_{XXZ} = \sum_{n=1}^{L-1} \left[\sigma_n^+ \sigma_{n+1}^- + \sigma_n^- \sigma_{n+1}^+ + \frac{\cos \eta}{2} \sigma_n^z \sigma_{n+1}^z \right] + i \frac{\sin \eta}{2} (\cot(\xi_+ + \eta/2) \sigma_1^z + \cot(\xi_- - \eta/2) \sigma_L^z)$$

by constructing a special monodromy matrix and using it to diagonalize Hamiltonian (3) and the corresponding transfer matrix by means of the algebraic Bethe ansatz.

Let $K_+(u) = K(u + \eta/2, \xi_+)$ and $K_-(u) = K(u - \eta/2, \xi_-)$, where

$$K(u, \xi) = \begin{bmatrix} \sin(u + \xi) & 0 \\ 0 & -\sin(u - \xi) \end{bmatrix}.$$

Then K_{\pm} satisfies the boundary Yang–Baxter equations [3, 6].

The Sklyanin monodromy matrix is defined [3] as

$$\Lambda(u) = \sigma^2 L^t(-u) \sigma^2 K_+(u) L(u) = \begin{bmatrix} \Lambda_1^1 & \Lambda_1^2 \\ \Lambda_2^1 & \Lambda_2^2 \end{bmatrix}.$$

Using (2) and the boundary Yang–Baxter equation, we can prove that $\Lambda(u)$ satisfies the same equation as $K_+(u)$. It gives the commutation relations between Λ_j^i . The Sklyanin transfer matrix is defined as

$$T_S(u) = \text{tr } \Lambda(u) K_-(u) = \sin(u - \eta/2 + \xi_-) \Lambda_1^1 - \sin(u - \eta/2 - \xi_-) \Lambda_2^2.$$

Explicit expressions for the elements of the Sklyanin monodromy matrix are

$$\begin{aligned}\Lambda_1^1(u) &= \sin(u + \eta/2 + \xi_+)A(u)D(-u) + \sin(u + \eta/2 - \xi_+)C(u)B(-u) \\ \Lambda_2^2(u) &= -\sin(u + \eta/2 + \xi_+)B(u)C(-u) - \sin(u + \eta/2 - \xi_+)D(u)A(-u) \\ \Lambda_1^2(u) &= \sin(u + \eta/2 + \xi_+)B(u)D(-u) + \sin(u + \eta/2 - \xi_+)D(u)B(-u) \\ \Lambda_2^1(u) &= -\sin(u + \eta/2 + \xi_+)A(u)C(-u) - \sin(u + \eta/2 - \xi_+)C(u)A(-u).\end{aligned}$$

Let $\xi_+ = v - \eta/2$ and $\xi_- = -v - \eta/2$. Then the following relations are satisfied:

$$B(v)T_S(u) = \widehat{T}_S(u)B(v) \quad (6)$$

where the explicit expression for the Sklyanin transfer matrix for these values ξ_{pm} is

$$\begin{aligned}T_S(u) &= \sin(u - v - \eta) \sin(u + v)A(u)D(-u) \\ &\quad + \sin(u - v - \eta) \sin(u - v + \eta)C(u)B(-u) \\ &\quad + \sin(u + v) \sin(u + v)B(u)C(-u) \\ &\quad + \sin(u + v) \sin(u - v + \eta)D(u)A(-u)\end{aligned} \quad (7)$$

and

$$\begin{aligned}\widehat{T}_S(u) &= \sin(u - v) \sin(u + v + \eta)A(u)D(-u) + \sin(u - v) \sin(u - v)C(u)B(-u) \\ &\quad + \sin(u + v + \eta) \sin(u + v - \eta)B(u)C(-u) \\ &\quad + \sin(u + v + \eta) \sin(u - v)D(u)A(-u).\end{aligned}$$

We have

$$B(v)\Lambda_1^2(u) = \widehat{\Lambda}_1^2(u)B(v) \quad (8)$$

where explicitly

$$\begin{aligned}\Lambda_1^2(u) &= \sin(u + v)B(u)D(-u) + \sin(u - v + \eta)D(u)B(-u) \\ \widehat{\Lambda}_1^2(u) &= \sin(u + v + \eta)B(u)D(-u) + \sin(u - v)D(u)B(-u).\end{aligned}$$

Equations (6) and (8) were verified by direct calculation.

The operators $T_S(u)$ and $\Lambda_1^2(u)$ thus satisfy the first condition, equation (4), for the restriction. We conjecture that they also satisfy the second one, equation (5), if q is a root of unity. This conjecture was explicitly verified by direct calculation for third-degree roots for the case $T_S(u)$ (but not for the case $\Lambda_1^2(u)$). It would be nice to find an elegant general proof.

The operators $T_S(u)$ and $\Lambda_1^2(u)$ depend on one parameter. There also exists a two-parameter family of operators satisfying (4) and (5).

By definition, let

$$T(x_{ij}; u_1, u_2) = x_{11}A(u_1)D(u_2) + x_{22}D(u_1)A(u_2) + x_{12}B(u_1)C(u_2) + x_{21}C(u_1)B(u_2).$$

Then

$$B(v)T(x_{ij}(v); u_1, u_2) = T(\widehat{x}_{ij}(v); u_1, u_2)B(v) \quad (9)$$

where $T(x_{ij}(v); u_1, u_2)$ and $T(\widehat{x}_{ij}(v); u_1, u_2)$ are obtained from $T(x_{ij}; u_1, u_2)$ by suitably substituting $x_{ij}(v)$ and $\widehat{x}_{ij}(v)$ for x_{ij} and

$$\begin{aligned}x_{11}(v) &= \sin(u_1 - \eta - v) \sin(u_2 - v) \\ x_{22}(v) &= \sin(u_1 + \eta - v) \sin(u_2 - v) \\ x_{12}(v) &= -\sin(u_2 - v) \sin(u_2 - v) \\ x_{21}(v) &= -\sin(u_1 + \eta - v) \sin(u_1 - \eta - v) \\ \widehat{x}_{11}(v) &= \sin(u_1 - v) \sin(u_2 - \eta - v) \\ \widehat{x}_{22}(v) &= \sin(u_1 - v) \sin(u_2 + \eta - v) \\ \widehat{x}_{12}(v) &= -\sin(u_2 - \eta - v) \sin(u_2 + \eta - v) \\ \widehat{x}_{21}(v) &= -\sin(u_1 - v) \sin(u_1 - v).\end{aligned}$$

We have

$$T(x'_{ij}(v); u_1, u_2)B(v + \eta)B(v + 2\eta) = B(v + \eta)B(v + 2\eta)T(\hat{x}'_{ij}(v); u_1, u_2) \quad (10)$$

where

$$\begin{aligned} x'_{11}(v) &= \sin(u_1 - \eta - v) \sin(u_2 + 3\eta - v) \\ x'_{22}(v) &= \sin(u_1 - 2\eta - v) \sin(u_2 - v) \\ x'_{12}(v) &= -\sin(u_2 - v) \sin(u_2 - 3\eta - v) \\ x'_{21}(v) &= -\sin(u_1 - \eta - v) \sin(u_1 - 2\eta - v) \\ \hat{x}'_{11}(v) &= \sin(u_1 - 3\eta - v) \sin(u_2 - \eta - v) \\ \hat{x}'_{22}(v) &= \sin(u_1 - v) \sin(u_2 - 2\eta - v) \\ \hat{x}'_{12}(v) &= -\sin(u_2 - \eta - v) \sin(u_2 - 2\eta - v) \\ \hat{x}'_{21}(v) &= -\sin(u_1 - v) \sin(u_1 - 3\eta - v). \end{aligned}$$

If we require that the operators $T(x_{ij}(v); u_1, u_2)$ and $T(x'_{ij}(v); u_1, u_2)$ coincide, we can verify that this requirement is satisfied if $\eta = \pi/3$ or $\eta = 2\pi/3$. As discussed above, it follows that the two-parameter family of operators $T(x_{ij}(v); u_1, u_2)$ can be restricted on $W(v)$.

We conjecture that $T(x_{ij}(v); u_1, u_2)$ satisfies restriction conditions (4) and (5) if $\eta = m\pi/p$, where m and p are coprime integers.

It is easy to verify that the relation

$$T_S(u) = T(x_{ij}(v); u, -u)$$

holds, where $T_S(u)$ is Sklyanin transfer matrix in (7).

4. Discussion

It was shown in [7] that the Sklyanin transfer matrix for the Pasquier–Saleur case ($v \rightarrow \infty$) after the restriction satisfies the truncated system of fusion functional equations. This system defines the spectrum $M(p/p + 1)$. This statement can also be generalized to finite v . It is remarkable that the spectrum of states surviving after the restriction is independent of v [8]. This fact was discovered numerically in [5].

The explicit construction for additional central elements of the elliptic Yang–Baxter algebra in roots of unity was given in [9]. It would be interesting to generalize the approach in this paper to the elliptic case.

Another important problem is to generalize the Kitanine–Maillet–Terras construction [10] of the local operators of the XXZ model in terms of elements of the monodromy matrix to the restricted models. Namely, it would be interesting to build explicit operators that simultaneously respect the restriction and have mutual locality (i.e. commutativity). Constructing such operators would allow explicit formulae to be obtained for the correlation functions in the restricted models.

Acknowledgments

I am indebted to M Jimbo, N Kitanine, and E Sklyanin for useful discussions and also to W Everett for editorial assistance. This work was presented at the conference ‘On Recent Advances in the Theory of Quantum Integrable Systems.’ I am indebted to the organizers of the conference and especially to P Sorba for the opportunity to participate in this nice and very interesting meeting. This work is supported in part by RFBR-01-02-16686, SSRF-20044.2003.2 and INTAS-00-00055.

References

- [1] Alcaraz F C, Barber M N and Batchelor M T 1987 *Phys. Rev. Lett.* **58** 771
- [2] Pasquier V and Saleur H 1990 Common structures between finite systems and conformal field theories through quantum groups *Nucl. Phys. B* **330** 523
- [3] Sklyanin E K 1988 Boundary conditions for integrable quantum systems *J. Phys. A: Math. Gen.* **21** 2375–89
- [4] Tarasov V 1992 *Int. J. Mod. Phys. A* **7** Suppl 1B 963
- [5] Alcaraz F C, Baake M, Grimm U and Rittenberg V 1989 *J. Phys. A: Math. Gen.* **22** L5–11
- [6] Wang Y S 2000 *J. Phys. A: Math. Gen.* **33** 4009
- [7] Belavin A and Stroganov Yu 1999 Minimal models of integrable lattice theory and truncated functional equations *Phys. Lett. B* **446** 281 (*Preprint hep-th/9908050*)
- [8] Belavin A unpublished
- [9] Belavin A and Jimbo M 2002 Central elements of the elliptic Yang–Baxter algebra at roots of unity *Preprint hep-th/0208224*
- [10] Kitanine N, Maillet J M and Terras V 1999 *Nucl. Phys. B* **554** 647