

Inhomogeneous loop models with open boundaries

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2005 J. Phys. A: Math. Gen. 38 6091

(<http://iopscience.iop.org/0305-4470/38/27/001>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.92

The article was downloaded on 03/06/2010 at 03:49

Please note that [terms and conditions apply](#).

Inhomogeneous loop models with open boundaries

P Di Francesco

Service de Physique Théorique de Saclay, CEA/DSM/SPhT, URA 2306 du CNRS,
F-91191 Gif sur Yvette Cedex, France

Received 8 April 2005, in final form 17 May 2005

Published 22 June 2005

Online at stacks.iop.org/JPhysA/38/6091

Abstract

We consider the crossing and non-crossing $O(1)$ dense loop models on a semi-infinite strip, with inhomogeneities (spectral parameters) that preserve the integrability. We compute the components of the ground-state vector and obtain a closed expression for their sum, in the form of Pfaffian and determinantal formulae.

PACS numbers: 02.10.Ab, 05.50.+q

Mathematics Subject Classification: 05A19, 82B20

1. Introduction

The interplay between statistical mechanics and combinatorics is an everlasting one, and takes many different guises as time goes by. Some activity has developed recently around conjectural observations in [1–3] on the ground-state vectors of some simple two-dimensional statistical models of loops, which may alternatively be viewed as one-dimensional quantum (spin) chains. As it turned out, and among other integer numbers, the total number of alternating sign matrices (ASM) popped out of the study of the ground-state vector of the integrable quantum spin chain corresponding to the dense $O(1)$ loop model on a semi-infinite cylinder of square lattice. This number counts the total number of configurations of the ice model on a square with domain wall boundary conditions. It also counts the configurations of the fully packed loop model on a square grid, yet another type of loop model, now with two kinds of loops crossing or touching at each vertex, and connecting by pairs the points at the periphery of the grid. This opened up the road to many more observations turned into conjectures, regarding correlation functions as well as other boundary conditions, and all involving integer sequences (see for instance [4–7]). An activity also developed in trying to relate some particular subsets of configurations of the fully packed loop model to rhombus tilings of planar domains with possible conic singularities [8–12].

The idea of considering inhomogeneous versions of the loop models came with trying to modify the boundary conditions of the loop model on a cylinder by introducing dislocations in the underlying lattice [13, 14], and it was realized and proved in [15] that the full multiparameter generalization of the loop model that preserves its integrability actually leads to a ground-state vector whose sum of suitably normalized components coincides with the so-called

Izergin–Korepin determinant, defined as the partition function of the inhomogeneous six vertex model on a square grid with domain wall boundary conditions [16, 17]. This allowed us, as a by-product, to prove the conjecture of [1] that the sum of suitably normalized entries of the ground-state vector of the $O(1)$ spin chain is the total number of alternating sign matrices. The general proof of [15] takes full advantage of the integrability of the model, and transforms intertwining relations for the transfer matrix of the loop model into local recursion relations for the ground-state vector's entries.

Another loop model, very similar in nature to the $O(1)$ loop model, also includes the possibility for loops of crossing one another. This is the so-called crossing or Brauer $O(1)$ loop model, for which many combinatorial conjectures were made in [18], surprisingly relating this quantum chain to degrees of components of the commuting variety, computed in [19]. The same techniques as those used in [15], making full use of the integrability of the loop model, were applied to this case in [20], allowing us to prove most of the conjectures of [18]. The algebro-geometric interpretation of these results was extended recently in [21].

The two works [20] and [15] are only concerned with loop models wrapped on a cylinder, i.e. with periodic boundary conditions. The aim of the present paper is to investigate the case of open boundary conditions, namely of (crossing or non-crossing) inhomogeneous loop models defined on a semi-infinite strip of square lattice. By adapting the techniques of [20, 15], we will derive sum rules for the components of the corresponding ground-state vectors. The main outcome will be some particularly simple Pfaffian or determinantal formulae for the state sum as an explicit function of the model's inhomogeneities (spectral parameters). In the case of crossing loops, we will obtain a 'reflected' generalization of the results of [20], while in the non-crossing case we will be able to identify the state sum with the partition function of so-called U-turn symmetric alternating sign matrices of [22]. Let us stress at this point that, as opposed to the crossing loop case where our proof is rigorous and complete, the non-crossing case relies on an assumption we make on the total degree of the vector's components, as functions of the spectral parameters. Although we have no doubt that this is true, proving it would certainly require a lot of effort, and we prefer to concentrate on the consequences of this property on the ground-state vector.

The paper is organized as follows. Section 2 is devoted to the case of crossing loops with open boundaries. After giving definitions in section 2.1, we show in section 2.2 the fundamental intertwining relations satisfied by the transfer matrix of the model. Solving for all the subsequent relations leads to the solution of section 2.3 in the form of an explicit recursion relation defining the 'fundamental' entry of the ground-state vector, out of which all others are iteratively constructed. The above relations are turned into recursion relations for all entries of the vector in section 2.4, as well as into symmetry properties of the entries in section 2.5. Using all these properties allows for proving two sum rules on the ground-state vector components in section 2.6. Section 3 is concerned with the case of non-crossing loops. We follow the same route: definitions (section 3.1), intertwining properties (section 3.2), solution (section 3.3), recursion relations (section 3.4), symmetries (section 3.5), and finally sum rule (section 3.6). A few concluding remarks are gathered in section 4, while samples of entries of the ground-state vector are given in appendices A and B respectively for the crossing and non-crossing case.

2. The inhomogeneous $O(1)$ crossing loop model with open boundaries

2.1. Transfer matrix and basic relations

We consider the open boundary version of the inhomogeneous $O(1)$ 'Brauer' crossing loop model considered in [20]. The latter was defined on a square lattice wrapped on a semi-infinite





Figure 1. A sample configuration of the Brauer loop model on a strip of width $N = 6$ (left). We have indicated the corresponding open crossing link pattern of connection of the points 1, 2, 3, 4, 5, 6 (right).

cylinder of even perimeter, thus giving rise to periodic boundary conditions. We now consider the same model on a square lattice that covers a semi-infinite strip of width N (even or odd), with centres of the lower edges labelled 1, 2, . . . , N . On each face of this domain of the square lattice, we draw at random, say with respective probabilities a_i, b_i, c_i in the i th column (at the vertical of the point labelled i) one of the three following configurations



The strip is moreover supplemented with the following pattern of fixed configurations of loops on the (left and right) boundaries:



In a given configuration, the points 1, 2, . . . , N are connected by pairs (except for one of them if N is odd, in which case it is connected to the infinity along the strip). Such a pattern of connection is called an *open crossing link pattern*. The set of open crossing link patterns on N points is denoted by CLP_N , and has cardinality $(2n - 1)!!$ for $N = 2n$ or $N = 2n - 1$. The open crossing link patterns $\pi \in CLP_N$ span a complex vector space of dimension $(2n - 1)!!$, with canonical basis $\{|\pi\rangle\}_{\pi \in CLP_N}$. An example of loop configuration together with its link pattern are depicted in figure 1.

One interesting question is to find for given probability weights a_i, b_i, c_i , the relative probabilities of the occurrence of the crossing link patterns $\pi \in CLP_N$. The crucial property of this loop model is that it is integrable for the following choice of probability weights:

$$a(u) = \frac{2(1 - u)}{(1 + u)(2 - u)}, \quad b(u) = \frac{u(1 - u)}{(1 + u)(2 - u)}, \quad c(u) = \frac{2u}{(1 + u)(2 - u)}. \quad (2.3)$$

More precisely, the so-called R -matrix of the model is an operator acting on a vector space of open crossing link patterns or any tensor product thereof, say at some points labelled i and j of the link patterns, as follows:

$$R_{i,j}(z, w) = \begin{array}{c} \text{---} \times \text{---} \\ | \\ \text{---} \times \text{---} \\ \uparrow \\ z \end{array} \begin{array}{c} w \\ \leftarrow \\ \text{---} \times \text{---} \\ | \\ \text{---} \times \text{---} \end{array} = a(z-w) \square + b(z-w) \square + c(z-w) \square \tag{2.4}$$

Here z and w are the (arbitrary complex) spectral parameters attached respectively to the points labelled i and j , and are carried, as well as the point labels, by the oriented straight lines in the pictorial representation on the left. Each of the three possible configurations of boxes on the right acts on open crossing link patterns as follows: the configuration must be connected by its lower end to the point i and by its right end to the point j of the link patterns, thus forming new patterns whose new points i and j are the upper and left ends of the box, respectively. Alternatively, the R -matrix may act locally at points $i, i + 1$ on open crossing link patterns with N points, via the permuted matrix $\check{R} = PR$:

$$\check{R}_{i,i+1}(z, w) = \begin{array}{c} \text{---} \times \text{---} \\ \uparrow \\ \text{---} \times \text{---} \\ | \\ \text{---} \times \text{---} \\ \uparrow \\ i \quad i+1 \end{array} \begin{array}{c} z-w \\ \leftarrow \\ \text{---} \times \text{---} \\ | \\ \text{---} \times \text{---} \end{array} = a(z-w) \square + b(z-w) \square + c(z-w) \square \tag{2.5}$$

$$= a(z-w)I \otimes I + b(z-w)f_i + c(z-w)e_i$$

for $i = 1, 2, \dots, N - 1$ and where P simply permutes the point labels, so that each label is conserved along the vertical direction. We have displayed the matrix \check{R} as a linear combination of the three local operators $I \otimes I, f_i, e_i, i = 1, 2, \dots, N - 1$, which form the generators of the Brauer algebra $B_N(1)$, subject to the relations:

$$e_i^2 = e_i, \quad f_i^2 = I, \quad e_i e_{i\pm 1} e_i = e_i, \quad f_i f_{i+1} f_i = f_{i+1} f_i f_{i+1}, \tag{2.6}$$

$$[e_i, e_j] = [e_i, f_j] = [f_i, f_j] = 0 \text{ if } |i - j| > 1, \quad f_i e_i = e_i f_i = e_i,$$

and a few others involving both e 's and f 's, all clear from the pictorial representation of the action on link patterns, namely: $I \otimes I$ leaves the link patterns unchanged, f_i crosses the links terminating at points i and $i + 1$, and e_i glues the two ends of links at i and $i + 1$ and adds up a new link connecting i to $i + 1$. If a loop is formed in the process, it must simply be erased (loops are given a weight 1 here, leading to the relation $e_i^2 = e_i$).

Following Sklyanin [23], we also introduce a boundary operator $K_i(z)$, whose action is diagonal at the points labelled $i = 1$ or N , with matrix element 1, but whose effect is to switch the spectral parameter $z \rightarrow -z$ attached to that point, with the pictorial representation

$$K_i(z) = \begin{array}{c} \text{---} \\ \uparrow \\ \bullet \\ \uparrow \\ \text{---} \\ i \end{array} \begin{array}{c} -z \\ \leftarrow \\ \text{---} \\ | \\ \text{---} \\ \uparrow \\ z \end{array} \tag{2.7}$$

In addition to the standard Yang–Baxter and unitarity relations (with additive spectral parameters), reading pictorially

$$\begin{array}{c} \text{---} \times \text{---} \\ \times \\ \text{---} \times \text{---} \end{array} = \begin{array}{c} \text{---} \times \text{---} \\ \times \\ \text{---} \times \text{---} \end{array} \quad \text{and} \quad \begin{array}{c} \text{---} \times \text{---} \\ \times \\ \text{---} \times \text{---} \end{array} = \begin{array}{c} \text{---} \times \text{---} \\ \times \\ \text{---} \times \text{---} \end{array} \tag{2.8}$$

The solution to these equations for which the \check{R} matrix is a linear combination of generators $I \otimes I, e_i, f_i$ is essentially unique (up to unimportant redefinitions) and takes form (2.4). Equation (2.8) is now supplemented by the boundary Yang–Baxter relations

(2.9)

and unitarity relation

$$K_i(z)K_i(-z) = I \quad \text{or} \quad \begin{array}{c} \xrightarrow{z} \bullet \xrightarrow{-z} \\ \xrightarrow{-z} \bullet \xrightarrow{z} \end{array} = \xrightarrow{z} \quad (2.10)$$

on both sides of the strip.

The transfer matrix $T(t|z_1, z_2, \dots, z_N)$ of our model reads pictorially

(2.11)

It acts from the vector space of open crossing link patterns with N points to itself. As a consequence of the Yang–Baxter and boundary Yang–Baxter equations, the transfer matrices at two distinct values of t commute.

We denote by $\Psi^{(N)}(z_1, z_2, \dots, z_N)$ the common ground-state vector of the T for fixed values of the z_i , namely such that

$$T(t|z_1, z_2, \dots, z_N)\Psi^{(N)}(z_1, z_2, \dots, z_N) = \Psi^{(N)}(z_1, z_2, \dots, z_N). \quad (2.12)$$

As T is a rational fraction of the z_i , we normalize Ψ_N so that all its entries are coprime polynomials of the z_i . Picking say $t = 0$, we may view the entries of $\Psi^{(N)}$ as the relative probabilities of open link pattern connections in random crossing loop configurations with inhomogeneous probabilities $(a_i, b_i, c_i) = (a(z_i), b(z_i), c(z_i))$ in each column i of the strip, and equation (2.12) expresses nothing but the invariance of probabilities under the addition of two rows to the semi-infinite cylinder (left and right boundaries are indeed invariant only under translations of two lattice spacings). This interpretation is *stricto sensu* only valid in the range of z_i leading to $a_i, b_i, c_i \in [0, 1]$, in which case $\Psi^{(N)}$ is the Perron–Frobenius eigenvector of T .

A last remark is in order. It turns out that the case of odd size $N = 2n - 1$ may always be recovered from that of even size $N = 2n$, upon taking $z_{2n} \rightarrow \infty$. Indeed, considering the transfer matrix (2.11) of size N , we see that when $z_N \rightarrow \infty$, the two rightmost R -matrix elements (acting at the point labelled N) both tend to f_N , hence the action at point N decouples from the transfer matrix, and we have the reduction $T(t|z_1, \dots, z_N) \rightarrow T(t|z_1, \dots, z_{N-1}) \otimes I$, reading pictorially

(2.13)


This implies that when $z_N \rightarrow \infty$, the eigenvector $\Psi^{(N)}(z_1, \dots, z_N)$ becomes proportional (at leading order in z_N) to $\Psi^{(N-1)}(z_1, \dots, z_{N-1})$. This allows for recovering the odd N case from the even N one. Henceforth, throughout the paper and unless otherwise specified, we will always assume that N is even, and write $N = 2n$.

2.2. Intertwining

As an immediate consequence of the Yang–Baxter equation, we have the intertwining property

$$T(t|z_1, \dots, z_i, z_{i+1}, \dots, z_N) \check{R}_{i,i+1}(z_i, z_{i+1}) = \check{R}_{i,i+1}(z_i, z_{i+1}) T(t|z_1, \dots, z_{i+1}, z_i, \dots, z_N) \tag{2.14}$$

for $i = 1, 2, \dots, N - 1$, also expressed pictorially as



Applying this to the eigenvector $\Psi^{(N)}$ results in the relation

$$\Psi^{(N)}(z_1, \dots, z_i, z_{i+1}, \dots, z_N) = \check{R}_{i,i+1}(z_i, z_{i+1}) \Psi^{(N)}(z_1, \dots, z_{i+1}, z_i, \dots, z_N). \tag{2.16}$$

When written in components, the latter translates into two sets of local relations for the entries of $\Psi^{(N)}$ on the basis of open crossing link patterns, namely

$$\Theta_i \Psi_\pi^{(N)} = \Psi_{f_i \pi}^{(N)} \tag{2.17}$$

for all π with no little arch connecting points $i, i + 1$ and

$$\Delta_i \Psi_\pi^{(N)} = \sum_{\substack{\pi' \neq \pi \\ e_i \pi' = \pi}} \Psi_{\pi'}^{(N)} \tag{2.18}$$

for all π with a little arch joining points i and $i + 1$, where Θ_i and $\Delta_i, i = 1, 2, \dots, N - 1$, are local divided difference operators acting on functions of (z_1, \dots, z_N) as

$$\begin{aligned} \Theta_i &= (1 + z_i - z_{i+1}) (2\partial_i - \tau_i) \frac{1}{1 + z_i - z_{i+1}} \\ \Delta_i &= (1 + z_i - z_{i+1}) \left(1 + \frac{z_{i+1} - z_i}{2} \right) \partial_i \end{aligned} \tag{2.19}$$

where ∂_i and τ_i act on functions $f(z_1, \dots, z_N)$ as

$$\partial_i f = \frac{\tau_i f - f}{z_i - z_{i+1}} \quad \tau_i f(z_1, \dots, z_i, z_{i+1}, \dots, z_N) = f(z_1, \dots, z_{i+1}, z_i, \dots, z_N). \tag{2.20}$$

With these definitions, it is clear that $\Theta_i^2 = I$, while $\Delta_i^2 = -\Delta_i$.

An important direct consequence of equation (2.17) is that $\Psi_\pi^{(N)}$ vanishes when $z_{i+1} = 1 + z_i$ if the link pattern π has no arch joining i and $i + 1$. This is easily deduced for instance from the relation (2.16) with R as in (2.4): indeed, when $z_{i+1} = 1 + z_i$, $\check{R}_{i,i+1} \propto e_i$, and therefore only components with an arch joining i to $i + 1$ may be nonzero. By taking appropriate products of \check{R} , this was straightforwardly extended in [20], and $\Psi^{(N)}$ actually has the general property:

(P1): For any pair $i < j$ of points such that, in the link pattern π , no arch connects any pair of points among $i, i + 1, \dots, j$, the component $\Psi_\pi^{(N)}$ vanishes when $z_j = 1 + z_i$.

The first set of relations (2.17) turns out to be sufficient to generate all the entries of $\Psi^{(N)}$ from say that corresponding to the maximally crossing link pattern, still denoted π_0 by a slight abuse of notation, and that connects points i and $i + n, i = 1, 2, \dots, n$: indeed, like in [20], we just have to follow ‘paths’ from π_0 to $\pi = f_{i_1} f_{i_2} \dots f_{i_k} \cdot \pi_0$ obtained by successive actions of the generators f_i , restricted in such a way that they do not act trivially (i.e. f_i never acts on a link pattern that connects points i and $i + 1$) and apply (2.17) accordingly. Any two such paths must be equivalent modulo the braid relations $f_i f_{i+1} f_i = f_{i+1} f_i f_{i+1}, f_i^2 = I$ and $f_i f_j = f_j f_i$ for $|i - j| > 1$. It is easy to show that the Θ also satisfy the braid relations, just like the ‘gauged’ operators

$$\delta_i = 2\partial_i - \tau_i \tag{2.21}$$

in terms of which $\Theta_i = (1 + z_i - z_{i+1})\delta_i / (1 + z_i - z_{i+1})$. However, as already observed in [20] in the periodic case, the representation of the symmetric group they form is not faithful (it has dimension $(2n - 1)!!$, to be compared with the order of symmetric group, $(2n)! = (2n - 1)!! \times 2^n n!$), and we must also implement the stabilizer relations

$$\Theta_i \Theta_{i+n} \Psi_{\pi_0}^{(N)} = \Psi_{\pi_0}^{(N)} \tag{2.22}$$

for $i = 1, 2, \dots, n - 1$. So, if $\Psi_{\pi_0}^{(N)}$ obeys relations (2.22), the result of the successive actions of Θ s yielding $\Psi_\pi^{(N)}$ out of $\Psi_{\pi_0}^{(N)}$ is independent of the path from π_0 to π , and all components of $\Psi^{(N)}$ are therefore determined by equation (2.17) without ambiguity.

Relations (2.22) however do not seem to determine $\Psi_0^{(N)} \equiv \Psi_{\pi_0}^{(N)}$ completely. The other set of relations (2.18) actually serves this purpose, as we shall see in the next section. To conclude this section, let us mention two more intertwining properties, one in the ‘bulk’, and the other on the boundary. The former will lead to the main recursion relation on entries of $\Psi^{(N)}$, while the latter will allow us to derive some boundary symmetry property of $\Psi^{(N)}$, both instrumental in eventually computing the sum on the entries of $\Psi^{(N)}$. Let us denote by φ_i the embedding of $\text{CLP}_{2n-2} \rightarrow \text{CLP}_{2n}$ that inserts a little arch between points $i - 1$ and i . We have the following restriction/projection property: if two neighbouring parameters z_i and z_{i+1} are such that $z_{i+1} = 1 + z_i$, then

$$T(t|z_1, \dots, z_i, z_{i+1} = 1 + z_i, \dots, z_{2n})\varphi_i = \varphi_i T(t|z_1, \dots, z_{i-1}, z_{i+2}, z_{2n}) \tag{2.23}$$

for $i = 1, 2, \dots, N - 1$. This is proved for instance in [20] by explicitly commuting φ_i through the product of two R matrices at points i and $i + 1$, and noting that when $z_{i+1} = 1 + z_i$, $\check{R}_{i,i+1}(z_i, z_{i+1}) \propto e_i$.

Finally, using the boundary operator K_1 at the leftmost point, and applying the boundary Yang–Baxter equation (2.9), we get

$$K_1(-z_1)T(t|-z_1, z_2, \dots, z_N) = T(t|z_1, z_2, \dots, z_N)K_1(-z_1) \tag{2.24}$$

or pictorially

$$(2.25)$$

This boundary intertwining relation, when applied on the vector $\Psi^{(N)}(-z_1, z_2, \dots, z_N)$, allows us to show that the latter is proportional to $\Psi^{(N)}(z_1, z_2, \dots, z_N)$, and we find for even $N = 2n$:

$$\Psi^{(N)}(-z_1, z_2, \dots, z_N) = \Psi^{(N)}(z_1, z_2, \dots, z_N). \tag{2.26}$$

The same reasoning at the other end with the point labelled N leads to the condition

$$\Psi^{(N)}(z_1, z_2, \dots, z_{N-1}, -z_N) = \Psi^{(N)}(z_1, z_2, \dots, z_N). \tag{2.27}$$

In both equations, the proportionality factors are fixed to be 1 by the fact that $\Psi^{(N)}$ is a polynomial.

Let us also mention that the system is invariant under reflection under which the points are reflected as $i \rightarrow N + 1 - i$, and the link patterns $\pi \rightarrow \rho(\pi)$ accordingly. Operatorwise, a global reflection reverts all orientations of lines, and therefore inverts all operators, which amounts to switching all $z_i \rightarrow -z_i$. As a result, the reflected eigenvector $\Psi^{(N)\rho}(-z_N, -z_{N-1}, \dots, -z_1)$, with components $(\Psi_{\pi}^{(N)\rho}(-z_N, -z_{N-1}, \dots, -z_1)) = \Psi_{\rho(\pi)}^{(N)} \times (-z_N, -z_{N-1}, \dots, -z_1)$, is proportional to $\Psi^{(N)}(z_1, z_2, \dots, z_N)$, and as $\Psi^{(N)}$ is a polynomial, we have

$$\Psi_{\rho(\pi)}^{(N)}(-z_N, -z_{N-1}, \dots, -z_1) = \Psi_{\pi}^{(N)}(z_1, z_2, \dots, z_N) \tag{2.28}$$

for all link patterns $\pi \in \text{CLP}_N$.

An important conclusive remark is in order. An alternative transfer matrix T' for the inhomogeneous crossing loop model with open boundary conditions may be written uniquely in terms of ‘bulk’ (\check{R}) and ‘boundary’ (K) operators, as shown for instance in figure 2. The commutation of T' with T is a consequence of the Yang–Baxter and boundary Yang–Baxter equations (2.8) and (2.9). This means in turn that the relations on $\Psi^{(N)}$ inherited from the intertwining properties involving \check{R} and K (namely equations (2.17)–(2.18) and (2.26)–(2.27)) completely determine $\Psi^{(N)}$ up to a global proportionality factor, as they produce an obvious (Perron–Frobenius) eigenvector for T' . The purpose of the next section is to exhibit a candidate for $\Psi_0^{(N)}$ for which all these relations will be satisfied: this in turn will prove, by a uniqueness argument, that the candidate for $\Psi_0^{(N)}$ is indeed the right value, thus solving our problem for all components of $\Psi^{(N)}$.

2.3. Solution for $\Psi_0^{(N)}$

As mentioned in the previous section, relations (2.16) together with the boundary symmetries (2.26)–(2.27) determine $\Psi^{(N)}$ completely up to a global normalization, which we have fixed by the coprimarity requirement, provided in addition the component $\Psi_{\pi_0}^{(N)} = \Psi_0^{(N)}$ satisfies the stabilizer conditions (2.22). The entry $\Psi_0^{(N)}$ must be further determined by relations (2.18) and (2.26)–(2.27). The former actually reduce to just one of them, as we may obtain any other relation in the list (2.18) by acting on a particular one with a succession of operators




Figure 2. The alternative transfer matrix T' , together with T . The two commute, as a consequence of the Yang–Baxter and boundary Yang–Baxter equations.

Θ_i : this simply amounts to generate any other relation by means of crossing/uncrossings of consecutive arches of the corresponding link patterns, via an action of the f_i operators that never hit little arches connecting i to $i + 1$. Let us therefore examine only the simplest (and generic) case of equation (2.18), corresponding to the link pattern $\pi = f_{n-1}f_{n-2} \cdots f_1\pi_0$, with a unique little arch, connecting points n and $n + 1$. In this case, equation (2.18) reads

$$\Delta_n \Theta_{n-1} \Theta_{n-2} \cdots \Theta_1 \Psi_0^{(N)} = (1 + \Theta_n) \sum_{j=1}^{n-1} \Theta_{n-1} \Theta_{n-2} \cdots \Theta_{j+1} \Theta_{j-1} \Theta_{j-2} \cdots \Theta_1 \Psi_0^{(N)} \quad (2.29)$$

or pictorially

$$\Delta_n \Psi \begin{array}{c} \text{arches from } n \text{ to } n+1 \\ \text{---} \\ n \quad n+1 \end{array} = \sum_{j=1}^{n-1} \Psi \begin{array}{c} \text{arches from } j \text{ to } j+n \\ \text{---} \\ j \quad n \quad n+1 \quad j+n \end{array} + \Psi \begin{array}{c} \text{arches from } j \text{ to } j+n \\ \text{---} \\ j \quad n \quad n+1 \quad j+n \end{array} \quad (2.30)$$

Noting that $1 + \Theta_i = \Delta_i \times 2/(1 + z_i - z_{i+1})$, we finally get the relation

$$\Delta_n \left(\Theta_{n-1} \Theta_{n-2} \cdots \Theta_1 - \frac{2}{1 + z_n - z_{n+1}} \sum_{j=1}^{n-1} \Theta_{n-1} \Theta_{n-2} \cdots \Theta_{j+1} \Theta_{j-1} \Theta_{j-2} \cdots \Theta_1 \right) \Psi_0^{(N)} = 0. \quad (2.31)$$

Due to simple commutation relations between Θ s and monomials of the form $1 + z_i - z_j$, this may be recast into

$$\Delta_n \Phi_n = 0 \quad (2.32)$$

where

$$\Phi_n = \left(\Theta_{n-1} - \frac{2}{1 + z_n - z_{n+1}} \right) \left(\Theta_{n-2} - \frac{2}{1 + z_{n-1} - z_{n+1}} \right) \cdots \left(\Theta_1 - \frac{2}{1 + z_2 - z_{n+1}} \right) \Psi_0^{(N)}. \quad (2.33)$$

As Δ_i is proportional to ∂_i , equation (2.32) simply expresses that Φ_n must be invariant under the interchange of z_n and z_{n+1} .

By explicit calculation of $\Psi_0^{(N)}$ and Φ_n from the eigenvector condition (2.12) for the first few values of $N = 2, 4, 6$, we have observed a particularly simple formula for Φ_n , which displays the desired invariance manifestly, namely,

$$\begin{aligned} \Phi_n &= \Psi_0^{(N-2)}(z_1, \dots, z_{n-1}, z_{n+2}, \dots, z_{2n}) \\ &\quad \times \prod_{i=1}^{n-1} (a_{i,n} b_{i,n} a_{i,n+1} b_{i,n+1} a_{n,n+i+1} c_{n,n+i+1} a_{n+1,n+i+1} c_{n+1,n+i+1}) \end{aligned} \quad (2.34)$$

where we have defined for convenience

$$a_{i,j} = 1 + z_i - z_j, \quad b_{i,j} = 1 - z_i - z_j, \quad c_{i,j} = 1 + z_i + z_j. \quad (2.35)$$

According to previous section, relation (2.34), if true, determines $\Psi_0^{(N)}$ completely, and therefore fixes the whole vector $\Psi^{(N)}$ as well. The validity of (2.34) for general N also implicitly states that no spurious overall polynomial divisor of the entries of $\Psi^{(N)}$ will occur, hence does not conflict with the coprimality requirement of its components. Before using them, let us first turn relations (2.33) and (2.34) into a recursion relation for $\Psi_0^{(N)}$. This is readily done upon using the inversion formula

$$\left(\Theta_j - \frac{2}{a_{j+1,n+1}} \right)^{-1} = \left(a_{j,n+1} \Theta_j \frac{1}{a_{j+1,n+1}} \right)^{-1} = a_{j+1,n+1} \Theta_j \frac{1}{a_{j,n+1}} = \Theta_j + \frac{2}{a_{j,n+1}} \quad (2.36)$$

where we have used $\Theta_j^2 = I$. This allows us to invert equation (2.33) into

$$\begin{aligned} \Psi_0^{(N)}(z_1, \dots, z_N) &= \left(\Theta_1 + \frac{2}{a_{1,n+1}} \right) \left(\Theta_2 + \frac{2}{a_{2,n+1}} \right) \times \dots \times \left(\Theta_{n-1} + \frac{2}{a_{n-1,n+1}} \right) \\ &\quad \times \left(\Psi_0^{(N-2)}(z_1, \dots, z_{n-1}, z_{n+2}, \dots, z_N) \right. \\ &\quad \left. \times \prod_{i=1}^{n-1} (a_{i,n} b_{i,n} a_{i,n+1} b_{i,n+1} a_{n,n+i+1} c_{n,n+i+1} a_{n+1,n+i+1} c_{n+1,n+i+1}) \right). \end{aligned} \quad (2.37)$$

We now state our main result: the entry $\Psi_0^{(N)}$ of the ground-state vector of the crossing loop model with open boundaries is given by the recursion relation (2.37), with the initial condition that $\Psi_0^{(2)} = 1$. To prove this statement, we must

- (i) check that $\Psi_0^{(N)}$ as given by (2.37) is indeed a polynomial;
- (ii) check that $\Psi_0^{(N)}$ satisfies property (P1) in order for guaranteeing the polynomiality of all other entries of $\Psi^{(N)}$, obtained via actions of the Θ_i on $\Psi_0^{(N)}$, and finally
- (iii) check that $\Psi^{(N)}$ thus constructed satisfies all relations (2.17)–(2.18), (2.26)–(2.27) and (2.22).

Actually in the latter step, only (2.26)–(2.27) and (2.22) must be checked, as equation (2.17) is used to generate the other entries of $\Psi^{(N)}$, and the main recursion relation also guarantees that equation (2.18) is satisfied.

To check (i), we will rearrange the various factors a, b, c in (2.37), using $\Theta_i + \frac{2}{a_{i,n+1}} = a_{i,i+1} a_{i+1,n+1} \delta_i \times 1/(a_{i,n+1} a_{i,n+1})$, with δ_i as in (2.21), and the fact that, like ∂_i and τ_i , δ_i commutes with the functions that are symmetric under the interchange $z_i \leftrightarrow z_{i+1}$. Let us

prove by induction, that for even $N = 2n$,

$$\Psi_0^{(N)}(z_1, \dots, z_N) = P_0^{(N)}(z_1, \dots, z_N) \times \prod_{1 \leq i < j \leq n} a_{i,j} b_{i,j} a_{i+n,j+n} c_{i+n,j+n} \times \prod_{\ell=2}^n \prod_{m=n+1}^{n+\ell-1} a_{\ell,m}, \tag{2.38}$$

where $P_0^{(N)}$ is a polynomial of the z . If we alternatively define the quantity $P_0^{(N)}$ via equation (2.38), we are simply left with proving that it is a polynomial. We apply the recursion relation (2.37) to equation (2.38) with $N \rightarrow N - 2$, and with the appropriate shifts of variables $z_i \rightarrow z_{i+2}$ for $i = n, n + 1, \dots, 2n - 2$. Commuting all possible a, b, c factors that are symmetric in z_i, z_{i+1} through the δ_i , we finally get that equation (2.38) is equivalent to the relation,

$$P_0^{(N)}(z_1, \dots, z_N) = \prod_{i=1}^{n-1} \delta_i a_{i+1,n+i+1} \prod_{j=1}^{n-1} b_{j,n+1} c_{n,n+j+1} P_0^{(N-2)}(z_1, \dots, z_{n-1}, z_{n+2}, \dots, z_N), \tag{2.39}$$

which proves the desired result, as the operators δ_i transform polynomials into polynomials of the same degree: we find that $P_0^{(N)}(z_1, \dots, z_N)$ is a polynomial of the z , with total degree $3n(n - 1)/2$.

Moreover, equation (2.38) also allows us to immediately check property (ii), namely that $\Psi_0^{(N)}$ has the expected vanishing properties of (P1) when $z_j = 1 + z_i$ (simply inspect the a factors). Note finally that $P_0^{(2)}(z_1, z_2) = 1$, as is readily seen from the explicit solution of the eigenvector equation (2.12), and therefore using iteratively (2.39), and commuting all operators δ as much as possible to the left, we arrive at the closed expression:

$$P_N(z_1, \dots, z_N) = \left(\prod_{r=1}^{n-1} \prod_{i=1}^{n-r} \delta_i a_{2,n+r+i} \right) \left(\prod_{s=1}^{n-1} \prod_{j=1}^{n-s} b_{j,n+s} c_{n+1-s,n+s+j} \right). \tag{2.40}$$

We are therefore left with the final task of checking that $\Psi_0^{(N)}$, defined via (2.38) and (2.40), indeed satisfies the stabilizer property (2.22) and the boundary symmetry properties (2.26)–(2.27), which will in turn be granted for any other component of $\Psi^{(N)}$, via actions with the Θ_i . Equation (2.22) is proved in its equivalent form $\Theta_i \Psi_0^{(N)} = \Theta_{i+n} \Psi_0^{(N)}$ by induction on n : assume it is satisfied by $\Psi_0^{(N-2)}$, for $n \rightarrow n - 1$. Going back to the original equivalent formulation of the recursion relation $D_n \Psi_0^{(N)} = \Phi_n$ with

$$D_n = \Theta_{n-1} \Theta_{n-2} \cdots \Theta_1 - \frac{2}{a_{n,n+1}} \sum_{j=1}^{n-1} \Theta_{n-1} \Theta_{n-2} \cdots \Theta_{j+1} \Theta_{j-1} \Theta_{j-2} \cdots \Theta_1 \tag{2.41}$$

and using the braid relations satisfied by the Θ , it is easy to show that

$$\Theta_i D_n = D_n \Theta_{i+1} \quad \Theta_{i+n+1} D_n = D_n \Theta_{i+n+1} \tag{2.42}$$

for $i = 1, 2, \dots, n - 2$. Moreover, in this range of indices,

$$\begin{aligned} \Theta_i \Phi_n &= \left(\prod_{i=1}^{n-1} a_{i,n} b_{i,n} a_{i,n+1} b_{i,n+1} a_{n,n+i+1} c_{n,n+i+1} a_{n+1,n+i+1} c_{n+1,n+i+1} \right) \\ &\quad \times \Theta_i \Psi_0^{(N-2)}(z_1, \dots, z_{n-1}, z_{n+2}, \dots, z_N) \\ &= \left(\prod_{i=1}^{n-1} a_{i,n} b_{i,n} a_{i,n+1} b_{i,n+1} a_{n,n+i+1} c_{n,n+i+1} a_{n+1,n+i+1} c_{n+1,n+i+1} \right) \end{aligned}$$

$$\begin{aligned} & \times \Theta_{i+n+1} \Psi_0^{(N-2)}(z_1, \dots, z_{n-1}, z_{n+2}, \dots, z_N) \\ & = \Theta_{i+n+1} \Phi_n, \end{aligned} \quad (2.43)$$

where we have used the explicit symmetry of the prefactor in z_i and z_{i+1} and the induction hypothesis. Combining equations (2.42) and (2.43), we immediately get that $\Theta_{i+1} \Psi_0^{(N)} = \Theta_{i+n+1} \Psi_0^{(N)}$, which amounts to equation (2.22) for $i = 2, 3, \dots, n-1$ as the Θ are involutions. The case $i = 1$ is more tedious, as no nice commutation relations like (2.42) are available. However the property $\Theta_1 \Psi_0^{(N)} = \Theta_{n+1} \Psi_0^{(N)}$ reduces to $\delta_1 a_{2,n+1} P_0^{(N)} = \delta_{n+1} a_{2,n+1} P_0^{(N)}$, which we now prove by using the explicit expression (2.40).

We will make extensive use of definition (2.21) of δ_i and of the following modified Leibniz rule

$$\partial_i(fg) = \tau_i(f)\partial_i(g) + g\partial_i(f) = \tau_i(g)\partial_i(f) + f\partial_i(g) \quad (2.44)$$

satisfied by the divided difference operator ∂_i of equation (2.20) acting on the product of functions f, g . When translated in terms of δ_i , upon noting that $\tau_i(fg) = \tau_i(f)\tau_i(g)$, this gives

$$\delta_i(fg) = \tau_i(f)\delta_i(g) + 2g\partial_i(f). \quad (2.45)$$

Isolating the first two terms in product (2.40), thus writing $P_0^{(N)} = \delta_1 a_{2,n+2} R_0^{(N)}$, we see that the condition $\delta_1 a_{2,n+1} P_0^{(N)} = \delta_{n+1} a_{2,n+1} P_0^{(N)}$ amounts to

$$\begin{aligned} \delta_1 a_{2,n+1} \delta_1 a_{2,n+2} R_0^{(N)}(z_1, \dots, z_N) &= (a_{1,n+1} \delta_1 + 2) \delta_1 a_{2,n+2} R_0^{(N)}(z_1, \dots, z_N) \\ &= \delta_{n+1} a_{2,n+1} \delta_1 a_{2,n+2} R_0^{(N)}(z_1, \dots, z_N) \\ &= (a_{2,n+2} \delta_{n+1} + 2) \delta_1 a_{2,n+2} R_0^{(N)}(z_1, \dots, z_N) \end{aligned} \quad (2.46)$$

where we have used relation (2.45) and $\partial_1(a_{2,k}) = 1 = \partial_k(a_{2,k})$ for $k > 2$. Equation (2.46) amounts to $a_{1,n+1} R_0^{(N)} = \delta_{n+1} \delta_1 a_{2,n+2} R_0^{(N)}$, or equivalently $P_0^{(N)} = \delta_1 a_{2,n+2} R_0^{(N)} = \delta_{n+1} a_{1,n+1} R_0^{(N)}$. To best illustrate the strategy of the proof, let us first treat the case $N = 6$. First, it is easy to check directly that

$$P_0^{(4)}(z_1, z_2, z_3, z_4) = \delta_1 a_{2,4} b_{1,3} c_{2,4} = \delta_3 a_{1,3} b_{1,3} c_{2,4}. \quad (2.47)$$

Then we write

$$P_0^{(6)} = \delta_1 a_{2,5} \delta_2 a_{3,6} \delta_1 a_{2,6} Q \quad Q = b_{1,4} c_{3,5} b_{2,4} c_{3,6} b_{1,5} c_{2,6} \quad (2.48)$$

and we have to prove that $P_0^{(6)} = S_0^{(6)}$, with

$$S_0^{(6)} = \delta_4 a_{1,4} \delta_2 a_{3,6} \delta_1 a_{2,6} Q = \delta_2 a_{3,6} \delta_4 a_{1,4} \delta_1 a_{2,6} Q. \quad (2.49)$$

We now wish to commute the operator δ_4 all the way to the right. For this, we apply formula (2.45) to rewrite $a_{1,4} \delta_1 = \delta_1 a_{2,4} - 2$, which yields

$$\begin{aligned} S_0^{(6)} &= (\delta_2 a_{3,6} \delta_1 a_{2,6} \delta_4 a_{2,4} - 2 \delta_2 a_{3,6} \delta_4 a_{2,6}) Q \\ &= (\delta_2 a_{3,6} \delta_1 a_{2,6} \delta_4 a_{2,4} - 2 \delta_2 a_{3,6} a_{2,6} \delta_4) Q. \end{aligned} \quad (2.50)$$

Now we note that

$$\begin{aligned} \delta_4 a_{2,4} Q &= b_{1,4} c_{3,6} b_{1,5} c_{2,6} \delta_4 a_{2,4} b_{2,4} c_{3,5} \\ &= b_{1,4} c_{3,6} b_{1,5} c_{2,6} P_0^{(4)}(z_2, z_3, z_4, z_5) \\ &= b_{1,4} c_{3,6} b_{1,5} c_{2,6} \delta_2 a_{3,5} b_{2,4} c_{3,5} \\ &= \delta_2 a_{3,5} Q \end{aligned} \quad (2.51)$$

where we have used property (2.47) with the substitution $(z_1, z_2, z_3, z_4) \rightarrow (z_2, z_3, z_4, z_5)$.

We now take back the operator δ_2 to the left:

$$\begin{aligned}
 S_0^{(6)} &= (\delta_2 \delta_1 a_{3,6} a_{2,6} \delta_2 a_{3,5} - 2 \delta_2 a_{3,6} a_{2,6} \delta_4) Q \\
 &= (\delta_2 \delta_1 \delta_2 a_{3,6} a_{2,6} a_{3,5} - 2 \delta_2 a_{3,6} a_{2,6} \delta_4) Q \\
 &= (\delta_1 \delta_2 \delta_1 a_{3,6} a_{2,6} a_{3,5} - 2 \delta_2 a_{3,6} a_{2,6} \delta_4) Q \\
 &= (\delta_1 \delta_2 a_{3,5} \delta_1 a_{3,6} a_{2,6} - 2 \delta_2 a_{3,6} a_{2,6} \delta_4) Q \\
 &= (\delta_1 (a_{2,5} \delta_2 + 2) \delta_1 a_{3,6} a_{2,6} - 2 \delta_2 a_{3,6} a_{2,6} \delta_4) Q \\
 &= (\delta_1 a_{2,5} \delta_2 a_{3,6} \delta_1 a_{2,6} + 2 \delta_2 a_{3,6} a_{2,6} (\delta_2 - \delta_4)) Q
 \end{aligned} \tag{2.52}$$

where we have used property (2.45) and the braid relation $\delta_2 \delta_1 \delta_2 = \delta_1 \delta_2 \delta_1$. Finally, we compute $(\delta_2 - \delta_4) Q = b_{1,4} c_{3,6} b_{1,5} c_{2,6} (\delta_2 - \delta_4) c_{3,5} b_{2,4} = 0$, again as a consequence of property (2.47) with the substitution $(z_1, z_2, z_3, z_4) \rightarrow (z_2, z_3, z_4, z_5)$, which reads $(\delta_2 a_{3,5} - \delta_4 a_{2,4}) c_{3,5} b_{2,4} = 0 = a_{2,5} (\delta_2 - \delta_4) c_{3,5} b_{2,4}$. Subsequently, equation (2.52) reduces to $S_0^{(6)} = P_0^{(6)}$, which completes the proof for $N = 6$. We now turn to the case of general N . The proof is by weak induction on N . We assume the property

$$\begin{aligned}
 P_0^{(N-2r)}(z_1, \dots, z_{N-2r}) &= \delta_1 a_{2,n-r+2} R_0^{(N-2r)}(z_1, \dots, z_{N-2r}) \\
 &= \delta_{n-r+1} a_{1,n-r+1} R_0^{(N-2r)}(z_1, \dots, z_{N-2r})
 \end{aligned} \tag{2.53}$$

for all $r = 1, 2, \dots, n - 2$. We start from the formula for $S_0^{(N)} = \delta_{n+1} a_{1,n+1} R_0^{(N)}$:

$$\begin{aligned}
 S_0^{(N)} &= \delta_{n+1} a_{1,n+1} U_{2,n-1}^{(1)} U_{1,n-2}^{(2)} U_{1,n-3}^{(3)} \cdots U_{1,2}^{(n-2)} U_{1,1}^{(n-1)} Q \\
 U_{s,t}^{(r)} &= \prod_{i=s}^t \delta_i a_{i+1,n+r+i} \quad Q = \prod_{r=1}^{n-1} \prod_{i=1}^{n-r} b_{i,n+r} c_{n+1-r,n+r+i},
 \end{aligned} \tag{2.54}$$

and commute the operator δ_{n+1} all the way to the right. Actually, as $\delta_{n+1} a_{1,n+1}$ commutes with $U_{2,n-1}^{(1)}$, we simply have to commute it through $U_{1,n-2}^{(2)}$, as δ_{n+1} then also commutes with the rest of the U on its right. We now use repeatedly formula (2.45) to commute δ_{n+1} all the way to the right:

$$\begin{aligned}
 \delta_{n+1} a_{1,n+1} U_{1,n-2}^{(2)} &= \delta_{n+1} (\delta_1 a_{2,n+1} - 2) a_{2,n+3} U_{2,n-2}^{(2)} \\
 &= (\delta_1 a_{2,n+3} \delta_{n+1} a_{2,n+1} - 2 a_{2,n+3} \delta_{n+1}) U_{2,n-2}^{(2)} \\
 &= U_{1,1}^{(2)} \delta_{n+1} (\delta_2 a_{3,n+1} - 2) a_{3,n+4} U_{3,n-2}^{(2)} - 2 a_{2,n+3} U_{2,n-2}^{(2)} \delta_{n+1} \\
 &= U_{1,2}^{(2)} \delta_{n+1} a_{3,n+1} U_{3,n-2}^{(2)} - 2 U_{1,1}^{(2)} a_{3,n+4} U_{3,n-2}^{(2)} \delta_{n+1} - 2 a_{2,n+3} U_{2,n-2}^{(2)} \delta_{n+1} \\
 &= U_{1,n-2}^{(2)} \delta_{n+1} a_{n-1,n+1} - 2 \sum_{r=1}^{n-2} U_{1,r-1}^{(2)} a_{r+1,n+r+2} U_{r+1,n-2}^{(2)} \delta_{n+1}.
 \end{aligned} \tag{2.55}$$

As the term $\delta_{n+1} a_{n-1,n+1}$ commutes with all $U_{1,n-r}^{(r)}$ for $r \geq 3$, we finally get

$$\begin{aligned}
 S_0^{(N)} &= U_{2,n-1}^{(1)} U_{1,n-2}^{(2)} \cdots U_{1,1}^{(n-1)} \delta_{n+1} a_{n-1,n+1} Q \\
 &\quad - 2 \sum_{r=1}^{n-2} U_{2,n-1}^{(1)} U_{1,r-1}^{(2)} a_{r+1,n+r+2} U_{r+1,n-2}^{(2)} U_{1,n-3}^{(3)} \cdots U_{1,1}^{(n-1)} \delta_{n+1} Q.
 \end{aligned} \tag{2.56}$$

We now note that

$$\begin{aligned}
 \delta_{n+1} a_{n-1,n+1} Q &= \frac{Q}{b_{n-1,n+1} c_{n,n+2}} \delta_{n+1} a_{n-1,n+1} b_{n-1,n+1} c_{n,n+2} \\
 &= \frac{Q}{b_{n-1,n+1} c_{n,n+2}} P_0^{(4)}(z_{n-1}, z_n, z_{n+1}, z_{n+2}) \\
 &= \frac{Q}{b_{n-1,n+1} c_{n,n+2}} \delta_{n-1} a_{n,n+2} b_{n-1,n+1} c_{n,n+2},
 \end{aligned} \tag{2.57}$$

where we have commuted δ_{n+1} through the piece of Q symmetric in z_{n+1}, z_{n+2} and used property (2.47) for $(z_1, z_2, z_3, z_4) \rightarrow (z_{n-1}, z_n, z_{n+1}, z_{n+2})$. We must now take the operator δ_{n-1} to the left. Again, it is readily seen to commute with $U_{1,n-r}^{(r)}$ for $r = n-1, n-2, \dots, 3$. We therefore concentrate on

$$\begin{aligned}
 U_{2,n-1}^{(1)} U_{1,n-2}^{(2)} \delta_{n-1} a_{n,n+2} &= U_{2,n-2}^{(1)} U_{1,n-3}^{(2)} \delta_{n-1} a_{n,2n} \delta_{n-2} a_{n-1,2n} \delta_{n-1} a_{n,n+2} \\
 &= U_{2,n-2}^{(1)} U_{1,n-3}^{(2)} \delta_{n-1} \delta_{n-2} \delta_{n-1} a_{n,2n} a_{n-1,2n} a_{n,n+2} \\
 &= U_{2,n-2}^{(1)} U_{1,n-3}^{(2)} \delta_{n-2} \delta_{n-1} a_{n,n+2} a_{n,2n} \delta_{n-2} a_{n-1,2n} \\
 &= U_{2,n-2}^{(1)} U_{1,n-3}^{(2)} \delta_{n-2} (a_{n-1,n+2} \delta_{n-1} + 2) a_{n,2n} U_{n-2,n-2}^{(2)} \\
 &= U_{2,n-2}^{(1)} U_{1,n-3}^{(2)} \delta_{n-2} a_{n-1,n+2} \delta_{n-1} a_{n,2n} U_{n-2,n-2}^{(2)} + 2 U_{2,n-2}^{(1)} U_{1,n-3}^{(2)} a_{n,2n} a_{n-1,2n} \delta_{n-1}^2 \\
 &= U_{2,n-2}^{(1)} U_{1,n-3}^{(2)} \delta_{n-2} a_{n-1,n+2} U_{n-1,n-1}^{(1)} U_{n-2,n-2}^{(2)} + 2 U_{2,n-1}^{(1)} U_{1,n-3}^{(2)} a_{n-1,2n} \delta_{n-1} \\
 &= \delta_1 a_{2,n+2} U_{2,n-1}^{(1)} U_{1,n-2}^{(2)} + 2 \sum_{r=1}^{n-2} U_{2,n-1}^{(1)} U_{1,n-r-2}^{(2)} a_{n-r,2n-r+1} \delta_{n-r} U_{n-r+1,n-2}^{(2)}
 \end{aligned} \tag{2.58}$$

where we have repeatedly used (2.45) and the braid relations, and also $\delta^2 = 1$ to rewrite $a_{n-r+1,2n-r+1} a_{n-r,2n-r+1} = \delta_{n-r} a_{n-r+1,2n-r+1} a_{n-r,2n-r+1} \delta_{n-r}$. Comparing equation (2.58) on one hand and (2.55) multiplied on the left by $U_{2,n-2}^{(1)}$ on the other hand, we find

$$\begin{aligned}
 \delta_{n+1} a_{1,n+1} U_{2,n-2}^{(1)} U_{1,n-2}^{(2)} &= \delta_1 a_{2,n+2} U_{2,n-1}^{(1)} U_{1,n-2}^{(2)} \\
 &+ 2 U_{2,n-1}^{(1)} \sum_{r=1}^{n-2} U_{1,r-1}^{(2)} a_{r+1,n+r+2} (\delta_{r+1} - \delta_{n+1}) U_{r+2,n-2}^{(2)}.
 \end{aligned} \tag{2.59}$$

When multiplied by all the remaining factors $U_{1,n-3}^{(3)} \cdots U_{1,1}^{(n-1)} Q$ on the right, equation (2.59) reads

$$S_0^{(N)} = P_0^{(N)} + 2 U_{2,n-1}^{(1)} \sum_{r=1}^{n-2} U_{1,r-1}^{(2)} a_{r+1,n+r+2} (\delta_{r+1} - \delta_{n+1}) U_{r+2,n-2}^{(2)} U_{1,n-3}^{(3)} \cdots U_{1,1}^{(n-1)} Q. \tag{2.60}$$

Each term in the sum is now computed by invoking the weak induction hypothesis (2.53) for the polynomials $P_0^{(N-2r)}(z_{r+1}, z_{r+2}, \dots, z_N)$, which yields

$$\begin{aligned}
 0 &= (\delta_{r+1} a_{r+2,n+2} - \delta_{n+1} a_{r+1,n+1}) R_0^{(N-2r)}(z_{r+1}, z_{r+2}, \dots, z_N) \\
 &= a_{r+1,n+1} (\delta_{r+1} - \delta_{n+1}) R_0^{(N-2r)}(z_{r+1}, z_{r+2}, \dots, z_N)
 \end{aligned} \tag{2.61}$$

by use of equation (2.45), and where

$$R_0^{(N-2r)}(z_{r+1}, z_{r+2}, \dots, z_N) = U_{r+2,n-2}^{(2)} U_{r+1,n-3}^{(3)} \cdots U_{r+1,r+1}^{(n-r-1)} Q' \tag{2.62}$$

with $Q' = \prod_{s=1}^{n-r-1} \prod_{i=1}^{n-r-s} b_{i+r,n+s} c_{n+1-s,n+s+i}$. Noting finally that the quantity $U_{r+2,n-2}^{(2)} U_{r+1,n-3}^{(3)} \cdots U_{r+1,r+1}^{(n-r-1)} Q'$ is exactly the piece of $U_{r+2,n-2}^{(2)} U_{1,n-3}^{(3)} \cdots U_{1,1}^{(n-1)} Q$ that does not commute with $(\delta_{r+1} - \delta_{n+1})$, we find that each term in the sum of (2.60) vanishes identically. We conclude that $S_0^{(N)} = P_0^{(N)}$, which completes the proof that $\Theta_1 \Psi_0^{(N)} = \Theta_{n+1} \Psi_0^{(N)}$.

To finally prove equations (2.26) and (2.27), we first note that $\Psi_0^{(N)} / P_0^{(N)}$, expressed through (2.38), is manifestly even in z_1 and z_{2n} , as the quantity $a_{1,j} b_{1,j}$ is invariant under $z_1 \rightarrow -z_1$, while $a_{j,2n} c_{j,2n}$ is invariant under $z_{2n} \rightarrow -z_{2n}$ for all j in the range of the product. Proceeding by induction on N and assuming that $P_0^{(N-2)}(z_1, \dots, z_{N-2})$ is even in z_1 and z_{N-2} , we immediately get from equation (2.39) that $P_0^{(N)}(z_1, \dots, z_{2n-1}, -z_{2n}) =$

$P_0^{(N)}(z_1, \dots, z_{2n-1}, z_{2n})$, as when $z_{2n} \rightarrow -z_{2n}$ the quantity $a_{n,2n}c_{n,2n}$, that carries the only dependence on z_{2n} in the prefactor, remains invariant. To prove the property for z_1 , we simply note that (2.26) is a consequence of equation (2.27) and the reflection symmetry (2.28), which we just have to prove for $\Psi_0^{(N)}$, whose link pattern is reflection-symmetric $\pi_0 = \rho(\pi_0)$. This is again done by induction on n . Denoting by $\tilde{z} = (-z_{2n}, -z_{2n-1}, \dots, -z_2, -z_1)$, we have the following property:

$$\Theta_i f(w)|_{w \rightarrow \tilde{z}} = \Theta_{2n-i} f(\tilde{z}) \tag{2.63}$$

as a direct consequence of the definition of Θ . Hence performing the substitution $z \rightarrow \tilde{z}$ in Φ_n results in $\Phi_n(\tilde{z}) = \tilde{D}_n \Psi_0^{(N)}(\tilde{z})$, where

$$\tilde{D}_n = \Theta_{n+2} \Theta_{n+3} \cdots \Theta_{2n-1} - \frac{2}{a_{n,n+1}} \sum_{j=1}^{n-1} \Theta_{n+2} \Theta_{n+3} \cdots \Theta_{n+j} \Theta_{n+j+2} \Theta_{n+j+3} \cdots \Theta_{2n-1}. \tag{2.64}$$

Letting this operator act on $\Psi_0^{(N)}(z)$ rather than on $\Psi_0^{(N)}(\tilde{z})$, we may use equation (2.22) repeatedly to rewrite the result as $\tilde{D}_n \Psi_0^{(N)}(z) = D_n \Psi_0^{(N)}(z)$: indeed, first replacing $\Theta_{2n-1} \rightarrow \Theta_{n-1}$ and commuting it all the way to the left, then repeating this for $\Theta_{2n-2} \rightarrow \Theta_{n-2}$, etc, until all original Θ factors are transformed, takes \tilde{D}_n back to D_n . By the induction hypothesis, we have $\Phi_n(\tilde{z}) = \Phi_n(z)$ as all prefactors of $\Psi_0^{(N-2)}$ are manifestly reflection invariant. We conclude that $\Phi_n(\tilde{z}) = \tilde{D}_n \Psi_0^{(N)}(\tilde{z})$ coincides with $\Phi_n(z) = D_n \Psi_0^{(N)}(z) = \tilde{D}_n \Psi_0^{(N)}(z)$, hence $\Psi_0^{(N)}(\tilde{z}) - \Psi_0^{(N)}(z)$ is annihilated by the invertible operator \tilde{D}_n , therefore vanishes identically, and we have proved the desired reflection symmetries.

This completes the proof that $\Psi_0^{(N)}$ is given by equations (2.38) and (2.40). By inspection, as the Θ are degree-preserving operators, we deduce immediately that $\Psi_0^{(2n)}$ is a polynomial of total degree $4n(n-1)$ while the partial degree in each variable is $4(n-1)$. In particular, in the reduction from size $N = 2n$ to size $N - 1 = 2n - 1$ mentioned in section 2.1, we have the following formula:

$$\Psi^{(N-1)}(z_1, \dots, z_{N-1}) = \lim_{z_N \rightarrow \infty} \frac{1}{z_N^{4(n-1)}} \Psi^{(N)}(z_1, \dots, z_N). \tag{2.65}$$

Recursion (2.39) may be implemented quite efficiently upon using the modified Leibniz rule (2.44). For $n = 2$, we have for instance

$$\begin{aligned} P_0^{(4)} &= (2\partial_1 - \tau_1)(a_{2,4}b_{1,3}c_{2,4}) \\ &= 2(b_{1,3}c_{2,4} + a_{1,4}c_{2,4} + a_{1,4}b_{2,3}) - a_{1,4}b_{2,3}c_{1,4} \\ &= b_{2,3}a_{1,4}b_{1,4} + 2c_{2,4}(a_{1,4} + b_{1,3}) \\ &= 5 + 3z_2 - 3z_3 - 2z_2z_3 - z_1^2 - z_4^2 + (z_1^2 - z_4^2)(z_2 + z_3) \end{aligned} \tag{2.66}$$

where we have used definition (2.19), the Leibniz rule (2.44), and the fact that $2 - c_{1,4} = b_{1,4}$. The explicit value of the whole vector $\Psi^{(4)}$ in terms of the z is given in appendix A, as well as that of $\Psi_0^{(6)}$. An important remark is in order. The action of the operators $2\partial_i$ on the a, b, c may only produce 2, -2 or 0 as an answer, henceforth starting from equations (2.38) and (2.39) and using the modified Leibniz formula (2.44) repeatedly ensures by induction on n that the final result for $\Psi_0^{(N)}$ may be written in general as an integer linear combination of products of a, b, c , the coefficients being only ‘ \pm ’ powers of 2, as is the case in the third line of equation (2.66). This in turn guarantees that $\Psi_0^{(N)}(0, 0, \dots, 0)$ is a (positive) integer in the homogeneous limit, and this property goes over to all other components of $\Psi^{(N)}$ via Θ actions. To compute the values of $\Psi_0^{(N)}(0, 0, \dots, 0)$ using the recursion relation (2.39), we only need to know the intermediate steps $P_0^{(N-2k)}$ up to terms of degree

$(n - k) + (n - k + 1) + \dots + (n - 1) = kn - k(k + 1)/2$. With this restriction, we have found the values:

$$1, 5, 129, 17\,369, 12\,275\,137, 45\,692\,809\,149, \dots \tag{2.67}$$

of $\Psi_0^{(N)}(0, 0, \dots, 0)$ for $N = 2, 4, 6, 8, 10, 12, \dots$ (the value 129 for $\Psi_0^{(6)}(0, 0, \dots, 0)$ may be read off equation (A.4) of appendix A).

Remarkably, a recursion relation similar to (2.37) may be derived in the case of a system with periodic boundary conditions, with even size $N = 2n$. In that case, the entry corresponding to the maximally crossing link pattern π_0 was shown to read [20]

$$\Psi_{0,\text{per}}^{(N)}(z_1, \dots, z_{2n}) = \prod_{i=1}^{2n} \prod_{k=1}^{n-1} a_{i,i+k} \tag{2.68}$$

where the indices are taken modulo N (with the convention that $i + N \equiv i$, for $i = 1, 2, \dots, N$). Using (2.68) it is easy to prove by induction that

$$\begin{aligned} \Psi_{0,\text{per}}^{(N)}(z_1, \dots, z_{2n}) &= \left(\Theta_1 + \frac{2}{1 + z_1 - z_{n+1}} \right) \left(\Theta_2 + \frac{2}{1 + z_2 - z_{n+1}} \right) \\ &\dots \left(\Theta_{n-1} + \frac{2}{1 + z_{n-1} - z_{n+1}} \right) \\ &\times \left(\Psi^{(N-2)}(z_1, \dots, z_{n-1}, z_{n+2}, \dots, z_{2n}) \prod_{i=1}^{n-1} a_{i,n} a_{i,n+1} a_{n,n+i+1} a_{n+1,n+i+1} \right). \end{aligned} \tag{2.69}$$

Actually, the ‘inverse’ formula analogous to (2.34) was obtained in [20], and was the keypoint of the proof in that case. So, in a certain sense, the recursion relation (2.37) is a natural extension of the recursion relation (2.69).

Another important property of $\Psi^{(N)}$ concerns its leading term, that is its piece of degree $4n(n - 1)$ in the z . We actually have the property:

(P2): The leading terms in $\Psi_\pi^{(N)}$ read

$$\Psi_\pi^{(N)} \sim (-1)^{c(\pi)} \frac{\prod_{1 \leq i < j \leq N} (z_i^2 - z_j^2)}{\prod_{\substack{\text{pairs } (i < j) \\ \text{connected in } \pi}} (z_i^2 - z_j^2)} \tag{2.70}$$

where $c(\pi)$ denotes the number of arch crossings in π .

To prove (P2), we first show by induction on n that it is satisfied by $\pi = \pi_0$. For this, we use (2.39) and note that at large z , $\delta_i \sim -\tau_i$, as the piece $2\partial_i$ lowers the degree, and $a_{i,j} \sim (z_i - z_j)$, $c_{i,j} \sim -b_{i,j} \sim (z_i + z_j)$. We get

$$P_0^{(N)} \sim P_0^{(N-2)}(z_1, \dots, z_{n-1}, z_{n+2}, \dots, z_N) \prod_{i=1}^{n-1} (z_1 - z_{n+i+1})(z_{i+1} + z_{n+1})(z_{n-1} + z_{n+i+1})$$

which, together with $P_0^{(2)} = 1$, gives the leading behaviour

$$P_0^{(N)}(z_1, \dots, z_N) \sim \prod_{i=2}^n \prod_{j=n+1}^{i+n-1} (z_i + z_j) \prod_{1 \leq k \leq \ell \leq n-1} (z_k^2 - z_{\ell+n+1}^2). \tag{2.72}$$

Once translated back in terms of $\Psi_0^{(N)}$ via (2.38), this yields the desired result (2.70) for $\pi = \pi_0$, with an overall sign $(-1)^{n(n-1)/2}$ from the b factors, where $n(n - 1)/2 = c(\pi_0)$ coincides with the number of crossings in π_0 . For an arbitrary link pattern π , we simply have

to act with operators Θ_i along a minimal path from π_0 to π . As we have $\Theta_i \sim \tau_i$ for large z , equation (2.71) follows immediately, as each transposition of neighbouring variables yields an overall minus sign, which parallels the fact that the number of crossings is decreased by 1 by the action of f_i on the corresponding link pattern.

To conclude this section, let us stress that we have now given a constructive definition of $\Psi_0^{(N)}$, leading to a vector $\Psi^{(N)}$ polynomial of total degree $4n(n-1)$ in the z . For it to match the other definition in section 2.1, we still have to check that no spurious non-trivial polynomial factor divides all entries of $\Psi^{(N)}$ (entries are coprime). This will actually be proved in section 2.6, when computing sum rules on the entries of $\Psi^{(N)}$.

2.4. Recursion relations

We now use the intertwining properties of section 2.2 to derive recursion relations for the entries of $\Psi^{(N)}$. Given a link pattern π , two situations may occur at a given pair of consecutive points $(i, i+1)$:

- (i) the pattern π has no arch connecting i to $i+1$, in which case property (P1) yields

$$\Psi_\pi^{(N)}(z_1, \dots, z_N)|_{z_{i+1}=1+z_i} = 0; \tag{2.73}$$

- (ii) the pattern π has a little arch joining i to $i+1$, in which case

$$\begin{aligned} \Psi_\pi^{(N)}(z_1, \dots, z_N)|_{z_{i+1}=1+z_i} &= \Psi_{\pi'}^{(N-2)}(z_1, \dots, z_{i-1}, z_{i+2}, \dots, z_N) \\ &\times \prod_{\substack{k=1 \\ k \neq i, i+1}}^N (2+z_i+z_k)(2+z_i-z_k)(1+z_k-z_i)(1-z_k-z_i) \end{aligned} \tag{2.74}$$

where π' is the link pattern π with the little arch $i, i+1$ removed ($\pi = \varphi_i \pi', \pi' \in \text{CLP}_{2n-2}$).

The latter is readily obtained by applying equation (2.23) to the vector $\Psi^{(N)}$ at $z_{i+1} = 1+z_i$, which shows proportionality between the restricted vector $\Psi^{(N)}$ and the projected one $\Psi^{(N-2)}$, and the polynomial proportionality factor in-between is fixed by the value of $\Theta_{n-1} \Theta_{n-2} \dots \Theta_1 \Psi_0^{(N)}$ extracted from equations (2.31), (2.33) and (2.34).

2.5. Symmetries

In the following section we will derive two sum rules for the components of $\Psi^{(N)}$. Before going into this let us display some symmetry properties of the sum over two particular sets of components of $\Psi^{(N)}$. We again concentrate on even $N = 2n$, unless otherwise specified.

By analogy with the case of periodic boundary conditions of [20], we may consider an interesting subset of the link patterns, which we call the *permutation sector*, in which each link pattern only connects points $1, 2, \dots, n$ to points among $n+1, n+2, \dots, 2n$. The name permutation sector is clear, as the connections may be encoded via a permutation $\sigma \in S_n$, namely $i \rightarrow n + \sigma(i)$ for instance. The simplest example of a link pattern in the permutation sector is the maximally crossing link pattern π_0 , which corresponds to the identity permutation.

Let b_N denote the indicator vector of the permutation sector, with entries equal to 1 in the sector, and 0 outside. Then we have the following relations:

$$b_N I = b_N, \quad b_N f_i = b_N, \quad b_N e_i = 0 \tag{2.75}$$

for $i \neq n$, which lead to

$$b_N \check{R}_{i,i+1}(z_i, z_{i+1}) = \frac{(1 - \frac{1}{2}(z_i - z_{i+1}))(1 + z_i - z_{i+1})}{(1 + \frac{1}{2}(z_i - z_{i+1}))(1 - z_i + z_{i+1})} b_N = \frac{(1 + a_{i+1,i})a_{i,i+1}}{(1 + a_{i,i+1})a_{i+1,i}} b_N. \tag{2.76}$$

Introducing

$$W^{(N)}(z_1, \dots, z_N) = b_n \cdot \Psi^{(N)}(z_1, \dots, z_N) \tag{2.77}$$

the sum over the entries of $\Psi^{(N)}$ in the permutation sector, let us act with b_N on both sides of equation (2.16). This immediately yields the symmetry relation

$$(1 + a_{i,i+1})a_{i+1,i}W^{(N)}(z_1, \dots, z_i, z_{i+1}, \dots, z_N) = (1 + a_{i+1,i})a_{i,i+1}W^{(N)}(z_1, \dots, z_{i+1}, z_i, \dots, z_N) \tag{2.78}$$

valid for $i \neq n$. More generally, replacing equation (2.16) with the action of a suitable chain-product of \check{R} allows us to express $\Psi^{(N)}$ as a product of consecutive \check{R} acting on $\Psi^{(N)}$ with z_i and z_j interchanged, for any pair of points i, j . This translates immediately into the generalized relation

$$(1 + a_{i,j})a_{j,i}W^{(N)}(\dots, z_i, \dots, z_j, \dots) = (1 + a_{j,i})a_{i,j}W^{(N)}(\dots, z_j, \dots, z_i, \dots) \tag{2.79}$$

valid only if both $i, j \leq n$ or both $i, j > n$. Using the boundary reflection symmetry (2.26), we also deduce that

$$W^{(N)}(-z_1, z_2, \dots, z_N) = W^{(N)}(z_1, z_2, \dots, z_N) = W^{(N)}(z_1, z_2, \dots, -z_N). \tag{2.80}$$

The same reasoning applies to the sum over all entries of $\Psi^{(N)}$. Let v_N denote the vector with all entries equal to 1, then it satisfies

$$v_N I = v_N, \quad v_N f_i = v_N, \quad v_N e_i = v_N, \quad v_N \check{R}_{i,i+1}(z_i, z_{i+1}) = v_N. \tag{2.81}$$

Then the sum over all components of $\Psi^{(N)}$,

$$Z^{(N)}(z_1, \dots, z_N) = v_N \cdot \Psi^{(N)}(z_1, \dots, z_N), \tag{2.82}$$

is symmetric in the z , as follows immediately from acting on both sides of equation (2.16) with v_N . $Z^{(N)}$ also satisfies the above-mentioned boundary reflection symmetries under $z_1 \rightarrow -z_1$ and $z_N \rightarrow -z_N$.

2.6. Sum rules

2.6.1. *Sum rule in the permutation sector.* We have, for even $N = 2n$ or odd $N = 2n - 1$,

$$W^{(N)}(z_1, \dots, z_N) = \prod_{1 \leq i < j \leq n} a_{i,j} b_{i,j} (1 + c_{i,j}) (1 + a_{j,i}) \prod_{n+1 \leq i < j \leq N} a_{i,j} c_{i,j} (1 + b_{i,j}) (1 + a_{j,i}). \tag{2.83}$$

The odd case is as a direct consequence of the even one, by application of equation (2.65).

To prove this for $N = 2n$, let us use the symmetry relation (2.79): the rhs of equation (2.79) vanishes if $z_j = 1 + z_i$ and also if $z_j = z_i - 2$, hence $W^{(N)}(z_1, \dots, z_N)$ must factor out a term $\prod_{1 \leq i < j \leq n} a_{i,j} (1 + a_{j,i}) a_{i+n,j+n} (1 + a_{j+n,i+n})$. Note that the factors $a_{i,j}$ here correspond to the simultaneous vanishings of all the components of $\Psi^{(N)}$ in the permutation sector, according to property (P1), as the only possible occurrence of a little arch connecting two consecutive points in a link pattern of the permutation sector is between points n and $n + 1$. Writing $W^{(N)}$ as

$$W^{(N)}(z_1, \dots, z_N) = X^{(N)}(z_1, \dots, z_N) \prod_{1 \leq i < j \leq n} a_{i,j} (1 + a_{j,i}) a_{i+n,j+n} (1 + a_{j+n,i+n}) \tag{2.84}$$

for some polynomial $X^{(N)}$, we deduce from equation (2.79) that $X^{(N)}$ is symmetric separately in z_1, z_2, \dots, z_n and in $z_{n+1}, z_{n+2}, \dots, z_{2n}$. Moreover, equation (2.80) implies that $W^{(N)}(-z_1, \dots, z_N) = W^{(N)}(z_1, \dots, z_N)$ has extra factors of $\prod_{j=2}^n b_{1,j} (1 + c_{1,j})$, and

similarly $W^{(N)}(z_1, \dots, -z_N) = W^{(N)}(z_1, \dots, z_N)$ has extra factors of $\prod_{i=n+1}^{2n-1} c_{j,2n}(1+b_{j,2n})$. These two quantities must therefore divide $X^{(N)}$, but by the above-mentioned symmetries of $X^{(N)}$, it must also be a multiple of $\prod_{1 \leq i < j \leq n} b_{i,j}(1+c_{i,j})$ and of $\prod_{n+1 \leq i < j \leq 2n} c_{i,j}(1+b_{i,j})$. This exhausts all factors in equation (2.83). We have finally found a total of $4n(n-1)$ factors for $W^{(N)}$, which is therefore entirely fixed to be given by (2.83) up to a constant, as it is a polynomial of degree $4n(n-1)$. The constant is now further fixed to be 1 by the leading term of $W^{(N)}$. Indeed, using property (P2) and equation (2.70), we may write the leading term in $W^{(N)}$ as

$$\begin{aligned} W^{(N)} &\sim \Delta(z_1^2, \dots, z_{2n}^2) \sum_{\sigma \in S_n} \frac{\text{sgn}(\sigma)}{\prod_{i=1}^n (z_i^2 - z_{n+\sigma(i)}^2)} \\ &= \Delta(z_1^2, \dots, z_{2n}^2) \det \left(\frac{1}{(z_i^2 - z_{n+i}^2)} \right)_{1 \leq i, j \leq n} \\ &= \Delta(z_1^2, \dots, z_{2n}^2) \frac{\Delta(z_1^2, \dots, z_n^2) \Delta(z_{n+1}^2, \dots, z_{2n}^2)}{\prod_{i,j=1}^n z_i^2 - z_{j+n}^2} \\ &= \Delta(z_1^2, \dots, z_n^2)^2 \Delta(z_{n+1}^2, \dots, z_{2n}^2)^2 \end{aligned} \tag{2.85}$$

where we have used the shorthand notation $\Delta(m_1, \dots, m_p)$ for the Vandermonde determinant $\prod_{1 \leq i < j \leq p} (m_i - m_j)$, applied the parametrization of link patterns in the permutation sector by the permutations $\sigma \in S_n$, interpreted (-1) to the number of crossings as the signature of the permutation, and finally applied the Cauchy determinant formula to re-express the resulting determinant as a product. The leading behaviour of the rhs of equation (2.83) is readily checked to coincide with this product.

As a side result of the sum rule (2.83), we conclude that the vector $\Psi^{(N)}$ constructed in section 2.3 indeed satisfies the coprimality constraint on its components, as its degree must be at least $4n(n-1)$ from the necessary factors of its sum rule within the permutation sector, leaving no place for overall spurious polynomial factors.

Note finally that in the homogeneous limit where all $z \rightarrow 0$, we simply get the integers

$$W^{(2n)}(0, 0, \dots, 0) = 2^{2n(n-1)} \quad W^{(2n-1)}(0, 0, \dots, 0) = 2^{2(n-1)^2} \tag{2.86}$$

as the sum of the integer entries of $\Psi^{(N)}(0, \dots, 0)$ in the permutation sector.

2.6.2. *Sum rule for all components of $\Psi^{(N)}$.* We have for even $N = 2n$:

$$\begin{aligned} Z^{(N)}(z_1, \dots, z_N) &= \prod_{1 \leq i < j \leq N} \frac{1 - (z_i - z_j)^2}{z_i - z_j} \frac{1 - (z_i + z_j)^2}{z_i + z_j} \\ &\times \text{Pf} \left(\frac{z_i - z_j}{1 - (z_i - z_j)^2} \frac{z_i + z_j}{1 - (z_i + z_j)^2} \right)_{1 \leq i < j \leq N} \end{aligned} \tag{2.87}$$

while for odd $N = 2n - 1$, we have

$$Z^{(N-1)}(z_1, \dots, z_{N-1}) = \lim_{z_N \rightarrow \infty} \frac{1}{z_N^{4(n-1)}} Z^{(N)}(z_1, \dots, z_N). \tag{2.88}$$

The latter relation is a consequence of equation (2.65).

Relation (2.87) is proved by induction on n . As mentioned in section 2.5, $Z^{(N)}$ is a symmetric polynomial of the z , and by a similar reasoning as above, we also conclude that $Z^{(N)}(z_1, \dots, z_{i-1}, -z_i, z_{i+1}, \dots, z_N) = Z^{(N)}(z_1, \dots, z_{i-1}, z_i, z_{i+1}, \dots, z_N)$, for all

$i = 1, 2, \dots, N$. Moreover, from properties (i)–(ii) of section 2.4, and upon summing over the entries of $\Psi^{(N)}$, we see that $Z^{(N)}$ satisfies for instance the recursion relation

$$Z^{(N)}(z_1 = z_2 - 1, z_2, z_3, \dots, z_N) = Z^{(N-2)}(z_3, z_4, \dots, z_N) \times \prod_{k=3}^N (1 + z_2 + z_k)(1 + z_2 - z_k)(2 + z_k - z_2)(2 - z_k - z_2). \tag{2.89}$$

By the above symmetries, this fixes the value of $Z^{(N)}$ for $z_1 = \pm(\pm z_k - 1), k = 2, 3, \dots, N$, hence a total of $4(n - 1)$ values, which fixes $Z^{(N)}$ as a function of z_1 up to a proportionality constant, as $Z^{(N)}$ is a polynomial of degree $4(n - 1)$ of z_1 . The latter is further fixed by writing the leading behaviour for large z of $Z^{(N)}$:

$$Z^{(N)} \sim \Delta(z_1^2, \dots, z_{2n}^2) \sum_{\pi \in \text{CLP}_{2n}} \frac{(-1)^{c(\pi)}}{\prod_{\substack{\text{pairs } (i < j) \\ \text{connected by } \pi}} (z_i^2 - z_j^2)} = \Delta(z_1^2, \dots, z_{2n}^2) \text{Pf} \left(\frac{1}{(z_i^2 - z_j^2)} \right)_{1 \leq i < j \leq 2n} \tag{2.90}$$

where we have directly identified the Pfaffian, upon interpreting the π as permutations of S_{2n} with only cycles of length 2, and $(-1)^{c(\pi)}$ as the signature of the corresponding permutation. The rhs of (2.87) is clearly a polynomial of the z , symmetric under interchange and sign reversal of the z , of total degree $4n(n - 1)$ and partial degree $4(n - 1)$ in each variable. Moreover, it clearly satisfies the recursion relation (2.89), as when $z_1 \rightarrow z_2 - 1$ the first two lines and columns of the matrix $A_{i,j} = (z_i^2 - z_j^2) / ((1 - (z_i - z_j)^2)(1 - (z_i + z_j)^2))$ are dominated by the terms $A_{1,2} = -A_{2,1}$, henceforth the determinant of A factors into that of A with the first two rows and columns deleted, and the proportionality factor coming from the prefactor in (2.87) matches that in (2.89). Moreover, for large z , the Pfaffian reduces to

$$\prod_{1 \leq i < j \leq N} (z_i^2 - z_j^2) \text{Pf} \left(\frac{1}{z_i^2 - z_j^2} \right)_{1 \leq i < j \leq N} \tag{2.91}$$

and matches exactly the sum over the leading terms of the components of $\Psi^{(N)}$, as given by equation (2.90). This completes the proof of the sum rule (2.87).

Note finally that in the homogeneous limit where all $z \rightarrow 0$, formula (2.87) reduces to

$$Z^{(2n)}(0, 0, \dots, 0) = \text{Pf} \left(\frac{1}{2} \left(\binom{2i + 2j + 1}{2j} - \binom{2i + 2j + 1}{2i} \right) \right)_{0 \leq i < j \leq 2n-1} \tag{2.92}$$

while in the odd $N = 2n - 1$ case, we have

$$Z^{(2n-1)}(0, 0, \dots, 0) = \text{Pf} \left(\frac{1}{2} \left(\binom{2i + 2j + 1}{2j} - \binom{2i + 2j + 1}{2i} \right) \right)_{1 \leq i < j \leq 2n-2}. \tag{2.93}$$

The numbers $Z^{(N)}(0, \dots, 0)$ read

$$1, 7, 39, 1771, 57\ 163, 16\ 457\ 953, 3125\ 503\ 009, 5643\ 044\ 005\ 273, 6357\ 601\ 085\ 989\ 209, \dots \tag{2.94}$$

for $N = 2, 3, 4, \dots, 10, \dots$

3. The inhomogeneous O(1) loop model with open boundaries

We now turn to the open boundary version of the inhomogeneous O(1) (non-crossing) loop model considered in [1–15]. Throughout this section and appendix B, we use the same

notations for transfer matrices, ground-state vectors, fundamental link patterns, etc as in the Brauer case, as there is no ambiguity that from now on we change the subject and deal with a different case. This allows for avoiding many repetitions, as many of the equations of the Brauer case still hold in the non-crossing one.

3.1. Transfer matrix and basic relations


Like in the Brauer case, the model of non-crossing loops was originally defined on a square lattice wrapped on a semi-infinite cylinder of even perimeter, giving rise to periodic boundary conditions. We now consider the same model on a square lattice that covers a semi-infinite strip of width N (even or odd), with centres of the lower edges labelled $1, 2, \dots, N$. We attach probabilities $(t_i, 1 - t_i)$ to the two face loop configurations



$$(3.1)$$


in the column above the edge labelled i . We moreover supplement the picture with the *same* patterns of fixed configurations of loops on the (left and right) boundaries as for the Brauer case, as depicted in equation (2.2). A given configuration now forms a planar pairing of the N labelled points (one of which is connected to infinity if N is odd), via $n = \lfloor N/2 \rfloor$ non-crossing arches. The set of such link patterns is denoted by LP_N , and has cardinality $c_n = \binom{2n}{n} / (n + 1)$ the n th Catalan number, for $N = 2n$ or $N = 2n - 1$. As before, we also consider the c_n -dimensional complex vector space with canonical basis indexed by elements of LP_N .

The transfer matrix for this system is built out of the basic R -matrix that acts on open link patterns or tensor products thereof via



$$(3.2)$$

where as before z and w are the spectral parameters attached to the points labelled i and j , respectively, and we use the same pictorial representation for the matrix elements of R (intersection between two oriented lines carrying the spectral parameters z and w). Alternatively, we have the permuted matrices



$$(3.3)$$

for $i = 1, 2, \dots, N - 1$ acting on the vector space of link patterns, where $e_i, i = 1, \dots, N - 1$ are now the generators of the Temperley–Lieb algebra $TL_N(1)$, subject to the relations

$$e_i^2 = e_i, \quad [e_i, e_j] = 0 \quad \text{if } |i - j| > 1, \quad e_i e_{i \pm 1} e_i = e_i. \quad (3.4)$$

While $I \otimes I$ leaves link patterns invariant, e_i glues the ends labelled i and $i + 1$ of the links and adds up a new link connecting points i and $i + 1$. If a loop is formed in the process, it must simply be erased (loops are given a weight 1 here, leading to the relation $e_i^2 = e_i$). Again, we use the integrable R -matrix, now corresponding to the choice

$$t(z, w) = \frac{qw - z}{qw - z} \quad q = e^{2i\pi/3}. \quad (3.5)$$

Following again Sklyanin [23], we also introduce a boundary operator $K_i(z)$, whose action is diagonal at the points labelled $i = 1$ or N , but whose effect is now to inverse the spectral

parameter $z \rightarrow 1/z$ attached to that point, represented pictorially like in the Brauer case (2.7) (except that $-z$ is now replaced by $1/z$). We still have the standard Yang–Baxter and unitarity relations (with multiplicative spectral parameters) that read pictorially as in (2.8), the boundary Yang–Baxter relation (2.9) (with $-z$ and $-w$ replaced by $1/z$ and $1/w$ respectively), and unitarity boundary relations at the leftmost and rightmost points:

$$K_1(z)K_1\left(\frac{1}{z}\right) = I \quad \text{and} \quad K_N(z)K_N\left(\frac{1}{z}\right) = I. \quad (3.6)$$

The transfer matrix $T(t|z_1, z_2, \dots, z_N)$ of our model reads pictorially exactly the same as in the Brauer case (2.11), only the intersection between two oriented lines carrying spectral parameters now correspond to definition (3.2). It now acts on the vector space of (non-crossing) link patterns with N points. As a consequence of the Yang–Baxter and boundary Yang–Baxter equations, the transfer matrices at two distinct values of t commute.

As before we denote by $\Psi^{(N)}(z_1, z_2, \dots, z_N)$ the common ground-state vector of the T for fixed values of the z_i , satisfying (2.12). As T is a rational fraction of the z_i , we normalize Ψ_N so that all its entries are coprime polynomials of the z_i . We may view the entries of $\Psi^{(N)}$ as relative probabilities of link pattern connections in random loop configurations with inhomogeneous probabilities $(t_i, 1 - t_i)$ in the i th column of the strip, and with $t_i = (qz_i - t)/(qt - z_i)$. This interpretation is again *stricto sensu* only valid in the range of z_i leading to $t_i \in [0, 1]$, in which case $\Psi^{(N)}$ is the Perron–Frobenius eigenvector of T .

The remainder of this note is based on an empirical observation, which we conjecture to be true, that for even $N = 2n$, $\Psi^{(N)}$ defined above is a polynomial of total degree $3n(n - 1)$ and partial degree $2(n - 1)$ in each variable. A similar property was proved in a rather indirect way in [15], involving the details of the Bethe ansatz solution of the corresponding integrable model. In the present case, we believe such a proof should be within reach, although technically tedious, but we will content ourselves with assuming the result. This property was the main difference between the strategies of proof in the crossing and non-crossing periodic boundary loop models of [20, 15], the former appearing as more straightforward, as it does not require any bound on the degree of the ground-state vector.

An illustration is given in appendix B, where the entries of $\Psi^{(N)}$ are listed for the case $N = 4$.

3.2. Intertwining properties

The intertwining relation (2.14) as well as its consequence (2.16) still hold in the non-crossing case, with the appropriate definition (3.2) of the R -matrix. When expressed in components, this translates into

$$\Delta_i \Psi_\pi^{(N)} = \sum_{\substack{\pi' \neq \pi \\ e_i \pi' = \pi}} \Psi_{\pi'}^{(N)} \quad (3.7)$$

where, for $i = 1, 2, \dots, N - 1$, the operator Δ_i acts on functions $f \equiv f(z_1, z_2, \dots, z_N)$ as

$$\Delta_i f = \frac{qz_i - z_{i+1}}{1 + q} \partial_i f \quad (3.8)$$

for $i = 1, 2, \dots, N - 1$, and with ∂_i acting as in (2.20). A first consequence of equation (2.16) is that the entries $\Psi_\pi^{(N)}$ satisfy a suitably modified property.

(P3): If $z_j = qz_i$, and if the link pattern π has no little arch connecting any pair of consecutive points between i and j , then $\Psi_\pi^{(N)}$ vanishes.

If $j = i + 1$, this is easily deduced from relation (2.16), by noting that $\check{R}_{i,i+1}(z_i, qz_i) \propto e_i$. It is easily generalized to more distant points $i < j$ by considering suitable products of R





Figure 3. A non-crossing link pattern (left) and its associated Dyck path (right). The box decomposition of the path is indicated, as well as the corresponding actions of e_i on the fundamental link pattern made of consecutive arches connecting points $2i - 1$ and $2i$.

matrices (see [15] for a detailed proof in the periodic boundary case; the adaptation to the open boundary case is straightforward).

For even $N = 2n$, these equations allow us to determine all the entries of $\Psi^{(N)}$ in terms of that corresponding to the link pattern π_0 with maximally nested arches that connects points i and $2n + 1 - i$. Indeed, we may decompose any link pattern canonically into successive actions of e_i on the ‘lowest’ one, made of n little arches connecting points $2i - 1$ to $2i$. This is best seen in the Dyck path formulation of link patterns, which are represented as paths on a square lattice as shown in figure 3. The path of a given link pattern is defined as follows. We visit the connected points say from left to right, and parallelly draw a path with the rule that if we encounter a new arch, the path goes up one step, and if we encounter an arch already opened earlier, the path goes down one step. The area below the path is then decomposed into square ‘boxes’, each of which corresponds to an action with an operator e_i , whose index is the horizontal coordinate of the box, while the vertical coordinate orders the successive actions. For instance, the decomposition of figure 3 corresponds to acting with $e_3e_2e_4e_6$ on the fundamental link pattern made of five successive little arches. With this formulation, it is easy to write down explicitly the antecedents $\pi' \neq \pi$ of a given link pattern π under the action of e_i . For these to exist, the Dyck path for π must necessarily have a maximum at horizontal position i . One obvious antecedent π'' is obtained by removing the box with this maximum. Others more subtle may arise from adding a whole row of boxes. The important property here is that we may order antecedents by strict inclusion. Indeed, the antecedent with the smallest number of boxes is π'' , and is strictly contained in all others. We may therefore express each new component in a triangular way with respect to strict inclusion. As an example let us treat the case $N = 6$, with five link patterns, explicitly. We label 1, 2, 3, 4, 5 the link patterns and their corresponding Dyck paths



(3.9)

Relations (3.7) allow us to express successively

$$\Psi_2 = \Delta_3 \Psi_1 \quad \Psi_3 = \Delta_4 \Psi_2 - \Psi_1 \quad \Psi_4 = \Delta_2 \Psi_2 - \Psi_1 \quad \Psi_5 = \Delta_2 \Psi_3 = \Delta_4 \Psi_4 \quad (3.10)$$

and we moreover have to write that

$$\begin{aligned} \Delta_5 \Psi_2 &= \Psi_2 & \Delta_1 \Psi_4 &= \Psi_2 & \Delta_1 \Psi_5 &= \Psi_1 + \Psi_3 \\ \Delta_3 \Psi_5 &= \Psi_3 + \Psi_4 & \Delta_5 \Psi_5 &= \Psi_1 + \Psi_4. \end{aligned} \tag{3.11}$$

The compatibility between these equations implies a number of relations to be satisfied by $\Psi_{\pi_0}^{(6)} \equiv \Psi_1$. This construction also applies to the periodic case, upon cutting the link patterns between points N and 1 and opening them. Note finally that Δ_i are degree-preserving operators, hence the total and partial degrees of $\Psi_{\pi_0}^{(N)}$ are shared by all other entries of $\Psi^{(N)}$.

Let us finally mention the non-crossing loop model counterparts of equations (2.23) and (2.24). We still denote by φ_i the embedding of $LP_{2n-2} \rightarrow LP_{2n}$ that acts on a link pattern with $n - 1$ arches by inserting a little arch between points $i - 1$ and i . We have the following restriction/projection property: if two neighbouring parameters z_i and z_{i+1} are such that $z_{i+1} = qz_i$, then

$$T(t|z_1, \dots, z_i, z_{i+1} = qz_i, \dots, z_{2n})\varphi_i = \varphi_i T(t|z_1, \dots, z_{i-1}, z_{i+2}, z_{2n}). \tag{3.12}$$

As in the Brauer case, this was proved in [15] by explicitly commuting φ_i through the product of two R matrices at lines i and $i + 1$, and noting that $\check{R}_{i,i+1}(z_i, qz_i) \propto e_i$.

Similarly, equation (2.24) immediately translates into

$$T(t|z_1, z_2, \dots, z_N)K_1\left(\frac{1}{z_1}\right) = K_1\left(\frac{1}{z_1}\right)T\left(t\left|\frac{1}{z_1}, z_2, \dots, z_N\right.\right) \tag{3.13}$$

with the same pictorial interpretation (2.25) with $-z_1$ replaced by $1/z_1$.

Finally, the reflection invariance of the system leads to the relation

$$(z_1 z_2 \dots z_N)^{2(n-1)} \Psi^{(N)\rho}\left(\frac{1}{z_N}, \frac{1}{z_{N-1}}, \dots, \frac{1}{z_1}\right) = \Psi^{(N)}(z_1, z_2, \dots, z_N) \tag{3.14}$$

also obtained by implementing the condition that $\Psi^{(N)}$ has partial degree $2(n - 1)$ in each variable z_i , and by using the reflection ρ of non-crossing link patterns.

3.3. Solution for $\Psi_{\pi_0}^{(N)}$

In the case of even $N = 2n$, applying condition (i) to the maximally nested pattern π_0 , with arches connecting points i and $2n - i$, we find that $\Psi_{\pi_0}^{(2n)} \equiv \Psi_0^{(2n)}$ must factor out the polynomial $\prod_{1 \leq i < j \leq n} (qz_i - z_j) \prod_{n+1 \leq i < j \leq 2n} (q^2 z_j - z_i)$. Moreover, the symmetry conditions (2.26)–(2.27) imply more vanishing conditions, and henceforth some extra polynomial factor $\prod_{2 \leq j \leq n} (q^2 z_1 z_j - 1)(qz_{n+j-1} z_N - 1)$. This exhausts the partial degree $2(n - 1)$ both in z_1 and z_N of $\Psi_0^{(N)}$, hence we may write for instance

$$\begin{aligned} \Psi_0^{(N)}(z_1, z_2, \dots, z_N) &= \prod_{j=2}^n (qz_1 - z_j)(q^2 z_1 z_j - 1) \\ &\times (q^2 z_N - z_{n+j-1})(qz_{n+j-1} z_N - 1) A^{(N)}(z_2, \dots, z_{N-1}). \end{aligned} \tag{3.15}$$

Now let us take $z_1 \rightarrow 0$. The R -matrix elements involving z_1 in the transfer matrix T reduce respectively to $-q^2 I \otimes I - qe_1$ and $-qI \otimes I - q^2 e_1$, whose product is $I \otimes I$. The net result of taking $z_1 \rightarrow 0$ is therefore to split T into the identity acting at the point labelled 1 and a similar transfer matrix acting on the points labelled $2, 3, \dots, N$. Taking then $z_N \rightarrow 0$ now reduces the transfer matrix to that of size $N - 2$, acting on the points labelled $2, 3, \dots, N - 1$. As a result, $A^{(N)}$ is proportional to the maximally nested entry of the ground-state vector at size $N - 2$, $\Psi_0^{(N-2)}(z_2, \dots, z_{N-1})$, and has the same partial and total degrees. We may now

apply condition (i) again, leading to more factors for $\Psi_0^{(N)}$. Iterating this process, we exhaust all factors and finally reach the total degree $3n(n-1)$, with, for even $N = 2n$:

$$\Psi_0^{(2n)}(z_1, z_2, \dots, z_{2n}) = \prod_{1 \leq i < j \leq n} (qz_i - z_j)(q^2 - z_i z_j) \prod_{n+1 \leq i < j \leq 2n} (q^2 z_j - z_i)(q - z_i z_j). \tag{3.16}$$

As explained before, this determines in turn all components of $\Psi^{(N)}$ to be polynomials of the same total and partial degrees.

As a by-product of the above discussion, the case of odd size $= 2n - 1$ is easily obtained from the case $N = 2n$ by simply taking $z_N \rightarrow 0$. We then have that

$$\Psi^{(N-1)}(z_1, z_2, \dots, z_{N-1}) = \lim_{z_N \rightarrow 0} \Psi^{(N)}(z_1, z_2, \dots, z_N). \tag{3.17}$$

Moreover, going from odd to even size gives

$$\Psi^{(N-2)}(z_1, z_2, \dots, z_{N-2}) = \lim_{z_{N-1} \rightarrow 0} \frac{\Psi^{(N-1)}(z_1, z_2, \dots, z_{N-1})}{z_1 z_2 \cdots z_{N-2}}. \tag{3.18}$$

3.4. Recursion relations

As a consequence of the above intertwining properties, given a link pattern $\pi \in LP_{2n}$, two situations may occur for a pair $(i, i + 1)$ of consecutive points:

- (i) the pattern π has no arch joining i to $i + 1$, in which case property (P1) implies that

$$\Psi_\pi^{(N)}(z_1, \dots, z_N)|_{z_{i+1}=qz_i} = 0; \tag{3.19}$$

- (ii) the pattern π has a little arch joining i to $i + 1$, in which case

$$\begin{aligned} \Psi_\pi^{(N)}(z_1, \dots, z_N)|_{z_{i+1}=qz_i} &= \Psi_{\pi'}^{(N-2)}(z_1, \dots, z_{i-1}, z_{i+2}, \dots, z_N) \\ &\times \prod_{\substack{k=1 \\ k \neq i, i+1}}^N (q^2 z_i - z_k)(q^2 z_i z_k - 1) \end{aligned} \tag{3.20}$$

where π' is the link pattern π with the little arch $i, i + 1$ removed ($\pi = \varphi_i \pi', \pi' \in LP_{2n-2}$). This is readily obtained by applying equation (3.12) to the vector $\Psi^{(N)}$ at $z_{i+1} = qz_i$. As a result, the restricted $\Psi^{(N)}$ must be proportional to the projected $\Psi^{(N-2)}$, and the proportionality factor is fixed by the value of $\Psi_0^{(N)}$ (3.16).

3.5. Symmetries

As before, the intertwining properties of previous section lead straightforwardly to symmetry properties for $\Psi^{(N)}$. Indeed, applying relation (3.12) on the vector $\Psi^{(N)}(\frac{1}{z_1}, z_2, \dots, z_N)$, we find that the latter must be proportional to $\Psi^{(N)}(z_1, z_2, \dots, z_N)$, with the result

$$z_1^{4(n-1)} \Psi^{(N)}\left(\frac{1}{z_1}, z_2, \dots, z_N\right) = \Psi^{(N)}(z_1, z_2, \dots, z_N). \tag{3.21}$$

The same reasoning at the other end with the space labelled N leads to the condition

$$z_N^{4(n-1)} \Psi^{(N)}\left(z_1, z_2, \dots, z_{N-1}, \frac{1}{z_N}\right) = \Psi^{(N)}(z_1, z_2, \dots, z_N). \tag{3.22}$$

In both equations, we have used the assumed fact that $\Psi^{(N)}$ has partial degree $2(n - 1)$ in each variable z_i .

As before, let us consider the sum over all components of $\Psi^{(N)}$, namely

$$Z^{(N)}(z_1, \dots, z_N) = \sum_{\pi \in LP_N} \Psi_{\pi}^{(N)}(z_1, z_2, \dots, z_N) = v_N \cdot \Psi^{(N)} \quad (3.23)$$

where v_N is the vector with all entries equal to 1. This vector also satisfies

$$v_N e_i = v_N \quad \text{and} \quad v_N \check{R}_{i,i+1}(z_i, z_{i+1}) = v_N \quad (3.24)$$

hence applying v_N to equation (2.16), we immediately get that $\tau_i Z^{(N)} = Z^{(N)}$, $i = 1, 2, \dots, N-1$, henceforth Z is symmetric in the z_i . Moreover, applying v_N to equation (3.21), we get

$$z_1^{4(n-1)} Z^{(N)}\left(\frac{1}{z_1}, z_2, \dots, z_N\right) = Z^{(N)}(z_1, z_2, \dots, z_N) \quad (3.25)$$

hence $Z^{(N)}$ is symmetric and reciprocal in each of the z .

3.6. Sum rule

For even $N = 2n$, we have

$$\begin{aligned} Z^{(N)}(z_1, \dots, z_{2n}) &= \frac{\prod_{i,j=1}^n (z_i^2 + z_i z_{j+n} + z_{j+n}^2) (1 + z_i z_{j+n} + z_i^2 z_{j+n}^2)}{\prod_{1 \leq i < j \leq n} (z_i - z_j) (1 - z_i z_j) (z_{i+n} - z_{j+n}) (1 - z_{i+n} z_{j+n})} \\ &\times \det \left(\frac{1}{z_i^2 + z_i z_{j+n} + z_{j+n}^2} \frac{1}{1 + z_i z_{j+n} + z_i^2 z_{j+n}^2} \right)_{1 \leq i, j \leq n}. \end{aligned} \quad (3.26)$$

Remarkably, this coincides with the partition function $Z_{UASM}(z_1, \dots, z_n; z_{n+1}, \dots, z_{2n})$ introduced in [22]. The proof of equation (3.26) parallels exactly that of equation (2.87), proceeding by induction on n , and makes use of the recursion relations (3.20), as well as of the symmetries of $Z^{(N)}$.

We also have

$$\begin{aligned} Z^{(N)}(z_1, \dots, z_{2n})^2 &= \prod_{1 \leq i < j \leq 2n} \frac{z_i^2 + z_i z_j + z_j^2}{z_i - z_j} \frac{1 + z_i z_j + z_i^2 z_j^2}{1 - z_i z_j} \\ &\times \text{Pf} \left(\frac{z_i - z_j}{z_i^2 + z_i z_j + z_j^2} \frac{1 - z_i z_j}{1 + z_i z_j + z_i^2 z_j^2} \right)_{1 \leq i, j \leq N} \end{aligned} \quad (3.27)$$

also proved by induction on n . The latter expression has the advantage of being explicitly symmetric in the z .

In the homogeneous limit where all z_i tend to 1, we find that for $N = 2n$, $Z^{(2n)}(1, 1, \dots, 1)$ is $3^{n(n-1)}$ times the total number of U-symmetric ASMs of size $(2n) \times (2n)$ discussed in [22], itself identical to that of vertically symmetric ASMs of size $(2n+1) \times (2n+1)$.

4. Conclusions

In this paper, we have derived sum rules for the ground-state vector of the inhomogeneous crossing and non-crossing $O(1)$ loop models on a semi-infinite strip. As opposed to the crossing case where the result is rigorous and proved completely, we have made in the non-crossing case a reasonable but crucial assumption on the degree of the ground-state vector as a polynomial of the inhomogeneities z_i . The completion of the latter proof would presumably involve invoking the algebraic Bethe ansatz solution of the XXZ spin chain with open boundaries, in

much the same spirit as in [15]. We have rather chosen here to concentrate on the various properties of this ground-state vector, for which we gave an explicit step-by-step construction by acting on a fundamental component with local divided difference operators, in order to generate all other entries of the vector. In this respect, it might be possible to unify both crossing and non-crossing cases by deriving a proof uniquely based on the main relations induced by equation (2.16), and that only involve the interplay between the symmetric group action on spectral parameters and the R -matrix of the integrable system. In order to do this, and by analogy with the crossing case, one should be able to check that the solution of the non-crossing case (3.16) actually satisfies all constraints inherited from the compatibility of all equations (3.7), as well as the boundary reflection properties (3.21)–(3.22). By a uniqueness argument, this would by-pass our approach, which assumes the value of the degree of $\Psi^{(N)}$. This is a problem for future work. Note finally that such a construction, both in the crossing and non-crossing cases, should be instrumental in trying to prove variants of the so-called Razumov–Stroganov conjectures [3]. It would also be interesting to extend the present work to other types of open boundary conditions, such as those considered in [24], and conjectured in the non-crossing case to be related to other symmetry classes of alternating sign matrices.

Our approach allows us, in particular, to compute the entries of the ground-state vector in the homogeneous limit, where it may be identified with the ground-state vector of the Hamiltonian of a suitable quantum chain, expressed as a particular weighted sum of generators of the Brauer (resp. Temperley–Lieb) algebra for the crossing (resp. non-crossing) case, acting on crossing (resp. non-crossing) link patterns (see [18, 3] for explicit expressions). As an outcome of our calculation, we show that these entries may be picked to be non-negative integers, summing to specific numbers as given by (2.92) and (2.93) for strips of even/odd size in the crossing case, and to the total number of vertically symmetric alternating sign matrices $A_V(2n+1)$ of [22] for strips of even size $N = 2n$. It is important to note that, as opposed to the standard case where the entries of the homogeneous ground-state vector are normalized so that the smallest one is 1 after division by their GCD (cf [15, 20]), the smallest entries in the open crossing case are not 1, but form themselves a quite intriguing sequence (2.67), as derived from the homogeneous limit of relation (2.40).

While the numbers $A_V(2n+1)$ have been given extensive combinatorial interpretations, that of numbers (2.92) and (2.93) is still elusive. Such an interpretation was suggested in [18] for the cylinder case, by noting and conjecturing that some entries of the homogeneous ground-state vector of the crossing loop model with periodic boundaries actually matched degrees of varieties related to the commuting variety [19]. This was further proved in [20] and extended in [21], where all the components of the ground-state vector were interpreted as the multidegrees of the components of a matrix variety. It is natural to hope that numbers (2.92) and (2.93) actually count the total degrees of some matrix varieties, still to be found. In this respect, the partial sum rule $W^{(N)}$ in the permutation sector (2.83) leading to the numbers (2.86) seems to indicate, like in the periodic case, that the corresponding components of the (yet unknown) matrix variety form a complete intersection, whose multidegree has the factorized form (2.83). We could also hope that the total multidegree, as given by equation (2.87), may alternatively be obtained like in [21] as the result of a ‘volume’ matrix integral over the putative matrix variety.

Acknowledgments

We acknowledge many interesting and stimulating discussions with P Zinn-Justin and J-B Zuber. This research was partly supported by the European network ENIGMA, grant MRTN-CT-2004-5652 and of the GEOCOMP project (ACI ‘Masse de données’).

Appendix A. Entries of the ground-state vector in the dense O(1) crossing loop model with open boundaries

In this appendix, we give the entries of $\Psi^{(N)}$ and their sum $Z^{(N)}(z_1, z_2, z_3, z_4)$ in the Brauer loop case for $N = 4$, as well as the entry $\Psi_{\pi_0}^{(N)}$ for $N = 6$. In the notations of section 2.3, and for $N = 4$, the three components of $\Psi^{(4)}$ read

$$\begin{aligned}
 \Psi^{(4)} &= a_{1,2}b_{1,2}a_{2,3}b_{3,4}c_{3,4}(5 + 3z_2 - 3z_3 - 2z_2z_3 - z_1^2 - z_4^2 + (z_1^2 - z_4^2)(z_2 + z_3)) \\
 \Psi^{(4)} &= a_{1,2}b_{1,2}a_{3,4}c_{3,4}(11 - 3z_1^2 + 8z_2 + z_2^2 - 8z_3 + 2z_1^2z_3 - 8z_2z_3 - 2z_2^2z_3 \\
 &\quad + z_3^2 + 2z_2z_3^2 - 3z_4^2 - 2z_2z_4^2 - (z_1^2 - z_3^2)(z_2^2 - z_4^2)) \\
 \Psi^{(4)} &= a_{2,3}(23 - 10z_1^2 + 3z_1^4 - 7z_2 + 18z_1^2z_2 - 3z_1^4z_2 - 11z_2^2 + 4z_1^2z_2^2 - z_1^4z_2^2 + 3z_3^2 \\
 &\quad - 4z_1^2z_3^2 + 7z_3 + 10z_1^2z_3 - z_1^4z_3 + 2z_2z_3 + 16z_1^2z_2z_3 - 2z_1^4z_2z_3 - z_2^2z_3 \\
 &\quad - 4z_1^2z_2^2z_3 - 4z_2^3z_3 - 11z_3^2 + 9z_1^2z_3^2 - 2z_1^4z_3^2 + z_2z_3^2 + z_1^2z_2z_3^2 + 3z_2^2z_3^2 \\
 &\quad + z_1^2z_2^2z_3^2 - z_2^3z_3^2 - 3z_3^3 - z_1^2z_3^3 - 4z_2z_3^3 - 2z_1^2z_2z_3^3 + z_2^2z_3^3 + 2z_2^3z_3^3 \\
 &\quad - 10z_4^2 - 11z_1^2z_4^2 + z_1^4z_4^2 - 10z_2z_4^2 + z_1^2z_2z_4^2 + 9z_2^2z_4^2 + z_1^2z_2^2z_4^2 + z_3^2z_4^2 \\
 &\quad - 18z_3z_4^2 - z_1^2z_3z_4^2 + 16z_2z_3z_4^2 + 6z_1^2z_2z_3z_4^2 - z_2^2z_3z_4^2 - 2z_2^3z_3z_4^2 + 4z_3^2z_4^2 \\
 &\quad + z_1^2z_3^2z_4^2 + 4z_2z_3^2z_4^2 + z_2^2z_3^2z_4^2 + 4z_3^3z_4^2 + 3z_4^4 + z_1^2z_4^4 + z_2z_4^4 - 2z_2^2z_4^4 + 3z_3z_4^4 \\
 &\quad - 2z_2z_3z_4^4 - z_3^2z_4^4 + (z_2 + z_3)(z_1^2 - z_3^2)(z_1^2 - z_4^2)(z_2^2 - z_4^2)). \tag{A.1}
 \end{aligned}$$

These may be obtained by explicitly solving the eigenvector equation (2.12). Alternatively, we have computed $\Psi_0^{(4)}$ in the text using (2.38)–(2.66), with the result $\Psi_0^{(4)} = a_{1,2}b_{1,2}a_{2,3}b_{3,4}c_{3,4}P_0^{(4)}$. The other components read simply

$$\begin{aligned}
 \Psi^{(4)} &= \Theta_1 \Psi^{(4)} \\
 \Psi^{(4)} &= \Theta_2 \Psi^{(4)}. \tag{A.2}
 \end{aligned}$$

The components (A.1) sum to

$$\begin{aligned}
 Z^{(4)}(z_1, z_2, z_3, z_4) &= 39 - 30z_1^2 + 7z_1^4 - 30z_2^2 + 14z_1^2z_2^2 - 4z_1^4z_2^2 + 7z_2^4 - 4z_1^2z_2^4 \\
 &\quad + z_1^4z_2^4 - 30z_3^2 + 14z_1^2z_3^2 - 4z_1^4z_3^2 + 14z_2^2z_3^2 + 12z_1^2z_2^2z_3^2 - z_1^4z_2^2z_3^2 \\
 &\quad - 4z_2^4z_3^2 - z_1^2z_2^4z_3^2 + 7z_3^4 - 4z_1^2z_3^4 + z_1^4z_3^4 - 4z_2^2z_3^4 - z_1^2z_2^2z_3^4 + z_2^4z_3^4 - 30z_4^2 \\
 &\quad + 14z_1^2z_4^2 - 4z_1^4z_4^2 + 14z_2^2z_4^2 + 12z_1^2z_2^2z_4^2 - z_1^4z_2^2z_4^2 - 4z_2^4z_4^2 - z_1^2z_2^4z_4^2 + 14z_3^2z_4^2 \\
 &\quad + 12z_1^2z_3^2z_4^2 - z_1^4z_3^2z_4^2 + 12z_2^2z_3^2z_4^2 + 6z_1^2z_2^2z_3^2z_4^2 - z_2^4z_3^2z_4^2 - 4z_3^4z_4^2 \\
 &\quad - z_1^2z_3^4z_4^2 - z_2^2z_3^4z_4^2 + 7z_4^4 - 4z_1^2z_4^4 + z_1^4z_4^4 - 4z_2^2z_4^4 - z_1^2z_2^2z_4^4 + z_2^4z_4^4 \\
 &\quad - 4z_3^2z_4^4 - z_1^2z_3^2z_4^4 - z_2^2z_3^2z_4^4 + z_3^4z_4^4. \tag{A.3}
 \end{aligned}$$

It takes only a few seconds for any formal manipulation software to check that this quantity indeed coincides with the Pfaffian expression of equation (2.87).

We also display the value of the component $\Psi_0^{(6)}$ for $N = 6$, as obtained from formula (2.40) and with extensive use of the modified Leibniz formula (2.44):

$$\begin{aligned}
 \Psi^{(6)} &= a_{1,2}b_{1,2}a_{1,3}b_{1,3}a_{2,3}b_{2,3}a_{2,4}a_{3,4}a_{3,5}a_{4,5}a_{4,6}a_{5,6}c_{5,6} \\
 &\quad \times (a_{1,6}b_{1,6}a_{2,6}b_{2,6}b_{3,4}b_{3,5}(a_{1,5}b_{1,5}b_{2,4} + 2c_{2,5}(1 + b_{5,4})) \\
 &\quad + 4a_{1,6}a_{2,6}b_{3,4}c_{3,6}(a_{1,5}b_{2,4}c_{2,6} + 2b_{1,6}(1 + b_{5,4}))
 \end{aligned}$$

$$\begin{aligned}
 &+ 4c_{3,5}c_{3,6}(1 + b_{4,6})(1 + b_{5,6})(a_{1,5}b_{1,6}b_{2,4} + 2c_{2,6}(1 + b_{5,4})) \\
 &+ 2c_{3,5}c_{3,6}(1 + b_{4,6})(a_{1,5}a_{2,6}b_{1,5}b_{2,4}c_{2,6} + 2a_{1,5}b_{1,5}b_{1,6}(a_{1,6} - b_{2,4})) \\
 &+ 2a_{1,6}b_{1,4}b_{1,6}(b_{2,5} - a_{1,5}) + 2c_{3,6}b_{3,4}(1 + b_{5,6})(a_{1,5}a_{1,6}b_{1,5}b_{1,6}b_{2,4} \\
 &+ 2b_{1,4}a_{2,6}c_{2,6}(b_{2,5} - a_{1,5}) + 2a_{1,5}b_{1,5}c_{2,6}(1 + b_{4,6})). \tag{A.4}
 \end{aligned}$$

As all a, b, c tend to 1 in the homogeneous limit where all $z \rightarrow 0$, we read off (A.4) that $\Psi_0^{(6)}(0, 0, 0, 0, 0, 0) = 129$. As expected for the case of arbitrary even N , the result for $\Psi_0^{(6)}$ is an integer linear combination of products of a, b, c , with coefficients \pm powers of 2.

Appendix B. Entries of the ground-state vector in the dense O(1) loop model with open boundaries

In this appendix, we give the entries of $\Psi^{(N)}$ and their sum $Z^{(N)}$ in the non-crossing loop case for $N = 4$. In the notations of section 3, and for $N = 4$, the two components of $\Psi^{(4)}$ read

$$\begin{aligned}
 \Psi^{(4)} \text{ (top diagram)} &= (qz_1 - z_2)(q^2z_1z_2 - 1)(q^2z_3 - z_4)(qz_3z_4 - 1) \\
 \Psi^{(4)} \text{ (bottom diagram)} &= (qz_2 - z_3)((q^2z_1 - z_4)(1 + z_1z_2z_3z_4) + (q^2z_4 - z_1)(z_1z_4 + z_2z_3) \\
 &\quad + (q^2 - 1)z_1z_4(z_2 + z_3)). \tag{B.1}
 \end{aligned}$$

These components sum to

$$\begin{aligned}
 Z^{(4)}(z_1, z_2, z_3, z_4) &= (1 + z_1z_2z_3z_4)(z_1z_2 + z_1z_3 + z_2z_3 + z_1z_4 + z_2z_4 + z_3z_4) \\
 &\quad + (z_1 + z_2)(z_3 + z_4)(z_1z_2 + z_3z_4) + (z_1z_3 + z_2z_4)(z_1z_4 + z_2z_3) + 3z_1z_2z_3z_4 \tag{B.2}
 \end{aligned}$$

which may easily be checked against equations (3.26) and (3.27).

References

[1] Batchelor M T, de Gier J and Nienhuis B 2001 The quantum symmetric XXZ chain at $\Delta = -1/2$, alternating sign matrices and plane partitions *J. Phys. A: Math. Gen.* **34** L265–L270 (Preprint cond-mat/0101385)

[2] Razumov A V and Stroganov Yu G 2001 Spin chains and combinatorics *J. Phys. A: Math. Gen.* **34** 3185 (Preprint cond-mat/0012141)

Razumov A V and Stroganov Yu G 2001 Spin chains and combinatorics: twisted boundary conditions *J. Phys. A: Math. Gen.* **34** 5335–40 (Preprint cond-mat/0102247)

[3] Razumov A V and Stroganov Yu G 2004 Combinatorial nature of ground state vector of O(1) loop model *Theor. Math. Phys.* **138** 333–7 (Preprint math.CO/0104216)

Razumov A V and Stroganov Yu G 2005 O(1) loop model with different boundary conditions and symmetry classes of alternating sign matrices *Theor. Math. Fiz.* **142** 273–43 (Preprint cond-mat/0108103)

[4] Pearce P A, Rittenberg V, de Gier J and Nienhuis B 2002 Temperley–Lieb stochastic processes *J. Phys. A: Math. Gen.* **35** L661–L668 (Preprint math-ph/0209017)

[5] de Gier J 2002 Loops, matchings and alternating-sign matrices *Preprint math.CO/0211285*

[6] Mitra S and Nienhuis B 2003 Osculating random walks on cylinders *Discrete Random Walks DRW'03* ed C Banderier and C Krattenthaler *Discrete Mathematics and Computer Science Proc. AC* pp 259–64 (Preprint math-ph/0312036)

[7] Mitra S, Nienhuis B, de Gier J and Batchelor M T 2004 Exact expressions for correlations in the ground state of the dense O(1) loop model *J. Stat. Mech.: Theor. Exp.* P09010 (Preprint cond-mat/0401245)

[8] Zuber J-B 2004 On the counting of fully packed loop configurations: some new conjectures *Electron. J. Combin.* **11** R13 (Preprint math-ph/0309057)

[9] Di Francesco P, Zinn-Justin P and Zuber J-B 2004 A bijection between classes of fully packed loops and plane partitions *Electron. J. Combin.* **11** R64 (Preprint math.CO/0311220)

- [10] Di Francesco P and Zuber J-B 2004 On FPL configurations with four sets of nested arches *J. Stat. Mech.: Theor. Exp.* P06005 (*Preprint cond-mat/0403268*)
- [11] Caselli F and Krattenthaler C 2004 Proof of two conjectures of Zuber on fully packed loop configurations *J. Combin. Theory A* **108** 123–46 (*Preprint math.CO/0312217*)
- [12] Caselli F, Krattenthaler C, Lass B and Nadeau P 2004 On the number of fully packed loop configurations with a fixed associated matching *Electron. J. Combin.* **11** R16 (*Preprint math.CO/0502392*)
- [13] Di Francesco P 2004 A refined Razumov–Stroganov conjecture *J. Stat. Mech.: Theor. Exp.* P08009 (*Preprint cond-mat/0407477*)
- [14] Di Francesco P 2004 A refined Razumov–Stroganov conjecture: II *J. Stat. Mech.: Theor. Exp.* P11004 (*Preprint cond-mat/0409576*)
- [15] Di Francesco P and Zinn-Justin P 2004 Around the Razumov–Stroganov conjecture: proof of a multi-parameter sum rule *Preprint math-ph/0410061*
- [16] Izergin A 1987 Partition function of the six-vertex model in a finite volume *Sov. Phys. Dokl.* **32** 878–9
- [17] Korepin V 1982 Calculation of norms of Bethe wave functions *Commun. Math. Phys.* **86** 391–418
- [18] De Gier J and Nienhuis B 2004 Brauer loops and the commuting variety *Preprint math.AG/0410392*
- [19] Knutson A 2003 Some schemes related to the commuting variety *Preprint math.AG/0306275*
- [20] Di Francesco P and Zinn-Justin P 2004 Inhomogeneous model of crossing loops and multidegree of some algebraic varieties *Preprint math-ph/0412031*
- [21] Knutson A and Zinn-Justin P 2005 A scheme related to the Brauer loop model *Preprint math.AG/0503224*
- [22] Kuperberg G 2002 Symmetry classes of alternating sign matrices under one roof *Ann. Math.* **156** 835–66 (*Preprint math.CO/0008184*)
- [23] Sklyanin E 1988 Boundary conditions for integrable quantum systems *J. Phys. A: Math. Gen.* **21** 2375–89
- [24] De Gier J and Rittenberg V 2004 Refined Razumov–Stroganov conjectures for open boundaries *J. Stat. Mech.: Theor. Exp.* P09009 (*Preprint math-ph/0408042*)