

High-precision numerical determination of eigenvalues for a double-well potential related to the Zinn-Justin conjecture

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2005 J. Phys. A: Math. Gen. 38 6785

(<http://iopscience.iop.org/0305-4470/38/30/012>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.92

The article was downloaded on 03/06/2010 at 03:52

Please note that [terms and conditions apply](#).

High-precision numerical determination of eigenvalues for a double-well potential related to the Zinn-Justin conjecture

H A Alhendi and E I Lashin¹

Department of Physics and Astronomy, College of Science, King Saud University, Riyadh, Saudi Arabia

E-mail: e_lashin@hotmail.com

Received 29 September 2004

Published 13 July 2005

Online at stacks.iop.org/JPhysA/38/6785

Abstract

A numerical method of high precision is used to calculate the energy eigenvalues and eigenfunctions for a symmetric double-well potential. The method is based on enclosing the system within two infinite walls with a large but finite separation and developing a power series solution for the Schrödinger equation. The obtained numerical results are compared with those obtained on the basis of the Zinn-Justin conjecture and found to be in excellent agreement.

PACS numbers: 03.65.Ge, 02.30.Hq

1. Introduction

Quantum mechanical tunnelling through finite barriers is a well-established phenomenon in theory and application. The symmetric double-well potential is one of the many examples exhibiting this phenomenon. In this case, the energy splitting generated by tunnelling can be estimated with the help of the well-known semi-classical WKB approximation and instanton techniques (see for example [1]). However, to calculate this splitting accurately, one needs an effective method of high precision.

In a series of papers, Zinn-Justin [2] developed a conjecture (to be termed ‘the Zinn-Justin conjecture’) to determine the energy levels of a quantum Hamiltonian H , in cases where the potential has degenerate minima. This conjecture takes the form of the generalized Bohr–Sommerfeld quantization formulae. It has been applied, among other potentials, to the case of the symmetric double well. In this case, the Hamiltonian is

$$H = -\frac{g}{2} \frac{\partial^2}{\partial q^2} + \frac{1}{g} V(q), \quad \text{where } V(q) = \frac{1}{2} q^2 (1 - q)^2. \quad (1)$$

¹ Permanent address: Department of Physics, Faculty of Science, Ain Shams University, Cairo, Egypt.

It is obvious that this Hamiltonian is invariant under the transformation ($q \rightarrow 1 - q$). The energy eigenvalues for this potential can be obtained by finding a solution to the Zinn-Justin conjecture equation:

$$\frac{1}{\sqrt{2\pi}} \Gamma\left(\frac{1}{2} - D(E, g)\right) \left(-\frac{2}{g}\right)^{D(E, g)} \exp[-A(E, g)/2] = \pm i. \quad (2)$$

The function $D(E, g)$ has a perturbative expansion in powers of g , of which the first few terms are

$$D(E, g) = E + g\left(3E^2 + \frac{1}{4}\right) + g^2\left(35E^3 + \frac{25}{4}E\right) + O(g^3). \quad (3)$$

The other function $A(E, g)$ receives contributions from the instanton expansion in the path integral and its first few terms are

$$A(E, g) = \frac{1}{3g} + g\left(17E^2 + \frac{19}{12}\right) + g^2\left(227E^3 + \frac{187}{4}E\right) + O(g^3). \quad (4)$$

The energy $E_{N, \pm}$ can be extracted from equation (2) by expanding in powers of g and in the two quantities

$$\lambda(g) = \ln\left(-\frac{2}{g}\right) \quad \text{and} \quad \xi(g) = \frac{\exp[-1/(6g)]}{\sqrt{\pi g}}. \quad (5)$$

The complete semi-classical expansion of $E_{N, \pm}$ has the form [4]

$$E_{\pm, N}(g) = \sum_{l=0}^{\infty} E_{N, l}^{(0)} g^l + \sum_{n=1}^{\infty} \left(\frac{2}{g}\right)^{Nn} \left(\mp \frac{e^{-1/6g}}{\sqrt{\pi g}}\right)^n \sum_{k=0}^{n-1} (\ln(-2/g))^k \sum_{l=0}^{\infty} \epsilon_{nkl}^{(N, \pm)} g^l. \quad (6)$$

The coefficients ϵ relevant to the numerical calculation have been explicitly calculated in [3]. The number N is the unperturbed quantum number which corresponds to

$$E_{\pm, N}(g) = N + 1/2 + O(g). \quad (7)$$

A detailed exposition of the above equations can be found in [4].

In [3], numerical calculations have been carried out and led to the energy eigenvalues for the ground and first excited states, respectively, for $g = 0.001$,

$$\begin{aligned} E_{0,+}(0.001) = & 0.498\ 995\ 454\ 862\ 109\ 171\ 689\ 130\ 839\ 481\ 921\ 636\ 820\ 947\ 240\ 208 \\ & 096\ 653\ 293\ 278\ 697\ 220\ 139\ 115\ 135\ 285\ 053\ 829\ 445\ 798\ 457\ 599\ 599 \\ & \underline{906\ 739\ 551\ 758\ 472\ 267\ 802\ 813\ 069\ 690\ 601\ 325\ 259\ 437\ 728\ 994\ 365} \\ & \underline{882\ 552\ 444\ 017\ 437\ 127\ 892\ 797\ 899\ 793}, \end{aligned} \quad (8)$$

$$\begin{aligned} E_{0,-}(0.001) = & 0.498\ 995\ 454\ 862\ 109\ 171\ 689\ 130\ 839\ 481\ 921\ 636\ 820\ 947\ 240\ 208 \\ & 096\ 653\ 293\ 278\ 697\ 220\ 139\ 129\ 839\ 929\ 595\ 580\ 370\ 812\ 277\ 499\ 244 \\ & \underline{848\ 259\ 367\ 436\ 475\ 768\ 328\ 848\ 353\ 551\ 134\ 663\ 063\ 098\ 233\ 151\ 885} \\ & \underline{233\ 080\ 862\ 284\ 780\ 527\ 221\ 010\ 367\ 282}. \end{aligned} \quad (9)$$

The above numerical results have been obtained by lattice extrapolation using a modified Richardson algorithm [3].

This tiny difference encourages us to seek for an independent but simple and direct method, which allows us to obtain the energy eigenvalues for the potential in equation (1) and compare them with the above numerical results. In addition, the present method allows us to obtain an accurate description for the corresponding wavefunctions. This method has been

previously applied to various potential functions with and without degenerate minima, leading to results with high accuracy [5].

The method, as will be described in the next two sections, is based on power series solution of the Schrödinger equation in a finite range. It has appeared from time to time in the literature [6–8], but has not been developed to its maximum efficiency. We shall show that, by using the computer algebra systems (for example, Mathematica) which can deal with exact numbers, the accuracy of the method can be substantially improved.

In the following section, for illustrative purpose, we explain our method by applying it to the well-known exactly solvable harmonic oscillator potential and then extend it to the symmetric double well.

2. Calculations and results

In this section we, first, consider the well-known exactly solvable harmonic oscillator. In this case, the Schrödinger equation reads ($\hbar = 1, m = 1$)

$$\left[-\frac{1}{2} \frac{d^2}{dq^2} + E - V(q) \right] \Psi(q) = 0, \quad (10)$$

where

$$V(q) = \frac{1}{2}q^2. \quad (11)$$

The exact energy eigenvalues and the corresponding eigenfunctions are

$$E_N = \left(N + \frac{1}{2} \right), \quad N = 0, 1, 2, \dots, \quad (12)$$

$$\Psi_N(q) = 2^{-\frac{N}{2}} (N!)^{-\frac{1}{2}} \pi^{-\frac{1}{4}} \exp\left(-\frac{q^2}{2}\right) H_N(q),$$

where $H_N(q)$ are the Hermite polynomials.

For the harmonic oscillator confined between two infinite walls at $q = \pm L$, we develop a power series solution in the form

$$\Psi(q) = \sum_{i=0}^{\infty} a_i q^i. \quad (13)$$

Substituting in equation (10), one gets the following recursion relation:

$$a_i = \frac{a_{i-4} - 2E a_{i-2}}{i(i-1)}, \quad i \neq 0, 1 \quad \text{and} \quad a_i = 0 \quad \text{when} \quad i < 0. \quad (14)$$

The symmetry of the potential implies that we have two types of solutions: the even solutions obtained by imposing (ignoring normalization) $a_0 = 1, a_1 = 0$ and the odd ones by imposing $a_0 = 0, a_1 = 1$. The energy eigenvalues are then obtained from the condition $\Psi(E, L) = 0$ for both cases.

For numerical calculations, we approximate the power series in equation (13) with a truncated one having a finite number of terms $\Psi_I(E, q)$, where I is the number of non-vanishing terms. The boundary condition for a specific value of L corresponds to $\Psi_I(E, L) = 0$. To get the zeros of $\Psi_I(E, L)$ with respect to E , we first plot a graph for $\Psi_I(L, E)$ as a function of E to locate where $\Psi_I(L, E)$ changes sign. We then can use two nearby points containing one single root as the initial iteration for the ‘bisection method’ to find the zeros. In doing this we have used Mathematica package version 3 and also have relied extensively on its ability




Figure 2. The normalized ground (left) and first excited (right) state wavefunctions for the bounded double-well potential with $g = 1/1000$.

in equation (16), one gets the following recurrence formula for the expansion coefficients, a_i :

$$a_i = \left(\frac{2}{g}\right) \frac{\frac{1}{2g} [a_{i-6} - \frac{1}{2}a_{i-4} + \frac{1}{16}a_{i-2}] - Ea_{i-2}}{i(i-1)}, \quad i \neq 0, 1 \quad \text{and} \quad a_i = 0 \quad \text{when} \quad i < 0. \tag{18}$$

For $L = 3$, the obtained eigenvalues are

$$E_{0,+}(0.001) = 0.498\ 995\ 454\ 862\ 109\ 171\ 689\ 130\ 839\ 481\ 921\ 636\ 820\ 947\ 240\ 208 \\ 096\ 653\ 293\ 278\ 697\ 220\ 139\ 115\ 135\ 285\ 053\ 829\ 445\ 798\ 457\ 599\ 599 \\ 906\ 739\ 551\ 758\ 472\ 267\ 802\ 813\ 069\ 690\ 601\ 325\ 259\ 437\ 728\ 994\ 365 \\ 882\ 552\ 444\ 017\ 437\ 127\ 892\ 797\ 899\ 793\ 989\ 220\ 053\ 606\ 978\ 041 \\ 386\ 525\ 573\ 028\ 377\ 235\ 024\ 167\ 171, \tag{19}$$

$$E_{0,-}(0.001) = 0.498\ 995\ 454\ 862\ 109\ 171\ 689\ 130\ 839\ 481\ 921\ 636\ 820\ 947\ 240\ 208 \\ 096\ 653\ 293\ 278\ 697\ 220\ 139\ 129\ 839\ 929\ 595\ 580\ 370\ 812\ 277\ 499\ 244 \\ 848\ 259\ 367\ 436\ 475\ 768\ 328\ 848\ 353\ 551\ 134\ 663\ 063\ 098\ 233\ 151\ 885 \\ 233\ 080\ 862\ 284\ 780\ 527\ 221\ 010\ 367\ 282\ 720\ 476\ 134\ 001\ 672\ 248 \\ 036\ 552\ 352\ 410\ 137\ 981\ 630\ 458\ 360. \tag{20}$$

These values agree with the ones obtained from the numerical calculations based on the Zinn-Justin conjecture. In figure 2 we present the ground and first excited state wavefunctions for the bounded double-well potential for $g = 1/1000$, $I = 4600$ and $L = 1$.

3. Discussion

It is important to note the following generic remarks. First, a reason for the capability of the present method is that for a bound state, the wavefunction is spatially localized which means that the probability density ($|\Psi|^2$) has appreciable values in a finite region of space behind which the probability density tends rapidly to zero. Thus, to a good approximation, it is, therefore, reasonable to consider the corresponding problem in a finite interval, with a suitable width, bounded by two infinite walls. The criteria for a suitable value of L can be quantitatively given by the condition $E \ll V(L)$. Second, from the WKB approximation, it can be made plausible that the zeros of $\Psi(E, L)$ provide upper bounds for the energy eigenvalues while the zeros of the derivative—with respect to q — $\Psi'(E', L)$ provide the lower ones; the same

finding can be proved in a rigorous way as shown in [9]. Thus, by matching the digits of the two zeros, one can get an accurate energy eigenvalue up to the number of coincident digits. As an example for the ground state of the bounded harmonic oscillator, with $L = 8$ and $I = 250$, one gets

$$\begin{aligned} E &= 0.500\,000\,000\,000\,000\,000\,000\,001\,436\,270\,705\,475\,576\,590\,375\,659\,826\,757\,972 \\ &\quad 824\,824\,621\,785\,332\,078\,167\,891\,514\,939\,744\,867\,648, \\ E' &= 0.499\,999\,999\,999\,999\,999\,999\,998\,540\,554\,357\,327\,868\,209\,274\,465\,258\,622\,103 \\ &\quad 903\,146\,216\,005\,437\,303\,539\,479\,001\,558\,808\,137\,418. \end{aligned} \quad (21)$$

The corresponding wavefunctions and their slopes are

$$\begin{aligned} \Psi(E, 8) &= 4.8 \times 10^{-49}, & \Psi'(E', 8) &= 8.1 \times 10^{-48}, \\ \Psi(E', 8) &= 2.6 \times 10^{-14}, & \Psi'(E, 8) &= -2 \times 10^{-13}. \end{aligned} \quad (22)$$

After matching the digits of the two numbers in equation (21), one gets the ground-state energy accurate up to 25 digits as shown in table 1. The remaining eigenvalues are obtained by the same procedure. However, one should pay attention that this accuracy is expected to be less than the accuracy of the bisection method. In this method, the accuracy estimation is $\varepsilon = (c - a)/2^n$ where n , here, is the number of iterations, and c and a are the two points enclosing only one root. In our case, we have taken for the ground state $n = 200$, $c = \frac{6}{10}$ and $a = 4/10$, giving $\varepsilon = 1.2 \times 10^{-61}$. Finally, according to the WKB approximation, the wave-function behaves for large q in the inaccessible region as

$$\Psi_{\text{WKB}}(q) \propto \frac{1}{(V(q) - E)^{\frac{1}{4}}} \exp\left(-\int_{q_t}^q \sqrt{V(q') - E} dq'\right), \quad (23)$$

where q_t is a turning point just left of the inaccessible region. The value of $\Psi_{\text{WKB}}(q = 8)$ is 6.5×10^{-14} , while for the truncated series solution of equation (13) it has the value 4.8×10^{-49} as given in equation (22). The reason for this huge difference is that the series solution is valid and convergent as long as q is finite [10]. In addition to this, the energy eigenvalues as extracted from the zeros of $\Psi(E, L)$ (for suitable L) result in a delicate cancellation between terms of opposite signs in the power series solution.

One may suspect that using a series solution in the form

$$\Psi(x) = \exp(-bx^2) \sum_j a_j x^j, \quad (24)$$

may help improve the rate of convergence for the obtained eigenvalues. In contrast, one needs more terms in the series expansion to achieve the same level of accuracy obtained by the series solution of the form given in equation (13). The reason behind this stems from the fact that any finite truncation for the series in the form given in equation (24) always decays, due to the exponential factor, as q becomes large making the determination of the energy eigenvalues less reliable, especially when the parameter b is large. As an example, when $b = \frac{1}{2}$ we can achieve the same accuracy reported in table 1 with the same number of non-vanishing terms in the truncated series, while for $b = 8$ we need 600 non-vanishing terms to achieve the same accuracy. Thus, the best thing which can be done is to work with the parameter b having zero value. However, it should be kept in mind that both the series in equations (13) and (24) are equivalent but only in the infinite sum limit.

We also study the effect of the parameter b in the case of the double-well potential given by

$$V(x) = -10x^2 + x^4 \quad (\text{in units } \hbar = 1, m = \frac{1}{2}). \quad (25)$$

Table 2. Precision versus I and the parameter b .

b	0	$\frac{1}{2}$	1	2	3	4	5	10
I	750	750	750	500	500	500	500	500
Precision	100	100	150	150	200	200	200	300

Table 3. The parameter b versus I (number of non-vanishing terms).

b	0	$\frac{1}{2}$	1	2	3	4	5	10
I	125	100	90	90	90	90	90	200

As an example, when we work with the precision 100 digits, then we find for $b = 0$, $I = 750$ and $L = 8$, that the ground-state energy has the value (accurate up to 69 digits)

$$E_0 = -20.633\,576\,702\,947\,799\,149\,958\,554\,837\,431\,508\,765\,315\,946\,057\,735\,513\,905\,710\,311\,428\,9292. \quad (26)$$

To achieve the same accurate energy determination for $b = 10$, we find that it is possible to use 500 terms which is not considerably less than the case of $b = 0$. However, this comes with the high cost of working with precision 300 digits. Working with such a high precision renders the calculation slow. At intermediate values of b like 2, 3 and 4, we can use less terms but with high precision as shown in table 2. According to our numerical investigations for the case of the double well, in the finite range, the choice $b = 0$ is the best compromise between the number of terms used and the degree of precision to get a more efficient calculation.

It is important to point out that in dealing with low-accuracy results (like nine digits), one cannot decide which is better, to work with or without the parameter b . Furthermore, employing the method in a non-efficient way may lead to wrong conclusions as in [8], where it is emphasized that setting a non-vanishing value for the parameter b greatly reduces the number of terms used. To clarify these points, we obtain for the potential given by equation (25) the four first energy levels ($E_0 = -20.633\,5767$, $E_1 = -20.633\,4568$, $E_3 = -12.379\,5437$, $E_4 = -12.375\,6738$) accurate up to 10 digits as presented in [8]; our results (using $L = 4.2$) are summarized in table 3. It is evident from table 3 that one cannot say it is a big advantage to use 90 terms (for $b = 2$) rather than 125 terms (for $b = 0$). However, numerical studies clearly indicate that the situation becomes worse when b increases (for $b = 10$ we need 200 terms). Another clear example is the pure quartic potential ($V(x) = x^4$) for which we get, for $b = 0$, $L = 3.5$ and $I = 75$, low-energy eigenvalues (the first five) determined accurately up to nine digits while obtaining the same results for the choice $b = 3$ and $I = 50$. Furthermore, the tenth eigenvalue is determined accurately up to nine digits, for $L = 3.9$, using $I = 75$ for $b = 3$, while $I = 125$ for $b = 0$. These findings are in contradiction with what has been claimed in [8], where it was mentioned that one should use about 2000 terms in the power series to determine the energy for the choice $b = 0$. Similar findings occur for the potential $V(x) = x^2 + x^8$. In such a situation, for $L = 2.5$, we can use 125 terms in the power series solution for $b = 0$ and 75 terms for $b = 5$, while getting the same accurate results up to nine digits.

The problem in the calculations found in [6, 8] comes from evaluating every term in the power series to a certain precision, and then summing the series which leads to an error accumulation, resulting in low-accuracy results despite using a large number of terms. In our approach, we sum all terms in the power series exactly, and then only in determining the roots

(energy) from the condition $\Psi_I(E, L) = 0$, do we resort to numerical calculation with a certain precision. Although the ability of the computer algebra system to deal with exact numbers was available from the early 1980s, it has not been used since then in such calculations.

4. Conclusion

In this paper we have presented an independent simple method leading to eigenvalues which agree well with the recently obtained numerical results based on the Zinn-Justin conjecture for the symmetric double-well potential. We have also included results with more significant digits than reported. It has been applied to some other potentials to illustrate its capability, and its precision has been compared with other calculations based on introducing an exponentially decaying factor (e^{-bx^2}). Several subtle points related to its precision have also been discussed and clarified. The method we opted for also enables us to get an accurate numerical determination of the corresponding wavefunctions.

Acknowledgment

This work was supported by the Research Center at College of Science, King Saud University, under project number Phys/1423/02.

References

- [1] Colemann S 1985 *Aspects of Symmetry* (Cambridge: Cambridge University Press)
- [2] Zinn-Justin J 1981 *J. Math. Phys.* **22** 511
Zinn-Justin J 1981 *Nucl. Phys. B* **192** 125
Zinn-Justin J 1983 *Nucl. Phys. B* **218** 333
Zinn-Justin J 1984 *J. Math. Phys.* **25** 549
- [3] Jentschura U D and Zinn-Justin J 2001 *J. Phys. A: Math. Gen.* **34** L253–8
- [4] Jentschura U D and Zinn-Justin J 2004 *Ann. Phys., NY* **313** 197
- [5] Alhendi H A and Lashin E I 2003 *Preprint* quant-ph/0306016
Alhendi H A and Lashin E I 2004 *Mod. Phys. Lett. A* **19** 2103
Alhendi H A and Lashin E I 2005 *Can. J. Phys.* **83** 541
- [6] Secrest D, Cashion K and Hirschfelder J O 1962 *J. Chem. Phys.* **37** 380
- [7] Barakat R and Rosner R 1981 *Phys. Lett. A* **83** 149
- [8] Killingbeck J 1981 *Phys. Lett. A* **84** 95
- [9] Taseli H and Eid R 1998 *J. Phys. A: Math. Gen.* **31** 3095
- [10] Coddington E A 1961 *An Introduction to Ordinary Differential Equations* (Englewood Cliffs, NJ: Prentice-Hall) pp 138–42