

Parafermionic derivation of Andrews-type multiple sums

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2005 J. Phys. A: Math. Gen. 38 8225

(<http://iopscience.iop.org/0305-4470/38/38/004>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.94

The article was downloaded on 03/06/2010 at 03:57

Please note that [terms and conditions apply](#).

Parafermionic derivation of Andrews-type multiple sums

P Jacob^{1,2} and P Mathieu^{1,2}

¹ Department of Mathematical Sciences, University of Durham, Durham, DH1 3LE, UK

² Département de Physique, Université Laval, Québec, G1K 7P4, Canada

E-mail: patrick.jacob@durham.ac.uk and pmathieu@phy.ulaval.ca

Received 10 May 2005, in final form 5 August 2005

Published 7 September 2005

Online at stacks.iop.org/JPhysA/38/8225

Abstract

A multi-parafermion basis of states for the \mathbb{Z}_k parafermionic models is derived. Its generating function is constructed by elementary steps. It corresponds to the Andrews multiple sum which enumerates partitions whose parts separated by the distance $k - 1$ differ by at least 2. Two analogous bases are derived for graded parafermions; one of these entails a new expression for their fermionic characters.

PACS numbers: 11.25.Hf, 02.10.Ox, 05.30.Pr, 11.10.-z

1. Introduction

1.1. Rogers–Ramanujan identities and the Andrews–Gordon generalization

The search for fermionic-type characters, that is, characters expressed as positive sums, has brought the topic of Rogers–Ramanujan identities within the framework of conformal field theory [1]³. The Rogers–Ramanujan identities are

$$\sum_{m \geq 0} \frac{q^{m^2 + (2-i)m} z^m}{(q)_m} = \prod_{n \neq 0, \pm i \pmod{5}} \frac{1}{1 - q^n}, \quad (i = 1, 2) \quad (1.1)$$

where

$$(a)_n = (a; q)_n = \prod_{i=0}^{n-1} (1 - aq^i). \quad (1.2)$$

This has various generalizations, the most relevant one being the Andrews–Gordon identity (see, e.g., [3]):

$$\sum_{m_1, \dots, m_{k-1} = 0}^{\infty} \frac{q^{N_1^2 + \dots + N_{k-1}^2 + N_1 + \dots + N_{k-1}}}{(q)_{m_1} \cdots (q)_{m_{k-1}}} = \prod_{n \neq 0, \pm i \pmod{2k+1}} \frac{1}{1 - q^n}, \quad (i = 1, \dots, k) \quad (1.3)$$

³ For further early references and a brief review of fermionic-type characters, see the introduction of [2].

with N_j defined as

$$N_j = m_j + \cdots + m_{k-1}. \quad (1.4)$$

The identity (1.3) has the following combinatorial interpretation: the lhs is the generating function for partitions (n_1, n_2, \dots) subject to the difference 2 condition

$$n_j \geq n_{j+k-1} + 2, \quad (1.5)$$

and containing at most $i - 1$ parts equal to 1, while the rhs is the generating function for partitions without parts equal to 0, $\pm i \pmod{2k + 1}$.

In the context of conformal field theory, we are mainly interested in the lhs, which is a fermionic-type expression. Granting that the two sets of partitions just described are equinumerous (which is the Gordon identity), the difficult part in establishing the analytic version (1.3) of this combinatorial identity is to demonstrate that the lhs is the proper generating function for partitions restricted by (1.5).

The point of this paper is to show that conformal field theory provides a simple method for constructing the sum-side of (1.3) and related extensions. But to put this statement in perspective, let us turn to some remarks concerning the Andrews multiple sum.

1.2. Remarks on the Andrews multiple sum

The generating function for partitions (n_1, \dots, n_m) with prescribed number of parts subject to the difference 2 condition (1.5), and containing at most $i - 1$ parts equal to 1, is

$$F_{k,i}(z; q) = \sum_{m_1, \dots, m_{k-1}=0}^{\infty} \frac{q^{N_1^2 + \cdots + N_{k-1}^2 + N_1 + \cdots + N_{k-1}} z^{N_1 + \cdots + N_{k-1}}}{(q)_{m_1} \cdots (q)_{m_{k-1}}}, \quad (1.6)$$

where the power of z gives the length of the partition. The standard proof of this result is based on the following indirect trick [4] (see also [3] chapter 7). One first shows that the number $f_{k,i}(m, n)$ of partitions of $n = \sum n_i$ with m parts subject to (1.5), and containing at most $i - 1$ parts equal to 1, satisfies a simple recurrence relation on i . This is then lifted to a recurrence relation for the generating function:

$$F_{k,i}(z; q) = \sum_{m, n \geq 0} z^m q^n f_{k,i}(m, n). \quad (1.7)$$

Finally, it is proved that the multiple sum on the right-hand side of (1.6) does satisfy this recurrence relation, with the same boundary conditions. The uniqueness of the solution of this recurrence problem completes the proof. But this is clearly a verification proof and not a constructive one⁴.

To our knowledge, there are no elementary constructive proofs of (1.6)⁵. To illustrate what is meant by such a proof, consider the case $k = 2$. The multiple sum reduces then to the sum-side of the Rogers–Ramanujan identity. As it is well known, the generating function $F_{2,i}$ is easily derived. Take $i = 2$. Looking for the generating function of partitions subject to the condition

$$n_j \geq n_{j+1} + 2, \quad (1.8)$$

one first counts those restricted partitions of length m and then sum over m . These restricted partitions can be described by the set of (standard) partitions of length at most m , whose

⁴ The original proof is based on the same recurrence relation for the generating function but the recurrence is not yet rooted to the restricted partitions, that is, to $f_{k,i}$ [5].

⁵ There are constructive proofs, using either Durfee dissections [6] or a bijection to lattice paths [7] (see also [8]), but (arguably) these are not quite elementary.

generating function is $(q)_m^{-1}$, to which we add the ‘staircase’ $(2m - 1, \dots, 5, 3, 1)$. Since the weight of the staircase is q^{m^2} , we end up with the following expression for $F_{2,2}$:

$$F_{2,2}(z; q) = \sum_{m \geq 0} \frac{q^{m^2} z^m}{(q)_m}, \quad (1.9)$$

where the variable z has been introduced to keep track of the length. For $i = 1$, there are no 1, so that the staircase is shifted to $(2m, \dots, 6, 4, 2)$ and this produces an extra term q^m within the sum. We thus recover the generating function $F_{2,i}(z; q)$ for $i = 1, 2$ by elementary steps.

How does this simple argument breaks for $k > 2$? Let us take $k = 3$ to illustrate the point and set $i = 3$. The ‘ground state’ that replaces the staircase of the previous example is now $(\dots, 7, 5, 5, 3, 3, 1, 1)$. To use the same strategy as for the $k = 2$ case would amount to trying to describe all partitions of length m with

$$n_j \geq n_{j+2} + 2, \quad (1.10)$$

in terms of the usual partitions of length at most m , to which we add the contribution of the ground state $(\dots, 7, 5, 5, 3, 3, 1, 1)$. But this simple description is simply not correct when $k > 2$. This can be seen plainly from a counter-example. There are three allowed partitions of length 3 and weight 7 satisfying (1.10): $(5, 1, 1)$, $(4, 2, 1)$ and $(3, 3, 1)$. Subtracting the ground-state contribution $(3, 1, 1)$, we are left with $(2, 0, 0)$, $(1, 1, 0)$ and $(0, 2, 0)$. But $(0, 2, 0)$ is not a genuine partition. This shows neatly that the argument used for $k = 2$ cannot be extended to higher value of k . This is a simple rationale justifying the non-elementary aspect of the proofs of (1.6).

1.3. The Andrews multiple sum in conformal field theory

We present here an elementary conformal-field-theoretical derivation of $F_{k,i}(z; q)$. As already mentioned, the multiple sum $F_{k,i}(z; q)$ has appeared in the description of the basis of states of some conformal field theories. In particular, with $z = 1$, it gives the irreducible (normalized) characters of the minimal models $\mathcal{M}(2, p)$ [9]. But more important for us here is that for a different specialization of z , one recovers the characters of the parafermionic \mathbb{Z}_k models in their fermionic form [10, 2]⁶.

But how does this function $F_{k,i}(z; q)$ actually appear in the parafermionic context? Using the generalized commutation relations between the modes of the basic parafermionic field and implementing the \mathbb{Z}_k invariance, we end up with a description of the basis of states formulated in terms of the condition (1.5), where parts at distance $k - 1$ differ by 2. More precisely, if $\mathcal{A}^{(1)}$ denotes the modes of the basic parafermionic field ψ_1 of dimension $1 - 1/k$, the descendent states are of the form⁷

$$\mathcal{A}_{-n_1}^{(1)} \cdots \mathcal{A}_{-n_m}^{(1)} |hws\rangle, \quad (1.11)$$

with n_i being positive integers subject to (1.5) and $|hws\rangle$ stands for a highest-weight state. There is in addition a boundary condition that specifies the irreducible module (the highest-weight state) under consideration. With the module labelled by an integer $1 \leq i \leq k$, this condition reads

$$n_{m-i+1} \geq 2. \quad (1.12)$$

This is clearly equivalent to the previously mentioned condition that specifies the maximal number of 1 that can appear at the right end of the associated partition (n_1, \dots, n_m) . At this

⁶ Note also that $F_{k,i}(z; q)$ is precisely the character of the principal subspace of the $\widehat{\mathfrak{su}}(2)_{k-1}$ affine algebra [11]—cf [12] for a readable account of these results and their implementation in a physical context.

⁷ Here the mode is defined up to a fractional part that is irrelevant for the present discussion.

point, i.e., having reached a description of the basis of states, the generating function (1.6) is invoked. Finally, by relating z to a power of q in order to adjust the total power of q to the proper conformal dimension of the states (taking thus due care of the omitted fractional parts in the modes), we recover the irreducible parafermionic characters⁸.

In this work, we present another basis of states for the parafermionic models. This basis is not formulated solely in terms of the basic parafermionic modes but involves rather the modes of the complete set of $k - 1$ parafermionic fields. The generating function of this basis of states turns out to be built by elementary steps, analogous to those that led to the sum-side of the Rogers–Ramanujan identity. The resulting expression is precisely the above function $F_{k,i}(z; q)$. Turning this around, the equivalence of the two bases of states for the parafermionic theories, the one exposed here and the previous one formulated in terms of partitions restricted by (1.5), entails a simple constructive proof of the Andrews multiple-sum identity.

Physically, this new derivation is quite appealing since each of the $k - 1$ sums on the rhs of (1.6) is linked to the counting of a given type of modes. In other words, the number m_j labels the number of parafermionic modes of type j .

1.4. The \mathbb{Z}_k multi-parafermion basis: combinatorial formulation

Let us state our result in a field-theoretical independent way. The multi-parafermion basis of states is equivalent to the set of $k - 1$ ordered partitions of respective lengths m_1, \dots, m_{k-1} , i.e.,

$$(n^{(1)}, n^{(2)}, \dots, n^{(k-1)}) \quad \text{with} \quad n^{(j)} = (n_1^{(j)}, \dots, n_{m_j}^{(j)}), \quad (1.13)$$

where the parts within a partition satisfy

$$n_l^{(j)} \geq n_{l+1}^{(j)} + 2j. \quad (1.14)$$

The different partitions are further subject to the boundary conditions:

$$n_{m_j}^{(j)} \geq j + \max(j - i + 1, 0) + 2j(m_{j+1} + \dots + m_{k-1}). \quad (1.15)$$

The length m and the weight n of the partitions enumerated by $f_{k,i}(m, n)$ are related to the above data as follows:

$$n = \sum_{j=1}^{k-1} \sum_{l=1}^{m_j} n_l^{(j)} \quad \text{and} \quad m = \sum_{j=1}^{k-1} j m_j. \quad (1.16)$$

Clearly, it is because we have a sequence of partitions with a difference condition at distance 1, i.e., the condition (1.14), that the generating function is so easily constructed.

1.5. A natural generalization

After deriving this ‘new’ basis of states, we have found that it has actually appeared previously in the literature on vertex operator algebras in [13] and in a much more general version⁹. Therefore, at the worse, we have provided a conformal-field-theoretical proof of a result already established by means of vertex-operator-algebra techniques. But we would like to stress the remarkable simplicity of our argument which, by itself, justifies its presentation.

In addition to be simple, our approach seems to have an important potential for generalization. This is illustrated here by the study of the graded parafermions (untreated

⁸ This construction could be rephrased in more Lie-algebraic terms in the language of vertex operator algebras following [10].

⁹ The basis in [13] pertains to all models of the form $\widehat{su}(r+1)_k/\widehat{u}(1)^r$. For $\widehat{su}(2)$, it reduces to the present basis.

in [13]) presented in section 4. In that case, two multi-parafermion bases are derived. One of the resulting generating function is new and it leads to a novel fermionic character formula for graded parafermions.

2. The \mathbb{Z}_k parafermionic models

The parafermionic conformal algebra is spanned by $k - 1$ parafermionic fields $\psi_r, r = 0, 1, \dots, k - 1$, with dimension

$$h_r = r \left(1 - \frac{r}{k}\right). \tag{2.1}$$

Note that $\psi_0 = I$, the identity field. For the present purpose, we will only need the following OPE [14]:

$$\psi_r(z)\psi_s(w) \sim \frac{c_{r,s}}{(z-w)^{2rs/k}} \left[\psi_{r+s}(w) + \frac{r}{r+s}(z-w)\partial\psi_{r+s} + \dots \right] \quad (r+s \leq k), \tag{2.2}$$

where the structure constants $c_{r,s}$ are fixed by associativity [14] (their explicit form will not be needed here).

Recall that the decomposition of the parafermionic field in modes depends upon the field on which it acts [14]. It is essentially fixed by the mutual locality, which is the phase that results from the substitution $z \rightarrow z e^{2\pi i}$, denoted by $e^{2\pi i\gamma}$. The OPE $\psi_r(z)\psi_s(w)$ indicates that the mutual locality coefficient of ψ_r and ψ_s , denoted by $\gamma_{r,s}$, is $\gamma_{r,s} = -2rs/k$. From the mutual locality coefficient, we can introduce a charge q , defined as

$$\gamma_{r,s} = -\frac{q_r q_s}{2k}. \tag{2.3}$$

The charge is normalized by setting $q_1 = 2$, so that $q_r = 2r$. The mutual locality coefficient of ψ_r and ϕ_q , a generic field of charge q , will then be $-rq/k$. Therefore, the mode decomposition of ψ_r acting on an arbitrary field ϕ_q reads

$$\psi_r(z)\phi_q(0) = \sum_{m=-\infty}^{\infty} z^{-rq/k-m-r} A_{r(r+q)/k+m}^{(r)} \phi_q(0), \tag{2.4}$$

the fractional power of z being fixed by the mutual locality.

In the following, and in agreement with our previous works [15, 2], the fractional part of the modes is omitted (being fixed unambiguously by the charge of the field or state on which it acts) and this is indicated by calligraphic symbols, i.e.,¹⁰

$$\mathcal{A}_n^{(r)} |\phi_q\rangle \equiv A_{n+r(r+q)/k}^{(r)} |\phi_q\rangle. \tag{2.5}$$

A form of the commutation relation between the $\mathcal{A}^{(r)}$ and $\mathcal{A}^{(s)}$ modes for $r+s \leq k$ follows from the computation of the integral

$$\frac{1}{(2\pi i)^2} \oint_{C_1} dw \oint_{C_2} dz z^{qr/k+n} w^{qs/k+m} (z-w)^{-2+2rs/k} \psi_r(z)\psi_s(w)\phi_q(0), \tag{2.6}$$

by standard contour deformation¹¹. The result is (omitting the state associated with $\phi_q(0)$ on which it acts):

$$\sum_{l=0}^{\infty} C_{2rs/k-2}^{(l)} [\mathcal{A}_{n-l-r-1}^{(r)} \mathcal{A}_{m+l-s+1}^{(s)} - \mathcal{A}_{m-l-s-1}^{(s)} \mathcal{A}_{n+l-r+1}^{(r)}] = ac_{r,s} \mathcal{A}_{n+m-r-s+1}^{(r+s)}, \tag{2.7}$$

¹⁰ This notation simplifies considerably the writing but it should be kept in mind that the conformal dimension of the mode is no longer given by minus its index. Note that here $|\phi_q\rangle$ stands for an arbitrary state of charge q .

¹¹ The integral for C_2 circulating around w while C_1 is a small contour around the origin is compared to the difference of two contours, one with $|z| > |w|$ and the other with $|z| < |w|$. Note that in the latter case, ψ_r passes over ψ_s and this produces a phase factor $(-1)^{-2rs/k}$ that is partly cancelled by the one coming from $(z-w)^{-2+2rs/k} \rightarrow (-1)^{2rs/k} (w-z)^{-2+2rs/k}$.

where

$$C_t^{(l)} = \frac{\Gamma(l-t)}{l!\Gamma(-t)}, \quad a = \left(\frac{ns - mr}{r+s} \right). \quad (2.8)$$

In the above integral, the power of $z - w$ is chosen in order to pick up precisely the first two non-vanishing terms of the OPE (in contradistinction with the usual presentation of the commutation relation where only the first non-vanishing term is picked out). We stress that this is made possible by the fact that in the module $A_{-r-s}^{(r+s)}|0\rangle$, there is a single descendant of relative charge 0 and relative level 1 and it is proportional to $L_{-1}A_{-r-s}^{(r+s)}|0\rangle$. Now the reason for which we pick up these two terms is to extract the maximal amount of constraint from the commutator without generating new types of fields, that is, fields other than ψ_{r+s} .¹²

Denote the parafermionic primary fields by $\{\varphi_\ell | \ell = 0, \dots, k-1\}$ [14, 15]. To each primary field, there corresponds a highest-weight state $|\varphi_\ell\rangle$. In particular, $|0\rangle = |\varphi_0\rangle$. The highest-weight conditions $\mathcal{A}_{-n-r}^{(r)}|\varphi_\ell\rangle = 0$ for $n < 0$ together with the singular vector $(\mathcal{A}_{-1}^{(1)})^{k-\ell+1}|\varphi_\ell\rangle = 0$ imply:

$$\mathcal{A}_{-n-r}^{(r)}|\varphi_\ell\rangle = 0 \quad \text{for } n < \max(r - k + \ell, 0). \quad (2.9)$$

Note that $\psi_r(0)|0\rangle = \mathcal{A}_{-r}^{(r)}|0\rangle \propto (\mathcal{A}_{-1}^{(1)})^r|0\rangle$.

3. A multi-parafermion basis of states

We look for a basis of states constructed out of the $k-1$ parafermionic modes, that is, a basis of the form

$$\mathcal{A}_{-n_1^{(1)}}^{(1)} \cdots \mathcal{A}_{-n_{m_1}^{(1)}}^{(1)} \mathcal{A}_{-n_1^{(2)}}^{(2)} \cdots \mathcal{A}_{-n_{m_2}^{(2)}}^{(2)} \cdots \mathcal{A}_{-n_1^{(k-1)}}^{(k-1)} \cdots \mathcal{A}_{-n_{m_{k-1}}^{(k-1)}}^{(k-1)} |\varphi_\ell\rangle. \quad (3.1)$$

The goal being to determine the set of independent states for a sequence of this type, one needs to find those conditions on the indices $n_l^{(j)}$ that would avoid over counting. These conditions are to be fixed by the commutation relations. In those relations, we can clearly set to zero those terms already considered. In particular, since each type of modes $\mathcal{A}_n^{(p)}$ for $0 \leq p \leq k-1$ is considered successively, we can drop their contribution on the rhs of the commutation relations (2.7) with $p = r+s$ and set

$$\sum_{l=0}^{\infty} C_{2rs/k-2}^{(l)} [\mathcal{A}_{n-l-r-1}^{(r)} \mathcal{A}_{m+l-s+1}^{(s)} - \mathcal{A}_{m-l-s-1}^{(s)} \mathcal{A}_{n+l-r+1}^{(r)}] \sim 0. \quad (3.2)$$

Let us now look at the consequences of these simplified relations. Consider first the string of $\mathcal{A}^{(1)}$ modes and set $r = s = 1$ in (3.2):

$$\sum_{l=0}^{\infty} C_{2/k-1}^{(l)} [\mathcal{A}_{n-l-2}^{(1)} \mathcal{A}_{m+l}^{(1)} - \mathcal{A}_{m-l-2}^{(1)} \mathcal{A}_{n+l}^{(1)}] \sim 0. \quad (3.3)$$

This shows that moving a $\mathcal{A}^{(1)}$ mode to the right of another $\mathcal{A}^{(1)}$ mode produces a shift Δ of its mode index by at least 2, that is,

$$\Delta = n+l - (n-l-2) = 2l+2 \geq 2. \quad (3.4)$$

Therefore, the $\mathcal{A}^{(1)}$ sequence of independent descendants takes the form

$$\mathcal{A}_{-n_1^{(1)}}^{(1)} \cdots \mathcal{A}_{-n_{m_1}^{(1)}}^{(1)} | \cdots \rangle \quad \text{with} \quad n_l^{(1)} \geq n_{l+1}^{(1)} + 2. \quad (3.5)$$

¹² For instance, the next subleading term of the OPE would involve the new field $(T\psi_{r+s})$.

In other words, because we have a shift by at least 2 in the commutation relation (3.3), we have a difference condition of 2 between adjacent parts. Moreover, the highest-weight condition requires $n_{m_1} \geq 1$. But this inequality on n_{m_1} is bounded to be modified by the presence of higher modes. Indeed, consider next the commutation of $\mathcal{A}^{(1)}$ and $\mathcal{A}^{(2)}$:

$$\sum_{l=0}^{\infty} C_{4/k-2}^{(l)} [\mathcal{A}_{n-l-2}^{(1)} \mathcal{A}_{m+l-1}^{(2)} - \mathcal{A}_{m-l-3}^{(2)} \mathcal{A}_{n+l}^{(1)}] \sim 0. \tag{3.6}$$

We see that by moving a $\mathcal{A}^{(1)}$ mode to the right of a $\mathcal{A}^{(2)}$ mode generates a shift of at least 2. Therefore, when the $\mathcal{A}^{(1)}$'s are preceded by a string of $m_2 \mathcal{A}^{(2)}$ modes, the $\mathcal{A}^{(1)}$ indices are shifted by the additional term $2m_2$. More generally, the relation

$$\sum_{l=0}^{\infty} C_{2r/k-2}^{(l)} [\mathcal{A}_{n-l-2}^{(1)} \mathcal{A}_{m+l-r+1}^{(r)} - \mathcal{A}_{m-l-r-1}^{(r)} \mathcal{A}_{n+l}^{(1)}] \sim 0 \tag{3.7}$$

shows that passing $\mathcal{A}^{(1)}$ over $\mathcal{A}^{(r)}$ (for any $r > 1$) generates a shift of at least 2. Therefore, the presence of higher modes to the right of the $\mathcal{A}^{(1)}$ ones induces a shift of all the $\mathcal{A}^{(1)}$ modes by $2(m_2 + \dots + m_{k-1})$. This reproduces (1.15) for $j = 1$ up to the ℓ -dependent boundary term.

Consider now the constraints on the $\mathcal{A}^{(2)}$ modes. The highest-weight condition requires $n_{m_2}^{(2)} \geq 2$. Now, since we have already taken into account the commutation of $\mathcal{A}^{(2)}$ with $\mathcal{A}^{(1)}$, it suffices to consider that of $\mathcal{A}^{(2)}$ with $\mathcal{A}^{(r)}$ modes for $r \geq 2$. But actually, the resulting constraints for those cases cannot be obtained by the commutation relations since the various types of modes have already been generated by the commutators that involve $\mathcal{A}^{(1)}$. To be explicit, we must take due care of the fact that, say $\mathcal{A}^{(1)} \mathcal{A}^{(3)} \sim \mathcal{A}^{(2)} \mathcal{A}^{(2)} \sim \mathcal{A}^{(4)}$. Instead, constraints on higher modes have to be determined by the associativity requirement. Since $\mathcal{A}^{(2)} \sim \mathcal{A}^{(1)} \mathcal{A}^{(1)}$, moving $\mathcal{A}^{(r)}$ past a $\mathcal{A}^{(2)}$ mode induces a shift of at least 4 (2 for each $\mathcal{A}^{(1)}$) for any $r \geq 2$. Therefore, within the string of $\mathcal{A}^{(2)}$ modes, we have a difference condition of 4 between adjacent modes (that follows by considering $r = 2$) and a global shift of four times the number of other type of modes at its right, that is, $4(m_3 + \dots + m_{k-1})$ (from the $r > 2$ cases). This yields (1.14) and (1.15) for $j = 2$ (again disregarding the ℓ -part of the boundary condition).

More generally, to extract the constraints for the commutation of $\mathcal{A}^{(i)}$ and $\mathcal{A}^{(j)}$ by associativity, in order to find the less restrictive conditions, we expand the mode with smallest index (i or j) in terms of $\mathcal{A}^{(1)}$ modes. We then find that the resulting shift is $2 \min(i, j)$ for the other mode (with index $\max(i, j)$). This readily shows that the parts $n_l^{(j)}$ satisfy

$$n_l^{(j)} \geq n_{l+1}^{(j)} + 2j, \tag{3.8}$$

together with

$$n_{m_j}^{(j)} \geq j + 2j(m_{j+1} + \dots + m_{k-1}). \tag{3.9}$$

Let us now construct the generating function for this basis of states, ignoring in the first step the boundary condition on ℓ . Let us first take into account the contribution of the $\mathcal{A}^{(1)}$ modes. It is given by enumerating ordinary partitions of length at most m_1 , all shifted by the staircase of weight:

$$\sum_{l=0}^{m_1-1} [2l + 1 + 2(m_2 + \dots + m_{k-1})] = m_1^2 + 2m_1(m_2 + \dots + m_{k-1}). \tag{3.10}$$

By introducing the dummy variable z_1 to keep track of the number of $\mathcal{A}^{(1)}$ modes, we have

$$\sum_{m_1 \geq 0} z_1^{m_1} \frac{q^{m_1^2 + 2m_1(m_2 + \dots + m_{k-1})}}{(q)_{m_1}}. \tag{3.11}$$

More generally, the contribution of the $\mathcal{A}^{(j)}$ modes is obtained by enumerating ordinary partitions of length at most m_j shifted by the staircase of step $2j$, whose weight, properly modified by the presence of the number of modes of higher type (i.e., $r > j$), is

$$j \sum_{l=0}^{m_j-1} [2l + 1 + 2(m_{j+1} + \dots + m_{k-1})] = jm_j^2 + 2jm_j(m_{j+1} + \dots + m_{k-1}). \tag{3.12}$$

This contributes to the factor

$$\sum_{m_j \geq 0} z_j^{m_j} \frac{q^{jm_j^2 + 2jm_j(m_{j+1} + \dots + m_{k-1})}}{(q)_{m_j}}. \tag{3.13}$$

Summing up all terms, we end up with the following generating function:

$$\sum_{m_1, \dots, m_{k-1} = 0}^{\infty} \frac{q^{N_1^2 + \dots + N_{k-1}^2} \prod_{j=1}^{k-1} z_j^{m_j}}{(q)_{m_1} \dots (q)_{m_{k-1}}}, \tag{3.14}$$

where the N_j are defined in (1.4). We can introduce a single variable to keep track of the relative charge of the descendant states instead of the length of its various parts by defining $z_j = z^j$. This leads to

$$\sum_{m_1, \dots, m_{k-1} = 0}^{\infty} \frac{q^{N_1^2 + \dots + N_{k-1}^2} z^{N_1 + \dots + N_{k-1}}}{(q)_{m_1} \dots (q)_{m_{k-1}}}. \tag{3.15}$$

Let us now take care of the boundary condition that characterizes the different modules, that is, the ℓ -dependence in the condition (2.9), which can be rewritten as:

$$n_{m_r}^{(r)} \geq r + \max(r - k + \ell, 0). \tag{3.16}$$

The second term in the bound (3.16) produces a further global shift for all the indices of type r such that $r - k + \ell > 0$. Summing their contribution generates the weight factor

$$\sum_{r=1}^{k-1} \max(r - k + \ell, 0) m_r = m_{k-\ell+1} + 2m_{k-\ell+2} + \dots + (\ell - 1)m_{k-1} = N_{k-\ell+1} + \dots + N_{k-1}. \tag{3.17}$$

This reproduces precisely the linear term in the exponent of q in (1.6) for $i = k - \ell + 1$. We have thus recovered the function $F_{k,i}(z; q) = F_{k,k-\ell+1}(z; q)$.

Note finally that by reinserting the fractional contribution of the modes (e.g., as described in section 5.2 of [2]) one recovers the Lepowsky–Primc character formula for the \mathbb{Z}_k parafermionic models [10].

4. New quasi-particle bases for graded parafermions

4.1. Preliminary remarks on graded parafermions

Graded parafermions [16] are associated with the coset $\widehat{\mathfrak{osp}}(1, 2)_k / \widehat{u}(1)$. The corresponding chiral algebra is generated by $2k - 1$ parafermions $\tilde{\psi}_r$, $r = 0, \frac{1}{2}, 1, \dots, k - \frac{1}{2}$, of dimension

$$\tilde{h}_r = r \left(1 - \frac{r}{k} \right) + \frac{\epsilon_r}{2}, \tag{4.1}$$

where $\epsilon_r = 0$ if r is integer and 1 otherwise. The conformal dimension of the lowest dimensional parafermion $\tilde{\psi}_{1/2}$ is thus $1 - 1/4k$. The defining OPE reads ($r + s \leq k$)

$$\tilde{\psi}_r(z) \tilde{\psi}_s(w) \sim \frac{\tilde{c}_{r,s}}{(z-w)^{2rs/k + \epsilon_r \epsilon_s}} [\tilde{\psi}_{r+s}(w) + \dots]. \tag{4.2}$$

Note that for $r + s$ half-integer, there are more than one descendant field at level 1. The mode decomposition is defined as

$$\tilde{\psi}_r(z)\phi_q(0) = \sum_{m=-\infty}^{\infty} z^{-rq/k-m-r-\epsilon_r/2} \tilde{A}_{r(r+q)/k+m}^{(r)} \phi_q(0). \tag{4.3}$$

As before, we will avoid writing the fractional part of the modes explicitly. The primary fields $\tilde{\varphi}_\ell$ are parametrized by an integer ℓ such that $0 \leq \ell \leq k$. The highest-weight conditions together with the singular vector $(\mathcal{A}_{-1}^{(1)})^{k-\ell+1} |\tilde{\varphi}_\ell\rangle = 0$ [17] imply:

$$\tilde{A}_{-r-\epsilon_r/2-n}^{(r)} |\tilde{\varphi}_\ell\rangle = 0 \quad \text{if } n < \max\left(r - \frac{\epsilon_r}{2} - k + \ell, 0\right). \tag{4.4}$$

4.2. A first graded multi-parafermion basis

The first basis we look for is of the form

$$\tilde{A}_{-n_1^{(0)}}^{(1/2)} \cdots \tilde{A}_{-n_{m_0}^{(0)}}^{(1/2)} \tilde{A}_{-n_1^{(1)}}^{(1)} \cdots \tilde{A}_{-n_{m_1}^{(1)}}^{(1)} \tilde{A}_{-n_1^{(2)}}^{(2)} \cdots \tilde{A}_{-n_{m_2}^{(2)}}^{(2)} \cdots \tilde{A}_{-n_1^{(k-1)}}^{(k-1)} \cdots \tilde{A}_{-n_{m_{k-1}}^{(k-1)}}^{(k-1)} |\tilde{\varphi}_\ell\rangle. \tag{4.5}$$

We then have to find the constraints on the different types of indices by considering the commutation relations. Consider first the commutator between two $\tilde{A}^{(1/2)}$ modes. For this, since the basis includes the $\tilde{A}^{(1)}$ modes and because the $\tilde{A}^{(1)}$ module has a single zero-charge descendant at level 1, we can pick up the first two non-vanishing terms in the OPE. This results into [16, 17]

$$\sum_{l \geq 0} C_{1/2k-1}^{(l)} [\tilde{A}_{n-l-1}^{(1/2)} \tilde{A}_{m+l}^{(1/2)} - \tilde{A}_{m-l-1}^{(1/2)} \tilde{A}_{n+l}^{(1/2)}] \sim 0. \tag{4.6}$$

This indicates a difference 1 between adjacent modes $\tilde{A}^{(1/2)}$:

$$n_l^{(0)} \geq n_{l+1}^{(0)} + 1. \tag{4.7}$$

The condition (4.4) yields $n_{m_0}^{(0)} \geq 1$. Next, we consider the commutator of $\tilde{A}^{(1/2)}$ with $\tilde{A}^{(r)}$ for r integer. Since we do not take into account the modes $\tilde{A}^{(r+1/2)}$ in this basis, we must avoid picking up any non-vanishing terms on the rhs of the corresponding OPE. The strongest constraint we get with this restriction is

$$\sum_{l \geq 0} C_{r/k+1}^{(l)} [\tilde{A}_{n-l+1}^{(1/2)} \tilde{A}_{m+l-r+1}^{(r)} - \tilde{A}_{m-l-r+1}^{(r)} \tilde{A}_{n+l+1}^{(1/2)}] = 0. \tag{4.8}$$

This implies that the smallest mode-shifting we can get when a $\tilde{A}^{(1/2)}$ mode is moved past a $\tilde{A}^{(r)}$ mode is zero. That indicates that the presence of higher modes at the right of the $\tilde{A}^{(1/2)}$ string does not affect the latter modes, that is, it does not modify the bound $n_{m_0}^{(0)} \geq 1$. As a result, there will be no interacting term of the type $m_0 m_r$ in the generating function.

For the other modes, the analysis is similar to the one pertaining to the non-graded case. Hence, (1.14) and (1.15) still hold for $j \geq 1$.

This basis has the following generating function:

$$\sum_{m_0, m_1, \dots, m_{k-1}=0}^{\infty} \frac{q^{m_0(m_0+1)/2 + N_1^2 + \dots + N_{k-1}^2 + N_{k-\ell+1} + \dots + N_{k-1}} z_0^{m_0} \prod_{j=1}^{k-1} z_j^{m_j}}{(q)_{m_0} (q)_{m_1} \cdots (q)_{m_{k-1}}}. \tag{4.9}$$

Setting $z_0 = z$, $z_j = z^{2j}$ and summing over the m_0 modes, this becomes

$$(-zq)_\infty \sum_{m_1, \dots, m_{k-1}=0}^{\infty} \frac{q^{N_1^2 + \dots + N_{k-1}^2 + N_{k-\ell+1} + \dots + N_{k-1}} z^{2(N_1 + \dots + N_{k-1})}}{(q)_{m_1} \cdots (q)_{m_{k-1}}} = (-zq)_\infty F_{k, k-\ell+1}(z^2; q), \tag{4.10}$$

which is precisely the result obtained in [18] (cf equation (3.24)). The simplicity of this derivation contrasts heavily with that in the later reference, which requires the enumeration of restricted jagged partitions [17, 19].

4.3. A second graded multi-parafermion basis

The second basis we consider involves all graded-parafermionic modes, that is,

$$\tilde{\mathcal{A}}_{-n_1^{(1/2)}}^{(1/2)} \cdots \tilde{\mathcal{A}}_{-n_{m_1/2}^{(1/2)}}^{(1/2)} \tilde{\mathcal{A}}_{-n_1^{(1)}}^{(1)} \cdots \tilde{\mathcal{A}}_{-n_{m_1}^{(1)}}^{(1)} \tilde{\mathcal{A}}_{-n_1^{(3/2)}}^{(3/2)} \cdots \tilde{\mathcal{A}}_{-n_{m_{3/2}}^{(3/2)}}^{(3/2)} \cdots \tilde{\mathcal{A}}_{-n_1^{(k-1/2)}}^{(k-1/2)} \cdots \tilde{\mathcal{A}}_{-n_{m_{k-1/2}}^{(k-1/2)}}^{(k-1/2)} |\tilde{\varphi}_\ell\rangle. \tag{4.11}$$

Let us note readily the boundary condition on each index $n_{m_j}^{(j)}$ that results from (4.4)

$$n_{m_j}^{(j)} \geq j + \frac{\epsilon_j}{2} + \max\left(j - \frac{\epsilon_j}{2} - k + \ell, 0\right). \tag{4.12}$$

Again we start by considering the commutation relation between the $\tilde{\mathcal{A}}^{(1/2)}$ and $\tilde{\mathcal{A}}^{(r)}$ modes, where now r can be both integer and half-integer. For $r = 1/2$, the analysis of the previous subsection still holds. Thus, here again, the $\tilde{\mathcal{A}}^{(1/2)}$ modes have to be distinct. For $r > 1/2$, if the produced module $\tilde{\mathcal{A}}^{(r+1/2)}$ has a single zero-relative-charge descendant at level 1, we can pick up two non-vanishing terms on the rhs of the OPE. This is the case when r is half-integer. The relevant commutation relation is then

$$\sum_{l \geq 0} C_{r/k-1}^{(l)} [\tilde{\mathcal{A}}_{n-l-1}^{(1/2)} \tilde{\mathcal{A}}_{m+l-r}^{(r)} - \tilde{\mathcal{A}}_{m-l-r-1}^{(r)} \tilde{\mathcal{A}}_{n+l}^{(1/2)}] \sim 0. \tag{4.13}$$

This implies a shift of 1 in $\tilde{\mathcal{A}}^{(1/2)}$ modes for each $\tilde{\mathcal{A}}^{(r)}$ modes at its right, with r half-integer. For r integer, it turns out that there are generically three zero-relative-charge descendant at level 1 in the module $\tilde{\mathcal{A}}^{(r+1/2)}$.¹³ Therefore, only the first non-vanishing term must be considered in the OPE $\psi_{1/2}(z)\psi_r(w)$. This gives

$$\sum_{l \geq 0} C_{r/k-1}^{(l)} [\tilde{\mathcal{A}}_{n-l-1}^{(1/2)} \tilde{\mathcal{A}}_{m+l-r+1}^{(r)} - \tilde{\mathcal{A}}_{m-l+r}^{(r)} \tilde{\mathcal{A}}_{n+l}^{(1/2)}] \sim 0, \tag{4.14}$$

and again this implies a shift of 1 in $\tilde{\mathcal{A}}^{(1/2)}$ mode for each $\tilde{\mathcal{A}}^{(r)}$ modes at its right, with r integer. Associativity (decomposition of higher modes into a product of $\tilde{\mathcal{A}}^{(1/2)}$ ones) shows that when $\tilde{\mathcal{A}}^{(r)}$ is passed over a $\tilde{\mathcal{A}}^{(s)}$, there is a difference $2 \min(r, s)$. This is thus a difference of $2r$ between the $\tilde{\mathcal{A}}^{(r)}$ modes and a shift of $2r$ for each higher modes at its right. When summing over the contribution of the r modes, this generates the weight

$$rm_r^2 + 2rm_r(m_{r+1/2} + m_{r+1} + \cdots + m_{k-1/2}). \tag{4.15}$$

The ℓ -dependent boundary term that has been ignored so far is evaluated as in the non-graded case:

$$\begin{aligned} & \sum_{r=1}^{k-1/2} \max\left(r - \frac{\epsilon_r}{2} - k + \ell, 0\right) m_r \\ &= (m_{k-\ell+1} + m_{k-\ell+3/2}) + \cdots + (\ell - 1)(m_{k-1} + m_{k-1/2}) \equiv \tilde{L}_{k-\ell+1}. \end{aligned} \tag{4.16}$$

¹³ To see this neatly, take k large. The character (normalized such that the leading term is 1) of the vacuum module of relative charge $2r$ (that is, the module of $\mathcal{A}_r^{(r)}|0\rangle$) is given by (for $r > 1/2$)

$$\chi_{2r}(q) \approx V_{2r}(q) - V_{2r+1}(q) = 1 + (1 + 2\epsilon_r)q + \cdots$$

where V_t denotes the Verma module of relative charge t (cf equations (5.4)–(5.6) and (5.12), (5.13) of [18]).

The resulting generating function is thus

$$\sum_{m_{1/2}, m_1, m_{3/2}, \dots, m_{k-1/2}=0}^{\infty} \frac{q^{\frac{1}{2}(\tilde{N}_{1/2}^2 + \tilde{N}_1^2 + \dots + \tilde{N}_{k-1/2}^2 + M_{1/2}) + \tilde{L}_{k-\ell+1}} \prod_{j=1/2}^{k-1/2} z_j^{m_j}}{(q)_{m_{1/2}} (q)_{m_1} \cdots (q)_{m_{k-1/2}}}. \tag{4.17}$$

where

$$\tilde{N}_j = m_j + m_{j+1/2} + \dots + m_{k-1/2} = M_j + M_{j+1/2}, \tag{4.18}$$

with

$$M_j = m_j + m_{j+1} + \dots + m_{k-1+\epsilon_j/2}, \tag{4.19}$$

and $\tilde{L}_{k-\ell+1}$ defined in (4.16). With $z_j = z^{2j}$, the z factor reduces to $z^{\tilde{N}}$ where $\tilde{N} = \sum 2jm_j$ and we have

$$G_{k,k-\ell+1}(z; q) = \sum_{m_{1/2}, m_1, m_{3/2}, \dots, m_{k-1/2}=0}^{\infty} \frac{q^{\frac{1}{2}(\tilde{N}_{1/2}^2 + \tilde{N}_1^2 + \dots + \tilde{N}_{k-1/2}^2 + M_{1/2}) + \tilde{L}_{k-\ell+1}} z^{\tilde{N}}}{(q)_{m_{1/2}} (q)_{m_1} \cdots (q)_{m_{k-1/2}}}. \tag{4.20}$$

4.4. A generalized Rogers–Ramanujan identity

The equivalence of the two new graded bases implies the equality (with $i = k - \ell + 1$):

$$G_{k,i}(z; q) = (-zq)_{\infty} F_{k,i}(z^2; q). \tag{4.21}$$

For $z = 1$, the rhs has a product form (cf [20] theorem 11). This and the above equality lead to the following generalization of the Rogers–Ramanujan identity¹⁴:

$$\sum_{m_{1/2}, \dots, m_{k-1/2}=0}^{\infty} \frac{q^{\frac{1}{2}(\tilde{N}_{1/2}^2 + \tilde{N}_1^2 + \dots + \tilde{N}_{k-1/2}^2 + M_{1/2}) + \tilde{L}_i}}{(q)_{m_{1/2}} (q)_{m_1} \cdots (q)_{m_{k-1/2}}} = \prod_{n=1}^{\infty} (1 + q^n) \prod_{n \neq 0, \pm i \pmod{2k+1}}^{\infty} (1 - q^n)^{-1}. \tag{4.22}$$

We stress that with the expression we had previously [18] for the specialized multi-sum, i.e., $(-q)_{\infty} F_{k,i}(1; q)$, the factor $(-q)_{\infty} = \prod_{n=1}^{\infty} (1 + q^n)$ would cancel on both sides of the ‘sum = product’ equality

$$(-q)_{\infty} F_{k,i}(1; q) = \prod_{n=1}^{\infty} (1 + q^n) \prod_{n \neq 0, \pm i \pmod{2k+1}}^{\infty} (1 - q^n)^{-1}, \tag{4.23}$$

reducing then to the usual Andrews–Gordon identity. But there is no such cancellation with (4.22) (except for the trivial case $k = 1$). In particular, for $k = 2$, it reads

$$\sum_{n,m,p=0}^{\infty} \frac{q^{\frac{1}{2}n^2 + m^2 + \frac{3}{2}p^2 + n(m+p) + 2mp + (2-i)(m+p) + \frac{1}{2}(n+p)}}{(q)_n (q)_m (q)_p} = \prod_{n=1}^{\infty} (1 + q^n) \prod_{n \neq 0, \pm i \pmod{5}} \frac{1}{1 - q^n}, \quad (i = 1, 2) \tag{4.24}$$

Because it involves the modulus 5 on the rhs, this identity could be viewed as the fermionic deformation of the original Rogers–Ramanujan identity (1.1).

¹⁴ Multiple sums similar but not identical to $G_{k,i}(z; q)$ have been conjectured in [21] as fermionic expressions for the Ramond characters of the superconformal minimal model $\mathcal{SM}(2, 4k)$. (These identities have been subsequently proved in [22]—see also theorem 4.4 of [23]). Note that if we relabel our m_j as m_{2j} and set $m_{2(k-\ell+1)} = m_s$ (so that s is even) together with $m_{2k-1} = 0$ in our formula, we recover the expression given in the second line of equation (2.6) in [21]. This signals an unexpected relation between the $\mathcal{SM}(2, 4k)$ models and the \mathbb{Z}_k graded parafermions. The present analysis, in the light of the recent work [24], provides a possible path for an alternative proof of these identities.

There is a striking similarity between (4.22) for $i = k$ and the identity of theorem 4.5 of [23]. In fact, Warnaar [25] has shown that these two relations are essentially equivalent. The sketch of the proof—which is an analytic counterpart of our conformal-field-theoretical proof of (4.21)—is reported in the appendix.

4.5. The \mathbb{Z}_k graded multi-parafermion bases: combinatorial formulation

Taken together, the results of [17] and the present ones have the following combinatorial interpretation. There is an equality between the number of partitions described by the following three sets.

- (1) The first set corresponds to the jagged partitions (n_1, \dots, n_m) defined as

$$n_j \geq n_{j+1} - 1, \quad n_j \geq n_{j+2}, \quad n_m \geq 1, \quad (4.25)$$

with at most $i = 1$ pairs of 01 and further subject to the following the k -restrictions:

$$n_j \geq n_{j+2k-1} + 1 \quad \text{or} \quad n_j = n_{j+1} - 1 = n_{j+2k-2} + 1 = n_{j+2k-1}, \quad (4.26)$$

for all values of $j \leq m - 2k + 1$, with $k > 1$.

- (2) The second set corresponds to a sequence of k ordered partitions $(n^{(0)}, n^{(1)}, n^{(2)}, \dots, n^{(k-1)})$ of respective lengths m_0, m_1, \dots, m_{k-1} , with

$$n_l^{(0)} \geq n_{l+1}^{(0)} + 1, \quad n_l^{(j)} \geq n_{l+1}^{(j)} + 2j, \quad (4.27)$$

with the different partitions being further subject to the boundary conditions

$$n_{m_0}^{(0)} \geq 1, \quad n_{m_j}^{(j)} \geq j + \max(j - i + 1, 0) + 2j(m_{j+1} + \dots + m_{k-1}), \quad (4.28)$$

with $j \geq 1$.

- (3) Finally, the third set corresponds to a sequence of $2k - 1$ ordered partitions $(n^{(1/2)}, n^{(1)}, n^{(3/2)}, \dots, n^{(k-1/2)})$ of respective lengths $m_{1/2}, m_1, \dots, m_{k-1/2}$, with

$$n_l^{(j)} \geq n_{l+1}^{(j)} + 2j, \quad (4.29)$$

and the boundary conditions

$$n_{m_j}^{(j)} \geq j + \frac{\epsilon_j}{2} \max\left(j - \frac{\epsilon_j}{2} + i + 1, 0\right) + 2j(m_{j+1/2} + m_{j+1} + \dots + m_{k-1/2}). \quad (4.30)$$

5. Conclusion

In this work, we have displayed a multi-parafermion basis of states for the \mathbb{Z}_k parafermionic models. The basis elements are in one-to-one correspondence with the set of $k - 1$ ordered partitions described in equations (1.13), (1.14) and (1.15). This is an alternative to the usual description of the basis in terms of partitions restricted by (1.5) [10, 2]. In the parafermionic context, the argued equivalence of the two bases leads us to the conclusion that the two sets of partitions, namely (1.13)–(1.15) and (1.5), are equinumerous. Clearly, finding a direct bijection would allow us to strip off this elementary derivation of $F_{k,i}$ from any parafermionic dressing. Moreover, such a bijection might point towards natural ‘higher-rank’ generalizations of the Andrews–Gordon identity.

As previously pointed out, the ‘new’ \mathbb{Z}_k basis has already been derived in [13]. We have thus emphasized here the novelty (and simplicity) of the conformal-field-theoretical derivation. As an original extension, two new bases of states for graded parafermions have been displayed. Each one leads to a distinct fermionic form of the graded-parafermion characters once the contribution of the fractional part of the parafermionic modes is reinserted. The expression

linked to the basis involving all parafermionic modes is new. It is interesting to see that for this basis, an unusual aspect of the representation theory of the graded parafermions (when compared to the standard \mathbb{Z}_k representation theory) plays a crucial role, which is that some graded-parafermionic modules have more than one level-1 descendant of relative charge 0.

This work offers another illustration of the non-uniqueness of the fermionic characters of the irreducible modules in a given model. Here, this is rooted in the non-uniqueness of the quasi-particle basis. There are indeed different choices for the spanning set of creation operators that are compatible with a description of the basis in terms of restriction rules akin to exclusion relations. For the \mathbb{Z}_k models, there are two choices: $\{\mathcal{A}^{(1)}\}$ and $\{\mathcal{A}^{(1)}, \dots, \mathcal{A}^{(k-1)}\}$. For the graded case, there are three such sets: $\{\tilde{\mathcal{A}}^{(1/2)}\}$, $\{\tilde{\mathcal{A}}^{(1/2)}, \tilde{\mathcal{A}}^{(1)}, \tilde{\mathcal{A}}^{(2)}, \dots, \tilde{\mathcal{A}}^{(k-1)}\}$ and $\{\tilde{\mathcal{A}}^{(1/2)}, \tilde{\mathcal{A}}^{(1)}, \tilde{\mathcal{A}}^{(3/2)}, \dots, \tilde{\mathcal{A}}^{(k-1/2)}\}$. These sets are not necessarily exhaustive since, for instance, one could possibly consider a choice where some parafermionic fields are ignored¹⁵, or even one involving a mixtures of selected parafermionic modes augmented by the addition the Virasoro or higher integer-spin field modes.

Acknowledgments

This work was supported by EPSRC (PJ) and NSERC (PM). We would like to thank O Warnaar for drawing to our attention some of the quoted references from the mathematical literature and, most importantly, for communicating to us the argument of the appendix.

Appendix. The analytic proof of (4.21)

The equality (4.21), that has been established here by a field-theoretical argument, can also be demonstrated by analytical methods [25]. The general argument would proceed by a simple extension of lemma A.1 of [23]. We will content ourselves with the consideration of the $k = 2$ case and indicate at the end how the analysis can be generalized to $k > 2$. Let us first replace m_j by m_{2j} :

$$G_{2,i}(z; q) = \sum_{m_1, m_2, m_3=0}^{\infty} \frac{q^{\frac{1}{2}(m_1+m_2+m_3)^2 + \frac{1}{2}(m_2+m_3)^2 + \frac{1}{2}m_3^2 + \frac{1}{2}(m_1+m_3) + (2-i)(m_2+m_3)} z^{m_1+2m_2+3m_3}}{(q)_{m_1}(q)_{m_2}(q)_{m_3}}. \tag{A.1}$$

Next, we replace m_2 by $m_2 - m_3$ and use the convention that $1/(q)_n = 0$ if $n < 0$

$$\begin{aligned} G_{2,i}(z; q) &= \sum_{m_1, m_2, m_3=0}^{\infty} \frac{q^{\frac{1}{2}(m_1+m_2)^2 + \frac{1}{2}m_2^2 + \frac{1}{2}m_3^2 + \frac{1}{2}(m_1+m_3) + (2-i)m_2} z^{m_1+2m_2+m_3}}{(q)_{m_1}(q)_{m_2-m_3}(q)_{m_3}} \\ &= \sum_{m_1, m_2=0}^{\infty} \frac{q^{\frac{1}{2}m_1(m_1+2m_2+1) + m_2^2 + (2-i)m_2} z^{m_1+2m_2}}{(q)_{m_1}(q)_{m_2}} \sum_{m_3=0}^{m_2} \frac{(q)_{m_2}}{(q)_{m_2-m_3}(q)_{m_3}} q^{\frac{1}{2}m_3(m_3+1)} z^{m_3}. \end{aligned} \tag{A.2}$$

Using the q -binomial theorem ([3], equation (3.3.6))

$$\sum_{j=0}^n \frac{(q)_n}{(q)_j(q)_{n-j}} q^{\frac{1}{2}j(j+1)} x^j = (-xq)_n, \tag{A.3}$$

we can perform the summation over m_3 and get

$$G_{2,i}(z; q) = \sum_{m_1, m_2=0}^{\infty} \frac{q^{\frac{1}{2}m_1(m_1+2m_2+1) + m_2^2 + (2-i)m_2} z^{m_1+2m_2}}{(q)_{m_1}(q)_{m_2}} (-zq)_{m_2}. \tag{A.4}$$

¹⁵ On the analytic side, the argument of the appendix shows clearly how to eliminate, from the generating function, an arbitrary set of modes associated with the graded parafermions ψ_r with r half-integer.

We next make use of the Euler relation ([3], equation (2.2.6)):

$$\sum_{n=0}^{\infty} \frac{q^{\frac{1}{2}n(n-1)} x^n}{(q)_n} = (-x)_{\infty}, \quad (\text{A.5})$$

to sum over m_1 with $x = zq^{m_2+1}$:

$$\begin{aligned} G_{2,i}(z; q) &= \sum_{m_2=0}^{\infty} (-zq^{m_2+1})_{\infty} (-zq)_{m_2} \frac{q^{m_2^2+(2-i)m_2} z^{2m_2}}{(q)_{m_2}} \\ &= (-zq)_{\infty} \sum_{m_2=0}^{\infty} \frac{q^{m_2^2+(2-i)m_2} z^{2m_2}}{(q)_{m_2}} = (-zq)_{\infty} F_{2,i}(z^2; q). \end{aligned} \quad (\text{A.6})$$

The generalization to $k > 2$ is straightforward. The odd modes m_{2j+1} for $j > 1$ are summed successively, starting from the largest one, by the q -binomial theorem, while the sum over m_1 is done by the Euler relation. The identity of theorem 4.5 in [23] is similarly related to the multiple sum of Andrews. With the suitable addition of a linear term, the latter is thus essentially equivalent to our (4.21).

References

- [1] Kedem R, Klassen T R, McCoy B M and Melzer E 1993 *Phys. Lett. B* **304** 263
Kedem R, Klassen T R, McCoy B M and Melzer E 1993 *Phys. Lett. B* **307** 68
Kedem R, McCoy B M and Melzer E 1995 *Recent Progress in Statistical Mechanics and Quantum Field Theory* ed P Bouwknegt *et al* (Singapore: World Scientific) p 195
- [2] Jacob P and Mathieu P 2002 *Nucl. Phys. B* **620** 351
- [3] Andrews G E 1984 *The Theory of Partitions* (Cambridge: Cambridge University Press)
- [4] Andrews G E 1981 *Houston J. Math.* **7** 11
- [5] Andrews G E 1974 *Proc. Natl Acad. Sci. USA* **71** 4082
- [6] Andrews G E 1977 *Am. J. Math.* **101** 735
- [7] Bressoud D 1987 Lattice paths and Rogers–Ramanujan identities *Number Theory, Madras 1987* ed K Alladi *Lect. Notes Math.* **1395** 140
- [8] Warnaar S O 1997 *Commun. Math. Phys.* **184** 203
- [9] Feigin B L, Nakanishi T and Ooguri H 1992 *Int. J. Mod. Phys. A* **7** (Suppl. 1A) 217
- [10] Lepowsky J and Primc M 1985 *Contemporary Mathematics* vol 46 (Providence, RI: American Mathematical Society)
- [11] Feigin B and Stoyanovsky A V 1993 Quasi-particles models for the representations of Lie algebras and geometry of flag manifold *Preprint hep-th/9308079*
Feigin B and Stoyanovsky A V 1994 *Funct. Anal. Appl.* **28** 55
- [12] Ardonne E, Kedem R and Stone M 2005 *J. Phys. A: Math. Gen.* **38** 617
- [13] Georgiev G 1996 *J. Pure Appl. Algebra* **112** 247
Georgiev G 1995 Combinatorial constructions of modules for infinite-dimensional Lie algebras: II. Parafermionic space *Preprint q-alg/9504024*
- [14] Zamolodchikov A B and Fateev V A 1985 *Sov. Phys.—JETP* **43** 215
- [15] Jacob P and Mathieu P 2000 *Nucl. Phys. B* **587** 514
- [16] Camino J M, Ramallo A V and Sanchez de Santos J M 1998 *Nucl. Phys. B* **530** 715
- [17] Jacob P and Mathieu P 2002 *Nucl. Phys. B* **630** 433
- [18] Bégin L, Fortin J-F, Jacob P and Mathieu P 2003 *Nucl. Phys. B* **659** 365
- [19] Fortin J-F, Jacob P and Mathieu P 2003 Jagged partitions *Preprint math.CO/0310079 (Ramanujan J. at press)*
- [20] Fortin J-F, Jacob P and Mathieu P 2005 *Electron. J. Comb.* **12** R12
- [21] Melzer E 1994 Supersymmetric analogs of the Gordon–Andrews identities, and related TBA systems *Preprint hep-th/9412154*
- [22] Bressoud D, Ismail M and Stanton D 2000 *Ramanujan J.* **4** 435
- [23] Warnaar S O 2003 *Discrete Math.* **272** 215
- [24] Fortin J-F, Jacob P and Mathieu P 2005 *J. Phys. A: Math. Gen.* **38** 1699
- [25] Warnaar S O 2005 Private communication