

## Supersymmetry for integrable hierarchies on loop superalgebras

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2005 J. Phys. A: Math. Gen. 38 9341

(<http://iopscience.iop.org/0305-4470/38/42/012>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.94

The article was downloaded on 03/06/2010 at 04:01

Please note that [terms and conditions apply](#).

# Supersymmetry for integrable hierarchies on loop superalgebras

H Aratyn<sup>1</sup>, J F Gomes<sup>2</sup>, G M de Castro<sup>2</sup>, M B Silka<sup>2</sup> and A H Zimerman<sup>2</sup>

<sup>1</sup> Department of Physics, University of Illinois at Chicago, 845 W. Taylor St., Chicago, IL 60607-7059, USA

<sup>2</sup> Instituto de Física Teórica-UNESP, Rua Pamplona 145, 01405-900 São Paulo, Brazil

Received 30 July 2005, in final form 1 August 2005

Published 5 October 2005

Online at [stacks.iop.org/JPhysA/38/9341](http://stacks.iop.org/JPhysA/38/9341)

## Abstract

An algebraic approach is employed to formulate  $N = 2$  supersymmetry transformations in the context of integrable systems based on loop superalgebras  $\widehat{\mathfrak{sl}}(p + 1, p)$ ,  $p \geq 1$ , with homogeneous gradation. We work with extended integrable hierarchies, which contain supersymmetric AKNS and Lund–Regge sectors. We derive the one-soliton solution for  $p = 1$  which solves positive and negative evolution equations of the  $N = 2$  supersymmetric model.

PACS numbers: 02.30.Ik, 11.10.Lm

## 1. Introduction

The well-known examples of the mKdV and sine-Gordon models [1] as well as the AKNS and the Lund–Regge (complex sine-Gordon) models [2] can be understood as positive and negative flows belonging to the same integrable hierarchy conveniently classified in terms of underlying algebraic structure. The basic objects in an algebraic approach to integrable models are the loop algebra  $\widehat{\mathcal{G}}$  endowed with a grading operator  $Q$  and a constant semisimple generator  $E$ . The grading decomposes the loop algebra into graded subspaces and specifies the space of physical fields.

There already exists an extensive literature devoted to construction of integrable models with extended supersymmetry based on superspace formalism, (see e.g. [3–12]). Here, a purely algebraic approach without superspace techniques is employed to construct a class of integrable hierarchies based on loop superalgebras  $\widehat{\mathfrak{sl}}(p + 1, p)$ ,  $p \geq 1$ , with homogeneous gradation. Within this framework the symmetry structure of the underlying model is realized as a graded subalgebra of  $\widehat{\mathfrak{sl}}(p + 1, p)$  defined as a centralizer  $\mathcal{K}$  of a semisimple element  $E$  ( $\mathcal{K} = \text{Ker}(E)$ ). The positive and negative grade elements in the centre of  $\mathcal{K}$  generate positive and negative evolution flows of the extended hierarchy. The positive flows are of the generalized AKNS type, while the first negative flow corresponds to the Lund–Regge

(relativistic) model. By comparing expressions for subsequent positive flows we derive the closed expression for the recursion operator  $\mathcal{R}$ .

The elements in  $\mathcal{K}$  not residing in the centre give rise to the non-abelian symmetry structure of the model. In particular,  $N = 2$  supersymmetry transformations arise in this context being generated by specific fermionic generators  $f_{\pm} \in \mathcal{K}$ . The supersymmetry algebra follows from the fact that the anti-commutator of  $f_{\pm}$  closes on  $E$  [3, 4].

Furthermore, the algebraic approach allows a systematic construction of Hamiltonians. We derive the Lagrangian density for the second flow and the canonical (Dirac) bracket which defines the first Poisson bracket structure of the model. Higher brackets can be obtained by successive applications of recursion operator.

We employ the dressing and vertex formalisms to construct soliton solutions. Explicit formulae for the one-soliton solution are found for  $p = 1$ . The formula is verified to be a solution of both the AKNS and Lund–Regge type of evolution equations.

This work completes the discussion started in [15], where integrable models based on loop superalgebras with principal gradation (like supersymmetric KdV model) were treated by algebraic methods.

Section 2 gives details of algebraic foundation of our construction. We write equations of motion governed by the second flow for the  $\widehat{\mathfrak{sl}}(p + 1, p)$  model for arbitrary  $p$  generalizing the results of [14]. Symmetries, conservation laws, recursion operator and the first Poisson bracket are presented within the algebraic framework in section 3. In section 4, we deal with the first negative flow of the extended hierarchy, which is shown to be invariant under  $N = 2$  supersymmetry. Next we write explicitly the Lund–Regge equations of motion (for the  $p = 1$ ) model using the Gauss decomposition of the zero grade subgroup element  $B$ . In section 5, we explicitly construct the one soliton solution by using the dressing and vertex operator techniques.

## 2. General construction of the model

Consider the graded Lie algebra  $\mathfrak{sl}(p + 1, s + 1)$  composed of bosonic (even)  $\alpha_i, \beta_t, i = 1, \dots, p, t = 1, \dots, s$ , and fermionic (odd),  $\gamma$ , simple roots:

$$\begin{aligned} \alpha_i &= e_i - e_{i+1}, & i &= 1, \dots, p, \\ \beta_t &= f_t - f_{t+1}, & t &= 1, \dots, s, \\ \gamma &= e_{p+1} - f_1 \end{aligned} \tag{1}$$

in terms of unit length vectors  $e_r, f_a$  such that  $e_r^2 = 1$  and  $f_a^2 = -1$  with  $r = 1, \dots, p + 1$  and  $a = 1, \dots, s + 1$ . The remaining bosonic roots can be written as

$$\pm(\alpha_r + \dots + \alpha_{q-1}) = \pm(e_r - e_q), \quad \pm(\beta_a + \dots + \beta_{b-1}) = \pm(f_a - f_b), \tag{2}$$

where  $r, q = 1, \dots, p + 1$  and  $a, b = 1, \dots, s + 1$ . The remaining fermionic roots have the form

$$\pm(\alpha_r + \dots + \alpha_{p+1} + \gamma + \beta_1 + \dots + \beta_{a-1}) = \pm(e_r - f_a). \tag{3}$$

Consider the homogeneous gradation with the grading operator  $Q = \lambda \frac{d\lambda}{d}$ . The affine super Lie algebra  $\widehat{\mathcal{G}} = \widehat{\mathfrak{sl}}(p + 1, s + 1)$  decomposes according to  $\widehat{\mathcal{G}} = \oplus \mathcal{G}_k$  with the graded subspaces  $\mathcal{G}_k$  that satisfy  $[Q, \mathcal{G}_k] = k\mathcal{G}_k$  with  $k \in \mathbb{Z}$  and consist of elements  $X^{(k)} = \lambda^k X$  for  $X \in \mathfrak{sl}(p + 1, s + 1)$ .

The key role in an algebraic approach to integrable models is played by the constant grade one generator

$$E = E^{(1)} \tag{4}$$

obtained by setting  $n = 1$  in the following expression for the semisimple generators of grade  $n$

$$E^{(n)} = \left( \sum_{i=1}^p e_i - \sum_{a=1}^{s+1} f_a \right) \cdot H^{(n)}.$$

The centralizer  $\mathcal{K} = \text{Ker}(E)$  is a subalgebra of  $\mathcal{G}$  spanned by the Cartan subalgebra of  $\text{sl}(p+1, s+1)$  and the step operators associated to roots

$$\begin{aligned} \pm(\alpha_i + \cdots + \alpha_j) &= \pm(e_i - e_j), & \pm(\beta_a + \cdots + \beta_{b-1}) &= \pm(f_a - f_b), \\ \pm(\alpha_i + \cdots + \alpha_p + \gamma + \beta_1 + \cdots + \beta_{a-1}) &= \pm(e_i - f_a), \\ i, j &= 1, \dots, p, & a, b &= 1, \dots, s+1. \end{aligned} \quad (5)$$

The image  $\mathcal{I} = \text{Im}(E)$  is then obtained as a linear combination of step operators of bosonic roots

$$\pm(\alpha_i + \cdots + \alpha_p) = \pm(e_i - e_{p+1}), \quad i = 1, \dots, p,$$

and fermionic roots

$$\pm(\gamma + \beta_1 + \cdots + \beta_{a-1}) = \pm(e_{p+1} - f_a), \quad a = 1, \dots, s+1,$$

respectively.

The fact that the element  $E$  is semisimple insures that  $\mathcal{G} = \mathcal{K} \oplus \mathcal{I}$ . In addition, it also holds that  $[\mathcal{I}, \mathcal{I}] \subset \mathcal{K}$ .

Because supersymmetry requires equal number of bosons and fermions, we impose from now on the condition  $p = s+1$  and denote  $\alpha_{p+1} = \gamma, \alpha_{p+2} = \beta_1, \alpha_{p+3} = \beta_2, \dots, \alpha_{2p} = \beta_{p-1}$ .

The superalgebra  $\widehat{\text{sl}}(p+1, p)$  gives rise to a supersymmetric integrable model specified by element  $E$  of equation (4) and the following Lax operator

$$L = \partial_x + E + A_0 = \partial_x + \mathcal{A}_x, \quad (6)$$

where  $\mathcal{A}_x = E + A_0$  and

$$\begin{aligned} A_0 &= \sum_{i=1}^p (\bar{b}_i E_{e_i - e_{p+1}}^{(0)} + b_i E_{-(e_i - e_{p+1})}^{(0)}) + \sum_{a=1}^p (\bar{F}_a E_{f_a - e_{p+1}}^{(0)} + F_a E_{-(f_a - e_{p+1})}^{(0)}) \\ &= \sum_{i=1}^p (\bar{b}_i E_{\alpha_i + \cdots + \alpha_p}^{(0)} + b_i E_{-(\alpha_i + \cdots + \alpha_p)}^{(0)}) + \sum_{a=1}^p (\bar{F}_a E_{-(\alpha_{p+1} + \cdots + \alpha_{p+a})}^{(0)} + F_a E_{\alpha_{p+1} + \cdots + \alpha_{p+a}}^{(0)}), \end{aligned} \quad (7)$$

lies in the grade zero sector of  $\mathcal{I}$ .

Now, we propose the zero curvature relation  $[\partial_{t_n} + \mathcal{A}_{t_n}, \partial_x + \mathcal{A}_x] = 0$ . We search for a solution of the form

$$\mathcal{A}_{t_n} = D^{(n)} + D^{(n-1)} + \cdots + D^{(0)}, \quad (8)$$

where  $D^{(k)} \in \mathcal{G}_k$ . The zero curvature relation decomposes grade by grade into the following chain of equations:

$$\begin{aligned} [E, D^{(n)}] &= 0, \\ -\partial_x D^{(n)} + [A_0, D^{(n)}] + [E, D^{(n-1)}] &= 0, \\ &\vdots \\ -\partial_x D^{(1)} + [A_0, D^{(1)}] + [E, D^{(0)}] &= 0, \\ -\partial_x D^{(0)} + \partial_{t_2} A_0 + [A_0, D^{(0)}] &= 0. \end{aligned} \quad (9)$$

The top equation in (9) implies that  $D^{(n)} \in \mathcal{K}$  and, consequently, we assume that  $D^{(n)} = E^{(n)}$ .

Solving equations (9) from the highest to the zero grade components we find for  $t_n = t_2$ , the following equations of motion for the coefficients of  $A_0$ :

$$\begin{aligned}
 \partial_{t_2} b_i + \partial_x^2 b_i - 2 \sum_{j=1}^p (b_j \bar{b}_j + F_j \bar{F}_j) b_i &= 0, \\
 \partial_{t_2} \bar{b}_i - \partial_x^2 \bar{b}_i + 2 \sum_{j=1}^p (b_j \bar{b}_j + F_j \bar{F}_j) \bar{b}_i &= 0, \\
 \partial_{t_2} F_i + \partial_x^2 F_i - 2 \sum_{j=1}^p (b_j \bar{b}_j + F_j \bar{F}_j) F_i &= 0, \\
 \partial_{t_2} \bar{F}_i - \partial_x^2 \bar{F}_i + 2 \sum_{j=1}^p (b_j \bar{b}_j + F_j \bar{F}_j) \bar{F}_i &= 0,
 \end{aligned} \tag{10}$$

where  $i = 1, \dots, p$ . This result generalizes equations of motion found in [13] in the  $p = 1$  case.

### 3. Symmetries and conservation laws

The symmetries of integrable models will be defined below in a systematic manner in the setting of a gauge transformation relating the bare Lax  $(\partial_x + E)$  to the dressed Lax operator  $L$  given by definition (6) [17].

The dressing operator is defined by

$$\Theta = e^{\theta^{(-1)} + \theta^{(-2)} + \dots}, \quad \theta^{(-k)} \in \mathcal{G}_{-k},$$

and satisfies the relation:

$$E = \Theta^{-1} (E + A_0) \Theta + \Theta^{-1} \partial_x \Theta. \tag{11}$$

Decomposing (11) according to the grading we find the following equations:

$$\begin{aligned}
 A_0 + [E, \theta^{(-1)}] &= 0, \\
 \partial_x \theta^{(-1)} + [E, \theta^{(-2)}] + \frac{1}{2} [A_0, \theta^{(-1)}] &= 0, \\
 \partial_x \theta^{(-2)} + [E^{(1)}, \theta^{(-3)}] + \frac{1}{2} [A_0, \theta^{(-2)}] + \frac{1}{12} [[A_0, \theta^{(-1)}], \theta^{(-1)}] &= 0, \\
 \partial_x \theta^{(-3)} + [E^{(1)}, \theta^{(-4)}] + \frac{1}{2} [A_0, \theta^{(-3)}] + \frac{1}{12} [[A_0, \theta^{(-1)}], \theta^{(-2)}] \\
 + \frac{1}{12} [[A_0, \theta^{(-2)}], \theta^{(-1)}] &= 0 \\
 \vdots &
 \end{aligned} \tag{12}$$

Decompose  $\theta^{(-i)}$  in its components in the centralizer  $\mathcal{K}$  and image  $\mathcal{I}$  according to  $\theta^{(-i)} = \theta_{\mathcal{K}}^{(-i)} + \theta_{\mathcal{I}}^{(-i)}$ . Then the top equation (12) yields

$$\theta_{\mathcal{I}}^{(-1)} = \sum_{i=1}^p (-\bar{b}_i E_{\alpha_i + \dots + \alpha_p}^{(0)} + b_i E_{-(\alpha_i + \dots + \alpha_p)}^{(0)} - \bar{F}_a E_{-(\alpha_{p+1} + \dots + \alpha_{p+i})}^{(0)} + F_a E_{\alpha_{p+1} + \dots + \alpha_{p+i}}^{(0)}). \tag{13}$$

The second equation (12) can be rewritten as

$$\partial_x \theta_{\mathcal{K}}^{(-1)} + \frac{1}{2} [A_0, \theta_{\mathcal{I}}^{(-1)}] = 0, \quad \partial_x \theta_{\mathcal{I}}^{(-1)} + [E, \theta_{\mathcal{I}}^{(-2)}] + \frac{1}{2} [A_0, \theta_{\mathcal{K}}^{(-1)}] = 0 \tag{14}$$

and can be used to determine a local expression for  $\theta_{\mathcal{I}}^{(-2)}$ .

An element  $X_m \in \text{Ker}(E)$  of  $m$ th grade generates a symmetry transformation via [17]:

$$\delta_{X_m} \Theta = (\Theta X_m \Theta^{-1})_- \Theta. \tag{15}$$

We will derive the corresponding transformation of  $A_0$  using that from the top relation in (12) which follows that

$$A_0 = [\theta^{(-1)}, E] = ([\Theta, E])_0, \quad (16)$$

where the right-hand side contains projection on the zero grade.

Employing the definition (15) of the symmetry transformations to the above relation we find

$$\begin{aligned} \delta_{X_m} A_0 &= ([(\Theta X_m \Theta^{-1})_{-1} \Theta, E])_0 = [(\Theta X_m \Theta^{-1})_{-1}, E] \\ &= ([\Theta X_m \Theta^{-1}, E])_0 = ([[\Theta, E] \Theta^{-1}, \Theta X_m \Theta^{-1}])_0. \end{aligned}$$

From equation (11) we find that

$$[\Theta, E] \Theta^{-1} = A_0 + \partial_x \Theta \Theta^{-1}$$

and, thus, transformation of  $A_0$  can be given by a general gauge transformation formula:

$$\delta_{X_n} A_0 = ([A_0 + \partial_x \Theta \Theta^{-1}, \Theta X_n \Theta^{-1}])_0, \quad (17)$$

where  $X_n$  is an element in  $\mathcal{K}$  of  $n$ th grade. For  $m = 0$ , we find

$$\delta_{X_0} A_0 = [A_0, X_0]. \quad (18)$$

For  $m = 1$ , equation (17) yields,

$$\delta_{X_1} A_0 = [A_0, [\theta^{(-1)}, X_1]] + [\partial_x \theta^{(-1)}, X_1]. \quad (19)$$

In particular, for  $X_1 = E$  we find that  $\delta_E A_0 = \partial_x A_0 = \partial A_0 / \partial t_1$ , in accordance with the fact that the centre of  $\mathcal{K}$  generates isospectral flows.

Define two odd elements of  $\text{Ker}(E)$ :

$$f_{\pm} = \sum_{i=1}^p (\epsilon^{(0)} E_{e_i - f_i} \pm \epsilon^{(1)} E_{-(e_i - f_i)}), \quad (20)$$

containing grade 0 and 1 generators and Grassmannian parameters  $\epsilon^{(0)}$  and  $\epsilon^{(1)}$ . These two odd elements satisfy the graded commutation relations:

$$[f_+, f_-] = -2\epsilon^{(0)} \epsilon^{(1)} E,$$

According to relation (16) the elements  $f_{\pm} \in \text{Ker}(E)$  give rise the symmetry transformations  $\delta_{f_{\pm}}$ . These transformations satisfy the basic  $N = 2$  supersymmetry relations

$$[\delta_{f_+}, \delta_{f_-}] A_0 = -2\epsilon^{(0)} \epsilon^{(1)} \partial_x A_0, \quad [\delta_{f_{\pm}}, \delta_{f_{\pm}}] A_0 = 0.$$

For the particular case of  $p = 1$ , i.e.  $\text{sl}(2, 1)$ , we find the supersymmetry transformations to be

$$\begin{aligned} \delta_{f_{\pm}} b_1 &= \pm \epsilon^{(1)} \left( \partial_x F_1 - b_1 \int \bar{b}_1 F_1 + F_1 \int (b_1 \bar{b}_1 + F_1 \bar{F}_1) \right), \\ \delta_{f_{\pm}} \bar{b}_1 &= -\epsilon^{(0)} \bar{F}_1 \pm \epsilon^{(1)} \bar{b}_1 \int \bar{b}_1 F_1, \\ \delta_{f_{\pm}} \bar{F}_1 &= \pm \epsilon^{(1)} \left( \partial_x \bar{b}_1 - \bar{b}_1 \int (b_1 \bar{b}_1 + F_1 \bar{F}_1) + \bar{F}_1 \int \bar{b}_1 F_1 \right), \\ \delta_{f_{\pm}} F_1 &= \epsilon^{(0)} b_1 \mp \epsilon^{(1)} F_1 \int \bar{b}_1 F_1. \end{aligned} \quad (21)$$

These are indeed symmetries of the equations of motion (10). For the  $\widehat{\text{sl}}(p+1, p)$  case, the supersymmetry transformation generated by  $f_{\pm}$  in (20) are explicitly listed in appendix A.

Let us now calculate the Hamiltonian densities. According to [17], the Hamiltonian densities are given by

$$\mathcal{H}_n = -\text{tr}(E^{(0)} A^{(-n)}) = \frac{1}{2} \sum_{k=0}^{n-1} \text{tr}(A^{(-k)} A^{(1+k-n)}), \tag{22}$$

where  $A^{(-n)}$  are defined via

$$\partial_x \Theta \Theta^{-1} = \sum_{k=1}^{\infty} A^{(-k)} \lambda^{-k},$$

where  $\partial_x \Theta \Theta^{-1}$  is the quantity, which entered equation (11). The symbol  $\text{tr}$  in expression (22) denotes a trace for the superalgebra  $\text{sl}(p+1, p)$ .

Now, we explicitly work out the first few expressions. Inserting  $n = 1$  in (22) we obtain

$$\mathcal{H}_1 = -\text{tr}(E^{(0)} A^{(-1)}) = \frac{1}{2} \text{tr}(A_0^2) = \sum_{i=1}^p (b_i \bar{b}_i + F_i \bar{F}_i), \tag{23}$$

where  $A_0$  is given by (7). In order to evaluate  $\mathcal{H}_2$  we use (13) to obtain

$$\mathcal{H}_2 = \text{tr}(A_0 \partial_x \theta^{(-1)}) = \sum_{i=1}^p (\bar{b}_i \partial_x b_i + \bar{F}_i \partial_x F_i). \tag{24}$$

For  $n = 3$ , we find

$$\mathcal{H}_3 = -\frac{1}{2} \text{tr}(2A_0 A^{(-2)} + A^{(-1)^2}), \tag{25}$$

where

$$A^{(-1)} = -\partial_x \theta^{(-1)}, \quad A^{(-2)} = \partial_x \theta^{(-2)} - \frac{1}{2} [\partial_x \theta^{(-1)}, \theta^{(-1)}] \tag{26}$$

and, therefore,

$$\frac{1}{3} \mathcal{H}_3 = \sum_{j=1}^p \partial_x \bar{b}_j \partial_x b_j + \sum_{a=1}^p \partial_x \bar{F}_a \partial_x F_a + \sum_{i,j=1}^p \bar{b}_j b_j \bar{b}_i b_i + 2 \sum_{i,b=1}^p \bar{b}_i b_i \bar{F}_b F_b + \sum_{a,b=1}^p \bar{F}_a F_a \bar{F}_b F_b. \tag{27}$$

### 3.1. Poisson structure of the second flow

The equations of motion (10) can be derived from the Lagrangian density

$$\begin{aligned} \mathcal{L}_{t_2} = & \frac{1}{2} \sum_{i=1}^p \partial_t b_i \bar{b}_i - \frac{1}{2} \sum_{i=1}^p b_i \partial_t \bar{b}_i - \frac{1}{2} \sum_{a=1}^p \partial_t F_a \bar{F}_a + \frac{1}{2} \sum_{a=1}^p F_a \partial_t \bar{F}_a - \sum_{j=1}^p \partial_x \bar{b}_j \partial_x b_j \\ & - \sum_{a=1}^p \partial_x \bar{F}_a \partial_x F_a - \sum_{i,j=1}^p \bar{b}_j b_j \bar{b}_i b_i - 2 \sum_{i,b=1}^p \bar{b}_i b_i \bar{F}_b F_b - \sum_{a,b=1}^p \bar{F}_a F_a \bar{F}_b F_b. \end{aligned} \tag{28}$$

The canonical momenta are given by

$$\begin{aligned} P_{b_i} &= \frac{\delta \mathcal{L}_{t_2}}{\delta \dot{b}_i} = \frac{1}{2} \bar{b}_i, & P_{\bar{b}_i} &= \frac{\delta \mathcal{L}_{t_2}}{\delta \dot{\bar{b}}_i} = -\frac{1}{2} b_i, \\ P_{F_a} &= \frac{\delta \mathcal{L}_{t_2}}{\delta \dot{F}_a} = -\frac{1}{2} \bar{F}_a, & P_{\bar{F}_a} &= \frac{\delta \mathcal{L}_{t_2}}{\delta \dot{\bar{F}}_a} = -\frac{1}{2} F_a, \end{aligned} \tag{29}$$

which satisfy the usual canonical Poisson brackets

$$\begin{aligned} \{P_{b_i}(x), b_j(y)\}_P &= \{P_{\bar{b}_i}(x), \bar{b}_j(y)\}_P = -\delta_{ij}\delta(x-y), \\ \{P_{F_a}(x), F_b(y)\}_P &= \{P_{\bar{F}_a}(x), \bar{F}_b(y)\}_P = -\delta_{ab}\delta(x-y). \end{aligned}$$

The momenta dependence on the fields in (29) define the primary constraints

$$\begin{aligned} \Phi_{b_i} &= P_{b_i} - \frac{1}{2}\bar{b}_i = 0, & \Phi_{\bar{b}_i} &= P_{\bar{b}_i} + \frac{1}{2}b_i = 0, \\ \Phi_{F_a} &= P_{F_a} + \frac{1}{2}\bar{F}_a = 0, & \Phi_{\bar{F}_a} &= P_{\bar{F}_a} + \frac{1}{2}F_a = 0, \end{aligned} \quad (30)$$

and accordingly the Poisson bracket algebra is given in terms of the Dirac brackets

$$\{b_i(x), \bar{b}_j(y)\}_D = -\delta_{ij}\delta(x-y), \quad \{F_a(x), \bar{F}_b(y)\}_D = -\delta_{ab}\delta(x-y), \quad (31)$$

which together with the total Hamiltonian  $H_3 = \int dx \mathcal{H}_3$  obtained from (27) reproduce the equations of motion (10) via formula  $\partial X/\partial t_2 = \{X, H_3\}_D$ .

A second bracket characterized by  $P_2$  can be obtained from the recursive relation

$$P_2 = \mathcal{R}P_1, \quad (32)$$

where  $P_1$  is the first bracket defined in (31) and  $\mathcal{R}$  denotes the recursion operator relating consecutive time evolutions, i.e.

$$\partial_{t_m} A_0 = [E, \partial_x \partial_{t_{m-1}} A_0] - [A_0, \partial_x^{-1} [A_0, [E, \partial_{t_{m-1}} A_0]]] = \mathcal{R} \partial_{t_{m-1}} A_0.$$

In components this recursion relation for flows reads as

$$\partial_{t_m} \xi_{i,l} = \sum_{k=1}^4 \sum_{j=1}^p \mathcal{R}_{i,k}^{l,j} \partial_{t_{m-1}} \xi_{k,j}, \quad i = 1, \dots, 4, \quad l = 1, \dots, p, \quad (33)$$

where the symbol  $\xi_{i,j}$  represents all the fields contained in  $A_0$  and is introduced in equation (13) through  $\xi_{1,j} = \bar{b}_j$ ,  $\xi_{2,j} = b_j$ ,  $\xi_{3,j} = F_j$  and  $\xi_{4,j} = \bar{F}_j$ . For the case of  $\mathfrak{sl}(2, 1)$  with  $p = 1$  one finds the pseudo-differential operator expressions for the recursion operator  $\mathcal{R}_{i,k} = \mathcal{R}_{i,k}^{1,1}$ :

$$\begin{aligned} \mathcal{R}_{11} &= \partial_x - 2\bar{b}\partial_x^{-1}b + \bar{F}\partial_x^{-1}F, & \mathcal{R}_{12} &= -2\bar{b}\partial_x^{-1}\bar{b}, \\ \mathcal{R}_{13} &= \bar{b}\partial_x^{-1}\bar{F} + \bar{F}\partial_x^{-1}\bar{b}, & \mathcal{R}_{14} &= -\bar{b}\partial_x^{-1}F, \\ \mathcal{R}_{21} &= 2b\partial_x^{-1}b, & \mathcal{R}_{22} &= -\partial_x + 2b\partial_x^{-1}\bar{b} + F\partial_x^{-1}\bar{F}, \\ \mathcal{R}_{23} &= b\partial_x^{-1}\bar{F}, & \mathcal{R}_{24} &= F\partial_x^{-1}b + b\partial_x^{-1}F, \\ \mathcal{R}_{31} &= F\partial_x^{-1}b + b\partial_x^{-1}F, & \mathcal{R}_{32} &= F\partial_x^{-1}\bar{b}, \\ \mathcal{R}_{33} &= -\partial_x + b\partial_x^{-1}\bar{b}, & \mathcal{R}_{34} &= 0, \\ \mathcal{R}_{41} &= -\bar{F}\partial_x^{-1}b, & \mathcal{R}_{42} &= -\bar{F}\partial_x^{-1}\bar{b} - \bar{b}\partial_x^{-1}\bar{F}, \\ \mathcal{R}_{43} &= 0, & \mathcal{R}_{44} &= \partial_x - \bar{b}\partial_x^{-1}b, \end{aligned}$$

where  $\partial_x$  and the pseudo-differential symbol  $\partial_x^{-1}$  are acting on all the fields appearing to their right.

In appendix B we give the general expression for  $\mathcal{R}$  in case of arbitrary  $\mathfrak{sl}(p+1, p)$ . These expressions allow an easy derivation of the second bracket.

#### 4. The relativistic model

Consider the negative grade time evolution equations given by

$$\partial_{t_{-j}} A_0 - \partial_x (D^{(-1)} + D^{(-2)} + \dots + D^{(-j)}) - [A_0 + E, D^{(-1)} + D^{(-2)} + \dots + D^{(-j)}] = 0. \quad (34)$$



Here, we only consider  $j = 1$ . We introduce notation  $z = t_{-1}$  and  $\bar{z} = -x$ . Thus,

$$\partial_z = \frac{\partial t_{-1}}{\partial}, \quad \bar{\partial} = \frac{\partial \bar{z}}{\partial} = -\partial_x,$$

and the above evolution equation becomes

$$\partial_z A_0 + \bar{\partial} D^{(-1)} - [A_0 + E, D^{(-1)}] = 0,$$

with the following solution:

$$D^{(-1)} = B E^{(-1)} B^{-1}, \quad A_0 = \bar{\partial} B B^{-1}, \tag{35}$$

where  $B$  is the group element of  $sl(p + 1, p)$  satisfying Leznov–Saveliev’s equations

$$\bar{\partial}(B^{-1} \partial_z B) + [E^{(-1)}, B^{-1} E B] = 0, \quad \partial_z(\bar{\partial} B B^{-1}) + [B E^{(-1)} B^{-1}, E] = 0. \tag{36}$$

It is customary to think about coordinates  $z, \bar{z}$  as light-cone coordinates. Such association allows us to interpret the model as being relativistic.

The group element  $B$  transforms under the symmetry transformations (15) as

$$\delta_{X_n} B = -(\Theta X_n \Theta^{-1})_0 B, \quad n \geq 0. \tag{37}$$

For  $n = 0$  and  $n = 1$  the above relation takes the form

$$\delta_{X_0} B = -X_0 B, \quad \delta_{X_1} B = [X_1, \theta^{(-1)}] B. \tag{38}$$

In particular, for  $X_1 = E$  the above definition yields

$$\delta_E B = -(\Theta E \Theta^{-1})_0 B = -[\theta^{(-1)}, E] B = -A_0 B.$$

Recalling that  $A_0 = -\partial_x B B^{-1}$  we obtain  $\delta_E B = \partial_x B$ , confirming again that  $E$  generates the first positive isospectral flow  $t_1 = x$ .

These transformations leave the above equations of motion (36) invariant. In particular, the equations of motion (36) remain unchanged under the  $N = 2$  supersymmetry transformations  $\delta_{f_{\pm}}$ .

Note that equations (36) yield chiral currents associated to the Kernel of  $E$ , i.e.  $\mathcal{K} = \{X \in sl(p + 1, p), [X, E] = 0\}$ . In order to make connection with the super AKNS model (7), i.e.  $A_0 \in \mathcal{I}$  in (35), we impose the following subsidiary constraints:

$$\text{tr}(X \bar{\partial} B B^{-1}) = \text{tr}(X B^{-1} \partial_z B) = 0, \quad X \in \mathcal{K}. \tag{39}$$

#### 4.1. The relativistic super Lund–Regge model

In order to define the relativistic member of the so-called super Lund–Regge hierarchy we consider the zero grade group element  $B \in G_0 = SL(2, 1)$  parametrized as

$$B = e^{\tilde{\chi} E_{-\alpha_1}} e^{\tilde{f}_1 E_{-\alpha_1 - \alpha_2}} e^{\tilde{f}_2 E_{\alpha_2}} e^{\frac{1}{2} \varphi_1 (\alpha_1 + \alpha_2) \cdot H + \frac{1}{2} \varphi_2 (\alpha_2) \cdot H} e^{\tilde{g}_2 E_{-\alpha_2}} e^{\tilde{g}_1 E_{\alpha_1 + \alpha_2}} e^{\tilde{\psi} E_{\alpha_1}}, \tag{40}$$

where we are using the basis  $E_{\pm \alpha_1}, E_{\pm \alpha_2}, E_{\pm(\alpha_1 + \alpha_2)}, \alpha_1 \cdot H$  and  $\alpha_2 \cdot H$ . Changing to the natural variables

$$\begin{aligned} \tilde{\psi} &= \psi e^{-\frac{\varphi_1 + \varphi_2}{2}}, & \tilde{g}_1 &= g_1 e^{-\frac{\varphi_2}{2}}, & \tilde{f}_1 &= f_1 e^{-\frac{\varphi_2}{2}} \\ \tilde{\chi} &= \chi e^{-\frac{\varphi_1 + \varphi_2}{2}}, & \tilde{g}_2 &= g_2 e^{-\frac{\varphi_1}{2}}, & \tilde{f}_2 &= f_2 e^{-\frac{\varphi_1}{2}} \end{aligned} \tag{41}$$

we find for the currents

$$\begin{aligned} J &= B^{-1} \partial_z B = J_{-\alpha_1} E_{\alpha_1} + J_{-\alpha_2} E_{\alpha_2} + J_{-\alpha_1 - \alpha_2} E_{\alpha_1 + \alpha_2} + J_{(\alpha_1 + \alpha_2) \cdot H} (\alpha_1 + \alpha_2) \cdot H \\ &\quad - J_{\alpha_2 \cdot H} \alpha_2 \cdot H + J_{\alpha_1} E_{-\alpha_1} + J_{\alpha_2} E_{-\alpha_2} + J_{\alpha_1 + \alpha_2} E_{-(\alpha_1 + \alpha_2)}, \\ \bar{J} &= \bar{\partial} B B^{-1} = \bar{J}_{-\alpha_1} E_{\alpha_1} + \bar{J}_{-\alpha_2} E_{\alpha_2} + \bar{J}_{-\alpha_1 - \alpha_2} E_{\alpha_1 + \alpha_2} + \bar{J}_{(\alpha_1 + \alpha_2) \cdot H} (\alpha_1 + \alpha_2) \cdot H \\ &\quad - \bar{J}_{\alpha_2 \cdot H} \alpha_2 \cdot H + \bar{J}_{\alpha_1} E_{-\alpha_1} + \bar{J}_{\alpha_2} E_{-\alpha_2} + \bar{J}_{\alpha_1 + \alpha_2} E_{-(\alpha_1 + \alpha_2)} \end{aligned} \tag{42}$$

with the coefficients  $J_{-\alpha_1}, \dots, \bar{J}_{\alpha_1 + \alpha_2}$  shown in appendix D.

The constraints (39)

$$J_{\pm(\alpha_1+\alpha_2)} = \bar{J}_{\pm(\alpha_1+\alpha_2)} = J_{(\alpha_1+\alpha_2)\cdot H} = \bar{J}_{(\alpha_1+\alpha_2)\cdot H} = J_{\alpha_2\cdot H} = \bar{J}_{\alpha_2\cdot H} = 0 \quad (43)$$

lead to the following equations for the non-local fields:

$$\begin{aligned} \partial_z f_1 &= \frac{1}{2} f_1 \partial_z \varphi_2 + g_2 \left[ \partial_z \chi - \frac{1}{2} \chi (\partial_z \varphi_1 + \partial_z \varphi_2) \right], \\ \partial_z g_1 &= \psi \partial_z f_2 + \frac{1}{2} g_1 \partial_z \varphi_2 - \frac{1}{2} \psi f_2 \partial_z \varphi_1, \\ \bar{\partial} f_1 &= \chi \bar{\partial} g_2 + \frac{1}{2} f_1 \bar{\partial} \varphi_2 - \frac{1}{2} \chi g_2 \bar{\partial} \varphi_1, \\ \bar{\partial} g_1 &= \frac{1}{2} g_1 \bar{\partial} \varphi_2 + f_2 \left[ \bar{\partial} \psi - \frac{1}{2} \psi (\bar{\partial} \varphi_1 + \bar{\partial} \varphi_2) \right], \\ \partial_z \varphi_1 &= \frac{\psi \left[ \partial_z \chi (1 + g_2 f_2) + \frac{1}{2} \chi g_2 \partial_z f_2 \right]}{1 + \psi \chi (1 + \frac{5}{4} g_2 f_2)}, \\ \partial_z \varphi_2 &= \frac{\psi \partial_z \chi (1 + \frac{3}{2} g_2 f_2) - g_2 \partial_z f_2 - \frac{1}{2} \psi \chi g_2 \partial_z f_2}{1 + \psi \chi (1 + \frac{5}{4} g_2 f_2)}, \\ \bar{\partial} \varphi_1 &= \frac{\chi \left[ \bar{\partial} \psi (1 + g_2 f_2) + \frac{1}{2} \psi \bar{\partial} g_2 f_2 \right]}{1 + \psi \chi (1 + \frac{5}{4} g_2 f_2)}, \\ \bar{\partial} \varphi_2 &= \frac{\chi \bar{\partial} \psi (1 + \frac{3}{2} g_2 f_2) + (\frac{1}{2} \psi \chi + 1) f_2 \bar{\partial} g_2}{1 + \psi \chi (1 + \frac{5}{4} g_2 f_2)}. \end{aligned} \quad (44)$$

The equations of motion are given in the Leznov–Saveliev form (36), after eliminating the non-local fields with the help of equation (44). In components they read

$$\begin{aligned} (1 + g_2 f_2) \left[ \bar{\partial} \partial_z \chi - \frac{1}{2} \bar{\partial} \chi (\partial_z \varphi_1 + \partial_z \varphi_2) - \frac{1}{2} \chi (\bar{\partial} \partial_z \varphi_1 + \bar{\partial} \partial_z \varphi_2) + \frac{1}{2} (\bar{\partial} \varphi_1 + \bar{\partial} \varphi_2) \right. \\ \left. \times (\partial_z \chi - \frac{1}{2} \chi (\partial_z \varphi_1 + \partial_z \varphi_2)) \right] + \bar{\partial} (g_2 f_2) \left[ \partial_z \chi - \frac{1}{2} \chi (\partial_z \varphi_1 + \partial_z \varphi_2) \right] + \chi = 0, \end{aligned} \quad (45)$$

$$\bar{\partial} \partial_z g_2 + \frac{1}{2} \bar{\partial} g_2 \partial_z \varphi_1 + \frac{1}{2} g_2 \bar{\partial} \partial_z \varphi_1 - \frac{1}{2} \partial_z g_2 \bar{\partial} \varphi_1 - \frac{1}{4} g_2 \bar{\partial} \varphi_1 \partial_z \varphi_1 + (1 + \psi \chi) g_2 = 0 \quad (46)$$

$$\begin{aligned} (1 + g_2 f_2) \left[ \partial_z \bar{\partial} \psi - \frac{1}{2} \partial_z \psi (\bar{\partial} \varphi_1 + \bar{\partial} \varphi_2) - \frac{1}{2} \psi (\partial_z \bar{\partial} \varphi_1 + \partial_z \bar{\partial} \varphi_2) + \frac{1}{2} (\partial_z \varphi_1 + \partial_z \varphi_2) \right. \\ \left. \times (\bar{\partial} \psi - \frac{1}{2} \psi (\bar{\partial} \varphi_1 + \bar{\partial} \varphi_2)) \right] + \partial_z (g_2 f_2) \left[ \bar{\partial} \psi - \frac{1}{2} \psi (\bar{\partial} \varphi_1 + \bar{\partial} \varphi_2) \right] + \psi = 0 \end{aligned} \quad (47)$$

$$\partial_z \bar{\partial} f_2 + \frac{1}{2} \partial_z f_2 \bar{\partial} \varphi_1 + \frac{1}{2} f_2 \partial_z \bar{\partial} \varphi_1 - \frac{1}{2} \bar{\partial} f_2 \partial_z \varphi_1 - \frac{1}{4} f_2 \partial_z \varphi_1 \bar{\partial} \varphi_1 + (1 + \psi \chi) f_2 = 0, \quad (48)$$

where the non-local fields  $\varphi_1$  and  $\varphi_2$  are given in equation (44).

#### 4.2. Connection between AKNS and Lund–Regge variables

On the basis of (35) we find the following non-local relations between the AKNS and the relativistic Lund–Regge variables:

$$\begin{aligned} \bar{b}_1 &= \bar{J}_{-\alpha_1} = \frac{e^{\frac{1}{2}(\varphi_1+\varphi_2)}}{1+f_2g_2} \left( \bar{\partial} \psi - \frac{1}{2} \psi (\bar{\partial} \varphi_1 + \bar{\partial} \varphi_2) \right), \\ F_1 &= \bar{J}_{-\alpha_2} = e^{-\frac{1}{2}\varphi_1} \left( \bar{\partial} f_2 + \frac{1}{2} f_2 \bar{\partial} \varphi_1 \right), \\ b_1 &= \bar{J}_{\alpha_1} = e^{-\frac{1}{2}(\varphi_1+\varphi_2)} \left( \bar{\partial} \chi + \frac{1}{2} \chi (\bar{\partial} \varphi_1 + \bar{\partial} \varphi_2) - \chi f_2 \bar{\partial} g_2 - \frac{1}{2} \chi \bar{\partial} \varphi_1 g_2 f_2 \right. \\ &\quad \left. + e^{\frac{1}{2}\varphi_1} f_1 \bar{J}_{-\alpha_2} - \chi^2 e^{-\frac{1}{2}(\varphi_1+\varphi_2)} \bar{J}_{-\alpha_1} \right), \\ \bar{F}_1 &= \bar{J}_{\alpha_2} = e^{\frac{1}{2}\varphi_1} \left( \bar{\partial} g_2 - \frac{1}{2} g_2 \bar{\partial} \varphi_1 + e^{-\frac{1}{2}(\varphi_1+\varphi_2)} f_1 \bar{J}_{-\alpha_1} \right). \end{aligned} \quad (49)$$

Note that they are all non-local fields in terms of the Lund–Regge variables due to the presence of  $\varphi_1, \varphi_2$  and  $f_1$  defined by (44). Also, we have seen that in terms of the AKNS variables, the models are invariant under supersymmetry transformations (21).

### 5. Soliton solutions

#### 5.1. Dressing transformations and vertex operators

The dressing transformation and the vertex operators method represent a powerful tool for the construction of solitons solutions of integrable models. The dressing transformation relates two solutions of the equations of motion written in the zero curvature representation. In particular, it relates the vacuum and the 1-soliton solutions by a gauge transformation,

$$A_\mu = \Theta_\pm A_\mu^{\text{vac}} \Theta_\pm^{-1} - (\partial_\mu \Theta_\pm) \Theta_\pm^{-1}, \tag{50}$$

where

$$\Theta_- = e^{t(-1)+\dots}, \quad \Theta_+ = e^{v(0)} e^{v(1)} \dots \tag{51}$$

The zero curvature representation implies for pure gauge solutions:

$$A_\mu^{\text{vac}} = -\partial_\mu T_0 T_0^{-1}, \quad A_\mu = -\partial_\mu T T^{-1} \tag{52}$$

which allows the following relation:

$$T = \Theta_\pm T_0, \quad \text{i.e. } \Theta_+ T = \Theta_- T_0 g, \tag{53}$$

where  $g \in \hat{G}$  is an arbitrary constant element of the corresponding affine group. Suppose  $T_0$  represents the vacuum solution,

$$T_0 = \exp(-t_n E^{(n)}) \exp(-x E), \quad n > 1, \tag{54}$$

i.e.,

$$A_{t_n}^{\text{vac}} = E^{(n)}, \quad A_x^{\text{vac}} = E. \tag{55}$$

As a consequence of (50) with (55) and (51) we can determine  $\Theta_\pm$ . Consider for instance equation (50) for  $A_x$  and  $\Theta_-$ . It determines its zero grade component  $t(-1)$  through

$$A_0 = [t(-1), E]. \tag{56}$$

Note that if we enlarge the loop algebra by adding central terms, then consistency requires the introduction of a additional field  $\nu$  associated with the extension  $A_0 \rightarrow A_0 + \partial_x \nu \hat{c}$  in constructing the Lax operator. For the specific  $\mathfrak{sl}(2, 1)$  model:

$$A_0 = \bar{b}_1 E_{\alpha_1}^{(0)} + b_1 E_{-\alpha_1}^{(0)} + F_1 E_{\alpha_2}^{(0)} + \bar{F}_1 E_{-\alpha_2}^{(0)} + \partial_x \nu \hat{c} \tag{57}$$

we find

$$t(-1) = -\bar{b}_1 E_{\alpha_1}^{(-1)} + b_1 E_{-\alpha_1}^{(-1)} + F_1 E_{\alpha_2}^{(-1)} - \bar{F}_1 E_{-\alpha_2}^{(-1)} + \partial_x \nu \frac{1}{2} (\alpha_1 + \alpha_2) \cdot H^{(-1)}. \tag{58}$$

From (53) we find the solution

$$\begin{aligned} \tau_0 &\equiv e^v = \langle \lambda_0 | T_0 g T_0^{-1} | \lambda_0 \rangle, \\ \tau_1 &\equiv \bar{b}_2 \tau_0 = \langle \lambda_0 | E_{-\alpha_1}^{(1)} T_0 g T_0^{-1} | \lambda_0 \rangle, \\ \tau_2 &\equiv b_2 \tau_0 = -\langle \lambda_0 | E_{\alpha_1}^{(1)} T_0 g T_0^{-1} | \lambda_0 \rangle, \\ \tau_3 &\equiv F_1 \tau_0 = \langle \lambda_0 | E_{-\alpha_2}^{(1)} T_0 g T_0^{-1} | \lambda_0 \rangle, \\ \tau_4 &\equiv \bar{F}_1 \tau_0 = \langle \lambda_0 | E_{\alpha_1}^{(1)} T_0 g T_0^{-1} | \lambda_0 \rangle. \end{aligned} \tag{59}$$

For the relativistic model let us consider equation (50) for  $A_x$  and  $\theta_+$ . We find

$$A_0 = -\partial_x e^{v(0)} e^{-v(0)}. \quad (60)$$

Since, for the affine case,  $A_0 = -\partial_x B B^{-1} - \partial_x v \hat{c}$ ,

$$e^{v(0)} = B e^{v \hat{c}}. \quad (61)$$

From (53) we find the solution

$$\langle \lambda_r | B e^{v \hat{c}} | \lambda_t \rangle = \langle \lambda_r | T_0 g T_0^{-1} | \lambda_t \rangle. \quad (62)$$

We now choose  $|\lambda_r\rangle, |\lambda_t\rangle$  to be the following weight states

$$|\rho\rangle = \left| \frac{1}{2} \alpha_1 \right\rangle, \quad |\epsilon\rangle = \left| -\frac{1}{2} \alpha_1 - \alpha_2 \right\rangle \quad (63)$$

and obtain

$$\begin{aligned} \langle \rho | B e^{v \hat{c}} | \rho \rangle &= \tau_0 e^{\frac{1}{2}(\varphi_1 + \varphi_2)} = \langle \rho | T_0 g T_0^{-1} | \rho \rangle, \\ \langle \epsilon | B e^{v \hat{c}} | \epsilon \rangle &= \tau_0 e^{\frac{1}{2}(\varphi_1 - \varphi_2)} = \langle \epsilon | T_0 g T_0^{-1} | \epsilon \rangle, \\ \langle \rho | B e^{v \hat{c}} E_{-\alpha_1 - \alpha_2} | \rho \rangle &= \frac{1}{2} \tilde{g}_1 \tau_0 e^{\frac{1}{2}(\varphi_1 + \varphi_2)} = \langle \rho | T_0 g T_0^{-1} E_{-\alpha_1 - \alpha_2} | \rho \rangle, \\ \langle \rho | B e^{v \hat{c}} E_{\alpha_2} | \rho \rangle &= -\frac{1}{2} \tilde{g}_2 \tau_0 e^{\frac{1}{2}(\varphi_1 + \varphi_2)} = \langle \rho | T_0 g T_0^{-1} E_{\alpha_2} | \rho \rangle, \\ \langle \rho | B e^{v \hat{c}} E_{-\alpha_1} | \rho \rangle &= \tau_0 e^{\frac{1}{2}(\varphi_1 + \varphi_2)} \left( \tilde{\psi} + \frac{1}{2} \tilde{g}_2 \tilde{g}_1 \right) = \langle \rho | T_0 g T_0^{-1} E_{-\alpha_1} | \rho \rangle, \\ \langle \rho | E_{\alpha_1 + \alpha_2} B e^{v \hat{c}} | \rho \rangle &= \frac{1}{2} \tilde{f}_1 \tau_0 e^{\frac{1}{2}(\varphi_1 + \varphi_2)} = \langle \rho | E_{\alpha_1 + \alpha_2} T_0 g T_0^{-1} | \rho \rangle, \\ \langle \rho | E_{-\alpha_2} B e^{v \hat{c}} | \rho \rangle &= -\frac{1}{2} \tilde{f}_2 \tau_0 e^{\frac{1}{2}(\varphi_1 + \varphi_2)} = \langle \rho | E_{-\alpha_2} T_0 g T_0^{-1} | \rho \rangle, \\ \langle \rho | E_{\alpha_1} B e^{v \hat{c}} | \rho \rangle &= \tau_0 e^{\frac{1}{2}(\varphi_1 + \varphi_2)} \left( \tilde{\psi} + \frac{1}{2} \tilde{g}_2 \tilde{g}_1 \right) = \langle \rho | E_{\alpha_1} T_0 g T_0^{-1} | \rho \rangle. \end{aligned} \quad (64)$$

### 5.2. Vertex operators and solutions

In order to evaluate the matrix elements in the r.h.s. of equations (59) and (64) it is instructive to introduce the eigenvectors  $(\mathcal{F}(\gamma))$  of  $E^m$ , i.e.,

$$[E^{(m)}, \mathcal{F}_i(\gamma)] = f_i^m(\gamma) \mathcal{F}_i(\gamma), \quad (65)$$

such that we can classify the soliton solutions in terms of the constant group element  $g = e^{\mathcal{F}_1(\gamma_1)} e^{\mathcal{F}_2(\gamma_2)} \dots$  and hence

$$T_0 g T_0^{-1} = \exp(\rho_1(\gamma_1) \mathcal{F}_1(\gamma_1) + \rho_2(\gamma_2) \mathcal{F}_2(\gamma_2) + \dots), \quad (66)$$

where  $\rho_i(\gamma_i) = \exp(-t_n f_i^n - x f_i^1)$ .

For  $\mathcal{G} = \mathfrak{sl}(2, 1)$  model,  $E^{(1)} = (\alpha_1 + \alpha_2) \cdot H^{(1)}$  and its eigenstates are given by

$$\begin{aligned} \mathcal{F}_1(\gamma) &= \sum_{n \in \mathbb{Z}} (E_{\alpha_1}^{(n)} + a_1 E_{-\alpha_2}^{(n)}) \gamma^{-n}, \\ \mathcal{F}_2(\gamma) &= \sum_{n \in \mathbb{Z}} (E_{-\alpha_1}^{(n)} + a_2 E_{\alpha_2}^{(n)}) \gamma^{-n}, \end{aligned} \quad (67)$$

with eigenvalues  $f_1^m(\gamma) = \gamma^m$ ,  $f_2^m(\gamma) = -\gamma^{-m}$ . We now consider soliton solutions associated to  $g = \exp(\mathcal{F}_1(\gamma_1)) \exp(\mathcal{F}_2(\gamma_2))$  and

$$T_0 g T_0^{-1} = (1 + \rho_1(\gamma_1) \mathcal{F}_1(\gamma_1)) (1 + \rho_2(\gamma_2) \mathcal{F}_2(\gamma_2)) \quad (68)$$

leading to the following matrix elements:

$$\begin{aligned}
 \tau_0 &= \langle 0|T_0gT_0^{-1}|0\rangle = 1 + \Gamma\rho'_1\rho'_2, \\
 \tau_1 &= \langle 0|E_{-\alpha_1}^{(1)}T_0gT_0^{-1}|0\rangle = -\gamma_2\rho'_2, \\
 \tau_2 &= \langle 0|T_0gT_0^{-1}E_{\alpha_1}^{(-1)}|0\rangle = \gamma_1\rho'_1, \\
 \tau_3 &= \langle 0|E_{-\alpha_2}^{(1)}T_0gT_0^{-1}|0\rangle = -a_2\gamma_2\rho'_2, \\
 \tau_4 &= \langle 0|T_0gT_0^{-1}E_{\alpha_2}^{(-1)}|0\rangle = a_1\gamma_1\rho'_1,
 \end{aligned}
 \tag{69}$$

where  $b = a_1a_2$ ,  $\Gamma_0 = \frac{\gamma_1\gamma_2}{(\gamma_1-\gamma_2)^2}$  and  $\Gamma = (1 - b)\Gamma_0$ . We therefore find the solution for the non-relativistic model (C.7),

$$\begin{aligned}
 \bar{b}_1 &= \frac{\gamma_1\rho'_1}{1 + \Gamma\rho'_1\rho'_2}, & b_1 &= -\frac{\gamma_2\rho'_2}{1 + \Gamma\rho'_1\rho'_2}, \\
 \bar{F}_1 &= -a_2\frac{\gamma_2\rho'_2}{1 + \Gamma\rho'_1\rho'_2}, & \bar{F}_1 &= a_1\frac{\gamma_1\rho'_1}{1 + \Gamma\rho'_1\rho'_2},
 \end{aligned}
 \tag{70}$$

where for the non-relativistic model  $\rho'_1(\gamma_1) = e^{-t_2\gamma_1^2 - \gamma_1x}$ ,  $\rho'_2(\gamma_1) = e^{t_2\gamma_2^2 + \gamma_2x}$ .

For the relativistic model described by equations of motion (45)–(48) we find the following matrix elements:

$$\begin{aligned}
 \langle \rho|T_0gT_0^{-1}|\rho\rangle &= 1 + \rho_1\rho_2\frac{\gamma_1}{(\gamma_1 - \gamma_2)^2} \left( \gamma_1 - \frac{1}{2}(\gamma_1 + \gamma_2)b \right), \\
 \langle \epsilon|T_0gT_0^{-1}|\epsilon\rangle &= 1 + \rho_1\rho_2\frac{\gamma_1}{(\gamma_1 - \gamma_2)^2} \left( \gamma_1 + \frac{1}{2}(\gamma_1 - 3\gamma_2)b \right), \\
 \langle \rho|T_0gT_0^{-1}E_{-\alpha_1}^{(0)}|\rho\rangle &= \rho_1, \\
 \langle \rho|E_{\alpha_1}^{(0)}T_0gT_0^{-1}|\rho\rangle &= \rho_2, \\
 \langle \rho|T_0gT_0^{-1}E_{\alpha_2}^{(0)}|\rho\rangle &= -\frac{1}{2}a_1\rho_1, \\
 \langle \rho|E_{-\alpha_2}^{(0)}T_0gT_0^{-1}|\rho\rangle &= -\frac{1}{2}a_2\rho_2, \\
 \langle \rho|T_0gT_0^{-1}E_{-\alpha_1-\alpha_2}^{(0)}|\rho\rangle &= \frac{1}{2}a_1\rho_1\rho_2\frac{\gamma_1}{(\gamma_1 - \gamma_2)}, \\
 \langle \rho|E_{\alpha_1+\alpha_2}^{(0)}T_0gT_0^{-1}|\rho\rangle &= \frac{1}{2}a_1\rho_2\frac{\gamma_1}{(\gamma_1 - \gamma_2)},
 \end{aligned}
 \tag{71}$$

which allow us to determine the solution

$$\begin{aligned}
 e^{\frac{1}{2}(\varphi_1+\varphi_2)} &= \frac{1 + \frac{\gamma_1}{\gamma_2}\Gamma_0\left(1 - b\frac{(\gamma_1+\gamma_2)}{2\gamma_1}\right)\rho_1\rho_2}{\tau_0}, \\
 e^{\frac{1}{2}(\varphi_1-\varphi_2)} &= \frac{1 + \Gamma_0\left(1 + b\frac{(\gamma_1-3\gamma_2)}{2\gamma_2}\right)\rho_1\rho_2}{\tau_0} = 1 - \frac{b(\gamma_2 - \gamma_1)\Gamma_0\rho_1\rho_2}{2\gamma_2(1 + \Gamma_0\rho_1\rho_2)}, \\
 \psi &= \frac{\rho_1}{\tau_0} \left( 1 - \frac{b\gamma_1\rho_1\rho_2}{2(\gamma_1 - \gamma_2)(1 + \frac{\gamma_1}{\gamma_2}\Gamma_0\rho_1\rho_2)} \right), \\
 \chi &= \frac{\rho_2}{\tau_0} \left( 1 - \frac{b\gamma_1\rho_1\rho_2}{2(\gamma_1 - \gamma_2)(1 + \frac{\gamma_1}{\gamma_2}\Gamma_0\rho_1\rho_2)} \right), \\
 g_1 &= a_2\frac{\gamma_1\rho_1\rho_2}{(\gamma_1 - \gamma_2)\tau_0} e^{-\frac{1}{2}\varphi_1}, & f_1 &= a_1\frac{\gamma_1\rho_1\rho_2}{(\gamma_1 - \gamma_2)\tau_0} e^{-\frac{1}{2}\varphi_1}, \\
 g_2 &= a_1\frac{\rho_1}{\tau_0} e^{-\frac{1}{2}\varphi_2}, & f_2 &= a_2\frac{\rho_2}{\tau_0} e^{-\frac{1}{2}\varphi_2}.
 \end{aligned}
 \tag{72}$$

From the solutions given in (72) it is easy to verify the consistency of the constraints (39) (or (44)). Let us take for instance

$$\bar{\partial} f_1 = \chi \bar{\partial} g_2 + \frac{1}{2} f_1 \bar{\partial} \varphi_2 - \frac{1}{2} g_2 \chi \bar{\partial} \varphi_1. \quad (73)$$

Substituting (72) in both sides of equation (73) and identifying  $t_{-1} = z$ ,  $x = -\bar{z}$  using the second equation (72) together with the fact that  $a_1 b = a_2 b = 0$  we find that the constraint (73) is satisfied. The same can be shown for all equations (44) (i.e. all constraints (39) are satisfied).

Finally we verify the connection between the non-relativistic and the relativistic variables given by equations (49). Substituting the solutions (72) in (49) and comparing with (70) we conclude that they agree if we associate

$$\gamma^2 t_2 \rightarrow \frac{z}{\gamma} = \frac{t_{-1}}{\gamma}, \quad x \rightarrow -\bar{z} \quad (74)$$

which implies  $\rho(\gamma) \rightarrow \rho'(\gamma)$ .

### Acknowledgments

HA, GMC and MBS acknowledge support from Fapesp. HA thanks the IFT-UNESP Institute for its hospitality. JFG and AHZ thank CNPq for a partial support.

### Appendix A. Supersymmetry transformation for $\widehat{\mathfrak{sl}}(p+1, p)$

Let  $f_{\pm}$  be the supersymmetry generators from equation (20). Using the symmetry transformations given in equations (18) and (19) and

$$\theta_I^{(-1)} = \sum_{i=1}^p (-\bar{b}_i E_{(e_i - e_{p+1})}^{(-1)} + b_i E_{-(e_i - e_{p+1})}^{(-1)}) + \sum_{a=1}^p (-\bar{F}_a E_{(f_a - e_{p+1})}^{(-1)} + F_a E_{-(f_a - e_{p+1})}^{(-1)}) \quad (A.1)$$

and

$$\begin{aligned} \partial_x \theta_K^{(-1)} = & - \sum_{j=1, j \neq i}^p \sum_{i=1}^p \bar{b}_j b_i E_{(e_i - e_j)}^{(-1)} - \sum_{j=1}^p \bar{b}_j b_j (e_j - e_{p+1}) \cdot H^{(-1)} \\ & - \sum_{j=1}^p \sum_{a=1}^p \bar{b}_j F_a E_{(e_i - f_a)}^{(-1)} - \sum_{j=1}^p \sum_{a=1}^p b_j \bar{F}_a E_{-(e_i - f_a)}^{(-1)} \\ & - \sum_{a=1, a \neq b}^p \sum_{b=1}^p \bar{F}_a F_b E_{(f_a - f_b)}^{(-1)} + \sum_{a=1}^p \bar{F}_a F_a (f_a - e_{p+1}) \cdot H^{(-1)} \end{aligned} \quad (A.2)$$

obtained from equation (12), we find

$$\begin{aligned} \delta_{f_{\pm}} b_r &= \pm \left( \partial_x F_r - \sum_{j=1}^p \left( b_j \int \bar{b}_j F_r - F_j \int (b_r \bar{b}_j + F_r \bar{F}_j) \right) \right) \epsilon^{(1)}, \\ \delta_{f_{\pm}} \bar{b}_r &= -\bar{F}_r \epsilon^{(0)} \pm \left( \sum_{j=1}^p \bar{b}_j \int \bar{b}_r F_j \right) \epsilon^{(1)}, \end{aligned}$$

$$\begin{aligned} \delta_{f_{\pm}} \bar{F}_r &= \pm \left( \partial_x \bar{b}_r - \left( \sum_{j=1}^p \bar{b}_j \int (b_j \bar{b}_r + F_j \bar{F}_r) \right) + \bar{F}_r \int \bar{b}_r F_r \right) \epsilon^{(1)}, \\ \delta_{f_{\pm}} F_r &= b_r \epsilon^{(0)} \mp \sum_{j=1}^p F_j \int \bar{b}_j F_r \epsilon^{(1)}. \end{aligned} \tag{A.3}$$

**Appendix B. Recursion operator for  $\widehat{\mathfrak{sl}}(p+1, p)$**

Using the same notation as in equation (33) we write down expression for the recursion pseudo-differential operator  $\mathcal{R}_{i,k}^{lj}$  valid for a general case of  $\widehat{\mathfrak{sl}}(p+1, p)$ . Here, the  $l, j = 1, \dots, p$  and indices  $i, k = 1, 2, 3, 4$  label the four modes  $\bar{b}_j, b_j, F_j, \bar{F}_j$ .

$$\begin{aligned} \mathcal{R}_{11}^{lj} &= \delta_{ij} \left( \partial_x - \sum_k \bar{b}_k \partial_x^{-1} b_k + \sum_b \bar{F}_b \partial_x^{-1} F_b \right) - \bar{b}_l \partial_x^{-1} b_j, \\ \mathcal{R}_{12}^{lj} &= -\bar{b}_j \partial_x^{-1} \bar{b}_l - \bar{b}_l \partial_x^{-1} \bar{b}_j, \\ \mathcal{R}_{13}^{lj} &= \bar{b}_l \partial_x^{-1} \bar{F}_j + \bar{F}_j \partial_x^{-1} \bar{b}_l, \\ \mathcal{R}_{14}^{lj} &= -\bar{b}_l \partial_x^{-1} F_j, \\ \mathcal{R}_{21}^{lj} &= b_j \partial_x^{-1} b_l + b_l \partial_x^{-1} b_j, \\ \mathcal{R}_{22}^{lj} &= \delta_{ij} \left( -\partial_x + \sum_k b_k \partial_x^{-1} \bar{b}_k + \sum_b F_b \partial_x^{-1} \bar{F}_b \right) + b_l \partial_x^{-1} \bar{b}_j, \\ \mathcal{R}_{23}^{lj} &= b_l \partial_x^{-1} \bar{F}_j, \\ \mathcal{R}_{24}^{lj} &= F_j \partial_x^{-1} b_l + b_l \partial_x^{-1} F_j, \\ \mathcal{R}_{31}^{lj} &= F_l \partial_x^{-1} b_j + b_j \partial_x^{-1} F_l, \\ \mathcal{R}_{32}^{lj} &= F_l \partial_x^{-1} \bar{b}_j, \\ \mathcal{R}_{33}^{lj} &= \delta_{ij} \left( -\partial_x + \sum_k b_k \partial_x^{-1} \bar{b}_k + \sum_b F_b \partial_x^{-1} \bar{F}_b \right) - F_l \partial_x^{-1} \bar{F}_j, \\ \mathcal{R}_{34}^{lj} &= F_l \partial_x^{-1} F_j - F_j \partial_x^{-1} F_l, \\ \mathcal{R}_{41}^{lj} &= -\partial_x^{-1} b_j \bar{F}_l, \\ \mathcal{R}_{42}^{lj} &= -\bar{F}_l \partial_x^{-1} \bar{b}_j - \bar{b}_j \partial_x^{-1} \bar{F}_l, \\ \mathcal{R}_{43}^{lj} &= -\bar{F}_j \partial_x^{-1} \bar{F}_l + \bar{F}_l \partial_x^{-1} \bar{F}_j, \\ \mathcal{R}_{44}^{lj} &= \delta_{ij} \left( \partial_x - \sum_k \bar{b}_k \partial_x^{-1} b_k + \sum_b \bar{F}_b \partial_x^{-1} F_b \right) - \bar{F}_l \partial_x^{-1} F_j. \end{aligned}$$

**Appendix C. Case of  $\widehat{\mathfrak{sl}}(3, 2)$**

Consider the  $\widehat{\mathfrak{sl}}(3, 2)$  case where

$$A_0 = b_1 E_{-\alpha_1 - \alpha_2}^{(0)} + \bar{b}_1 E_{\alpha_1 + \alpha_2}^{(0)} + b_2 E_{-\alpha_2}^{(0)} + \bar{b}_2 E_{\alpha_2}^{(0)} + F_1 E_{\alpha_3}^{(0)} + \bar{F}_1 E_{-\alpha_3}^{(0)} + F_2 E_{\alpha_3 + \alpha_4}^{(0)} + \bar{F}_2 E_{-\alpha_3 - \alpha_4}^{(0)} \tag{C.1}$$

and propose the zero curvature representation

$$\partial_{t_2} A_0 - \partial_x (D^{(0)} + D^{(1)} + D^{(2)}) - [A_0 + E, D^{(0)} + D^{(1)} + D^{(2)}] = 0 \quad (\text{C.2})$$

from where we find the following solutions for equations (9):

$$\begin{aligned} D^{(2)} &= (D^{(2)})_{\text{Ker}} = (e_1 + e_2 - f_1 - f_2) \cdot H^{(2)}, \\ D^{(1)} &= (D^{(1)})_{\text{IM}} = b_1 E_{-\alpha_1 - \alpha_2}^{(1)} + b_2 E_{-\alpha_2}^{(1)} + \bar{b}_1 E_{\alpha_1 + \alpha_2}^{(1)} + \bar{b}_2 E_{\alpha_2}^{(1)} \\ &\quad + F_1 E_{\alpha_3}^{(1)} + F_2 E_{\alpha_3 + \alpha_4}^{(1)} + \bar{F}_1 E_{-\alpha_3}^{(1)} + \bar{F}_2 E_{-\alpha_3 - \alpha_4}^{(1)} \\ (D^{(0)})_{\text{IM}} &= -\partial_x b_1 E_{-\alpha_1 - \alpha_2}^{(0)} - \partial_x b_2 E_{-\alpha_2}^{(0)} + \partial_x \bar{b}_1 E_{\alpha_1 + \alpha_2}^{(0)} + \partial_x \bar{b}_2 E_{\alpha_2}^{(0)} \\ &\quad - \partial_x F_1 E_{\alpha_3}^{(0)} - \partial_x F_2 E_{\alpha_3 + \alpha_4}^{(0)} + \partial_x \bar{F}_1 E_{-\alpha_3}^{(0)} + \partial_x \bar{F}_2 E_{-\alpha_3 - \alpha_4}^{(0)} \\ (D^{(0)})_{\text{Ker}} &= -(b_1 \bar{b}_1) h_1^{(0)} - (b_1 \bar{b}_1 + b_2 \bar{b}_2) h_2^{(0)} + (F_1 \bar{F}_1 + F_2 \bar{F}_2) h_3^{(0)} + (F_2 \bar{F}_2) h_4^{(0)} - (b_1 \bar{b}_2) E_{-\alpha_1}^{(0)} \\ &\quad - (b_1 \bar{F}_1) E_{-\alpha_1 - \alpha_2 - \alpha_3}^{(0)} - (b_1 \bar{F}_2) E_{-\alpha_1 - \alpha_2 - \alpha_3 - \alpha_4}^{(0)} - (b_2 \bar{b}_1) E_{\alpha_1}^{(0)} - (b_2 \bar{F}_1) E_{-\alpha_2 - \alpha_3}^{(0)} \\ &\quad - (b_2 \bar{F}_2) E_{-\alpha_2 - \alpha_3 - \alpha_4}^{(0)} - (\bar{b}_1 F_1) E_{\alpha_1 + \alpha_2 + \alpha_3}^{(0)} - (\bar{b}_1 F_2) E_{\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4}^{(0)} - (\bar{b}_2 F_1) E_{\alpha_1 + \alpha_2}^{(0)} \\ &\quad - (\bar{b}_2 F_2) E_{\alpha_2 + \alpha_3 + \alpha_4}^{(0)} - (F_1 \bar{F}_2) E_{-\alpha_4}^{(0)} - (F_2 \bar{F}_1) E_{\alpha_4}^{(0)} \end{aligned} \quad (\text{C.3})$$

and the equations of motion

$$\begin{aligned} \partial_{t_2} b_1 + \partial_x^2 b_1 - 2(b_1 \bar{b}_1 + b_2 \bar{b}_2 + F_1 \bar{F}_1 + F_2 \bar{F}_2) b_1 &= 0 \\ \partial_{t_2} b_2 + \partial_x^2 b_2 - 2(b_1 \bar{b}_1 + b_2 \bar{b}_2 + F_1 \bar{F}_1 + F_2 \bar{F}_2) b_2 &= 0 \\ \partial_{t_2} \bar{b}_1 - \partial_x^2 \bar{b}_1 + 2(b_1 \bar{b}_1 + b_2 \bar{b}_2 + F_1 \bar{F}_1 + F_2 \bar{F}_2) \bar{b}_1 &= 0 \\ \partial_{t_2} \bar{b}_2 - \partial_x^2 \bar{b}_2 + 2(b_1 \bar{b}_1 + b_2 \bar{b}_2 + F_1 \bar{F}_1 + F_2 \bar{F}_2) \bar{b}_2 &= 0 \\ \partial_{t_2} F_1 + \partial_x^2 F_1 - 2(b_1 \bar{b}_1 + b_2 \bar{b}_2 + F_2 \bar{F}_2) F_1 &= 0 \\ \partial_{t_2} F_2 + \partial_x^2 F_2 - 2(b_1 \bar{b}_1 + b_2 \bar{b}_2 + F_1 \bar{F}_1) F_2 &= 0 \\ \partial_{t_2} \bar{F}_1 - \partial_x^2 \bar{F}_1 + 2(b_1 \bar{b}_1 + b_2 \bar{b}_2 + F_2 \bar{F}_2) \bar{F}_1 &= 0 \\ \partial_{t_2} \bar{F}_2 - \partial_x^2 \bar{F}_2 + 2(b_1 \bar{b}_1 + b_2 \bar{b}_2 + F_1 \bar{F}_1) \bar{F}_2 &= 0. \end{aligned} \quad (\text{C.4})$$

Consider

$$\begin{aligned} X_0 &= \epsilon_{-\alpha_2 - \alpha_3} E_{\alpha_2 + \alpha_3}^{(0)} + \epsilon_{\alpha_2 + \alpha_3} E_{-\alpha_2 - \alpha_3}^{(0)} + \epsilon_{-\alpha_2 - \alpha_3 - \alpha_4} E_{\alpha_2 + \alpha_3 + \alpha_4}^{(0)} \\ &\quad + \epsilon_{\alpha_2 + \alpha_3 + \alpha_4} E_{-\alpha_2 - \alpha_3 - \alpha_4}^{(0)} + \epsilon_{-\alpha_1 - \alpha_2 - \alpha_3} E_{\alpha_1 + \alpha_2 + \alpha_3}^{(0)} + \epsilon_{\alpha_1 + \alpha_2 + \alpha_3} E_{-\alpha_1 - \alpha_2 - \alpha_3}^{(0)} \\ &\quad + \epsilon_{-\alpha_1 - \alpha_2 - \alpha_3 - \alpha_4} E_{\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4}^{(0)} + \epsilon_{\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4} E_{-\alpha_1 - \alpha_2 - \alpha_3 - \alpha_4}^{(0)} \end{aligned} \quad (\text{C.5})$$

leading from (18) to the supersymmetry transformations

$$\begin{aligned} \delta_{X_0} b_1 &= \epsilon_{\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4} F_2 + \epsilon_{\alpha_1 + \alpha_2 + \alpha_3} F_1, \\ \delta_{X_0} b_2 &= \epsilon_{\alpha_2 + \alpha_3 + \alpha_4} F_2 + \epsilon_{\alpha_2 + \alpha_3} F_1, \\ \delta_{X_0} \bar{b}_1 &= \epsilon_{-\alpha_1 - \alpha_2 - \alpha_3 - \alpha_4} \bar{F}_2 + \epsilon_{-\alpha_1 - \alpha_2 - \alpha_3} \bar{F}_1, \\ \delta_{X_0} \bar{b}_2 &= \epsilon_{-\alpha_2 - \alpha_3 - \alpha_4} \bar{F}_2 + \epsilon_{-\alpha_2 - \alpha_3} \bar{F}_1, \\ \delta_{X_0} F_1 &= -\epsilon_{-\alpha_2 - \alpha_3} b_2 - \epsilon_{-\alpha_1 - \alpha_2 - \alpha_3} b_1, \\ \delta_{X_0} F_2 &= -\epsilon_{-\alpha_1 - \alpha_2 - \alpha_3 - \alpha_4} b_1 - \epsilon_{-\alpha_2 - \alpha_3 - \alpha_4} b_2, \\ \delta_{X_0} \bar{F}_1 &= \epsilon_{\alpha_2 + \alpha_3} \bar{b}_2 + \epsilon_{\alpha_1 + \alpha_2 + \alpha_3} \bar{b}_1, \\ \delta_{X_0} \bar{F}_2 &= \epsilon_{\alpha_1 + \alpha_2 + \alpha_3 + \alpha_4} \bar{b}_1 + \epsilon_{\alpha_2 + \alpha_3 + \alpha_4} \bar{b}_2. \end{aligned} \quad (\text{C.6})$$



Of course when  $b_1 = \bar{b}_1 = 0$  and  $F_2 = \bar{F}_2 = 0$  we recover the  $p = 1$  or  $SL(2, 1)$  case proposed in [14] corresponding to  $A_0 = \bar{b}_2 E_{\alpha_1} + b_2 E_{-\alpha_1} + F_1 E_{\alpha_2} + \bar{F}_1 E_{-\alpha_2}$ , i.e.,

$$\begin{aligned} \partial_{t_2} b_2 + \partial_x^2 b_2 - 2(b_2 \bar{b}_2 + F_1 \bar{F}_1) b_2 &= 0 \\ \partial_{t_2} \bar{b}_2 - \partial_x^2 \bar{b}_2 + 2(b_2 \bar{b}_2 + F_1 \bar{F}_1) \bar{b}_2 &= 0 \\ \partial_{t_2} F_1 + \partial_x^2 F_1 - 2b_2 \bar{b}_2 F_1 &= 0 \\ \partial_{t_2} \bar{F}_1 - \partial_x^2 \bar{F}_1 + 2b_2 \bar{b}_2 \bar{F}_1 &= 0 \end{aligned} \tag{C.7}$$

**Appendix D.  $SL(2, 1)$  super-currents**

The components of the  $sl(2, 1)$  currents  $J = B^{-1} \partial B$  and  $\bar{J} = \bar{\partial} B B^{-1}$  given in equation (42) read in terms of the fields defined in (40) and (41):

$$\begin{aligned} J_{\alpha_1} &= e^{\frac{1}{2}(\varphi_1 + \varphi_2)} (\partial_z \chi - \frac{1}{2} \chi (\partial_z \varphi_1 + \partial_z \varphi_2) + \partial_z f_1 f_2 - \frac{1}{2} f_1 f_2 \partial_z \varphi_2), \\ J_{\alpha_2} &= e^{-\frac{1}{2}\varphi_1} (\partial g_2 + \frac{1}{2} g_2 \partial \varphi_1 + \partial f_1 \psi - \frac{1}{2} f_1 \partial \varphi_2 \psi - \psi g_2 e^{-\frac{1}{2}(\varphi_1 + \varphi_2)} J_{\alpha_1}), \\ J_{\alpha_1 + \alpha_2} &= e^{\frac{1}{2}\varphi_2} (\partial f_1 - \frac{1}{2} f_1 \partial \varphi_2 - g_2 e^{-\frac{1}{2}(\varphi_1 + \varphi_2)} J_{\alpha_1}), \\ J_{\alpha_2 \cdot H} &= \partial \varphi_2 - \partial f_2 g_2 + \frac{1}{2} f_2 g_2 \partial \varphi_1 - \psi e^{-\frac{1}{2}(\varphi_1 + \varphi_2)} J_{\alpha_1}, \\ J_{(\alpha_1 + \alpha_2) \cdot H} &= \partial \varphi_1 + \partial f_1 g_1 - \frac{1}{2} f_1 g_1 \partial \varphi_2 - (\psi + g_2 g_1) e^{-\frac{1}{2}(\varphi_1 + \varphi_2)} J_{\alpha_1}, \\ J_{-\alpha_1 - \alpha_2} &= e^{\frac{1}{2}\varphi_2} (\partial g_1 - \frac{1}{2} g_1 \partial \varphi_2 - \psi \partial f_2 + \frac{1}{2} \psi \partial \varphi_1 f_2 + g_1 J_{\alpha_2 \cdot H}), \\ J_{-\alpha_1} &= e^{-\frac{1}{2}(\varphi_1 + \varphi_2)} (\partial \psi + \frac{1}{2} \psi (\partial \varphi_1 + \partial \varphi_2) + \psi g_2 \partial f_2 \\ &\quad - \frac{1}{2} \psi \partial \varphi_1 g_2 f_2 + g_1 e^{\frac{1}{2}\varphi_1} J_{\alpha_2} - \psi^2 e^{-\frac{1}{2}(\varphi_1 + \varphi_2)} J_{\alpha_1}), \\ J_{-\alpha_2} &= e^{\frac{1}{2}\varphi_1} (\partial f_2 - \frac{1}{2} f_2 \partial \varphi_1 + g_1 e^{\frac{1}{2}\varphi_1} J_{\alpha_1}), \end{aligned}$$

and

$$\begin{aligned} \bar{J}_{-\alpha_1} &= e^{\frac{1}{2}(\varphi_1 + \varphi_2)} (\bar{\partial} \psi - \frac{1}{2} \psi (\bar{\partial} \varphi_1 + \bar{\partial} \varphi_2) + g_2 \bar{\partial} g_1 - \frac{1}{2} g_2 g_1 \partial_z \varphi_2), \\ \bar{J}_{-\alpha_2} &= e^{-\frac{1}{2}\varphi_1} (\bar{\partial} f_2 + \frac{1}{2} f_2 \bar{\partial} \varphi_1 + \bar{\partial} g_1 \chi - \frac{1}{2} g_1 \bar{\partial} \varphi_2 \chi - \chi f_2 e^{-\frac{1}{2}(\varphi_1 + \varphi_2)} \bar{J}_{-\alpha_1}), \\ \bar{J}_{-\alpha_1 - \alpha_2} &= e^{\frac{1}{2}\varphi_2} (\bar{\partial} g_1 - \frac{1}{2} g_1 \bar{\partial} \varphi_2 - f_2 e^{-\frac{1}{2}(\varphi_1 + \varphi_2)} \bar{J}_{-\alpha_1}), \\ \bar{J}_{\alpha_2 \cdot H} &= \bar{\partial} \varphi_2 + \bar{\partial} g_2 f_2 + \frac{1}{2} f_2 g_2 \bar{\partial} \varphi_1 - \chi e^{-\frac{1}{2}(\varphi_1 + \varphi_2)} \bar{J}_{-\alpha_1}, \\ \bar{J}_{(\alpha_1 + \alpha_2) \cdot H} &= \bar{\partial} \varphi_1 - \bar{\partial} g_1 f_1 - \frac{1}{2} f_1 g_1 \bar{\partial} \varphi_2 - (\chi + f_1 f_2) e^{-\frac{1}{2}(\varphi_1 + \varphi_2)} \bar{J}_{-\alpha_1}, \\ \bar{J}_{\alpha_1 + \alpha_2} &= e^{-\frac{1}{2}\varphi_2} (\bar{\partial} f_1 - \frac{1}{2} f_1 \bar{\partial} \varphi_2 - \chi \bar{\partial} g_2 + \frac{1}{2} \chi \bar{\partial} \varphi_1 g_2 + f_1 \bar{J}_{\alpha_2 \cdot H}), \\ \bar{J}_{\alpha_1} &= e^{-\frac{1}{2}(\varphi_1 + \varphi_2)} (\bar{\partial} \chi + \frac{1}{2} \chi (\bar{\partial} \varphi_1 + \bar{\partial} \varphi_2) - \chi f_2 \bar{\partial} g_2 - \frac{1}{2} \chi \bar{\partial} \varphi_1 g_2 f_2 \\ &\quad + f_1 e^{\frac{1}{2}\varphi_1} \bar{J}_{-\alpha_2} - \chi^2 e^{-\frac{1}{2}(\varphi_1 + \varphi_2)} \bar{J}_{-\alpha_1}), \\ \bar{J}_{\alpha_2} &= e^{\frac{1}{2}\varphi_1} (\bar{\partial} g_2 - \frac{1}{2} g_2 \bar{\partial} \varphi_1 + f_1 e^{-\frac{1}{2}(\varphi_1 + \varphi_2)} \bar{J}_{-\alpha_1}). \end{aligned} \tag{D.1}$$

**References**

[1] Chodos A 1980 *Phys. Rev. D* **21** 2818  
 Tracy C A and Widom H 1996 *Commun. Math. Phys.* **179** 1 (Preprint solv-int/9506006)  
 Ferreira L A, Miramontes J L and Guillén J S 1997 *J. Math. Phys.* **38** 882 (Preprint hep-th/9606066)  
 Dorfmeister J, Gradl H and Szmigielski J 1998 *Acta Applicandae Math.* **53** 1  
 Fioravanti D and Stanishkov M 2000 *Nucl. Phys. B* **591** 685 (Preprint hep-th/0005158)

- [2] Aratyn H, Ferreira L A, Gomes J F and Zimerman A H 2000 *J. Phys. A: Math. Gen.* **33** L331 (Preprint nlin.si/0007002)  
Kamchatnov A M and Pavlov M V 2002 *Phys. Lett. A* **301** 269 (Preprint nlin.si/0208025)
- [3] Delduc F and Gallot L 1998 *J. Math. Phys.* **39** 4729 (Preprint solv-int/9802013)
- [4] Madsen J O and Miramontes J L 2001 *Commun. Math. Phys.* **217** 249 (Preprint hep-th/9905103)
- [5] Popowicz Z 1996 *J. Phys. A: Math. Gen.* **29** 1281 (Preprint hep-th/9510185)
- [6] Sorin A S and Kersten P H M 2002 *Lett. Math. Phys.* **60** 135 (Preprint nlin.si/0201026)
- [7] Delduc F and Sorin A S *Proc. 2002 Workshop on Integrable Theories, Solitons and Duality* ed L A Ferreira, J F Gomes and A H Zimerman *J. High Energy PRHEP-unesp2002/044* (Preprint nlin.si/0206037)
- [8] Nissimov E and Pacheva S 2002 *J. Math. Phys.* **43** 2547 (Preprint nlin.si/0103055)
- [9] Ivanov E, Krivonos S and Toppan F 1999 *Mod. Phys. Lett. A* **14** 2673 (Preprint solv-int/9912003)
- [10] Delduc F, Gallot L and Sorin A 1999 *Nucl. Phys. B* **558** 545 (Preprint solv-int/9907004)
- [11] Delduc F and Gallot L 1997 *Commun. Math. Phys.* **190** 395 (Preprint solv-int/9609008)
- [12] Ivanov E and Krivonos S 1997 *Phys. Lett. A* **231** 75 (Preprint hep-th/9609191)
- [13] Aratyn H and Rasinariu C 1997 *Phys. Lett. B* **391** 99 (Preprint hep-th/9608107)  
Aratyn H, Das A and Rasinariu C 1997 *Mod. Phys. Lett. A* **12** 2623 (Preprint solv-int/9704119)
- [14] Aratyn H, Das A, Rasinariu C and Zimerman A H 1998 *Supersymmetry and Integrable Models* ed H Aratyn *et al* (Berlin: Springer) (*Lecture Notes in Physics* vol 502)
- [15] Aratyn H, Gomes J F and Zimerman A H 2004 *Nucl. Phys. B* **676** 537 (Preprints hep-th/0309099, hep-th/0409171)  
Aratyn H, Gomes J F and Zimerman A H *Contribution to the 11th Int. Conf. on Symmetry Methods in Physics (SYMPHYS-11) (Prague)* (Preprint hep-th/0408231)
- [16] Inami T and Kanno H 1991 *Commun. Math. Phys.* **136** 519
- [17] Aratyn H, Gomes J F, Nissimov E, Pacheva S and Zimerman A H 2001 Symmetry flows, conservation laws and dressing approach to integrable models *Integrable Hierarchies and Modern Physical Theories* ed H Aratyn and A Sorin (Dordrecht: Kluwer) pp 243–75