

Fractal diffusion coefficient from dynamical zeta functions

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2006 J. Phys. A: Math. Gen. 39 L151

(<http://iopscience.iop.org/0305-4470/39/10/L01>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.101

The article was downloaded on 03/06/2010 at 04:13

Please note that [terms and conditions apply](#).

LETTER TO THE EDITOR

Fractal diffusion coefficient from dynamical zeta functions

Giampaolo Cristadoro

Max Planck Institute for the Physics of Complex Systems, Nöthnitzer Str. 38, D 01187 Dresden, Germany

E-mail: giampo@mpipks-dresden.mpg.de

Received 28 September 2005, in final form 31 January 2006

Published 22 February 2006

Online at stacks.iop.org/JPhysA/39/L151

Abstract

Dynamical zeta functions provide a powerful method to analyse low-dimensional dynamical systems when the underlying symbolic dynamics is under control. On the other hand, even simple one-dimensional maps can show an intricate structure of the grammar rules that may lead to a non-smooth dependence of global observables on parameters changes. A paradigmatic example is the fractal diffusion coefficient arising in a simple piecewise linear one-dimensional map of the real line. Using the Baladi–Ruelle generalization of the Milnor–Thurston kneading determinant, we provide the exact dynamical zeta function for such a map and compute the diffusion coefficient from its smallest zero.

PACS number: 05.45.–a

(Some figures in this article are in colour only in the electronic version)

Chaotic dynamic forces move the attention from individual trajectory to evolution of densities. The study of the spectrum of the evolution operator, the Perron–Frobenius operator, is a well-established method for the analytic treatment of hyperbolic dynamical systems. In particular, dynamical averages can be extracted from the spectra of the generalized transfer operator: this approach was successfully used to compute physical quantities like, e.g., the diffusion coefficient for extended hyperbolic dynamical systems or escape rates for open systems [1]. Relaxing the assumption of full hyperbolicity is the natural step forward: lots of efforts are still devoted to understanding how to extend the assumptions of strong chaoticity to include the presence of regions of regular motion into a meaningful periodic orbit expansion. Along this line, it was possible to show how to treat systems with marginally stable periodic orbits [2]. In such a case, marginality has a deep effect on the analytic structure of zeta functions and the system shows peculiar behaviour like anomalous diffusion and power law decay of correlations [3].

Even without relaxing the hyperbolicity assumption a strong limitation on the practical implementation of the periodic orbits programme relies on the need of control over periodic orbits proliferation: the number of periodic orbits grows exponentially with the period (at a rate given by the topological entropy) and an analytical treatment requires the control over such a huge amount of information. Cycle expansions technique prevents the explosion of the theory organizing cycles into fundamental terms and curvature corrections (constructed with longer cycles and their shadowing by shorter orbits) that comes out to be a good perturbative expansion [1]. The main ingredient is then a good understanding of the cycles that build the fundamental and curvature terms that implies a good control over the symbolic dynamic of the system. Unfortunately, the missing of such a control is the normal situation rather than the exception. Even for a very simple one-dimensional map without complete branches such a control is typically impossible. As a paradigmatic example, we will use a simple one-dimensional linear map on the real line where it was shown [4] that the diffusion coefficient is a fractal function of the parameter defining it. Such a complex behaviour is the result of the intricacy of the symbolic dynamics for a generic parameter value. Even if transport properties of such maps were already derived in [5] any attempt to analytically organize the cycle expansions for the full parameter range has failed so far: to our knowledge until now it was only possible to identify a family of parameter values where symbolic dynamics become tractable and dynamical zeta function can be derived [6]. Here, we show how it is possible to derive the diffusion coefficient for the full range of parameter values using an alternative formulation of the dynamical zeta functions presented by Baladi–Ruelle [8] (their approach generalize the Milnor–Thurston work [7] originally devoted to topological zeta function only).

1. Invariant coordinates and kneading determinants

For $\Lambda \geq 2$ let us define the map $g_\Lambda : \mathbb{R} \rightarrow \mathbb{R}$ as a lift of the map $\hat{g}_\Lambda : [0, 1] \rightarrow \mathbb{R}$:

$$g_\Lambda(x+n) = \hat{g}_\Lambda(x) + n, \quad n \in \mathbb{Z} \quad (1)$$

$$\hat{g}_\Lambda(x) = \frac{1}{2} + \Lambda \left(x - \frac{1}{2}\right). \quad (2)$$

Let us define $f_\Lambda(x) : [0, 1] \rightarrow [0, 1]$ to be the restriction on the unit interval of $\hat{g}_\Lambda(x)$, i.e.,

$$f_\Lambda(x) = \hat{g}_\Lambda(x), \quad \text{mod } 1. \quad (3)$$

A set of initial conditions for the map $g_\Lambda(x)$ will spread out diffusively on the real line and the transport coefficients (for example, the drift or the diffusion coefficient) can be computed with the help of a generalized dynamical zeta function [1] of the form

$$\zeta^{-1}(z, \beta) = \prod_{\{p\}} \left(1 - \frac{z^{n_p} e^{\beta \sigma_p}}{|\Lambda_p|} \right) \quad (4)$$

where $\{p\}$ denotes the set of prime periodic orbits of the reduced map $f(x)$ and the only quantities that enter the definition are the period of the orbit n_p , the instability $\Lambda_p = \prod_{i=0}^{n_p-1} f'(f^i(x))$ and the integer jumping number $\sigma_p = g(x) - f(x)$ (that recalls the number of cells the trajectory jumps once unfolded on the real line). The cycle expansions technique [1] provides an organization of the infinite set of periodic orbits $\{p\}$ that produces (for fully hyperbolic systems) a rapidly convergent power expansion of the dynamical zeta function (4). Transport properties of the system are encoded into the behaviour of its smallest




Figure 1. Example of the map $f_\Lambda(x)$ with $\Lambda = 3.23$.

zero $z_0(\beta)$ around its value $z_0(0) = 1$. For example, the drift J and the diffusion coefficient D can be computed by

$$J = \left. \frac{dz_0(\beta)}{d\beta} \right|_{\beta=0}, \quad D = \frac{1}{2} \left. \frac{d^2z_0(\beta)}{d\beta^2} \right|_{\beta=0}. \tag{5}$$

For odd integer Λ , the map consists of complete branches and the symbolic dynamic reduces to a full shift on an n -letter alphabet. In such a case, the dynamical zeta function can be written in closed form and the diffusion coefficient is easily computed. The situation is similar for even integer. For a specific set of the parameter Λ , it was possible [6] to control the symbolic dynamic and construct the dynamical zeta function, while for the generic case we will expect an infinite number of pruning rules (changing discontinuously under parameter variation) and no simple expression for the cycle expansions organization of periodic orbits. On the other hand, Milnor and Thurston [7] have shown how it is possible to control the symbolic dynamic of maps of the unit interval following the iteration of critical points. In particular, they were able to relate the topological entropy to the determinant of a finite matrix (the kneading determinant) where the entries are formal power series (with coefficients determined by the kneading trajectories). Later Baladi and Ruelle [8] have generalized the result to constant weight (see also [9] and references therein for more general results). We will closely follow the notation of Baladi–Ruelle [8] to construct the dynamical zeta function for the family of maps $f_\Lambda(x)$ defined above.

Fix Λ and rename for simplicity $f(x) := f_\Lambda(x)$. Call $a_0 < a_1 < \dots < a_N$ the ordered sequence of end points of each branch (see figure 1). Define $\epsilon(x) = \pm 1$, whether $f(x)$ is increasing or decreasing and $t(x)$ as a constant weight for $x \in [a_{i-1}, a_i]$. In our specific case, the restriction of these functions on the i -interval will take the constant values:

$$\epsilon_i = 1 \tag{6}$$

$$t_i = \frac{z e^{\beta\sigma_i}}{\Lambda} \tag{7}$$

where σ_i is the jumping number associated with the i th branch.

Let us associate to each point x the address vector in \mathbb{Z}^{N-1}

$$\vec{\alpha}(x) = [\text{sgn}(x - a_1), \dots, \text{sgn}(x - a_{N-1})] \quad (8)$$

and define the invariant coordinate of x by the formal series:

$$\vec{\theta}(x) = \sum_{n=0}^{\infty} \left[\prod_{k=0}^{n-1} (\epsilon t)(f^k(x)) \right] \vec{\alpha}(f^n(x)) \quad (9)$$

with the convention that the product is equal to 1 if $n = 0$ (the invariant coordinate θ is single valued once we put $\epsilon(a_i) = 0$).

Defining $\phi(a^\pm) = \lim_{x \rightarrow a^\pm} \phi(x)$, we compute the discontinuity vector at the critical points a_i for $i = 1, \dots, N - 1$:

$$\vec{K}_i(z, \beta) = \frac{1}{2} [\vec{\theta}(a_i^+) - \vec{\theta}(a_i^-)]. \quad (10)$$

The kneading matrix $K(z, \beta)$ is defined as the $(N - 1) \times (N - 1)$ matrix with $\vec{K}_i, i = 1, \dots, N - 1$ as rows. Let us call $\Delta(z, \beta) = \det K(z, \beta)$ the kneading determinant.

2. Relation between kneading determinant and dynamical zeta function

Without entering into many details, we restrict to the case under study and present the formula that relates the kneading determinant and the dynamical zeta function for the map $f(x)$ defined above. Let us call $\{\bar{p}\}$ the set of prime periodic orbits that have a critical point as periodic point (i.e., $f^{n_{\bar{p}}}(a_i) = a_i$). We note explicitly that there is at most a finite number of such prime orbits (for finite number of a_i). It is possible to show that the dynamical zeta function is equal to the kneading determinant up to a rational function (see [8] for more general results and a proof of relation (11)):

$$\Delta(z, \beta) = R(z, \beta) \zeta^{-1}(z, \beta) \quad (11)$$

$$R(z, \beta) = \left[1 - \frac{1}{2} (\epsilon_1 t_1 + \epsilon_N t_N) \right] \prod_{\{\bar{p}\}} [1 - t_{\bar{p}}(z, \beta)]^{-1}. \quad (12)$$

In their original work, Baladi and Ruelle [8] inserted the product over the set $\{\bar{p}\}$ into a definition of a reduced zeta function where they do not count such (repelling) cycles. Here, we prefer to remain with the usual zeta function and explicitly factorize out the critical cycles. In particular, formula (11) shows that the smallest zero of the kneading determinant coincides with the smallest zero of the dynamical zeta function for $\beta \rightarrow 0$.

3. Fractal diffusion coefficient

Let us start explicitly computing the kneading determinant for the map f_Λ . It is easy to see that, for $i = 1, \dots, N - 1$,

$$\vec{\theta}(a_i^+) = (\dots, +, +, -, \dots) + t_{i+1} \vec{\theta}(a_0^+) \quad (13)$$

$$\vec{\theta}(a_i^-) = (\dots, +, -, -, \dots) + t_i \vec{\theta}(a_N^-) \quad (14)$$

where the change of sign in the component of the first addend appears at the i th position. From the choice (6), we have $t_{i+1} = e^\beta t_i$ and so

$$\vec{K}_i(z, \beta) = (0, \dots, 0, 1, 0, \dots, 0) + \frac{1}{2} t_i e^{\frac{\beta}{2}} [e^{\frac{\beta}{2}} \vec{\theta}(a_0^+) - e^{-\frac{\beta}{2}} \vec{\theta}(a_N^-)]. \quad (15)$$

Simplifying the notation, we get that the kneading matrix is of the form

$$K_{ij} = \delta_{ij} - W_{ij}, \quad W_{ij} = p_i q_j \tag{16}$$

and the kneading determinant can be easily computed to be

$$\Delta = \det(K) = \exp \left[- \sum_{n=1}^{\infty} \frac{\text{Tr}(W^n)}{n} \right] \tag{17}$$

$$= 1 - \text{Tr} W \tag{18}$$

where in the last passage we have used the fact that for matrix W of the form (16) we have $\text{Tr}(W^n) = (\text{Tr} W)^n$. Using relations (16) and (18), the kneading determinant simplifies to

$$\Delta(z, \beta) = 1 + \frac{z}{2\Lambda} \sum_{i=1}^{N-1} e^{\beta(\sigma_i+1/2)} [e^{\frac{\beta}{2}\vec{\theta}(a_0^+)} - e^{-\frac{\beta}{2}\vec{\theta}(a_N^-)}]_i. \tag{19}$$

Due to the symmetry under $x \rightarrow -x$, the map $g(x)$ presents no net drift. In order to show it, we compute the first derivative of the smallest zero of the kneading determinant (5). Let us relabel the intervals by the associated jumping numbers: call it b_j , $j = -M, \dots, 0, \dots, M$, and call $\vec{\alpha}(j)$ the address (8) of a point in the interval b_j . Let us define the function $s_j(z, \beta)$ by the relation

$$e^{\beta/2\vec{\theta}(a_0^+)} =: \sum_{j=-M}^M s_j(z, \beta) \vec{\alpha}(j). \tag{20}$$

In the symmetric case, iterations of the critical point a_0 can be mapped to iterations of the other ‘external’ critical point a_N and so

$$e^{-\beta/2\vec{\theta}(a_N^-)} = \sum_{j=-M}^M s_j(z, -\beta) \vec{\alpha}(-j). \tag{21}$$

With the above definition, the function $s_j(z, \beta)$ receives contributions only when the trajectory of the critical point a_0 traverses the interval b_j .

Each of $\vec{\alpha}(j)$ has $2M$ components (see 8) that, with abuse of notation, we can label by $k = [(-M + \frac{1}{2}), \dots, (-2 + \frac{1}{2}), -\frac{1}{2}, \frac{1}{2}, (2 - \frac{1}{2}), \dots, (M - \frac{1}{2})]$. The kneading determinant can be rewritten as

$$\Delta(z, \beta) = 1 + \frac{z}{2\Lambda} \sum_k \sum_j e^{\beta k} [s_j(z, \beta) \vec{\alpha}(j) - s_j(z, -\beta) \vec{\alpha}(-j)]_k \tag{22}$$

and the first derivative is then

$$\frac{\partial \Delta(z, \beta)}{\partial \beta} \Big|_{z=1, \beta=0} = \frac{z}{2\Lambda} \sum_{k,j} [s_j(1, 0) k \vec{x}_k(j) + \partial_\beta s_j(1, 0) \vec{y}_k(j)] \Big|_{z=1, \beta=0} \tag{23}$$

$$= 0, \tag{24}$$

where we have used

$$\vec{x}(j) = [\vec{\alpha}(j) - \vec{\alpha}(-j)]; \quad \vec{y}(j) = [\vec{\alpha}(j) + \vec{\alpha}(-j)] \tag{25}$$

$$[\vec{x}(j)]_k = [\vec{x}(j)]_{-k}; \quad [\vec{y}(j)]_k = -[\vec{y}(j)]_{-k}. \tag{26}$$

For $\beta = 0$, the leading zero of the dynamical zeta function must be $z(0) = 1$ and then the kneading determinant must satisfy




Figure 2. Fractal diffusion coefficient as a function of the slope of the map $f_\Lambda(x)$ computed from the smallest zero of the dynamical zeta function. The inset is a blow up of a part of the main figure.

$$\Delta(1, 0) = 1 + \frac{1}{2\Lambda} \sum_{k,j} s_j(1, 0) [\vec{x}(j)]_k \quad (27)$$

$$= 1 + \frac{2}{\Lambda} \sum_{j=-M}^M j s_j(1, 0) = 0, \quad (28)$$

where in the last passage we have used the fact that $\sum_k [\vec{x}(j)]_k = 4j$. In order to clarify equation (28), let us check it for the simplest case of odd integer Λ . In such a case, the map consists of complete branches and $a_0 = 0$ becomes a fixed point. The function $s_j(z, \beta)$ vanishes for all indexes but $j = -M$ where

$$s_{-M}(z, \beta) = \sum_{n=0}^{\infty} \left[\frac{z e^{-\beta M}}{\Lambda} \right]^n e^{-\frac{\beta}{2}}. \quad (29)$$

Using the fact that $M = (\Lambda - 1)/2$, the relation (28) becomes

$$\Delta(1, 0) = 1 - \frac{1}{\Lambda} (\Lambda - 1) \frac{\Lambda}{\Lambda - 1} = 0. \quad (30)$$

Similar relations are found for even integer Λ where again the address of a_0 can be written in closed form due to the fact that $f^n(0) = 1/2, \forall n > 0$.

In the same spirit, it is possible to compute the diffusion coefficient for a generic value of Λ , rewriting equation (5) as

$$D = -\frac{1}{2} \left(\frac{\partial^2 \Delta(z, \beta)}{\partial \beta^2} \Big/ \frac{\partial \Delta(z, \beta)}{\partial z} \right) \Big|_{z=1, \beta=0} \quad (31)$$

where

$$\frac{\partial^2 \Delta(z, \beta)}{\partial \beta^2} \Big|_{1,0} = \frac{1}{2\Lambda} \sum_{k,j} \left[k^2 s_j(z, \beta) \vec{x}(j)_k + 2k \frac{\partial s_j(z, \beta)}{\partial \beta} \vec{y}(j)_k + \frac{\partial^2 s_j(z, \beta)}{\partial \beta^2} \vec{x}(j)_k \right] \Big|_{1,0} \quad (32)$$

$$\frac{\partial \Delta(z, \beta)}{\partial z} \Big|_{1,0} = -1 + \frac{1}{2\Lambda} \sum_{k,j} \frac{\partial s_j(z, \beta)}{\partial z} \vec{x}(j)_k \Big|_{1,0}. \quad (33)$$

The evaluation of expression (31) for a parameter choice $\Lambda \in [2, 3]$ is shown in figure 2 and can be compared, for example, with figure 10 in [4].

4. Conclusions

The fractal diffusion coefficient that appears in some simple piecewise linear map is the consequence of the sensitivity of the topological entropy to parameter variation: even arbitrary small parameter change creates and destroys infinitely many periodic orbits and the cycle expansions approach to dynamical zeta function becomes hard to follow. Avoiding periodic orbits and using the Baladi–Ruelle generalization of the Milnor–Thurston kneading determinant, we were able to construct the dynamical zeta function for arbitrary parameter values and compute from it the fractal diffusion coefficient.

Acknowledgments

I would like to thank R Artuso for helpful comments and suggestions and R Klages for interesting discussions.

References

- [1] Artuso R, Aurell E and Cvitanović P 1990 *Nonlinearity* **3** 325
Artuso R, Aurell E and Cvitanović P 1990 *Nonlinearity* **3** 361
Cvitanović P, Artuso R, Dahlqvist P, Mainieri R, Tanner G, Vattay G, Whelan N and Wirzba A 2003 *Chaos: Classical and Quantum* (www.nbi.dk/ChaosBook)
- [2] Prellberg T 2003 *J. Phys. A: Math. Gen.* **36** 2455–61
Isola S 2002 *Nonlinearity* **15** 1521–39
Artuso R, Cvitanović P and Tanner G 2003 *Prog. Theor. Phys. Suppl.* **150** 1
- [3] Artuso R, Casati G and Lombardi R 1993 *Phys. Rev. Lett.* **71** 62–4
Dahlqvist P and Artuso R 1996 *Phys. Lett. A* **219** 212
Dahlqvist P 1999 *Phys. Rev. E* **60** 6639–44
Artuso R and Cristadoro G 2004 *J. Phys. A: Math. Gen.* **37** 85–103
- [4] Klages R and Dorfman J R 1999 *Phys. Rev. E* **59** 5361–83
- [5] Groeneveld J and Klages R 2002 *J. Stat. Phys.* **109** 821–61
- [6] Tseng H-C, Chen H-J, Li P-C, Lai W-Y, Chou C-H and Chen H-W 1994 *Phys. Lett. A* **195** 74–80
Chen C-C 1995 *Phys. Rev. E* **51** 2815–22
Tseng H-C and Chen H-J 1996 *Int. J. Mod. Phys. B* **10** 1913–34
- [7] Milnor J and Thurston W 1988 *Lect. Notes Math.* **1342** 465–563
- [8] Baladi V and Ruelle D 1994 *Ergodic Theor. Dyn. Syst.* **14** 621–32
- [9] Baladi V and Ruelle D 1996 *Invent. Math.* **123** 553–74
Baillif M and Baladi V 2004 *Preprint* [math.DS/0211343](https://arxiv.org/abs/math/0211343)