

## Factorized steady states in mass transport models on an arbitrary graph

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2006 J. Phys. A: Math. Gen. 39 4859

(<http://iopscience.iop.org/0305-4470/39/18/006>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.104

The article was downloaded on 03/06/2010 at 04:26

Please note that [terms and conditions apply](#).

# Factorized steady states in mass transport models on an arbitrary graph

M R Evans<sup>1,2</sup>, Satya N Majumdar<sup>3</sup> and R K P Zia<sup>4</sup>

<sup>1</sup> SUPA, School of Physics, University of Edinburgh, Mayfield Road, Edinburgh EH9 3JZ, UK

<sup>2</sup> Isaac Newton Institute for Mathematical Sciences, 20 Clarkson Road, Cambridge, CB3 0EH, UK

<sup>3</sup> Laboratoire de Physique Théorique et Modèles Statistiques, Université Paris-Sud, Bat 100, 91405, Orsay-Cedex, France

<sup>4</sup> Department of Physics and Center for Stochastic Processes in Science and Engineering, Virginia Tech, Blacksburg, VA 24061-0435, USA

Received 24 February 2006

Published 19 April 2006

Online at [stacks.iop.org/JPhysA/39/4859](http://stacks.iop.org/JPhysA/39/4859)

## Abstract

We study a general mass transport model on an arbitrary graph consisting of  $L$  nodes each carrying a continuous mass. The graph also has a set of directed links between pairs of nodes through which a stochastic portion of mass, chosen from a site-dependent distribution, is transported between the nodes at each time step. The dynamics conserves the total mass and the system eventually reaches a steady state. This general model includes as special cases various previously studied models such as the zero-range process and the asymmetric random average process. We derive a general condition on the stochastic mass transport rules, valid for arbitrary graph and for both parallel and random sequential dynamics, that is sufficient to guarantee that the steady state is factorizable. We demonstrate how this condition can be achieved in several examples. We show that our generalized result contains as a special case the recent results derived by Greenblatt and Lebowitz for  $d$ -dimensional hypercubic lattices with random sequential dynamics.

PACS numbers: 05.40.-a, 02.50.Ey, 64.60.-i

## 1. Introduction

Diverse physical phenomena such as traffic flow [1], force propagation in granular media [2, 3], clustering of buses [4], aggregation and fragmentation of clusters [5], phase separation dynamics [6], shaken granular gases [7, 8] and sandpile dynamics [9] share one common feature—their microscopic dynamics involves stochastic transport of ‘mass’, or some conserved quantity, from one point in space to another. To simplify analysis, continuous space is typically replaced by (or ‘binned’ into) discrete sites. Several such lattice models

of stochastic mass transport have been introduced and studied, most notably the zero-range process (ZRP) [10–12], and the asymmetric random average process (ARAP) [13, 14]. These models are defined by specifying the microscopic dynamics, i.e. the basic stochastic rules for mass transport. Given the dynamics, there are two principal theoretical issues: (i) to identify the steady state if there is any, i.e. to find the invariant measure and (ii) once the steady state is known, to understand various physical properties in the steady state, e.g. the phenomenon of ‘condensation’ that happens when a finite fraction of the total mass condenses onto a single site [15].

It turns out that step (i) itself is often very difficult and the exact steady state is known in only very few cases [11]. In many of these known cases, the steady state is *factorizable*. This means that the steady state probability  $P(\{m_i\})$  of finding the system with mass  $m_1$  at site 1, mass  $m_2$  at site 2 etc is given by a product of (scalar) factors  $f(m_i)$ —one factor for each of the  $L$  sites of the system—e.g. for a homogeneous system where all sites  $i$  have equivalent connectivities

$$P(\{m_i\}) = Z(M, L)^{-1} \prod_{i=1}^L f(m_i) \delta\left(\sum_{i=1}^L m_i - M\right), \quad (1)$$

where  $Z(M, L)$  is a normalization which ensures that the integral of the probability distribution over all configurations containing the total mass  $M$  is unity; hence

$$Z(M, L) = \prod_{i=1}^L \left[ \int_0^\infty dm_i f(m_i) \right] \delta\left(\sum_{i=1}^L m_i - M\right). \quad (2)$$

Here, the  $\delta$ -function has been introduced to guarantee that we only include those configurations containing mass  $M$  in the integral. The single-site weights,  $f(m)$ , are determined by the details of the mass transfer rules and for a heterogeneous system may depend on the site  $i$ .

The advantage of having a factorizable steady state is that step (ii) mentioned above is often easier to carry out explicitly. This has been demonstrated recently by an exact analysis of the condensation phenomenon that occurs in a general class of mass transport models [15, 16]. This raises a natural question: when does the steady state in these mass transport models factorize? This issue was recently addressed in the context of a sufficiently general ‘mass transport model’ in one dimension, that includes, as special cases, the previously studied ZRP, ARAP and the chipping model [5]. In this model a mass  $m_i$  resides at each site  $i$  of a one-dimensional lattice with periodic boundary conditions. At each time, a stochastic portion,  $\tilde{m}_i \leq m_i$  of the mass  $m_i$  at site  $i$ , chosen from a distribution  $\varphi(\tilde{m}_i|m_i)$ , is chipped off to site  $i + 1$ . The distribution  $\varphi(\tilde{m}|m)$  was called the chipping kernel and by choosing its form appropriately one can recover the ZRP, the ARAP and the chipping model of [5]. Even though the model above is defined in discrete time where all sites are updated in parallel, by appropriately choosing the chipping kernel it is easy to study the continuous time limit, which corresponds to a random sequential update sequence, as a special case [17]. Similarly, one can also recover, as a special case, the model with only discrete masses as in ZRP. Thus the discrete time dynamics generalizes continuous time dynamics but a continuous mass variable generalizes discrete mass.

A natural question, first addressed in [17], is what should be the form of the chipping kernel  $\varphi(\tilde{m}|m)$  for the final steady state to be factorizable. In that study, it was proved that the *necessary and sufficient* condition for a factorized steady state in the one-dimensional directed case defined above is that the chipping kernel is of the form

$$\varphi(\tilde{m}|m) = \frac{u(\tilde{m})v(m - \tilde{m})}{\int_0^m d\tilde{m} u(\tilde{m})v(m - \tilde{m})}, \quad (3)$$

where  $u(z)$  and  $v(z)$  are arbitrary non-negative functions. Then the single-site weight in equation (1) is given by

$$f(m) = \int_0^m d\tilde{m} u(\tilde{m})v(m - \tilde{m}). \quad (4)$$

Furthermore, given a chipping kernel  $\varphi(\tilde{m}|m)$ , sometimes it is hard to verify explicitly that it is of the form (3) and thereby to identify the functions  $u(m)$  and  $v(m)$  in order to construct the weight  $f(m)$  in equation (4). This problem was circumvented by devising a test [18] to check if a given explicit  $\varphi(\tilde{m}|m)$  satisfies the condition (3) or not. Further, if it ‘passes this test,’ the weight  $f(m)$  can be found explicitly by a simple quadrature [18]. Finally, for any desired function  $f(m)$ , one can construct dynamical rules (i.e.,  $\varphi$ ) that will yield  $f(m)$  in a factorized steady state.

It was further demonstrated in [17] that the corresponding necessary and sufficient condition in the case of random sequential dynamics in continuous time can be easily obtained, by taking a suitable limit, from the condition for the parallel dynamics manifest in equation (3). This is done by choosing the chipping kernel as

$$\varphi(\tilde{m}|m) = \left[ 1 - dt \int_0^m \alpha(\tilde{m}|m) d\tilde{m} \right] \delta(\tilde{m}) + \alpha(\tilde{m}|m) dt \quad (5)$$

for small time increment  $dt$  and  $\delta(z)$  is the Dirac delta function. The function  $\alpha(\tilde{m}|m)$  denotes the ‘rate’ at which a mass  $\tilde{m}$  is transferred from a site with mass  $m$  to its right neighbour. Note that the form in equation (5) ensures the normalization,  $\int_0^m \varphi(\tilde{m}|m) d\tilde{m} = 1$ . Then, the necessary and sufficient condition for the factorizable steady state, derived from the more general condition in equation (3), is that the rate  $\alpha(\tilde{m}|m)$  must be of the following form [17]:

$$\alpha(\tilde{m}|m) = \frac{x(\tilde{m})v(m - \tilde{m})}{v(m)}, \quad (6)$$


where  $x(z)$  and  $v(z)$  are arbitrary non-negative functions. The corresponding steady state weight is then simply,  $f(m) = v(m)$ .

The condition (3) for factorizability in the mass transport model was derived only in one dimension and also only for unidirectional mass transport (from site  $i$  to site  $i + 1$ ). A natural question, therefore, is whether one can generalize this condition to higher-dimensional lattices, or to arbitrary graphs where mass transport can take place, in general, between any pair of sites  $i$  and  $j$ . Recently, Greenblatt and Lebowitz were able to derive a sufficiency condition [19] for factorizability in the mass transport model with nearest-neighbour mass transport on a regular  $d$ -dimensional lattice with periodic boundary conditions, but considered only the case of random sequential dynamics. They showed that if  $\alpha_q(\tilde{m}|m) dt$  is the probability of mass  $\tilde{m} \leq m$  being chipped off a site with mass  $m$  to a nearest neighbour in the direction  $q$  (there being  $2^d$  nearest neighbours on a hypercubic lattice in  $d$  dimensions), then the sufficient condition for factorizability is a direct generalization of the condition in equation (6), namely that the rate function must be of the form

$$\alpha_q(\tilde{m}|m) = \frac{x_q(\tilde{m})v(m - \tilde{m})}{v(m)} \quad (7)$$

for each  $q$ , where  $x_q(z)$  for each  $q$  and  $v(z)$  are arbitrary functions. The steady state weight is simple,  $f(m) = v(m)$ . However, it was not possible to prove that the condition (7) is also necessary [19], except in the case of generalized zero-range processes.

The purpose of this paper is to derive a more general sufficiency condition, valid for arbitrary graphs where the mass transport takes place not necessarily between nearest neighbours and for the more general case of parallel dynamics. Our results boil down to



**Figure 1.** An example graph with four nodes labelled 1, 2, 3 and 4 and directed links between certain pairs of nodes. The associated  $(4 \times 4)$  mass-transfer matrix  $[\mu]$  is shown, whose element  $\mu_{ij}$  denotes the stochastic mass transferred from site  $i$  to site  $j$  in one single update, provided there is a directed link between the two sites. If there is no directed link, the corresponding matrix element is always identically zero. The diagonal element  $\mu_{ii}$  is the amount of mass that stays at site  $i$  during the update. The row sum and the column sum associated with any node  $i$ ,  $\sum_j \mu_{ij} = m_i$  and  $\sum_j \mu_{ji} = m'_i$ , represent respectively the mass at  $i$  before and after the update.

equations (29) and (31). The former yields a sufficiency condition and the latter an additional consistency condition which must be satisfied.

In the special case of random sequential dynamics on a regular hypercubic lattice with nearest-neighbour mass transport, our sufficiency condition reduces to the one derived by Greenblatt and Lebowitz. Our results, however, are considerably more general.

The paper is organized as follows. In section 2, we define the mass transport model on an arbitrary graph. In section 3, we derive a sufficient condition on the chipping kernels, valid for parallel dynamics on an arbitrary graph  $\mathcal{G}$ , that would guarantee that the steady state is factorizable. We show that there are some additional consistency conditions that need to be satisfied in general and we demonstrate explicitly how these conditions are satisfied in several examples. In section 4, we extend our approach to random sequential dynamics on an arbitrary graph. We conclude with a summary and discussion in section 5.

## 2. The mass transport model on an arbitrary graph

We consider a fixed arbitrary graph  $\mathcal{G}$  consisting of  $L$  nodes labelled  $i = 1, 2, 3, \dots, L$  and a set of directed links or channels between certain pairs of nodes. At a given time  $t$ , a node  $i$  has mass  $m_i \geq 0$  where  $m_i$  are continuous variables. We consider discrete time dynamics where at each step the masses at all the nodes are updated in parallel according to the following rules. We first define a  $(L \times L)$  mass-transfer matrix  $[\mu]$  as follows. An element  $\mu_{ij}$  of the matrix  $[\mu]$  is identically zero at all times if there is no directed link from site  $i$  to site  $j$  on  $\mathcal{G}$ . If there is a directed link from  $i$  to  $j$ , then  $\mu_{ij} \geq 0$  is a non-negative stochastic variable that represents the mass transferred from site  $i$  to site  $j$  during one update. In figure 1 we give an example which we refer to for illustrative purposes throughout this section. The diagonal element  $\mu_{ii}$  represents the mass that stays at site  $i$  at the end of the single update. We assume that the dynamics of mass transport conserves the total mass. Thus if  $\{m_1, m_2, \dots, m_L\}$  represents the masses before the update, by virtue of mass conservation, the row sums of the matrix  $[\mu]$

are given by  $\sum_j \mu_{ij} = m_i$  (see figure 1). Similarly, the column sum corresponding to a node  $i$ ,  $\sum_j \mu_{ji} = m_i$  represents the mass at site  $i$  just after the update.

Note that, by definition, some elements of the matrix  $[\mu]$  are permanently zero (when there is no directed channel available for mass transfer between a pair of sites). On the other hand, when there is an available channel from  $i$  to  $j$ , the matrix element  $\mu_{ij}$  is a stochastic variable which is chosen in the following way. For each node  $i$ , we define a generalized ‘chipping kernel’  $\varphi_i(\{\mu_{ij}\}|m_i)$  that represents the joint distribution of the masses transported from site  $i$  in a single update. Here the set  $\{\mu_{ij}\}$  runs over only those sites  $j$  which are connected to  $i$  via a directed link, and in addition it includes the diagonal element  $\mu_{ii}$ . In other words,  $\{\mu_{ij}\}$  is simply the set of non-zero elements in the  $i$ th row of the mass-transfer matrix  $[\mu]$ . For example, in figure 1 where we have a graph of four nodes along with the directed links, we need to define four chipping kernels as follows:  $\varphi_1(\mu_{11}, \mu_{12}, \mu_{14}|m_1)$ ,  $\varphi_2(\mu_{21}, \mu_{22}, \mu_{23}|m_2)$ ,  $\varphi_3(\mu_{31}, \mu_{33}|m_3)$  and  $\varphi_4(\mu_{43}, \mu_{44}|m_4)$ , respectively. The chipping kernels must be normalized to unity at all sites  $i$ , i.e.

$$\int \varphi_i(\{\mu_{ij}\}|m_i) \delta\left(\sum_j \mu_{ij} - m_i\right) \prod_j d\mu_{ij} = 1, \tag{8}$$

where it is implied that the index  $j$  in the sum as well as in the product runs over all the sites that  $i$  feeds into, including  $i$  itself. Thus in this model the amount of mass transported from a given site  $i$  in one update depends only on that site  $i$ , and not, e.g. on the destination sites to which the mass is transported. Also, we assume that the chipping kernels  $\varphi_i(\{\mu_{ij}\}|m_i)$  do not contain the time  $t$  explicitly. Note also that this chipping kernel  $\varphi_i(\{\mu_{ij}\}|m_i)$  generalizes that of [17, 18] sufficiently to include hypercubic lattices[19] or more complicated graphs.

The chipping kernels thus specify the dynamics, i.e. the mass update rules. Given these kernels, we next ask what is the steady state joint probability distribution of masses  $\underline{m} \equiv \{m_1, m_2, \dots, m_L\}$ , i.e.  $P(\underline{m}, t \rightarrow \infty)$ . In particular, our goal is to determine the properties of the chipping kernels  $\varphi_i(\{\mu_{ij}\}|m_i)$  required in order to guarantee that the steady state joint probability distribution is factorizable, i.e. of the form

$$P(\underline{m}, t \rightarrow \infty) = Z(M, L)^{-1} \left[ \prod_{i=1}^L f_i(m_i) \right] \delta\left(\sum_{i=1}^L m_i - M\right), \tag{9}$$

where the normalization constant is given by

$$Z(M, L) = \prod_{i=1}^L \left[ \int_0^\infty dm_i f_i(m_i) \right] \delta\left(\sum_{i=1}^L m_i - M\right). \tag{10}$$

Besides, if the steady state factorizes as in equation (9), we would also like to know the single-site weights  $f_i(m_i)$  in terms of the prescribed chipping kernels  $\varphi_i(\{\mu_{ij}\}|m_i)$ . Note that on an arbitrary graph  $\mathcal{G}$ , the single-site weights  $f_i(m_i)$  are, in general, different from site to site. Hence there is an additional subscript  $i$  in  $f_i(m_i)$ . On a homogeneous graph, where all sites have equivalent connectivities, the weight function  $f(m)$  does not depend on the site  $i$  explicitly as in equation (1).

### 3. Factorizable steady state on an arbitrary graph

Since the dynamics conserves the total mass, at any time  $t$  we can write  $P(\underline{m}, t) \propto F(\underline{m}, t) \delta(\sum_i m_i - M)$  where  $M$  is the total mass and  $F(\underline{m}, t)$  is the unnormalized weight at time  $t$ . Below, we will first write down the general evolution equation of the weight  $F(\underline{m}, t)$

under the mass transport rules prescribed by the chipping kernels. While the notations that we will use for a general graph  $\mathcal{G}$  may seem a bit complicated, it is instructive to keep the simple example in figure 1 in mind and use it as a guide to the general notations.

Let us consider a single update from time  $t$  to time  $t + 1$ . Let  $\underline{m} \equiv \{m_1, m_2, \dots, m_L\}$  denote the masses at time  $t$  before the update and  $\underline{m}' \equiv \{m'_1, m'_2, \dots, m'_L\}$  denote the masses at time  $t + 1$  after the update. In terms of the elements of the mass-transfer matrix  $[\mu]$ , we thus have,  $m_i = \sum_k \mu_{ik}$  and  $m'_i = \sum_k \mu_{ki}$  for all  $i$ . The master equation for the evolution of the weight then reads

$$F(\underline{m}', t + 1) = \prod_{i=1}^L \left[ \int_0^\infty dm_i \prod_j \int d\mu_{ij} \varphi_i(\{\mu_{ij}\} | m_i) \delta\left(m'_i - \sum_k \mu_{ki}\right) \delta\left(m_i - \sum_k \mu_{ik}\right) \right] F(\underline{m}, t), \quad (11)$$

where the product over  $j$  runs over only the sites to which the site  $i$  feeds into, i.e., when there is a directed link between sites  $i$  and  $j$  (note that this set includes the site  $i$  itself). For other sites that are not connected to  $i$  by a directed link, the corresponding matrix element  $\mu_{ij} = 0$  identically (see figure 1) and hence they are not integration variables in equation (11).

Our strategy now is to assume that the steady state factorizes and to determine a sufficient condition for this assumption to hold. First, we take the  $t \rightarrow \infty$  limit on both sides of equation (11) and assume that the steady state weight factorizes

$$F(\underline{m}) = \prod_i f_i(m_i) \quad (12)$$

and write

$$f_i(m_i) \varphi_i(\{\mu_{ij}\} | m_i) = \mathcal{P}_i(\{\mu_{ij}\}) \delta\left(m_i - \sum_j \mu_{ij}\right), \quad (13)$$

thus

$$\prod_{i=1}^L f_i(m'_i) = \prod_{i=1}^L \left[ \int_0^\infty dm_i \prod_j \int d\mu_{ij} \mathcal{P}_i(\{\mu_{ij}\}) \delta\left(m'_i - \sum_k \mu_{ki}\right) \delta\left(m_i - \sum_k \mu_{ik}\right) \right]. \quad (14)$$

Now we Laplace transform this equation to obtain

$$\prod_{i=1}^L g_i(s_i) = \prod_{i=1}^L \left[ \int_0^\infty dm_i \prod_j \int d\mu_{ij} \mathcal{P}_i(\{\mu_{ij}\}) \delta\left(m_i - \sum_k \mu_{ik}\right) e^{-s_i \sum_k \mu_{ki}} \right], \quad (15)$$

where we have defined  $g_i(s_i) = \int_0^\infty f_i(m) e^{-s_i m} dm$ . Next we trivially perform the integration over  $m_i$  variables on the rhs of equation (15) and rearrange  $s_i$  to get

$$\prod_{i=1}^L g_i(s_i) = \prod_{i=1}^L \left[ \prod_j \int d\mu_{ij} \mathcal{P}_i(\{\mu_{ij}\}) e^{-\sum_k s_k \mu_{ik}} \right] \equiv \prod_i X_i(\{s_k\}_i), \quad (16)$$

where  $\{s_k\}_i$  indicates that  $k$  runs over the sites to which  $i$  is connected by a directed link, including the site  $i$  itself. For example, for the graph in figure 1, we have

$$g_1(s_1) g_2(s_2) g_3(s_3) g_4(s_4) = X_1(s_1, s_2, s_4) X_2(s_1, s_2, s_3) X_3(s_1, s_3) X_4(s_3, s_4). \quad (17)$$

One solution (not the most general as we shall discuss in section 5) to (16) is

$$X_i(\{s_k\}_i) = \prod_j K_{ij}(s_j) \quad (18)$$

so that

$$g_i(s_i) = \prod_j K_{ji}(s_i). \quad (19)$$

As before, the product in equation (18) runs over only those sites  $j$  that the site  $i$  feeds into (including itself), whereas in equation (19) the product runs over only those sites  $j$  that feed onto  $i$  (including the site  $i$ ). For example, for the graph in figure 1, we will have

$$\begin{aligned} X_1(s_1, s_2, s_4) &= K_{11}(s_1)K_{12}(s_2)K_{14}(s_4) \\ X_2(s_1, s_2, s_3) &= K_{21}(s_1)K_{22}(s_2)K_{23}(s_3) \\ X_3(s_1, s_3) &= K_{31}(s_1)K_{33}(s_3) \\ X_4(s_3, s_4) &= K_{43}(s_3)K_{44}(s_4) \end{aligned} \quad (20)$$

and correspondingly

$$\begin{aligned} g_1(s_1) &= K_{11}(s_1)K_{21}(s_1)K_{31}(s_1) \\ g_2(s_2) &= K_{12}(s_2)K_{22}(s_2) \\ g_3(s_3) &= K_{23}(s_3)K_{33}(s_3)K_{43}(s_3) \\ g_4(s_4) &= K_{14}(s_4)K_{44}(s_4). \end{aligned} \quad (21)$$

Since the rhs of equation (19) is a product of Laplace transforms, its inverse Laplace transform  $f_i(m_i)$  therefore has the convolution form

$$f_i(m) = \left[ \prod_{*j} v_{ji} \right] (m), \quad (22)$$

where the notation indicates a product over sites  $j$  feeding into  $i$  which is a multiple convolution and the function  $v_{ji}(m)$  is the inverse Laplace transform of  $K_{ji}(s_i)$ , i.e.

$$K_{ji}(s_i) = \int_0^\infty v_{ji}(\sigma) e^{-s_i\sigma} d\sigma. \quad (23)$$

Again, going back to figure 1, we will have, for example, by inverting equation (21) the following convolution for the site labelled 1:

$$f_1(m_1) = \int v_{11}(m_1 - \sigma_{21} - \sigma_{31})v_{21}(\sigma_{21})v_{31}(\sigma_{31}) d\sigma_{21} d\sigma_{31} \quad (24)$$

and similarly for the sites labelled 2, 3 and 4.

Having obtained the form of  $f_i(m)$ , let us ask what does this imply for the chipping kernels. To see that, we go back to the definitions of  $X_i$  in equation (16). Substituting the ansatz for  $X_i$  in equation (18) on the rhs of equation (16) we see that for each  $i$

$$\prod_j \int d\mu_{ij} \mathcal{P}_i(\{\mu_{ij}\}) \exp\left(-\sum_k s_k \mu_{ik}\right) = \prod_j K_{ij}(s_j). \quad (25)$$

For example, for the site 1 in figure 1, we have

$$\int d\mu_{11} d\mu_{12} d\mu_{14} P_1(\mu_{11}, \mu_{12}, \mu_{14}) e^{-s_1\mu_{11}-s_2\mu_{12}-s_4\mu_{14}} = K_{11}(s_1)K_{12}(s_2)K_{14}(s_4), \quad (26)$$



and similarly for the sites 2, 3 and 4. Equation (25) immediately implies for the general case

$$\mathcal{P}_i(\{\mu_{ij}\}) = \prod_j v_{ij}(\mu_{ij}), \quad (27)$$

where  $j$  runs over the sites which the site  $i$  feeds into and  $v_{ij}$  is the Laplace inversion of  $K_{ij}$  as in equation (23). For example, for the site 1 in figure 1 we have

$$P_1(\mu_{11}, \mu_{12}, \mu_{14}) = v_{11}(\mu_{11})v_{12}(\mu_{12})v_{14}(\mu_{14}). \quad (28)$$

Substituting equation (27) into equation (13) we get the following expression for the chipping kernel:

$$\varphi_i(\{\mu_{ij}\}|m_i) = \frac{\prod_j v_{ij}(\mu_{ij})}{[\prod_{*k} v_{ki}](m_i)}. \quad (29)$$

In the numerator the product is over sites  $j$  which the site  $i$  feeds into via a directed link connecting  $i$  to  $j$  (this includes site the  $i$  itself), and one has the constraint  $\sum_j \mu_{ij} = m_i$ . In the denominator, on the other hand, the product is over sites  $k$  feeding into site  $i$  (including once again the site  $i$  itself). Going back again to the example in figure 1, we have for the site 1 the following chipping kernel:

$$\varphi_1(\mu_{11}, \mu_{12}, \mu_{14}|m_1) = \frac{v_{11}(\mu_{11})v_{12}(\mu_{12})v_{14}(\mu_{14})}{\int v_{11}(m_1 - \sigma_{21} - \sigma_{31})v_{21}(\sigma_{21})v_{31}(\sigma_{31}) d\sigma_{21} d\sigma_{31}}, \quad (30)$$

where  $\mu_{11} + \mu_{12} + \mu_{14} = m_1$ . Similarly one can easily express the chipping kernels for the other sites 2, 3 and 4 in terms of the functions  $v_{ij}$ .

However this is not quite the end of the story since the chipping kernel  $\varphi_i(\{\mu_{ij}\}|m_i)$  must obey a key consistency condition, namely its normalization: equation (8). This implies from equation (29) that for each site  $i$

$$\left[ \prod_{*j} v_{ji} \right] (m) = \left[ \prod_{*j} v_{ij} \right] (m). \quad (31)$$

Again, the convolution on the lhs is over sites  $j$  feeding into the site  $i$  but the convolution on the rhs is over sites  $j$  which  $i$  feeds into. For example, for the site 1 in figure 1, this consistency condition reads

$$\begin{aligned} & \int v_{11}(m_1 - \sigma_{21} - \sigma_{31})v_{21}(\sigma_{21})v_{31}(\sigma_{31}) d\sigma_{21} d\sigma_{31} \\ &= \int v_{11}(m_1 - \mu_{12} - \mu_{14})v_{12}(\mu_{12})v_{14}(\mu_{14}) d\mu_{12} d\mu_{14}. \end{aligned} \quad (32)$$

One can write down similar consistency conditions for sites 2, 3 and 4 also.

In summary, a sufficient condition for factorization is that the chipping kernels are of product form as in equation (29) where  $v_{ij}$  are non-negative functions that satisfy the consistency conditions in equation (31), but otherwise are arbitrary. If this condition holds, then the steady state will factorize with single-site weight  $f_i(m)$  given by the convolution formula in equation (22). These conditions form the central result of this paper. The conditions arise from making the ansatz in equation (18) which immediately implies equation (22) for  $f_i(m_i)$  and equation (27) for  $\mathcal{P}_i(\{\mu_{ij}\})$ . These two equations, when substituted in equation (13), give equation (29) for the chipping kernel. Finally, equation (31) arises from the requirement that the chipping kernel in equation (29) is normalized to unity.

Conditions (29) and (31) certainly form a sufficient condition. Whether it is also a necessary condition remains a non-trivial issue. In the special case of a one-dimensional graph with unidirectional transport, it was explicitly proved in [17] that this is also a necessary

condition. Similarly, this is true (see below) in the case of a complete graph where every site is connected to every other site. However, it is unlikely, in general, that (29) is also necessary. In section 5, we provide an explicit example showing solutions to (16) which are not of the form (29). Although we are unable, for a technical reason to be discussed later, to prove that this class of solutions are also ‘legitimate,’ we feel that valid solutions different from (29) should exist. Beyond this simple example, finding the most general valid solution to equation (16) for the arbitrary graph seems to pose significant challenges.

### 3.1. Specific examples

Let us now discuss several special cases where one can directly verify that the consistency conditions in equation (31) are satisfied automatically by the functions  $v_{ij}$  due to the nature of the underlying graph  $\mathcal{G}$ , thereby guaranteeing factorizability provided the chipping kernels are of the product form as in equation (29).

**Example 1** (Mass transport in one dimension over extended range): Consider a one-dimensional lattice with periodic boundary conditions where the mass from a given site can get transported up to a range  $l$  to the left or to the right. In this case

$$\varphi_i(\{\mu_{ij}\}|m_i) = \varphi_i(\mu_{i,j-l} \dots \mu_{i,j+l}|m_i) \tag{33}$$

and we have a translationally invariant chipping kernel i.e.  $\varphi_i(\cdot|m)$  is the same function of  $(2l + 1)$  arguments for each site. Then the consistency condition (31) reads

$$[v_{i,i-l} * v_{i,i-l+1} * \dots * v_{i,i+l-1} * v_{i,i+l}](m_i) = [v_{i-l,i} * v_{i-l+1,i} * \dots * v_{i+l-1,i} * v_{i+l,i}](m_i) \tag{34}$$

which is automatically satisfied due to the translational invariance  $v_{i,i+d}(\mu) = v_{i-d,i}(\mu)$ .

**Example 2** (Symmetric chipping rule): It is easy to see that if the chipping rules are symmetric, i.e.  $v_{ij}(\mu) = v_{ji}(\mu)$ , then the consistency condition (31) holds automatically.

**Example 3** (A more general chipping rule): A more general way to meet the condition (31) is to assume that the chipping rules are such that at each site  $i$ , for every directed link  $(i \rightarrow j)$  out of site  $i$ , there is an incoming directed link  $(k \rightarrow i)$  to  $i$  from some other site  $k$  with  $v_{ij}(\mu) = v_{ki}(\mu)$ . For example, in the 1-d example above; this is achieved through  $v_{i,i+d}(\mu) = v_{i-d,i}(\mu)$ . Similarly, one can achieve this on a hypercubic lattice in arbitrary dimensions and on any graph where the number of links out of a site equals the number of links into it.

**Example 4** (A complete graph with permutationally invariant chipping kernel): So far we have proved that equation (29) is a sufficient condition for factorizability on any arbitrary graph provided we can satisfy the consistency condition (31) for each site. As discussed before, to prove that this condition is also necessary seems hard for a general graph. However, one can prove this for a complete graph where every site is connected to every other site via a directed link *and* the chipping kernel is permutationally invariant. This can be proved as follows. On a complete homogeneous graph the functions  $g_i(s) = g(s)$  in equation (16) do not depend explicitly on the site index  $i$ . Moreover, since every site is connected to every other sites and using the fact that the chipping kernel is permutationally invariant equation (16) simply becomes

$$g(s_1)g(s_2) \dots g(s_L) = [X(s_1, s_2, \dots, s_L)]^L. \tag{35}$$

Clearly  $X(s_1, s_2, \dots, s_L) = K(s_1)K(s_2) \cdots K(s_L)$  with  $K(s)$  being arbitrary, is *one* solution to equation (35) and thereby  $g(s) = [K(s)]^L$ . As in the general case, this will then lead to the sufficiency condition (29). To prove that  $X(s_1, s_2, \dots, s_L) = K(s_1)K(s_2) \cdots K(s_L)$  is also the most general form of the solution that one can write down for equation (35), we take logarithm on both sides of equation (35) and then take derivatives with respect to  $s_i$  and  $s_j$  with  $i \neq j$ . This gives

$$\frac{\partial^2 \ln X}{\partial s_i \partial s_j} = 0, \quad (36)$$

for any  $i \neq j$ . It is then easy to see that the most general solution of the partial differential equation (36) is of the form,  $X(s_1, s_2, \dots, s_L) = K_1(s_1)K_2(s_2) \cdots K_L(s_L)$  where  $K_i(s)$  are arbitrary functions. Since the graph is homogeneous, we also have  $K_i(s) = K(s)$  independent on the site index  $i$ . Thus  $X(s_1, s_2, \dots, s_L) = K(s_1)K(s_2) \cdots K(s_L)$  with  $K(s)$  being an arbitrary function is the only solution of equation (35) that respects homogeneity. Since this solution finally leads to the sufficiency condition, the uniqueness of its form guarantees then that equation (29) is both necessary and sufficient. Note that the consistency condition (31) is automatically satisfied in this case.

#### 4. Random sequential dynamics

The sufficiency condition in equation (29) and the associated consistency condition in equation (31), derived above for parallel update dynamics in discrete time, can be easily extended to the case of random sequential dynamics. This can be achieved by letting the probability of the chipping event in a time step  $\propto dt$  so that, to leading order in  $dt$  for small  $dt$ , at most one chipping event can occur in the whole graph  $\mathcal{G}$  per update, i.e. the chipping events occur sequentially one per update. In addition, taking  $dt \rightarrow 0$  one can obtain the continuous time limit where chipping events occur with ‘rates’ per unit time. Thus, the corresponding sufficiency condition for the random sequential dynamics will be specified in terms of the chipping ‘rate’ kernels, rather than the chipping probability kernels  $\varphi_i$  in parallel dynamics. To translate the sufficiency condition in equation (29) valid for the probability kernels into one that is valid for ‘rate’ kernels, we first redefine the functions  $v_{ij}(\mu_{ij})$ , for all  $i \neq j$ , in the following way:

$$v_{ij}(\mu_{ij}) = \delta(\mu_{ij}) + x_{ij}(\mu_{ij}) dt, \quad (37)$$

where  $x_{ij}(\mu_{ij})$  are arbitrary functions. The diagonal functions  $v_{ii}(\mu_{ii})$  are left unchanged.

With this re-definition of  $v_{ij}$ , the steady state weight in equation (22) becomes

$$f_i(m) = v_{ii}(m) + dt \left[ \sum_{j \neq i} \int_0^m v_{ii}(m - \mu_{ji}) x_{ji}(\mu_{ji}) d\mu_{ji} \right] + O(dt^2), \quad (38)$$

where the sum over  $j$  in the second term runs over all sites  $j \neq i$  that feed into the site  $i$  on  $\mathcal{G}$ . Using the re-defined  $v_{ij}$  in equation (37) one can similarly rewrite the chipping kernels in equation (29). Since the diagonal elements play a special role, it is convenient to redefine the chipping kernel only in terms of non-diagonal elements, i.e.,  $\varphi_i(\{\mu_{ij}\}|m_i) \equiv \varphi_i(\{\mu_{ij}\}'|m_i)$  where  $\{\mu_{ij}\}'$  denotes the set of matrix elements in the row  $i$  without the diagonal element  $\mu_{ii}$ . We are allowed to get rid of the diagonal element using the row sum,  $\sum_j \mu_{ij} = m_i$ . For example, for the graph in figure 1, we will rewrite  $\varphi_1(\mu_{11}, \mu_{12}, \mu_{14}|m_1) \equiv \varphi_1(\mu_{12}, \mu_{14}|m_1)$ .

Substituting equation (37) in equation (29) and taking the limit  $dt \rightarrow 0$ , one gets

$$\begin{aligned} \varphi_i(\{\mu_{ij}\}'|m_i) &= \left[ 1 - \frac{dt}{v_{ii}(m_i)} \sum_{j \neq i} [x_{ji} * v_{ii}](m_i) \right] \prod_{j \neq i} \delta(\mu_{ij}) \\ &+ dt \left[ \sum_{j \neq i} \frac{x_{ij}(\mu_{ij})v_{ii}(m_i - \mu_{ij})}{v_{ii}(m_i)} \prod_{k \neq j} \delta(\mu_{ik}) \right] + O(dt^2), \end{aligned} \tag{39}$$

where  $[x * y](m) = \int_0^m x(\sigma)y(m - \sigma) d\sigma$  denotes the convolution integral. The notation in equation (39) may look a bit complicated, but actually it is rather simple. For example, for the graph in figure 1, the kernel in equation (39) for the site labelled 1 reads

$$\begin{aligned} \varphi_1(\mu_{12}, \mu_{14}|m_1) &= \left[ 1 - \frac{dt}{v_{11}(m_1)} ([v_{11} * x_{31}](m_1) + [v_{11} * x_{21}](m_1)) \right] \delta(\mu_{12})\delta(\mu_{14}) \\ &+ dt \left[ \frac{x_{12}(\mu_{12})v_{11}(m_1 - \mu_{12})}{v_{11}(m_1)} \delta(\mu_{14}) + \frac{x_{14}(\mu_{14})v_{11}(m_1 - \mu_{14})}{v_{11}(m_1)} \delta(\mu_{12}) \right]. \end{aligned} \tag{40}$$

Taking  $dt \rightarrow 0$  limit in equation (39) we then obtain the continuous time limit where a portion  $\mu_{ij}$  gets chipped off the site  $i$  with mass  $m_i$ , to another site  $j$  which is connected to  $i$  by a directed link from  $i$  to  $j$  on  $\mathcal{G}$ , with a rate  $x_{ij}(\mu_{ij})v_{ii}(m_i - \mu_{ij})/v_{ii}(m_i)$ . Also, we notice that in the limit  $dt \rightarrow 0$ , the steady state weight  $f_i(m_i)$  is given simply from equation (38)

$$f_i(m_i) = v_{ii}(m_i). \tag{41}$$

However, as in the case of parallel update, we need the functions  $x_{ij}$  to satisfy certain additional consistency conditions. This is because the chipping kernel  $\varphi_i(\{\mu_{ij}\}'|m_i)$ , when integrated over its arguments  $\{\mu_{ij}\}'$  must give unity. Integrating equation (39) then gives the required consistency condition that must be satisfied for each node  $i$ ,

$$\sum_{j \neq i} [x_{ij} * v_{ii}](m_i) = \sum_{j \neq i} [x_{ji} * v_{ii}](m_i), \tag{42}$$

where the sum on the lhs runs over all sites  $j$  that the site  $i$  feeds into (excluding the site  $i$  itself), but on the rhs the sum runs over all sites  $j$  that feed into the site  $i$  (excluding the site  $i$  itself). Furthermore, taking the Laplace transform, cancelling the factors involving  $v_{ii}$  and then inverting back, equation (42) reduces to a simpler relation,

$$\sum_{j \neq i} x_{ij}\mu = \sum_{j \neq i} x_{ji}\mu. \tag{43}$$

As an example of this condition for the site labelled 1 in figure 1, we have

$$x_{12}(\mu) + x_{14}(\mu) = x_{21}(\mu) + x_{31}(\mu). \tag{44}$$

Similar consistency conditions can be written down for sites 2, 3 and 4 in figure 1.

Thus, in summary, for the mass transport model on an arbitrary graph  $\mathcal{G}$  with random sequential dynamics specified by the rates  $\alpha_{ij}(\mu|m_i)$  of mass  $\mu$  to be transferred from site  $i$  to site  $j$  (provided there is a directed link between  $i$  and  $j$  on  $\mathcal{G}$ ), a sufficiency condition for the steady state to be factorizable with weights  $f_i(m_i)$  is that  $\alpha_{ij}(\mu|m_i)$  is of the form

$$\alpha_{ij}(\mu|m_i) = \frac{x_{ij}(\mu)v_{ii}(m_i - \mu)}{v_{ii}(m_i)} \tag{45}$$

for all  $i$ , where  $x_{ij}$  and  $v_{ii}$  are functions that must satisfy the consistency conditions in equation (42), but otherwise arbitrary. Also, the corresponding steady state weight is then given simply by  $f_i(m_i) = v_{ii}(m_i)$ . Note that though formally equation (45) looks like a direct

generalization of the one-dimensional condition in equation (6), the additional consistency conditions in equation (42) are nontrivial to satisfy. For the one-dimensional example in equation (6), the corresponding consistency condition,  $[x_{i,i+1} * v_i](m_i) = [x_{i-1,i} * v_i](m_i)$  is automatically satisfied due to the translational invariance,  $x_{i-1,i}(\mu) = x_{i,i+1}(\mu)$ .

#### 4.1. Specific example of hypercubic lattice

It is easy to verify that the consistency condition in equation (42) is automatically satisfied in the four example cases of section 3.1 (as it should be since random sequential dynamics is just a limit of the discrete time case). Let us do this explicitly for the case where the graph  $\mathcal{G}$  is a homogeneous hypercubic lattice with periodic boundary conditions and mass transfer takes place only between nearest neighbours. For a hypercubic lattice, from each site  $i$  there are  $2^d$  outgoing links to the  $2^d$  nearest neighbours of  $i$ . Similarly, there are exactly  $2^d$  incoming links to site  $i$  from its nearest neighbours. Then the sufficiency condition in equation (45) can be written in a simplified notation

$$\alpha_{i,q}(\mu|m_i) = \frac{x_{i,q}(\mu)v_{ii}(m_i - \mu)}{v_{ii}(m_i)}, \quad (46)$$

where the index  $q = \pm 1$  runs over the  $2^d$  directions  $\{\pm e_1, \pm e_2, \dots, \pm e_d\}$ ,  $\alpha_{i,q}(\mu|m_i)$  denotes the rate of transfer of mass  $\mu$  from site  $i$  with mass  $m_i$  in the direction  $q$  and  $x_{i,q}$  denotes the function associated with the link  $(i, i + q)$ . Similarly, the consistency condition in equation (43) can be rewritten as

$$\sum_q x_{i,q}(\mu) = \sum_{j \in \text{neighbours of } i} x_{j,-q}(\mu). \quad (47)$$

We next use the fact that the lattice is homogeneous, i.e. it is translationally invariant in all directions. Clearly then  $x_{i-q,q}(z) = x_{i,i+q} = g_q(z)$  due to translational invariance in the  $q$ th direction. In that case the condition in equation (47) is clearly satisfied at all  $i$ . Also, due to the translational invariance, the rate function  $\alpha_{i,q}(\mu|m_i) = \alpha_q(\mu|m_i)$  does not depend on the site index  $i$ . By the same requirement,  $v_{ii}(m_i) = v(m_i)$ . Thus, the sufficiency condition in equation (45) simply reads

$$\alpha_q(\mu|m_i) = \frac{g_q(\mu)v(m_i - \mu)}{v(m_i)}, \quad (48)$$

where  $g_q(z)$  and  $v(z)$  are arbitrary non-negative functions. The consistency conditions are automatically satisfied as proved above. The steady state single site weight  $f(m_i)$  is simply  $f(m_i) = v(m_i)$ , and naturally it does not depend on the site index  $i$  explicitly. The condition in equation (48) is precisely that derived by Greenblatt and Lebowitz. Thus, our general sufficiency condition, valid for an arbitrary graph  $\mathcal{G}$ , recovers this special case when  $\mathcal{G}$  is a homogeneous hypercubic lattice with nearest-neighbour mass transport.

#### 4.2. ZRP on arbitrary graph

We can also check that the steady state for the continuous time zero-range process on an arbitrary graph is recovered. The zero-range process involves discrete masses and is specified by the rates for a unit of mass to hop from site  $i$  to  $j$ . In the case where these rates are given by

$$\alpha_{ij}(1|m_i) = y_{ij}w_i(m_i), \quad (49)$$

where  $w_i(m_i)$  is the total rate for the unit mass leaving site  $i$  and  $y_{ij}$  is the probability that the random destination for a hop from site  $i$  is  $j$ , the steady state factorizes with single-site weight

$$f_i(m_i) = \frac{P_i^{m_i}}{\prod_{n=1}^{m_i} w_i(n)}, \quad (50)$$

where  $p_i = \sum_{j \neq i} y_{ji} p_j$  is the steady state probability of a single random walker moving from site  $i$  to  $j$  with probability  $y_{ij}$  [11, 12].

To make the connection between the forms (49) and (45) we identify

$$x_{ij}(1) = y_{ij} p_i \quad \text{and} \quad w_i(m_i) = \frac{p_i v_{ii}(m_i - 1)}{v_{ii}(m_i)}. \tag{51}$$

Inverting the latter equality to express  $v_{ii}(m_i)$  in terms of  $w_i$  yields, by virtue of (41), the single-site weight (50). Finally the consistency condition (42) becomes

$$\sum_{j \neq i} x_{ij}(1) v_{ii}(m - 1) = \sum_{j \neq i} x_{ji}(1) v_{ii}(m - 1), \tag{52}$$

and is satisfied with  $x_{ij}(1) = y_{ij} p_i$  since we have

$$\sum_{j \neq i} x_{ij}(1) = \sum_{j \neq i} x_{ji}(1). \tag{53}$$

### 5. Conclusion

In this work we have derived the sufficient condition (29) along with a consistency condition (31) for factorization of the general (discrete time, continuous mass) mass transport model on an arbitrary graph. In this case the single-site weight is given by (22). We gave in section 3.1 specific examples of geometries where the additional consistency condition associated with the sufficient condition is automatically fulfilled. Moreover on a complete graph with permutationally invariant chipping functions we showed that the sufficient condition is, in fact, also necessary.

Of course a significant improvement would be to generalize condition (29),(31) to a necessary and sufficient condition. To illustrate the challenges involved in accomplishing this task, we offer another simple example where we can derive a condition that is both necessary and sufficient.

This example, a seemingly trivial generalization of the one in [17, 18], involves a one-dimensional lattice with chipping to both nearest-neighbour sites and we also assume that the chipping kernel is translationally invariant ( $l = 1$  in example 1 of section 3.1). Then, due to the translational invariance, (16) becomes

$$\prod_i g(s_i) = \prod_i X(s_{i-1}, s_i, s_{i+1}). \tag{54}$$

Using a similar procedure to that employed in Example 4 of section 3.1—taking the logarithm of (54) then successive derivatives with respect to  $s_i$  and  $s_j$ —we find that the general solution to (54) is

$$X(s_{i-1}, s_i, s_{i+1}) = K_{ii-1}(s_{i-1}) K_{ii}(s_i) K_{ii+1}(s_{i+1}) \frac{H(s_{i-1}, s_i)}{H(s_i, s_{i+1})}, \tag{55}$$

where  $K_{ii-1}, K_{ii}, K_{ii+1}$  and  $H$  are arbitrary functions, independent of  $i$ . Inverting the Laplace transforms to give  $\mathcal{P}(\mu_{ii-1}, \mu_{ii}, \mu_{ii+1})$  would yield a considerably more complicated form for the chipping kernel  $\varphi(\mu_{ii-1}, \mu_{ii}, \mu_{ii+1})$  than (29). In principle, therefore, there could be a whole family of chipping kernels, generated by the choice of  $H$ , which give rise to the same steady state (i.e. the same single-site weight  $f(m)$ ) as that for  $H = 1$ . Thus,  $H$  can be thought of as a ‘gauge function’.

In addition, for each choice of  $H$ , one has to ensure the consistency condition, namely that

$$\varphi(\mu_{ii-1}, \mu_{ii}, \mu_{ii+1}) = \frac{\mathcal{P}(\mu_{ii-1}, \mu_{ii}, \mu_{ii+1})}{f(m_i)} \delta(m_i - (\mu_{ii-1} + \mu_{ii} + \mu_{ii+1})) \tag{56}$$

is correctly normalized to unity. On the face of it, it may appear that this imposes a formidable constraint on the choice of  $H$ . However, we now show that this consistency condition does not impose any additional constraint on  $H$ . In other words, if the consistency condition is ensured for  $H = 1$ , then it is automatically satisfied for all other choices of  $H$ . To see this, consider the expression of the chipping kernel (56). Now, the denominator  $f(m_i)$  is, of course, independent of the choice of  $H$ . So, to prove that the consistency condition of normalization of  $\varphi$  does not impose any additional constraint on  $H$ , one has to prove that the integral

$$I(m) \equiv \int d\mu_{ii-1} d\mu_{ii} d\mu_{ii+1} \mathcal{P}(\mu_{ii-1}, \mu_{ii}, \mu_{ii+1}) \delta(m - \mu_{ii-1} - \mu_{ii} - \mu_{ii+1}) \quad (57)$$

is independent of the choice of  $H$ . Now, taking the Laplace transform of equation (57) one obtains

$$\begin{aligned} \int dm e^{-sm} I(m) &= \int d\mu_{ii-1} d\mu_{ii} d\mu_{ii+1} e^{-s(\mu_{ii-1} + \mu_{ii} + \mu_{ii+1})} \mathcal{P}(\mu_{ii-1}, \mu_{ii}, \mu_{ii+1}) \\ &= X(s, s, s) \\ &= K_{ii-1}(s) K_{ii}(s) K_{ii+1}(s), \end{aligned} \quad (58)$$

where we have used definition (16) in going from the first line to the second and equation (55) in going from the second line to the third. But this expression on the rhs does not contain the ‘gauge function’  $H$  thereby proving that the integral of  $\mathcal{P}$  is independent of the choice of  $H$ . Thus the consistency condition is automatically fulfilled for arbitrary choices of  $H$  as long as it is ensured for  $H = 1$ , which is indeed the case as shown in section 3.1 example 1.

A more serious constraint on  $H$  is that the inverse Laplace transform of (55) must be *non-negative*. The case  $H = 1$  in (55) obviously imposes the trivial constraint that the inverse Laplace transform of the  $K$ ’s be non-negative. It remains to be shown whether there is also a class of  $H \neq 1$  which lead to valid chipping kernels. If it could be shown that this class is non-empty, the one-dimensional  $l = 1$  example we have discussed would show explicitly that it is *not* necessary for  $\varphi$  to be of the form (29). Mapping the extent of this class of  $H$  (if indeed it is non-empty) remains a daunting task.

For an arbitrary graph, we have made some in-roads with similar considerations. Unfortunately, the most general solution to (16) remains difficult to formulate.

Finally we note that having a factorized steady state opens the door for the study of condensation as in [15, 16]. Thus one should be able to analyse condensation in various geometries or even on scale-free networks [20].

## Acknowledgments

RKPZ acknowledges the support of the US National Science Foundation through DMR-0414122. We thank Peter Sollich for pointing out that equation (43) follows from equation (42).

## References

- [1] Chowdhury D, Santen L and Schadschneider A 2000 *Phys. Rep.* **329** 199
- [2] Liu C-h, Nagel S R, Schecter D A, Coppersmith S N, Majumdar S, Narayan O and Witten T A 1995 *Science* **269** 513
- [3] Coppersmith S N, Liu C-h, Majumdar S, Narayan O and Witten T A 1996 *Phys. Rev. E* **53** 4673
- [4] O’Loan O J, Evans M R and Cates M E 1998 *Phys. Rev. E* **58** 1404
- [5] Majumdar S N, Krishnamurthy S and Barma M 1998 *Phys. Rev. Lett.* **81** 3691  
Majumdar S N, Krishnamurthy S and Barma M 2000 *J. Stat. Phys.* **99** 1
- [6] Kafri Y, Levine E, Mukamel D, Schütz G M and Török J 2002 *Phys. Rev. Lett.* **89** 035702
- [7] van der Meer D, van der Weele K and Lohse D 2004 *J. Stat. Mech.: Theor. Exp.* P04004

- [8] Torok J 2005 *Physica A* **355** 374–82
- [9] Jain K 2005 *Phys. Rev. E* **72** 017105
- [10] Spitzer F 1970 *Adv. Math.* **5** 246
- [11] Evans M R 2000 *Braz. J. Phys.* **30** 42
- [12] Evans M R and Hanney T 2005 *J. Phys. A: Math. Gen.* **38** R195
- [13] Krug J and Garcia J 2000 *J. Stat. Phys.* **99** 31
- [14] Rajesh R and Majumdar S N 2000 *J. Stat. Phys.* **99** 943  
Rajesh R and Majumdar S N 2001 *Phys. Rev. E* **64** 036103
- [15] Majumdar S N, Evans M R and Zia R K P 2005 *Phys. Rev. Lett.* **94** 180601
- [16] Evans M R, Majumdar S N and Zia R K P 2005 *Preprint cond-mat/0510512*
- [17] Evans M R, Majumdar S N and Zia R K P 2004 *J. Phys. A: Math. Gen.* **37** L275
- [18] Zia R K P, Evans M R and Majumdar S N 2004 *J. Stat. Mech.: Theor. Exp.* **L10001**
- [19] Greenblatt R L and Lebowitz J L 2006 *J. Phys. A: Math. Gen.* **39** 1505 (*Preprint cond-mat/0506776*)
- [20] Noh J D, Shim G M and Lee H 2005 *Phys. Rev. Lett.* **94** 198701