

Renormalization-group analysis of repulsive three-body systems

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2006 J. Phys. A: Math. Gen. 39 L49

(<http://iopscience.iop.org/0305-4470/39/2/L01>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.104

The article was downloaded on 03/06/2010 at 04:28

Please note that [terms and conditions apply](#).

LETTER TO THE EDITOR

Renormalization-group analysis of repulsive three-body systems

Michael C Birse

Theoretical Physics Group, School of Physics and Astronomy, The University of Manchester, Manchester, M13 9PL, UK

Received 30 September 2005

Published 14 December 2005

Online at stacks.iop.org/JPhysA/39/L49

Abstract

A coordinate space approach, based on that used by Efimov, is applied to three-body systems with contact interactions between pairs of particles. In systems with nonzero orbital angular momentum or with asymmetric spatial wavefunctions, the hyperradial equation contains a repulsive $1/r^2$ potential. The resulting wavefunctions are used in a renormalization group analysis. This confirms Griesshammer's power counting for short-range three-body forces in these systems. The only exceptions are ones like the ^4S channel for three nucleons, where any derivatives needed in the interaction are found to be already counted by the scaling with the cut-off.

PACS numbers: 03.65.Nk, 11.10.Hi, 21.45.+v, 21.30.Fe

Effective field theories (EFTs) are now being widely applied to few-nucleon systems, see [1, 2]. The starting point is usually an organization of the terms in an effective potential according to naive dimensional analysis (NDA), as originally suggested by Weinberg [3]. This classifies terms according to number of powers of low-energy scales they contain. In some cases, most notably S-wave nucleon–nucleon scattering, the leading-order (LO) terms turn out to be unnaturally large. This is a consequence of low-energy bound or virtual states. It means that the LO terms need to be iterated to all orders in solving the Schrödinger or Lippmann–Schwinger equation [4–6].

However it is now clear that NDA is not valid in all systems. There can be nonperturbative effects associated with strong long-range potentials that significantly change the power counting for short-range interactions. This was first noted in the context of attractive three-body systems (such as three bosons, or the ^2S channel for three nucleons) [7, 8], where the leading three-body forces must be promoted to LO. More recently, failures of NDA have been observed in repulsive three-body systems [9], and for nucleon–nucleon scattering in spin-triplet waves [10, 11]. In the first example, short-range three-body forces in most low

partial waves are demoted to higher orders than naively expected; in the second, short-range forces are promoted to lower orders in P- and D-waves. In both cases, the terms in potentials scale with noninteger anomalous dimensions, and so the standard classification of terms as LO, next-to-leading order (NLO) etc is no longer convenient.

In this letter I examine repulsive three-body systems using the renormalization group (RG) approach developed in [12–14]. This provides an independent confirmation of results of Griesshammer for the power counting in these systems [9]. That work solved the Skorniakov–Ter-Martirosian equation [15] in momentum space, whereas here I work in coordinate space following the approach developed by Efimov for attractive three-body systems [16] and recently extended by Gasaneo and Macek to cases with nonzero angular momentum [17]. I also comment on differences between the counting for derivative interactions in systems with strong long-range forces compared with the pure short-range case. This corrects the counting in [9] for the leading three-body force in the 4S channel for three nucleons.

If particles interact only through zero-range forces, then their wavefunctions satisfy the free Schrödinger equation, except where two of them coincide. The two-body forces can then be represented by boundary conditions at these points. These boundary conditions form the basis for Efimov’s approach [16] as well as more recent work in [17–19]. In particular, Gasaneo and Macek have used this method to find solutions for systems with symmetric spatial wavefunctions. Here I generalize their results to cover asymmetric cases, such as the spin-quartet channels for three nucleons. It is convenient to work in hyperspherical coordinates since the boundary conditions are separable in the limit of infinite two-body scattering length. The resulting hyperradial equation then has the form of a free radial Schrödinger equation with a centrifugal-like $1/r^2$ term whose strength is given by the hyperangular eigenvalue. This potential determines the form of three-body wavefunctions at small hyperradii, and hence it controls RG flow of the short-range three-body forces [13, 14].

Sets of relative coordinates for three particles with equal masses are

$$\mathbf{x}_i = \mathbf{r}_k - \mathbf{r}_j, \quad \mathbf{y}_i = \mathbf{r}_i - \frac{1}{2}(\mathbf{r}_j + \mathbf{r}_k), \quad (1)$$

where i, j, k are a cyclic permutation of 1, 2, 3, and I have used the traditional ‘odd-man-out’ notation to label the sets. Hyperspherical coordinates can then be defined in terms of these as

$$r = \sqrt{x_i^2 + \frac{4}{3}y_i^2}, \quad \alpha_i = \arctan\left(\frac{\sqrt{3}x_i}{2y_i}\right). \quad (2)$$

The three-body wavefunction can be decomposed into Faddeev components [20] as

$$\Psi(r, \Omega) = \sum_{i=1}^3 \frac{2}{r^2 \sin(2\alpha_i)} \phi_i(r, \alpha_i, \hat{\mathbf{x}}_i, \hat{\mathbf{y}}_i), \quad (3)$$

where a factor of $1/(x_i y_i)$ has been taken out to simplify the radial parts of the Hamiltonian. Away from the configurations where two particles coincide, each of these components satisfies a free Schrödinger equation. In hyperspherical coordinates, this has the form

$$-\frac{1}{M} \left[\frac{\partial^2}{\partial r^2} + \frac{1}{2} \frac{\partial}{\partial r} + \frac{1}{r^2} \frac{\partial^2}{\partial \alpha_i^2} \right] \phi_i + \frac{1}{Mr^2} \left[\frac{\mathbf{L}_i^2}{\cos^2 \alpha_i} + \frac{\mathbf{L}_{jk}^2}{\sin^2 \alpha_i} \right] \phi_i = E \phi_i. \quad (4)$$

Here \mathbf{L}_{jk} denotes the relative angular momentum of the pair jk , and \mathbf{L}_i the angular momentum of particle i relative to that pair. If the problem is separable, we can write ϕ_i in the form

$$\phi_i(r, \alpha_i, \hat{\mathbf{x}}_i, \hat{\mathbf{y}}_i) = F_i(r) u_i(\alpha_i) Y_{l_i m_i}(\hat{\mathbf{x}}_i) Y_{l_i m_i}(\hat{\mathbf{y}}_i), \quad (5)$$

where $F_i(r)$ and $u_i(\alpha_i)$ satisfy the ordinary differential equations

$$-\frac{1}{M} \left[\frac{d^2}{dr^2} + \frac{1}{r} \frac{d}{dr} - \frac{v^2}{r^2} \right] F_i = p^2 F_i, \quad -\frac{d^2 u_i}{d\alpha_i^2} + \left[\frac{l_i(l_i + 1)}{\cos^2 \alpha_i} + \frac{l'_i(l'_i + 1)}{\sin^2 \alpha_i} \right] u_i = v^2 u_i. \quad (6)$$

This hyperradial equation looks just like a free radial Schrödinger equation in two dimensions, with the hyperangular eigenvalue ν^2 determining the strength of the centrifugal-like $1/r^2$ term. In the cases of interest, where pairs of particles interact only in S-waves, we can simplify the hyperangular equations by setting $l'_i = 0$ and $l_1 = l_2 = l_3 \equiv l$. The component ϕ_i is then independent of $\hat{\mathbf{x}}_i$.

From the definition of the reduced Faddeev components ϕ_i in equation (3), they must vanish at $x_i = 0$. In hyperspherical coordinates these boundary conditions are

$$\phi_i\left(r, \frac{\pi}{2}, \hat{\mathbf{y}}_i\right) = 0. \quad (7)$$

In the limit of infinite two-body scattering length, the logarithmic derivative of the reduced wavefunction must vanish whenever two particles coincide and so can interact via the two-body force [16–19]:

$$\frac{1}{x_i \Psi} \frac{\partial}{\partial x_i} (x_i \Psi) \Big|_{x_i=0} = 0. \quad (8)$$

The points where $x_i = 0$ correspond to $\mathbf{x}_j = \mathbf{y}_i$, $\mathbf{y}_j = -\frac{1}{2}\mathbf{y}_i$ and $\mathbf{x}_k = -\mathbf{y}_i$, $\mathbf{y}_k = -\frac{1}{2}\mathbf{y}_i$ in the other relative coordinate systems. In terms of the hyperangular coordinates these are $\alpha_j = \alpha_k = \frac{\pi}{3}$. The resulting boundary conditions on the Faddeev components are

$$\frac{\partial \phi_i}{\partial \alpha_i} \Big|_{\alpha_i=0} + \frac{2}{\sin(2\alpha_j)} \phi_j \Big|_{\alpha_j=\pi/3, \hat{\mathbf{y}}_j=-\hat{\mathbf{y}}_i} + \frac{2}{\sin(2\alpha_k)} \phi_k \Big|_{\alpha_k=\pi/3, \hat{\mathbf{y}}_k=-\hat{\mathbf{y}}_i} = 0. \quad (9)$$

These are separable and lead to the hyperangular conditions

$$\frac{du_i}{d\alpha_i} \Big|_{\alpha_i=0} + \frac{4}{\sqrt{3}} (-1)^l \left[u_j\left(\frac{\pi}{3}\right) + u_k\left(\frac{\pi}{3}\right) \right] = 0. \quad (10)$$

The symmetries of the spatial wavefunction can be used to simplify these conditions further. There are two cases of physical interest. The first is a spatial wavefunction that is symmetric under exchange of any pair of particles. In this case the Faddeev components are equal:

$$\phi_1 = \phi_2 = \phi_3 \equiv \phi. \quad (11)$$

This describes three identical bosons, or three fermions with different quantum numbers (spin, isospin, etc) whose intrinsic state is completely antisymmetric. Most importantly for nuclear physics this corresponds to three nucleons with total spin $\frac{1}{2}$. The second is where the spatial wavefunction is antisymmetric under exchange of one pair of particles, but symmetric under exchange of either of the others. The components are then related by

$$\phi_1 = 0, \quad \phi_2 = -\phi_3 \equiv \phi. \quad (12)$$

This describes three nucleons with total spin $\frac{3}{2}$. Generically we may write the boundary conditions in the form

$$\frac{du}{d\alpha} \Big|_{\alpha=0} + \lambda (-1)^l \frac{8}{\sqrt{3}} u\left(\frac{\pi}{3}\right) = 0, \quad (13)$$

where $\lambda = +1$ for completely symmetric spatial wavefunctions and $\lambda = -\frac{1}{2}$ for cases with one antisymmetric pair.

As noted by Gasaneo and Macek [17], the hyperangular equation for $l' = 0$ can be solved in terms of hypergeometric functions. After defining the new variable $z = \cos^2 \alpha$ and writing $u(z) = z^{(l+1)/2} v(v)$, the equation takes the form of the hypergeometric equation [21],

$$z(1-z) \frac{d^2 v}{dz^2} + [c - (a+b+1)z] \frac{dv}{dz} - abv = 0, \quad (14)$$

with

$$a = \frac{l+1-\nu}{2}, \quad b = \frac{l+1+\nu}{2}, \quad c = l + \frac{3}{2}. \quad (15)$$

The hyperangular eigenfunctions are thus¹

$$u(\alpha) = (\cos \alpha)^{l+1} {}_2F_1\left(\frac{l+1-\nu}{2}, \frac{l+1+\nu}{2}, l + \frac{3}{2}; \cos^2 \alpha\right). \quad (16)$$

The corresponding radial solutions are just Bessel functions of order ν and so we get

$$\phi(r, \alpha, \hat{\mathbf{y}}_i) = N J_\nu(pr) (\cos \alpha)^{l+1} {}_2F_1\left(\frac{l+1-\nu}{2}, \frac{l+1+\nu}{2}, l + \frac{3}{2}; \cos^2 \alpha\right) Y_{lm}(\hat{\mathbf{y}}_i). \quad (17)$$

These vanish at $\alpha = \frac{1}{2}$, as required by the first boundary condition. The other condition, arising from the contact interactions at $x_i = 0$, then provides an equation for the eigenvalues ν^2 . The hypergeometric functions have the properties [21],

$$\frac{\partial}{\partial z} {}_2F_1(a, b, c; z) = \frac{ab}{c} {}_2F_1(a+1, b+1, c+1; z), \quad (18)$$

and for z equal to or close to 1

$$\begin{aligned} {}_2F_1(a, b, c; 1) &= \frac{\Gamma(c)\Gamma(c-a-b)}{\Gamma(c-a)\Gamma(c-b)} && \text{if } c-a-b > 0, \\ {}_2F_1(a, b, c; z) &\sim \frac{\Gamma(c)\Gamma(a+b-c)}{\Gamma(a)\Gamma(b)} && \text{if } c-a-b < 0. \end{aligned} \quad (19)$$

Using these, equation (13) can be expressed in the form

$$1 = \lambda \left(-\frac{1}{2}\right)^l \frac{2}{\sqrt{3\pi}} \frac{\Gamma(\frac{l+1-\nu}{2})\Gamma(\frac{l+1+\nu}{2})}{\Gamma(l+\frac{3}{2})} {}_2F_1\left(\frac{l+1-\nu}{2}, \frac{l+1+\nu}{2}, l + \frac{3}{2}; \frac{1}{4}\right), \quad (20)$$

which matches equation (2.18) of [9], with the substitution of s by ν . It also agrees with the results of [17] if $\lambda = 1$. For symmetric systems ($\lambda = 1$) with $l = 0$ this is the equation first derived by Danilov [22], which has an imaginary solution for ν . The corresponding hyperradial wavefunctions show oscillatory behaviour at small distances, and this is responsible for Thomas [23] and Efimov [16] effects. For $l \geq 1$ or systems with one antisymmetric pair, the roots of the equation are real and can be found in table 2 of [9]. In these cases the $1/r^2$ potential is repulsive and so there is no Efimov effect.

Having constructed the wavefunctions for the long-range forces in these systems, we can now use the methods of [13] to find the RG eigenvalues, which give the power counting for terms in the short-range potential. In fact for real values of ν , we can immediately use the results in equation (54) of that paper if we multiply the hyperradial solutions by $\sqrt{\pi}/(2pr)$ to get functions that satisfy a three-dimensional radial Schrödinger equation. A term in the rescaled potential proportional to p^{2n} (n powers of the energy) varies with the cut-off Λ as $\Lambda^{2(n+\nu)} p^{2n}$ and so its RG eigenvalue is

$$\rho = 2(n + \nu). \quad (21)$$

If we assign Λ -independent terms to LO in our expansion of the EFT, then ρ also labels the order of a term. The leading term in each channel is thus of order 2ν , in agreement with the results in table 3 of [9], except for the ^4S and Wigner-antisymmetric ^2S channels, where Griesshammer adds two extra powers of low-energy scales.

¹ Note that I have chosen to write these in the form that will make most direct contact with the results of [9], rather than the equivalent form given in [17]. Also, the factor of 2 in equation (13) of [17] is incorrect and should be omitted.

The motivation for adding these two powers is the antisymmetry of the wavefunction in these channels which prevents all three particles from coinciding. As a result a pure δ -function interaction has no effect on them; one with at least two derivatives would be needed. For any finite cut-off, however, a contact interaction becomes nonlocal and so can contribute. In [9], this happens implicitly through the momentum cut-off. In contrast, [13] does it explicitly by using a δ -shell form for the short-distance interactions. This was done to ensure that the interaction has an effect even though the wavefunctions vanish as $r \rightarrow 0$ as a result of the $1/r^2$ potential. For example, the same RG analysis can be applied to two-body scattering with nonzero angular momentum L by setting $\nu = L + \frac{1}{2}$ [13]. It shows that the leading short-distance interaction in this partial wave is of order $2L$, as expected from the fact that $2L$ derivatives of a δ -function are needed to form a contact interaction that acts in this wave. Note that these derivatives are already counted by the RG eigenvalue, $\rho = 2L + 1$.

The wavefunctions in two-body channels with nonzero angular momentum, or in three-body channels with $\nu > -\frac{1}{2}$, satisfy a radial equation of the form

$$\frac{1}{r} \frac{d^2}{dr^2} (r\psi) = -p^2\psi + \frac{\nu^2 - \frac{1}{4}}{r^2}\psi. \quad (22)$$

Acting on one of the wavefunctions where the long-range $1/r^2$ interaction has been iterated to all orders, an interaction with two derivatives thus gives rise to two contributions. One is just proportional to two powers of the low-energy scale p , and so is two orders higher in the power counting. However the other, proportional to $(1/r^2)\psi$, is of the same order as the term without those derivatives, since at small distances $1/r$ is not a low-energy scale. This second piece is absent if $L = 0$ (or equivalently $\nu = \frac{1}{2}$). In that more familiar case, additional derivatives do indeed increase the order of the interaction.

The bottom line is that any derivatives needed to construct appropriate interactions for the repulsive $1/r^2$ potentials are already counted in the RG eigenvalue (or by the superficial degree of divergence in [9]), without any need to add additional powers. But, apart from this rather minor amendment, the present analysis confirms Griesshammer's results for the power counting in repulsive three-body systems [9]. The leading term in each channel has RG eigenvalue 2ν , where ν^2 is the hyperangular eigenvalue. These eigenvalues are not integers and so the usual classification of terms as NLO etc in the EFT becomes inconvenient. In most cases this counting demotes short-distance three-body interactions to higher orders than predicted by NDA, although in some channels there is a small degree of promotion.

Acknowledgments

I am grateful to the organizers of the ECT* Workshop on 'Nuclear forces and QCD: never the twain shall meet?' where this work was started. I also thank H Griesshammer for useful discussions and correspondence.

References

- [1] Beane S R, Bedaque P F, Haxton W C, Phillips D R and Savage M J 2001 *At the Frontier of Particle Physics: Handbook of QCD* vol 1 ed M Shifman (Singapore: World Scientific) p 133 (Preprint [nucl-th/0008064](#))
- [2] Bedaque P F and van Kolck U 2002 *Ann. Rev. Nucl. Part. Sci.* **52** 339 (Preprint [nucl-th/0203055](#))
- [3] Weinberg S 1990 *Phys. Lett. B* **251** 288
Weinberg S 1991 *Nucl. Phys. B* **363** 3
- [4] Bedaque P F and van Kolck U 1998 *Phys. Lett. B* **428** 221 (Preprint [nucl-th/9710073](#))
- [5] van Kolck U 1999 *Nucl. Phys. A* **645** 273 (Preprint [nucl-th/9808007](#))

-
- [6] Kaplan D B, Savage M J and Wise M B 1998 *Phys. Lett. B* **424** 390 (Preprint [nucl-th/9801034](#))
Kaplan D B, Savage M J and Wise M B 1998 *Nucl. Phys. B* **534** 329 (Preprint [nucl-th/9802075](#))
 - [7] Bedaque P F, Hammer H-W and van Kolck U 1999 *Phys. Rev. Lett.* **82** 463 (Preprint [nucl-th/9809025](#))
Bedaque P F, Hammer H-W and van Kolck U 1999 *Nucl. Phys. A* **646** 444 (Preprint [nucl-th/9811046](#))
Bedaque P F, Hammer H-W and van Kolck U 2000 *Nucl. Phys. A* **676** 357 (Preprint [nucl-th/9906032](#))
 - [8] Bedaque P F, Griesshammer H W, Hammer H-W and Rupak G 2003 *Nucl. Phys. A* **714** 589 (Preprint [nucl-th/0207034](#))
 - [9] Griesshammer H 2005 *Nucl. Phys. A* **760** 110 (Preprint [nucl-th/0502039](#))
 - [10] Nogga A, Timmermans R G E and van Kolck U 2005 Preprint [nucl-th/0506005](#)
 - [11] Birse M C 2005 Preprint [nucl-th/0507077](#)
 - [12] Birse M C, McGovern J A and Richardson K G 1999 *Phys. Lett. B* **464** 169 (Preprint [hep-ph/9807302](#))
 - [13] Barford T and Birse M C 2003 *Phys. Rev. C* **67** 064006 (Preprint [hep-ph/0206146](#))
 - [14] Barford T and Birse M C 2005 *J. Phys. A: Math. Gen.* **38** 697 (Preprint [nucl-th/0406008](#))
 - [15] Skorniakov G V and Ter-Martirosian K A 1957 *Sov. Phys.—JETP* **4** 648
 - [16] Efimov V N 1971 *Sov. J. Nucl. Phys.* **12** 589
Efimov V N 1979 *Sov. J. Nucl. Phys.* **29** 546
 - [17] Gasaneo G and Macek J H 2002 *J. Phys. B: At. Mol. Opt. Phys.* **35** 2239
 - [18] Fedorov D V and Jensen A S 1993 *Phys. Rev. Lett.* **71** 4103
Fedorov D V and Jensen A S 2001 *J. Phys. A: Math. Gen.* **34** 6003 (Preprint [quant-ph/0106039](#))
Fedorov D V and Jensen A S 2002 *Nucl. Phys. A* **697** 783 (Preprint [nucl-th/0107027](#))
 - [19] Gasaneo G, Ovchinnikov S and Macek J H 2001 *J. Phys. A: Math. Gen.* **34** 8941
 - [20] Faddeev L D 1961 *Sov. Phys.—JETP* **12** 1014
 - [21] Abramowitz M and Stegun I A 1970 *Handbook of Mathematical Functions* (New York: Dover)
 - [22] Danilov G S 1961 *Sov. Phys.—JETP* **13** 349
 - [23] Thomas L H 1935 *Phys. Rev.* **47** 903