

Classification of integrable hydrodynamic chains and generating functions of conservation laws

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2006 J. Phys. A: Math. Gen. 39 10803

(<http://iopscience.iop.org/0305-4470/39/34/014>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.106

The article was downloaded on 03/06/2010 at 04:48

Please note that [terms and conditions apply](#).

Classification of integrable hydrodynamic chains and generating functions of conservation laws

Maxim V Pavlov

Department of Mathematical Sciences, Loughborough University, Loughborough, LE11 3TU UK

Received 31 March 2006, in final form 20 June 2006

Published 9 August 2006

Online at stacks.iop.org/JPhysA/39/10803

Abstract

A new approach to classification of integrable hydrodynamic chains is established. Generating functions of conservation laws are classified by the method of hydrodynamic reductions. The N parametric family of explicit hydrodynamic reductions allows us to reconstruct corresponding hydrodynamic chains. Many new hydrodynamic chains are found.

PACS numbers: 02.30.Ik, 02.30.Jr

Mathematics Subject Classification: 35L40, 35L65, 37K10

1. Introduction

The first integrable hydrodynamic chain

$$A_t^k = A_x^{k+1} + kA^{k-1}A_x^0, \quad k = 0, 1, 2, \dots \quad (1)$$

was derived in [1]. The integrability of this hydrodynamic chain was developed in a set of publications [2–5]. The ‘integrability’ means the existence of infinitely many conservation laws

$$\partial_t \mathbf{H}_k(A^0, A^1, \dots, A^k) = \partial_x G_k(A^0, A^1, \dots, A^{k+1}), \quad k = 0, 1, 2, \dots \quad (2)$$

and infinitely many commuting flows (see [2])

$$A_{t^m}^k = [kA^{k+n-1}\partial_x + n\partial_x A^{k+n-1}] \frac{\partial \mathbf{H}_{m+1}}{\partial A^n}, \quad k, n, m = 0, 1, 2, \dots,$$

where $x \equiv t^0$, $t \equiv t^1$, $\mathbf{H}_0 = A^0$, $\mathbf{H}_1 = A^1$, $\mathbf{H}_2 = A^2 + (A^0)^2$, $\mathbf{H}_3 = A^3 + 3A^0A^1$, \dots

The first commuting flow is

$$A_y^k = A_x^{k+2} + A^0A_x^k + (k+1)A^kA_x^0 + kA^{k-1}A_x^1, \quad k = 0, 1, 2, \dots, \quad (3)$$

where $y \equiv t^2$. Eliminating the moments A^1 and A^2 from the equations

$$A_t^0 = A_x^1, \quad A_t^1 = \partial_x[A^2 + (A^0)^2/2], \quad A_y^0 = \partial_x[A^2 + (A^0)^2]$$

one can obtain the Khohlov–Zabolotzkaya equation

$$A_{tt}^0 = \partial_x [A_y^0 - A^0 A_x^0], \quad (4)$$

which is the dispersionless limit of the KP equation. The compatibility condition $\partial_t(p_y) = \partial_y(p_t)$ of two generating functions of conservation laws for the Benney hydrodynamic chain (1)

$$p_t = \partial_x \left(\frac{p^2}{2} + A^0 \right) \quad (5)$$

and for its first commuting flow (3)

$$p_y = \partial_x \left(\frac{p^3}{3} + A^0 p + A^1 \right) \quad (6)$$

yields (4). Thus, the integrability of this equation is equivalent to the integrability of the Benney hydrodynamic chain (see [6–9]). Substituting the series

$$p = \lambda - \frac{\mathbf{H}_0}{\lambda} - \frac{\mathbf{H}_1}{\lambda^2} - \frac{\mathbf{H}_2}{\lambda^3} - \dots \quad (7)$$

into both the above equations one can obtain (2) and the infinite series of conservation laws for (3),

$$\partial_y \mathbf{H}_k(A^0, A^1, \dots, A^k) = \partial_x \mathbf{Q}_k(A^0, A^1, \dots, A^{k+2}), \quad k = 0, 1, 2, \dots$$

Later many integrable hydrodynamic chains

$$A_t^k = \sum_{n=0}^{k+1} F_n^k(\mathbf{A}) A_x^n, \quad k = 0, 1, 2, \dots, \quad \frac{\partial F_n^k}{\partial A^m} = 0, \quad m > k + 1 \quad (8)$$

and corresponding 2+1 quasilinear equations (see, for instance, (4)) were found in [10–16]. Recently, some of these integrable hydrodynamic chains have been rediscovered (see [17]) and studied in [17–22].

At this moment we have several tools allowing us to find a complete classification of the integrable hydrodynamic chains (8) and more complicated hydrodynamic chains

$$A_t^k = \sum_{n=0}^{N_k} F_n^k(\mathbf{A}) A_x^n, \quad k = 0, 1, 2, \dots, \quad \frac{\partial F_n^k}{\partial A^m} = 0, \quad m > M_k, \quad (9)$$

where N_k and M_k are some integers.

- (1) The existence of an extra commuting flow (an *external* method), successfully used for the Egorov hydrodynamic chains (see [12]);
- (2) The existence of an extra conservation law (an *external* method), successfully used for the Kupershmidt Poisson bracket (see [10] and [34]) and for (an incomplete classification) the Kupershmidt–Manin Poisson bracket (see [10]);
- (3) Vanishing of the Haantjes tensor (an *internal* method), successfully (unfinished classification) used for the integrable hydrodynamic chains (8) and successfully (complete classification) for the Kupershmidt–Manin Poisson bracket (see [16]).

In this paper, we establish the *combined* method based on three key tools, previously successfully utilized in the theory of 2+1 hydrodynamic-type systems and 2+1 quasilinear equations (see [11, 13–15]). These are method of hydrodynamic reductions (see [5, 13]) and the method of pseudopotentials (see [11–13]) combined with the ‘concept’ of the Gibbons equation (see [23]).

This paper is organized in the following order. In the second section the concept of the Gibbons equation is introduced. In comparison with [5] hydrodynamic reductions are described in a conservative form. In the third section a new approach of the classification of integrable hydrodynamic chains via their generating functions of conservation laws is established. In the fourth section explicit hydrodynamic reductions associated with local Hamiltonian structures are discussed. In the fifth section the simple (but nontrivial) generalization of the Benney hydrodynamic chain is investigated. N parametric family of the Hamiltonian hydrodynamic reductions is found. The corresponding Riemann surface is constructed. Thus, N series of conservation laws and commuting flows can be found (see [23]), then infinitely many particular solutions can be obtained by the generalized hodograph method (see [24]). In the sixth section another integrable hydrodynamic chain is described, whose hydrodynamic reductions coincide with hydrodynamic reductions of the hydrodynamic chain described in the fifth section. One particular case of this hydrodynamic chain is connected with 2+1 quasilinear equation determined by the ‘integrable’ Lagrangian considered in [15]. In the seventh section three different approaches allowing us to construct commuting flows are presented. The Miura-type transformation is used for links between some well-known hydrodynamic chains and corresponding 2+1 quasilinear systems via generating functions of conservation laws. In the eighth section we briefly mention the general case of generating functions of conservation laws connected with arbitrary integrable hydrodynamic chains. This general case is closely connected with the Hamiltonian hydrodynamic chains considered in [25].

2. The Gibbons equation

The Benney hydrodynamic chain (1) is connected with the formal series

$$\lambda = p + \frac{A^0}{p} + \frac{A^1}{p^2} + \frac{A^2}{p^3} + \dots \tag{10}$$

by the Gibbons equation (see details below)

$$\lambda_t - p\lambda_x = \frac{\partial \lambda}{\partial p} \left[p_t - \partial_x \left(\frac{p^2}{2} + A^0 \right) \right].$$

The *method of hydrodynamic reductions* suggested in [5] (and developed in [13]) means the existence of infinitely many sub-systems (which are called the ‘hydrodynamic reductions’)

$$r_t^i = p^i(\mathbf{r})r_x^i, \quad i = 1, 2, \dots, N, \tag{11}$$

where all moments A^k are functions of N Riemann invariants r^k . This hydrodynamic-type system must be consistent with the generating function of conservation laws (5). Then one can obtain

$$\partial_i p = \frac{\partial_i A^0}{p^i - p}. \tag{12}$$

The compatibility conditions $\partial_i(\partial_k p) = \partial_k(\partial_i p)$ yield the *system in involution*

$$\partial_i p^k = \frac{\partial_i A^0}{p^i - p^k}, \quad \partial_{ik} A^0 = 2 \frac{\partial_i A^0 \partial_k A^0}{(p^i - p^k)^2}, \quad i \neq k, \tag{13}$$

which we call the *Gibbons–Tsarev system*.

Any hydrodynamic reduction (11) can be written via first moments A^k ($k = 0, 1, \dots, N - 1$), where all other moments A^n ($n = N, N + 1, \dots$) are functions of the first N moments A^k (see details in [5]).

Also, any hydrodynamic reduction (11) can be written in the conservative form (see (5))

$$a_t^i = \partial_x \left(\frac{(a^i)^2}{2} + A^0(\mathbf{a}) \right), \quad i = 1, 2, \dots, N, \quad (14)$$

where the function A^0 satisfies the Gibbons–Tsarev system (written via field variables a^k)

$$(a^i - a^k) \partial_{ik} A^0 = \partial_k A^0 \partial_i \left(\sum \partial_n A^0 \right) - \partial_i A^0 \partial_k \left(\sum \partial_n A^0 \right), \quad i \neq k, \quad (15)$$

$$(a^i - a^k) \frac{\partial_{ik} A^0}{\partial_i A^0 \partial_k A^0} + (a^k - a^j) \frac{\partial_{jk} A^0}{\partial_j A^0 \partial_k A^0} + (a^j - a^i) \frac{\partial_{ij} A^0}{\partial_i A^0 \partial_j A^0} = 0, \quad i \neq j \neq k,$$

which is a consequence of the compatibility conditions $\partial_i(\partial_k p) = \partial_k(\partial_i p)$

$$\frac{p - a^i}{\partial_i A^0} \sum \frac{\partial_{in} A^0}{p - a^n} - \frac{p - a^k}{\partial_k A^0} \sum \frac{\partial_{kn} A^0}{p - a^n} + \frac{(a^i - a^k) \partial_{ik} A^0}{\partial_i A^0 \partial_k A^0} \left(\sum \frac{\partial_n A^0}{p - a^n} - 1 \right) = 0, \quad (16)$$

where $\partial_i \equiv \partial/\partial a^i$ and (cf (12))

$$\partial_i p = \frac{\partial_i A^0}{p - a^i} \left(\sum \frac{\partial_n A^0}{p - a^n} - 1 \right)^{-1}. \quad (17)$$

Example. The simplest hydrodynamic reduction is given by $A^0 = \sum \varepsilon_k a^k$. This is the so-called waterbag reduction (see [6, 26]). Then (17)

$$\partial_i p = \frac{\varepsilon_i}{p - a^i} \left(\sum \frac{\varepsilon_n}{p - a^n} - 1 \right)^{-1}$$

can be integrated. The equation of the Riemann surface

$$\lambda = p - \sum \varepsilon_k \ln(p - a^k), \quad (18)$$

where λ is an integration factor, can be expanded (at the infinity $\lambda \rightarrow \infty, p \rightarrow \infty$) in the formal series (10), where $A^k = \sum \varepsilon_i (a^i)^{k+1}/(k+1)$, if $\sum \varepsilon_i = 0$. If $\sum \varepsilon_i \neq 0$, then first the parameter λ in the equation of the Riemann surface must be re-scaled,

$$\lambda - \sum \varepsilon_k \ln \lambda = p - \sum \varepsilon_k \ln(p - a^k).$$

Then all other moments A^k are some non-homogeneous polynomials of $\sum \varepsilon_i (a^i)^k$.

This is not a unique choice. For instance, any hydrodynamic reductions can be written in a conservative form in infinitely many ways. Let us mention just two simplest choices here (all other are discussed in [23]):

1. $a_t^i = \partial_x \left(\frac{(a^i)^2}{2} + A^0(\mathbf{a}, \mathbf{b}) \right), \quad b_t^i = \partial_x(a^i b^i) \quad i = 1, 2, \dots, N,$
2. $a_t^i = \partial_x \left(\frac{(a^i)^2}{2} + A^0 \right), \quad A_t^0 = \partial_x A^1(\mathbf{a}) \quad i = 1, 2, \dots, N.$

(19)

For the first choice, the simplest hydrodynamic reduction is given by $A^0 = \sum b^k$; for the second choice $A^1 = \sum b^k$.

The Riemann invariants are most suitable coordinates to prove such general properties like the existence of infinitely many hydrodynamic reductions parameterized by N arbitrary functions of a single variable (see [5]), but they are very inconvenient for a search of particular and explicit hydrodynamic reductions. Vice versa, it is very difficult to derive the Gibbons–Tsarev system in the coordinates a^k (see (17)), but explicit particular hydrodynamic reductions can be found naturally in many cases (see a lot of examples in [22]).

The *phenomenological* algebro-geometric approach for integrability of symmetric hydrodynamic-type systems

$$a_t^i = \partial_x \psi(a^1, a^2, \dots, a^N; p)|_{p=a^i} \tag{20}$$

was formulated in [23].

Statement 1. *If the symmetric hydrodynamic-type system (20) is integrable, then this system has the generating function of conservation laws*

$$p_t = \partial_x \psi(a^1, a^2, \dots, a^N; p). \tag{21}$$

Statement 2. *Then some function $\lambda(a^1, a^2, \dots, a^N; p)$ satisfies the Gibbons equation*

$$\lambda_t - \frac{\partial \psi}{\partial p} \lambda_x = \frac{\partial \lambda}{\partial p} [p_t - \partial_x \psi(\mathbf{a}; p)]. \tag{22}$$

We call the function $\lambda(a^1, a^2, \dots, a^N; p)$ the equation of the Riemann surface. This function is a solution of the set of *linear* PDE's

$$A_i^k \frac{\partial \lambda}{\partial u^k} + \frac{\partial \psi}{\partial u^i} \frac{\partial \lambda}{\partial p} = 0,$$

where the matrix A_i^k is given by

$$A_i^k(\mathbf{u}; p) = \left(\frac{\partial \psi}{\partial p} \Big|_{p=u^i} - \frac{\partial \psi}{\partial p} \right) \delta_i^k + \frac{\partial \psi}{\partial u^i} \Big|_{p=u^k}.$$

The Gibbons equation (22) describes a deformation of the Riemann surface $\lambda(\mathbf{a}; p)$. This equation has three distinguish features:

- (1) if one fixes $\lambda = \text{const}$ (free parameter), then one obtains (21),
- (2) if one fixes $p = \text{const}$ (free parameter), then one obtains the kinetic equation (a collisionless Vlasov equation) written in the so-called Lax form

$$\lambda_{t^i} = \{\lambda, \hat{\mathbf{H}}\} = \frac{\partial \lambda}{\partial x} \frac{\partial \hat{\mathbf{H}}}{\partial p} - \frac{\partial \lambda}{\partial p} \frac{\partial \hat{\mathbf{H}}}{\partial x},$$

where $\hat{\mathbf{H}} = \psi(\mathbf{a}; p)$.

- (3) if one chooses coordinates, which are the Riemann invariants determined by the condition $\partial \lambda / \partial p = 0$, then the corresponding hydrodynamic-type system (20) can be written in the diagonal form (cf (11))

$$r_{t^i}^i = \mu^i(\mathbf{r}) r_x^i, \quad i = 1, 2, \dots, N, \tag{23}$$

where the characteristic velocities

$$\mu^i = \frac{\partial \psi}{\partial p} \Big|_{p=p^i}$$

can be found from the algebraic system $\det A_i^k(\mathbf{u}; p) = 0$; then the corresponding values p^i can be expressed via the Riemann invariants r^k . In this algebro-geometric construction the Riemann invariants are the branch points $r^i = \lambda|_{\partial \lambda / \partial p = 0}$ of the Riemann surface (exactly as it is in the Whitham theory, see [8, 27]).

Example. The dispersionless limit of the vector NLS (see (19) and [3])

$$a_t^i = \partial_x \left(\frac{(a^i)^2}{2} + \sum b^n \right), \quad b_t^i = \partial_x (a^i b^i), \quad i = 1, 2, \dots, N \tag{24}$$

is the first known hydrodynamic reduction of the Benney hydrodynamic chain (1) determined by the moment decomposition $A^k = \sum (a^i)^k b^i$. In such a case the formal series (10) reduces to the equation of the Riemann surface

$$\lambda = p + \sum \frac{b^k}{p - a^k}. \quad (25)$$

The Zakharov reduction (24) written in the Riemann invariants (23) (see (11))

$$r_t^i = p^i(\mathbf{r})r_x^i, \quad i = 1, 2, \dots, 2N,$$

where the characteristic velocities $p^i(\mathbf{r})$ can be found from $(\partial\lambda/\partial p = 0)$

$$1 = \sum \frac{b^k}{(p - a^k)^2}$$

has infinitely many conservation laws, which can be obtained from (25) with the aid of the Bürmann–Lagrange series (see, for instance, [28] and some details in [23]).

The compatibility conditions $\partial_i(\partial_k p) = \partial_k(\partial_i p)$ must be valid *identically* for any symmetric hydrodynamic-type system (20), but a whole set (parameterized by N arbitrary functions of a single variable) of such symmetric hydrodynamic-type systems (for each fixed function $\psi(\mathbf{a}; p)$) is *described* by these compatibility conditions $\partial_i(\partial_k p) = \partial_k(\partial_i p)$.

Thus, the problem of the description of semi-Hamiltonian symmetric hydrodynamic-type system is the problem of the classification of integrable hydrodynamic chains.

3. The Gibbons–Tsarev system

The *main claim* of this paper is that *the classification of integrable hydrodynamic chains (9) is equivalent to the classification of the generating functions of conservation laws (21) by the method of hydrodynamic reductions (see [5, 13])*.

In this section we derive integrability conditions, which are a nonlinear PDE's system in involution. This *extended Gibbons–Tsarev system* generalizes the Gibbons–Tsarev system obtained in [5] and describes a *set* of generating functions of conservation laws together with their hydrodynamic reductions parameterized by N arbitrary functions of a single variable.

The simplest case is

$$p_t = \partial_x \psi(u; p). \quad (26)$$

Let us first introduce the new notations

$$f_i \equiv f(u, p^i, p) = \frac{\psi_u}{\psi_p|_{p=p^i} - \psi_p}, \quad f_{ik} \equiv f(u, p^i, p^k) = \frac{\psi_u|_{p=p^k}}{\psi_p|_{p=p^i} - \psi_p|_{p=p^k}}, \quad i \neq k, \quad (27)$$

$$\varphi_{ik}(u, p^i, p^k, p) \equiv \frac{f_{ik}\partial_p f_k - f_{ki}\partial_{p^i} f_i + f_i\partial_p f_k - f_k\partial_p f_i - \partial_u(f_i - f_k)}{f_i - f_k}, \quad i \neq k. \quad (28)$$

Differentiating the generating function of conservation laws (26) with respect to the Riemann invariants r^i (see (23)), one obtains

$$\partial_i p = f_i \partial_i u. \quad (29)$$

If $p = p^k$ ($k \neq i$), then (29) reduces to

$$\partial_i p^k = f_{ik} \partial_i u. \quad (30)$$

The compatibility condition $\partial_k(\partial_i p) = \partial_i(\partial_k p)$ yields

$$\partial_{ik}u = \varphi_{ik}\partial_i u \partial_k u. \tag{31}$$

Thus, the nonlinear PDE's system (30), (31) is in involution iff the functions φ_{ik} do not depend on p and the compatibility conditions $\partial_j(\partial_i p^k) = \partial_i(\partial_j p^k)$ and $\partial_j(\partial_{ik}u) = \partial_i(\partial_{jk}u)$ are fulfilled *identically*. We call the system (30), (31) the Gibbons–Tsarev system (cf (13)).

However, in general case the functions φ_{ik} depend on p . The compatibility conditions $\partial_k(\partial_i p^j) = \partial_i(\partial_k p^j)$ yield (cf (31))

$$\partial_{ik}u = \bar{\varphi}_{ik}\partial_i u \partial_k u, \tag{32}$$

where (cf (28))

$$\bar{\varphi}_{ik}(u, \mathbf{p}) \equiv \varphi_{ik}(u, p^i, p^k, p)|_{p=p^j}, \quad i \neq j \neq k.$$

Thus, the Gibbons–Tsarev system is determined by (30) and (32).

Finally, one must check the compatibility conditions $\partial_j(\partial_{ik}u) = \partial_i(\partial_{jk}u)$

$$\begin{aligned} \partial_u(\varphi_{ik} - \varphi_{jk}) + \varphi_{ij}(\varphi_{ik} - \varphi_{jk}) + f_{ji}\partial_{p^i}\varphi_{ik} + f_{jk}\partial_{p^k}\varphi_{ik} + f_j\partial_p\varphi_{ik} \\ = f_{ij}\partial_{p^j}\varphi_{jk} + f_{ik}\partial_{p^k}\varphi_{jk} + f_i\partial_p\varphi_{jk} \end{aligned}$$

following from (31), and the compatibility conditions $\partial_j(\partial_{ik}u) = \partial_i(\partial_{jk}u)$ following from (32). Such an over-determined nonlinear PDE system is said to be the *extended* Gibbons–Tsarev system.

Let me emphasize that the extended Gibbons–Tsarev system is a system on the sole function $\psi(u; p)$ only. *The general solution of this system yields a classification of integrable hydrodynamic chains.*

Remark. Any N hydrodynamic reduction (23) can be written in the conservative form (see (26))

$$a_t^i = \partial_x \psi(u(\mathbf{a}); a^i), \quad i = 1, 2, \dots, N. \tag{33}$$

If the function $\psi(u(\mathbf{a}); p)$ determines the integrable hydrodynamic-type system (33), then the compatibility conditions $\partial_i(\partial_k p) = \partial_k(\partial_i p)$ satisfy *identically*, where $\partial_k \equiv \partial/\partial a^k$ and (cf (17))

$$\partial_i p = \frac{\psi_u \partial_i u}{\psi_p|_{p=a^i} - \psi_p} \left[1 + \sum \frac{\psi_u|_{p=a^k} \partial_k u}{\psi_p|_{p=a^k} - \psi_p} \right]^{-1}. \tag{34}$$

If the function $\psi(u; p)$ determines the integrable hydrodynamic chain, then the compatibility conditions $\partial_i(\partial_k p) = \partial_k(\partial_i p)$ describe its N component hydrodynamic reductions parameterized by N arbitrary functions of a single variable. If the function $\psi(u; p)$ is unknown, then the compatibility conditions $\partial_i(\partial_k p) = \partial_k(\partial_i p)$ and the compatibility conditions $\partial_j(\partial_{ik}u) = \partial_i(\partial_{jk}u)$ yield the extended Gibbons–Tsarev system.

Let us finally emphasize: the Gibbons–Tsarev system describes N component hydrodynamic reductions parameterized by N arbitrary functions of a single variable for any *a priori* given function $\psi(\mathbf{u}; p)$; the extended Gibbons–Tsarev system describes all possible functions $\psi(\mathbf{u}; p)$ and their N component hydrodynamic reductions parameterized by N arbitrary functions of a single variable.

4. Explicit Hamiltonian hydrodynamic reductions

Suppose the hydrodynamic reductions (14) of the Benney hydrodynamic chain (1) (simultaneously, they are hydrodynamic reductions of the Khohlov–Zabolotzkaya equation (4)) are Hamiltonian (see details in [29])

$$a_t^i = \frac{1}{\varepsilon_i} \partial_x \frac{\partial \mathbf{h}}{\partial a^i},$$

where ε_i are arbitrary constants. Then the Hamiltonian density $\mathbf{h} = \Sigma \varepsilon_k (a^k)^3 / 6 + f(\Delta)$, where $\Delta = \Sigma \varepsilon_k a^k$ and $f'(\Delta) = A^0(\Delta)$. Substitution $A^0(\Delta)$ into (16) yields the choice $A^0(\Delta) = \Delta$ only (up to an insufficient constant factor). This is exactly the so-called waterbag hydrodynamic reduction (see (18)).

Thus, the *main claim* of this section is that the Hamiltonian hydrodynamic reductions (cf (33))

$$a_t^i = \partial_x \psi(\mathbf{u}; a^i)$$

for any hydrodynamic chain determined by the the generating function of conservation laws (see (22) and (26))

$$p_t = \partial_x \psi(\mathbf{u}; p),$$

one can seek in the form

$$a_t^i = \partial_x \left(\delta_i \frac{\partial \mathbf{h}}{\partial a^i} + \gamma_i \sum_{k \neq i} \gamma_k \frac{\partial \mathbf{h}}{\partial a^k} \right), \quad (35)$$

where δ_i and γ_i are some constants (see examples in [23]).

The second step is a reconstruction of the Riemann surface determined by the equation $\lambda(\mathbf{u}; p)$ by virtue of (35) and a computation of the moments $A^k = \Sigma f_n^k(a^n)$, where the functions $f_n^k(a^n)$ are determined from the expansion of the equation of the Riemann surface at infinity ($\lambda \rightarrow \infty$, $p \rightarrow \infty$). Couple of examples are considered below.

5. Generalized Benney hydrodynamic chain

In this section, we restrict our consideration on the first simple and very important particular case (cf (5))

$$p_t = \partial_x [U(p) + u]. \quad (36)$$

Theorem. *N component hydrodynamic reductions (23)*

$$r_t^i = U'(p^i) r_x^i$$

are described by solutions of the Gibbons–Tsarev system (cf (13))

$$\begin{aligned} \partial_{ik} u &= \frac{2\alpha U'(p^i) U'(p^k) + \beta [U'(p^i) + U'(p^k)] + 2\gamma}{(U'(p^i) - U'(p^k))^2} \partial_i u \partial_k u, \\ \partial_i p &= \frac{\partial_i u}{U'(p^i) - U'(p)}, \quad \partial_i p^k = \frac{\partial_i u}{U'(p^i) - U'(p^k)}, \end{aligned} \quad (37)$$

where the function $U(p)$ is a solution of the second-order ODE with respect to the independent variable p

$$U''(p) = \alpha U'^2(p) + \beta U'(p) + \gamma. \quad (38)$$

The compatibility conditions $\partial_j(\partial_{ik}A^0) = \partial_i(\partial_{jk}A^0)$ are identically satisfied for any constants α, β, γ . In general case ($\alpha \neq 0$) the function $U(p)$ can be written in the parametric form only,

$$U = \frac{1}{\alpha(q_1 - q_2)} [q_1 \ln(q - q_1) - q_2 \ln(q - q_2)],$$

$$p = \frac{1}{\alpha(q_1 - q_2)} [\ln(q - q_1) - \ln(q - q_2)].$$

If $\alpha = 0$, but $\beta \neq 0$, then without lost of generality $U = e^p$; if $\beta = 0$, then $U = p^2/2$; if $\alpha \neq 0$, but $\beta = 0$ and $\gamma = 0$, then $U = \ln p$; if $\alpha \neq 0$, but $\beta = 0$, then $U = \ln \sinh p$ (or $U = \ln \cosh p$).

The first step in the reconstruction of integrable hydrodynamic chains is an extracting of any explicit hydrodynamic reduction (see (33))

$$a_t^i = \partial_x [U(a^i) + u(\mathbf{a})]. \tag{39}$$

Then the compatibility conditions $\partial_i(\partial_k p) = \partial_k(\partial_i p)$ yield (cf (15), (16)) (38) two distinct index relationships

$$[U'(a^i) - U'(a^k)](\partial_{iku} + \alpha \partial_i u \partial_k u) = \partial_k u \partial_i \left(\sum \partial_n u \right) - \partial_i u \partial_k \left(\sum \partial_n u \right), \tag{40}$$

and three distinct index relationships

$$[U'(a^i) - U'(a^k)]\partial_j u \partial_{ik} u + [U'(a^k) - U'(a^j)]\partial_i u \partial_{jk} u + [U'(a^j) - U'(a^i)]\partial_k u \partial_{ij} u = 0,$$

where (see (34))

$$\partial_i p = \frac{\partial_i u}{U'(a^i) - U'(p)} \left(\sum \frac{\partial_k u}{U'(a^k) - U'(p)} + 1 \right)^{-1}. \tag{41}$$

Following the previous section we are looking for the Hamiltonian hydrodynamic reductions (39)

$$a_t^i = \partial_x [U(a^i) + u(\mathbf{a})] = \frac{1}{\varepsilon_i} \partial_x \frac{\partial \mathbf{h}}{\partial a^i}. \tag{42}$$

Theorem. *The Hamiltonian hydrodynamic reductions (42) are determined by the Hamiltonian density $\mathbf{h} = \sum \varepsilon_k \int U(a^k) da^k + f(\Delta)$, where $\Delta = \sum \varepsilon_k a^k$, $f'(\Delta) = u(\Delta)$ and $u(\Delta) = \Delta$, if $\alpha = 0$; $u(\Delta) = \ln(\Delta)$, if $\alpha \neq 0$.*

Proof. It can be obtained by the substitution (42) into (40). □

The equation of the Riemann surface (see (41)) can be found in quadratures

$$d\lambda = \frac{e^{\beta p + 2\alpha U(p)}}{u'(\Delta)} dp + e^{\beta p + 2\alpha U(p)} \sum \frac{\varepsilon_k d(p - a^k)}{U'(a^k) - U'(p)} \tag{43}$$

for this Hamiltonian hydrodynamic reduction.

Let us introduce the moments $A^k = \sum \varepsilon_n \int U^k(a^n) da^n$, then the hydrodynamic-type system (42) can be rewritten as the Hamiltonian hydrodynamic chain

$$A_t^k = \sum_{n=0}^{k+2} F_n^k(\mathbf{A}) A_x^n, \quad k = 0, 1, 2, \dots$$

determined by the Hamiltonian density $\mathbf{h}(A^0, A^1) = A^1 + f(A^0)$ (where $f(A^0) = (A^0)^2/2$, if $\alpha = 0$ and $f(A^0) = A^0 \ln A^0$) and by the *Dorfman* Poisson bracket (see [30])

$$\{A^0, A^0\} = \sum \varepsilon_n \delta'(x - x'),$$

$$\{A^k, A^n\} = [k(\alpha A^{k+n+1} + \beta A^{k+n} + \gamma A^{k+n-1})\partial_x + n\partial_x(\alpha A^{k+n+1} + \beta A^{k+n} + \gamma A^{k+n-1})]\delta(x - x').$$

For instance, this hydrodynamic chain is

$$A_t^0 = \partial_x \left[\alpha A^2 + \beta A^1 + \left(\gamma + \sum \varepsilon_n \right) A^0 \right]$$

$$A_t^k = (\alpha A^{k+2} + \beta A^{k+1} + \gamma A^k)_x + k(\alpha A^{k+1} + \beta A^k + \gamma A^{k-1})A_x^0, \quad k = 1, 2, \dots,$$

if the Hamiltonian density is $\mathbf{h}(A^0, A^1) = A^1 + (A^0)^2/2$.

6. The Gibbons–Tsarev system and hydrodynamic chains

In this section we consider the second simple and very important particular case

$$p_t = \partial_x [V(p)v]. \quad (44)$$

Then N component hydrodynamic reductions (23)

$$r_t^i = v V'(p^i) r_x^i$$

are compatible with the above generating function of conservation laws iff (see (29))

$$\partial_i p = V(p) \frac{\partial_i \ln v}{V'(p^i) - V'(p)}.$$

It is easy to see that under re-scaling

$$\frac{dp}{V(p)} = d\tilde{p}, \quad V'(p) = U'(\tilde{p}),$$

one gets the main formula from the previous example (see (37)), where $u = \ln v$. The corresponding equation is (see (38))

$$V V'' = \alpha V'^2 + \beta V' + \gamma, \quad (45)$$

where

$$V(p) = \exp U(\tilde{p}), \quad dp = \exp U(\tilde{p}) d\tilde{p}. \quad (46)$$

In general case ($\alpha \neq 0$) the function $V(p)$ can be written in the parametric form only

$$V(p) = (q - q_1)^{\frac{q_1}{\alpha(q_1 - q_2)}} (q - q_2)^{-\frac{q_2}{\alpha(q_1 - q_2)}},$$

$$p = \frac{1}{\alpha} \int (q - q_1)^{\frac{q_1}{\alpha(q_1 - q_2)} - 1} (q - q_2)^{-\frac{q_2}{\alpha(q_1 - q_2)} - 1} dq.$$

The corresponding degenerate cases (see the above example) are $V(p) = p^k$, where k is an arbitrary constant, $V(p) = e^p$, $V(p) = \cosh p$ (or $V(p) = \sinh p$) and two others given in the implicit form

$$p = \int \frac{dV}{\ln V}, \quad p = \int \frac{dV}{\sqrt{\ln V}}.$$

Since the equations describing hydrodynamic reductions of both these examples (36) and (44) are equivalent, one can recalculate any hydrodynamic reduction of one example to a corresponding hydrodynamic reduction of the second example (the Hamiltonian reductions of the generating function of conservation laws (44) are found in [23]). The relationship (46) is

nontrivial. For instance, $U(\tilde{p}) = \tilde{p}^2/2$ for the Benney hydrodynamic chain (see (5)), but $V(p)$ cannot be expressed via known elementary or special functions. Nevertheless, hydrodynamic reductions of the Benney hydrodynamic chain can be recalculated. One can compare the formulae (34) for (36) and (44)

$$\frac{\partial p}{\partial a^i} = \frac{V(p)\partial v/\partial a^i}{V'(a^i) - V'(p)} \left[v + \sum \frac{V(a^k)\partial v/\partial a^k}{V'(a^k) - V'(p)} \right]^{-1},$$

$$\frac{\partial \tilde{p}}{\partial c^i} = \frac{\partial u/\partial c^i}{U'(c^i) - U'(p)} \left[1 + \sum \frac{\partial u/\partial c^k}{U'(c^k) - U'(p)} \right]^{-1}.$$

They are equivalent under the transformation (46) and under the same transformation for the field variables $a^k \leftrightarrow c^k$,

$$V(a^i) = \exp U(c^i), \quad da^i = \exp U(c^i) dc^i.$$

The equation of the Riemann surface (43) also can be recalculated,

$$d\lambda = (V(p))^{2\alpha-1} \exp \left[\beta \int \frac{dp}{V(p)} \right] \left(\frac{v(\Delta) dp}{v'(\Delta)} + \sum \frac{\varepsilon_k}{V(a^k)} \frac{V(a^k) dp - V(p) da^k}{V'(a^k) - V'(p)} \right),$$

where $\Delta = \sum \varepsilon_k \int da^k / V(a^k)$, $v(\Delta) = \exp \Delta$, if $\alpha = 0$; $v(\Delta) = \Delta$, if $\alpha \neq 0$.

Example. The Lagrangian quasilinear equation associated with an elliptic curve.

The Lagrangian

$$L = \int z_x z_y z_t dx dy dt$$

creates the Euler–Lagrange equation

$$z_t z_{xy} + z_y z_{xt} + z_x z_{yt} = 0, \tag{47}$$

which is an integrable 2+1 quasilinear equation (see [15]). The pseudopotentials (cf (4), (5) and (6); see also [11, 13]) are

$$\frac{S_x}{z_x} = \zeta(\sigma), \quad \frac{S_y}{z_y} = \zeta(\sigma) + \frac{\wp'(\sigma) + \varepsilon}{2\wp(\sigma)}, \quad \frac{S_t}{z_t} = \zeta(\sigma) + \frac{\wp'(\sigma) - \varepsilon}{2\wp(\sigma)}, \tag{48}$$

where $\zeta(\sigma)$ and $\wp'(\sigma)$ are Weiershtrass elliptic functions ($\zeta'(\sigma) = -\wp(\sigma)$, $\wp'^2(\sigma) = 4\wp^3(\sigma) + \varepsilon^2$). Introducing the new functions $a = z_x$, $b = z_y$, $c = z_t$, (47) can be written in the form

$$a_t = c_x, \quad a_y = b_x, \quad b_t = c_y, \quad ca_y + ba_t + ab_t = 0.$$

Correspondingly, (48) can be written as the couple of generating functions of conservation laws

$$p_y = \partial_x \left[\left(\frac{p}{a} + \frac{\wp'(\sigma) + \varepsilon}{2\wp(\sigma)} \right) b \right], \quad p_t = \partial_x \left[\left(\frac{p}{a} + \frac{\wp'(\sigma) - \varepsilon}{2\wp(\sigma)} \right) c \right], \tag{49}$$

where $p = S_x$ and σ can be found from $\zeta(\sigma) = p/a$.

The reciprocal transformation (see [31, 32])

$$dz = a dx + b dy + c dt, \quad d\tilde{t} = dt, \quad d\tilde{y} = dy$$

reduces the above Lagrangian to

$$\tilde{L} = \int \frac{x_{\tilde{y}} x_{\tilde{t}}}{x_z^2} dz d\tilde{y} d\tilde{t}$$

and (49) to the couple of generating functions of conservation laws

$$\tilde{p}_y = \partial_z \left[\frac{\wp'(\sigma) + \varepsilon}{2\wp(\sigma)} b \right], \quad \tilde{p}_t = \partial_z \left[\frac{\wp'(\sigma) - \varepsilon}{2\wp(\sigma)} c \right], \quad (50)$$

where $\tilde{p} = \zeta(\sigma)$. However, these generating functions of conservation laws are considered in this section (see (44)). Indeed, equation (45)

$$V \frac{\partial^2 V}{\partial \tilde{p}^2} = 3 \left(\frac{\partial V}{\partial \tilde{p}} \right)^2 + 9 \frac{\partial V}{\partial \tilde{p}} + 6$$

has the general solution given in the parametric form

$$V(\tilde{p}) = \frac{\wp'(\sigma) \pm \varepsilon}{2\wp(\sigma)}, \quad \tilde{p} = \zeta(\sigma).$$

7. Commuting flows and 2+1 quasilinear systems

In this section three different approaches in construction of commuting flows for any given hydrodynamic chain are presented.

- (1) Suppose all generating functions of conservation laws (26) are enumerated. Then generating functions of conservation laws for commuting flows one should seek in the forms

$$p_{t^i} = \partial_x \psi_i(u^1; p), \quad p_{t^2} = \partial_x \psi_2(u^1, u^2; p), \dots \quad (51)$$

The compatibility conditions $\partial_t(p_{t^i}) = \partial_{t^i}(p_t)$, $\partial_t(p_{t^2}) = \partial_{t^2}(p_t)$, ... allow us to reconstruct functions ψ_k in quadratures (the corresponding example is given by the Benney hydrodynamic chain (1) and the Khohlov–Zabolotzkaya equation (4); see (5) and (6)).

Example. Let us consider the couple of commuting generating functions of conservation laws (cf (50))

$$p_y = \partial_x [V(p)b], \quad p_t = \partial_x [W(p)c].$$

The compatibility condition $\partial_t(p_y) = \partial_y(p_t)$ yields (45)

$$V V'' = \left(1 + \frac{\beta}{\alpha} \right) V'^2 + \left[\gamma - \delta - 2 \frac{\beta\delta}{\alpha} \right] V' + \frac{\delta}{\alpha} (\beta\delta - \alpha\gamma),$$

$$W W'' = \left(1 + \frac{\delta}{\gamma} \right) W'^2 + \left[\alpha - \beta - 2 \frac{\beta\delta}{\gamma} \right] W' + \frac{\beta}{\gamma} (\beta\delta - \alpha\gamma),$$

where 2+1 quasilinear system is ($\alpha, \beta, \gamma, \delta$ are arbitrary constants)

$$b_t = \alpha b c_x + \beta c b_x, \quad c_y = \gamma c b_x + \delta b c_x.$$

- (2) Let us replace $u^1 \rightarrow p(\zeta)$ and $t^1 \rightarrow \tau(\zeta)$ in the first equation (51)

$$\partial_{\tau(\zeta)} p(\lambda) = \partial_x \psi_1(p(\zeta), p(\lambda)). \quad (52)$$

Definition. Equation (52) is called the generating function of conservation laws and commuting flows.

All generating functions of conservation laws (51) can be obtained by the expansion of (52) in the series according to the Bürmann–Lagrange series of $p(\zeta)$, where $\partial_{\tau(\zeta)}$ is the formal series, whose coefficients ∂_{t^k} enumerate different commuting flows.

Example. The Benney hydrodynamic chain has the generating function of conservation laws (5) and the generating function of conservation laws and commuting flows is

$$\partial_{\tau(\zeta)} p(\lambda) = \partial_x \ln[p(\lambda) - p(\zeta)], \tag{53}$$

where $A^0_{\tau(\zeta)} = \partial_x p(\zeta)$. Substituting the expansion (7) for $p(\lambda)$ into the above formula, one can obtain the generating function of commuting flows in the form

$$A^0_{\tau(\zeta)} = \partial_x p(\zeta), \quad A^1_{\tau(\zeta)} = \partial_x \left(\frac{(A^0)^2}{2} + p(\zeta) \right), \dots$$

Substituting the expansion (7) for $p(\lambda)$ into the above formula, one can obtain the generating functions of conservation laws in the form (see (5) and (6))

$$p_{t^1} = \partial_x \left(\frac{p^2}{2} + A^0 \right), \quad p_{t^2} = \partial_x \left(\frac{p^3}{3} + A^0 p + A^1 \right), \dots$$

- (3) Suppose for any given function $\psi(u; p)$ (see (26)) we already know the corresponding Hamiltonian hydrodynamic reductions

$$a^i_{t^1} = \partial_x \left(\bar{g}^{ik} \frac{\partial \mathbf{h}_1}{\partial a^k} \right),$$

where \bar{g}^{ik} is a constant non-degenerate symmetric matrix. Then one can seek the *higher* conservation law density \mathbf{h}_2 for this hydrodynamic-type system. This conservation law density determines the higher commuting flow

$$a^i_{t^2} = \partial_x \left(\bar{g}^{ik} \frac{\partial \mathbf{h}_2}{\partial a^k} \right).$$

The generating function of conservation laws for this hydrodynamic-type system (as well as for the corresponding hydrodynamic chain) can be found by the replacement $a^i \rightarrow p$ (see details in [23]). The generating functions of commuting flows are given by

$$a^i_{\tau(\zeta)} = \partial_x \left(\bar{g}^{ik} \frac{\partial p(\zeta)}{\partial a^k} \right).$$

Let us replace $\partial_{\tau(\zeta)} \rightarrow \partial_{t^{-1}}$ and, correspondingly, $p(\zeta) \rightarrow A^{-1}$, then the generating function of conservation laws (53) can be written in the form

$$\tilde{p}_x = \partial_{t^{-1}} (e^{\tilde{p}} + A^{-1}), \tag{54}$$

where the *generating function of the Miura-type transformations* (see [33]) is given by $\tilde{p} = \ln(p - A^{-1})$. This generating function (54) determines the continuum limit of the discrete KP hierarchy (see [6])

$$B_x^k = B_z^{k+1} + k B^k B_z^0, \quad k = 0, 1, 2, \dots,$$

where $z \equiv t^{-1}$ and $B^0 \equiv A^{-1}$.

Remark. All moments A^k are connected with the moments B^k by the *Miura-type transformations*

$$A^0 = A^0(B^0, B^1), \quad A^1 = A^1(B^0, B^1, B^2), \quad A^2 = A^2(B^0, B^1, B^2, B^3), \dots,$$

which can be obtained by a substitution the generating function of the Miura-type transformations $p = A^{-1} + \exp \tilde{p}$ into (10) and comparing with (see [6])

$$\lambda = e^{\tilde{p}} + B^0 + B^1 e^{-\tilde{p}} + B^2 e^{-2\tilde{p}} + \dots$$

Let us substitute the expansion $\partial_{\tau(\zeta)} = \partial_{t^{-1}} + \zeta \partial_{t^{-2}} + \dots$ and $p(\zeta) \rightarrow A^{-1} + \zeta A^{-2} + \dots$ into (53). Then the compatibility condition $\partial_{t^{-2}}(\partial_{t^{-1}} p) = \partial_{t^{-1}}(\partial_{t^{-2}} p)$ yields the 2+1 shallow water system (see [11])

$$\partial_{t^{-2}} A^0 = \partial_x A^{-2}, \quad \partial_{t^{-1}} A^{-1} = \partial_x \left(\frac{(A^{-1})^2}{2} + A^0 \right), \quad \partial_{t^{-1}} A^{-2} = \partial_x (A^{-1} A^{-2}), \quad (55)$$

where

$$\partial_{t^{-2}} p = \partial_x \frac{A^{-2}}{A^{-1} - p}.$$

The compatibility condition $\partial_{t^{-2}}(\partial_{t^{-1}} p) = \partial_{t^{-1}}(\partial_{t^{-2}} p)$ yields the famous Boyer–Finley equation

$$\partial_{t^{-1}} A^{-1} = \partial_x \ln A^{-2}, \quad \partial_{t^{-1}} A^{-2} = \partial_{t^{-2}} A^{-1}. \quad (56)$$

In both cases (55) and (56) N component hydrodynamic reductions (11) and (see (23))

$$r_{t^{-1}}^i = \frac{1}{p^i - A^{-1}} r_x^i, \quad r_{t^{-2}}^i = \frac{A^{-2}}{(p^i - A^{-1})^2} r_x^i$$

are compatible (i.e. $\partial_{t^{-1}}(r_{t^{-2}}^i) = \partial_{t^{-2}}(r_{t^{-1}}^i)$; see [13, 24])

$$\frac{\partial_k p^i}{p^k - p^i} = \frac{\partial_k \left(\frac{1}{p^i - A^{-1}} \right)}{\frac{1}{p^k - A^{-1}} - \frac{1}{p^i - A^{-1}}} = \frac{\partial_k \left(\frac{A^{-2}}{(A^{-1} - p^i)^2} \right)}{\frac{A^{-2}}{(A^{-1} - p^k)^2} - \frac{A^{-2}}{(A^{-1} - p^i)^2}},$$

iff the functions p^i and A^0 satisfy the Gibbons–Tsarev system (13), where

$$\partial_i A^{-1} = \frac{\partial_i A^0}{p^i - A^{-1}}, \quad \partial_i \ln A^{-2} = \frac{\partial_i A^0}{(p^i - A^{-1})^2}.$$

It is not easy to verify, if did not take into account, that all these 2+1 quasilinear systems (4), (55) and (56) are members of the unique Benney hydrodynamic *lattice* (see [22]).

8. General case

The *main statement* of this paper is that all integrable hydrodynamic chains described by the generating functions of conservation laws (21) can be split into sub-classes according to the number M of the functions u^m . The first such a case (26) was considered in the previous section ($M = 1$). The second case ($M = 2$) is

$$p_t = \partial_x \psi(u, v; p).$$

Then N component hydrodynamic reductions (23) are compatible with this generating function if the compatibility conditions $\partial_i(\partial_k p) = \partial_k(\partial_i p)$ are fulfilled, where

$$\partial_i p = \frac{\psi_u \partial_i u + \psi_v \partial_i v}{\psi_p|_{p=p^i} - \psi_p}.$$

To complete this computation, we need some relationship between u and v (except trivial link $v(u)$). Without loss of generality we can choose these field variables u and v as a conservation law density and a flux, respectively:

$$u_t = v_x.$$

Since the hydrodynamic-type system (23) has this conservation law,

$$\partial_i v = \psi_p|_{p=p^i} \partial_i u,$$

and (cf (27), (29))

$$\partial_i p = \frac{\psi_u + \psi_v \psi_p|_{p=p^i}}{\psi_p|_{p=p^i} - \psi_p} \partial_i u.$$

However, in such a case the Gibbons–Tsarev system can be derived from the extended compatibility conditions $\partial_i(\partial_k p) = \partial_k(\partial_i p)$ and $\partial_i(\partial_k v) = \partial_k(\partial_i v)$.

9. Conclusion and outlook

In this paper, a new look on a classification of the integrable hydrodynamic chains is presented. The main object which should be under an investigation is the generating function of conservation laws

$$p_t = \partial_x \psi(\mathbf{u}; p),$$

where all distinct cases are separated by the number M of independent functions u^m . The simplest case (26) is considered in detail. Each such case has infinitely many sub-cases enumerated by the number K of independent functions v^k in the commuting generating functions of conservation laws

$$p_y = \partial_x \varphi(v; p).$$

Thus, all integrable hydrodynamic chains can be split into sub-classes by virtue of the two numbers M and K only.

Suppose all these functions $\psi(\mathbf{u}; p)$ are found. Then the Gibbons–Tsarev system describing N component hydrodynamic reductions for every function $\psi(\mathbf{u}; p)$ can be derived automatically in the Riemann invariants and in the field variables a^k (which are conservation law densities) simultaneously. Then corresponding hydrodynamic-type systems with a local Hamiltonian structure (as well as with *a priori* prescribed any nonlocal Hamiltonian structure) can be extracted. The equation of the Riemann surface $\lambda(\mathbf{u}; p)$ can be found in quadratures for the hydrodynamic reductions, whose characteristic velocities are invariant with respect to any Lie group symmetry. Asymptotic of the equation of the Riemann surface in the vicinity of any singular point (usually $\lambda \rightarrow \infty$, $p \rightarrow \infty$) yields explicit expressions of corresponding integrable hydrodynamic chains. The ‘flat’ hydrodynamic reductions can be used for a construction of a large class of particular solutions for these hydrodynamic chains and related 2+1 quasilinear equations.

Acknowledgments

I thank Eugeni Ferapontov, John Gibbons, Yuji Kodama, Boris Kupershmidt and Sergey Tsarev for their stimulating and clarifying discussions. I am grateful to the Institute of Mathematics in Taipei (Taiwan) where some part of this work has been done, and especially to Derchyi Wu, Jen-Hsu Chang, Ming-Hien Tu and Jyh-Hao Lee for fruitful discussions.

References

- [1] Benney D J 1973 Some properties of long non-linear waves *Stud. Appl. Math.* **52** 45–50
- [2] Kuperschmidt B A and Manin Yu I 1977 Long wave equation with free boundaries: I. Conservation laws *Funct. Anal. Appl.* **11** (no 3) 188–97
- Kuperschmidt B A and Manin Yu I 1978 Long wave equations with a free surface: II. The Hamiltonian structure and the higher equations *Funct. Anal. Appl.* **12** (no 1) 20–9
- Lebedev D R and Manin Yu I 1979 Conservation laws and representation of Benney’s long wave equations *Phys. Lett. A* **74** (no 3, 4) 154–6

- Lebedev D R 1979 Benney's long wave equations: Hamiltonian formalism *Lett. Math. Phys.* **3** 481–8
- [3] Zakharov V E 1980 Benney's equations and quasi-classical approximation in the inverse problem method *Funct. Anal. Appl.* **14** 89–98
- Zakharov V E 1981 On the Benney's equations *Physica D* **3** 193–200
- [4] Gibbons J 1981 Collisionless Boltzmann equations and integrable moment equations *Physica D* **3** 503–11
- [5] Gibbons J and Tsarev S P 1996 Reductions of Benney's equations *Phys. Lett. A* **211** 19–24
- Gibbons J and Tsarev S P 1999 Conformal maps and reductions of the Benney equations *Phys. Lett. A* **258** 263–70
- [6] Kodama Yu 1988 A method for solving the dispersionless KP equation and its exact solutions *Phys. Lett. A* **129** 223–6
- Kodama Yu 1988 A solution method for the dispersionless KP equation *Prog. Theor. Phys. Suppl.* **94** 184
- [7] Gibbons J and Kodama Yu 1994 Solving dispersionless Lax equations *Singular Limits of Dispersive Waves (NATO ASI Series B vol 320)* ed N Ercolani (New York: Plenum) p 61
- [8] Krichever I M 1992 The dispersionless equations and topological minimal models *Commun. Math. Phys.* **143** 415–29
- Krichever I M 1994 The τ -function of the universal Whitham hierarchy, matrix models and topological field theories *Commun. Pure Appl. Math.* **47** 437–75
- [9] Bogdanov L V and Konopelchenko B G 2004 Symmetry constraints for dispersionless integrable equations and systems of hydrodynamic type *Phys. Lett. A* **330** 448–59
- [10] Kupershmidt B A 1983 Deformations of integrable systems *Proc. R. Irish Acad. Sect. A* **83** 45–74
- Kupershmidt B A 1984 Normal and universal forms in integrable hydrodynamical systems *Proc. Berkeley-Ames Conf. on Nonlinear Problems in Control and Fluid Dynamics (Berkeley, CA, 1983) (Lie Groups: Hist., Frontiers and Appl. Ser. B: Systems Inform. Control, II)* (Brookline, MA: Math Sci Press) pp 357–78
- [11] Zakharov V E 1994 Dispersionless limit of integrable systems in 2+1 dimensions *Singular Limits of Dispersive Waves* ed N M Ercolani *et al* (New York: Plenum) pp 165–74
- [12] Pavlov M V 2004 Classification of the Egorov hydrodynamic chains *Theor. Math. Phys.* **138** 55–71
- [13] Ferapontov E V and Khusnutdinova K R 2004 On integrability of (2+1)-dimensional quasilinear systems *Commun. Math. Phys.* **248** 187–206
- Ferapontov E V and Khusnutdinova K R 2004 The characterization of 2-component (2+1)-dimensional integrable systems of hydrodynamic type *J. Phys. A: Math. Gen.* **37** 2949–63
- [14] Ferapontov E V, Khusnutdinova K R and Pavlov M V 2005 Classification of integrable (2+1) dimensional quasilinear hierarchies *Theor. Math. Phys.* **144** 35–43
- [15] Ferapontov E V, Khusnutdinova K R and Tsarev S P 2006 On a class of three dimensional integrable lagrangians *Commun. Math. Phys.* **261** 225–43
- [16] Ferapontov E V and Marshall D G 2005 Differential-geometric approach to the integrability of hydrodynamic chains: the Haanties tensor *Preprint nlin.SI/0505013*
- [17] Błaszak M and Szablikowski B M 2002 Classical R -matrix theory of dispersionless systems: I. (1+1)-dimension theory *J. Phys. A: Math. Gen.* **35** 10325–44
- Błaszak M and Szablikowski B M 2002 Classical R -matrix theory of dispersionless systems: II. (2+1)-dimension theory *J. Phys. A: Math. Gen.* **35** 10345–64
- [18] Pavlov M V 2003 Integrable hydrodynamic chains *J. Math. Phys.* **44** 4134–56
- [19] Chang J H 2005 Hydrodynamic reductions of dispersionless Harry dym hierarchy *J. Phys. A: Math. Gen.* **38** 6505–15
- [20] Chen Y T and Tu M H 2003 A note on the dispersionless dym hierarchy *Lett. Math. Phys.* **63** 125–39
- [21] Manas M, Alonso L M and Medina E 2002 Reductions of the dispersionless KP hierarchy *Theor. Math. Phys.* **133** 1712–21
- Konopelchenko B, Alonso L M and Medina E 2002 Quasiconformal mappings and solutions of the dispersionless KP hierarchy *Theor. Math. Phys.* **133** 1529–38
- [22] Pavlov M V 2006 The Kupershmidt hydrodynamic chains and lattices *Int. Math. Res. Not.* submitted
- [23] Pavlov M V 2006 Algebro-geometric approach in the theory of integrable hydrodynamic type systems *Comm. Math. Phys.* at press
- [24] Tsarev S P 1985 On poisson brackets and one-dimensional hamiltonian systems of hydrodynamic type *Sov. Math. Dokl.* **31** 488–91
- Tsarev S P 1991 The geometry of hamiltonian systems of hydrodynamic type. The generalized hodograph method *Math. USSR Izvestiya* **37** 397–419
- [25] Pavlov M V 2006 The Hamiltonian approach in the classification and the integrability of hydrodynamic chains *Physica D* submitted
- [26] Gibbons J and Yu L A 2000 The initial value problem for reductions of the Benney equations *Inverse Problems* **16** 605–18

- Yu L A 2000 Waterbag reductions of the dispersionless discrete KP hierarchy *J. Phys. A: Math. Gen.* **33** 8127–38
- [27] Dubrovin B A 1992 Hamiltonian formalism of Whitham-type hierarchies and topological landau-ginsburg models *Commun. Math. Phys.* **145** 195–207
- Dubrovin B A 1996 *Geometry of 2D Topological Field Theories (Lecture Notes in Mathematics vol 1620)* (Berlin: Springer) pp 120–348
- [28] Lavrentiev M A and Shabat B V 1965 *Metody teorii funktsii kompleksnogo peremennogo* (Russian) (Methods of the Theory of Functions of a Complex Variable) 3rd corrected edn Izdat. (Moscow: Nauka) 716 pp
- [29] Dubrovin B A and Novikov S P 1983 Hamiltonian formalism of one-dimensional systems of hydrodynamic type and the bogolyubov-whitham averaging method *Sov. Math. Dokl.* **27** 665–9
- Dubrovin B A and Novikov S P 1989 Hydrodynamics of weakly deformed soliton lattices. Differential geometry and Hamiltonian theory *Russ. Math. Surveys* **44** 35–124
- [30] Dorfman I Ya 1993 Dirac structures and integrability of nonlinear evolution equations *Nonlinear Science: Theory and Applications* (New York: Wiley) p 176
- [31] Rogers C and Shadwick W F 1982 *Bäcklund Transformations and their Applications* (New York: Academic)
- [32] Rozhddestvenski B L and Yanenko N N 1983 Systems of quasilinear equations and their applications to gas dynamics *Translations of Mathematical Monographs* vol 55 Translated from the second Russian edition by J R Schulenberger (Providence, RI: American Mathematical Society) (Moscow: Nauka, 1968, Russian edn.)
- [33] Pavlov M V 2006 Hydrodynamic chains and the classification of their Poisson brackets *J. Math. Phys.* submitted
- [34] Ferapontov E V, Khusnutdinova K, Marshall M V and Pavlov M V 2006 Classification of integrable Hamiltonian hydrodynamic chains associated with Kuperschmidt's brackets *J. Math. Phys.* submitted