

A unification in the theory of linearization of second-order nonlinear ordinary differential equations

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2006 J. Phys. A: Math. Gen. 39 L69

(<http://iopscience.iop.org/0305-4470/39/3/L01>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.105

The article was downloaded on 03/06/2010 at 04:42

Please note that [terms and conditions apply](#).

LETTER TO THE EDITOR

A unification in the theory of linearization of second-order nonlinear ordinary differential equations

V K Chandrasekar, M Senthilvelan and M Lakshmanan

Centre for Nonlinear Dynamics, Department of Physics, Bharathidasan University,
Tiruchirapalli-620 024, India

Received 18 October 2005, in final form 25 November 2005

Published 21 December 2005

Online at stacks.iop.org/JPhysA/39/L69

Abstract

In this letter, we introduce a new generalized linearizing transformation (GLT) for second-order nonlinear ordinary differential equations (SNODEs). The well-known invertible point (IPT) and non-point transformations (NPT) can be derived as sub-cases of the GLT. A wider class of nonlinear ODEs that cannot be linearized through NPT and IPT can be linearized by this GLT. We also illustrate how to construct GLTs and to identify the form of the linearizable equations and propose a procedure to derive the general solution from this GLT for the SNODEs. We demonstrate the theory with two examples which are of contemporary interest.

PACS numbers: 02.30.Hq, 05.45.–a

Linearizing nonlinear ordinary differential equations (NODEs) is still an open problem in the theory of differential equations [1–3]. If one raises the question whether a given arbitrary nonlinear ODE is linearizable or not, no definitive answer can be given in general. Three main points which need attention for further understanding of this problem are: (i) there is still no comprehensive literature available on the types of transformations that can linearize the ODEs, (ii) the general form of linearizable equation also differs from transformation to transformation and (iii) higher-order ODEs possess a greater variety of linearizing transformations than the lower-order ODEs. Due to these reasons no general treatment on linearizing transformations or linearizable equations has been formulated so far.

In this letter, we make an attempt to unify the linearizing transformations known for the case of second-order nonlinear ODEs (SNODEs) and extend their scope. As far as the SNODEs are concerned it has been shown that, in general, one can linearize them through two different kinds of transformations. One is the well-known invertible point transformation (IPT) and the other is the non-point transformation (NPT). As far as the IPT is concerned it has been shown [3–8] that the most general SNODE that can be linearized through such a transformation,

$$X = F(t, x), \quad T = G(t, x), \quad (1)$$

is of the form

$$\ddot{x} = D(t, x)\dot{x}^3 + C(t, x)\dot{x}^2 + B(t, x)\dot{x} + A(t, x), \quad (2)$$

where over dot denotes differentiation with respect to t and the functions A , B , C and D should satisfy the following two equations:

$$\begin{aligned} 3D_{tt} + 3BD_t - 3AD_x + 3DB_t + B_{xx} - 6DA_x + CB_x - 2CC_t - 2C_{tx} &= 0, \\ C_{tt} + 6AD_t - 3AC_x + 3DA_t - 2B_{tx} - 3CA_x + 3A_{xx} + 2BB_x - BC_t &= 0. \end{aligned} \quad (3)$$

The transformation (1) converts equation (2) into the linear ‘free particle’ equation,

$$\frac{d^2X}{dT^2} = 0. \quad (4)$$

On the other hand, it has also been shown that one can consider NPTs of the form

$$X = \hat{F}(t, x), \quad dT = \hat{G}(t, x) dt, \quad (5)$$

and linearize the given SNODE. The most general SNODE that can be linearized through the transformation (5) possess the form [9]

$$\ddot{x} + A_2(t, x)\dot{x}^2 + A_1(t, x)\dot{x} + A_0(t, x) = 0. \quad (6)$$

The set of relations between the functions A_i 's, $i = 0, 1, 2$, and the transformation (5) is given by

$$\begin{aligned} A_2 &= (\hat{G}\hat{F}_{xx} - \hat{F}_x\hat{G}_x)/K, \\ A_1 &= (2\hat{G}\hat{F}_{xt} - \hat{F}_x\hat{G}_t - \hat{F}_t\hat{G}_x)/K, \\ A_0 &= (\hat{G}\hat{F}_{tt} - \hat{F}_t\hat{G}_t)/K \end{aligned} \quad (7)$$

with $K = \hat{F}_x\hat{G} \neq 0$. The NPT also transforms equation (6) to the free particle equation (4). The functions A_i 's, $i = 0, 1, 2$, should satisfy the following relations [9]:

(i)

$$S_1(t, x) = A_{1x} - 2A_{2t} = 0, \quad (8)$$

$$S_2(t, x) = 2A_{0xx} - 2A_{1tx} + 2A_0A_{2x} - A_{1x}A_1 + 2A_{0x}A_2 + 2A_{2tt} = 0. \quad (9)$$

(ii) If $S_1(t, x) \neq 0$ and $S_2(t, x) \neq 0$, then

$$S_2^2 + 2S_{1t}S_2 - 2S_1^2A_{1t} + 4S_1^2A_{0x} + 4S_1^2A_0A_2 - 2S_1S_{2t} - S_1^2A_1^2 = 0, \quad (10)$$

$$S_{1x}S_2 + S_1^2A_{1x} - 2S_1^2A_{2t} - S_1S_{2x} = 0. \quad (11)$$

The NPT is also called a generalized Sundman transformation, see for example [10, 11].

Even though both the IPT and NPT transform the second-order nonlinear ODE to the free particle equation (4), the NPT has some disadvantages over the former. For example, in the case of IPT one can unambiguously invert the free particle solution and deduce the solution of the associated nonlinear equation, whereas in the case of NPT it is not so straightforward due to the non-local nature of the independent variable.

In this work, we unearth a more general transformation,

$$X = F(t, x), \quad dT = G(t, x, \dot{x}) dt, \quad (12)$$

and show that this transformation can be utilized to linearize a wider class of SNODEs and, in particular, certain equations which cannot be linearized by the NPT and IPT. We designate this transformation as the *generalized linearizing transformation* (GLT). If the function G in (12) is independent of the variable \dot{x} then it becomes an NPT (vide equation (5)). On the other hand, if G is a perfect differentiable function then it becomes an IPT, that is $G(t, x, \dot{x}) = \frac{d}{dt}\hat{G}(t, x)$,

then $dT = \frac{d\hat{G}}{dt} dt \Rightarrow T = \hat{G}(t, x)$. We stress here that (12) is a unified transformation as it includes IPT and NPT as special cases.

We demonstrate our above assertion with the case where G is a polynomial function in \dot{x} and in particular where it is linear in \dot{x} with coefficients which are arbitrary functions of t and x . Indeed, even such a simple case leads to interesting results as we see below. To be specific, we focus here on the case

$$X = F(t, x), \quad dT = (G_1(t, x)\dot{x} + G_2(t, x)) dt. \quad (13)$$

We note that in equation (13) even if we consider $X = F(t, x, \dot{x})$ and $dT = (G_1\dot{x} + G_2) dt$, after substitution into (16), we deduce that $F_{\dot{x}} = 0$ and so the form (13) is taken. Generalizations involving higher degree polynomials in \dot{x} for $G(t, x, \dot{x})$ will be dealt with elsewhere.

Substituting the transformation (13) into the free particle equation (4), the most general SODE that can be linearized through the GLT (13) can be shown to be of the form

$$\ddot{x} + A_3(t, x)\dot{x}^3 + A_2(t, x)\dot{x}^2 + A_1(t, x)\dot{x} + A_0(t, x) = 0 \quad (14)$$

and the functions A_i 's $i = 0, 1, 2, 3$, are connected to the transformation functions F and G through the relations

$$\begin{aligned} A_3 &= (G_1 F_{xx} - F_x G_{1x})/M, \\ A_2 &= (G_2 F_{xx} + 2G_1 F_{xt} - F_x G_{2x} - F_t G_{1x} - F_x G_{1t})/M, \\ A_1 &= (2G_2 F_{xt} + G_1 F_{tt} - F_x G_{2t} - F_t G_{2x} - F_t G_{1t})/M, \\ A_0 &= (G_2 F_{tt} - F_t G_{2t})/M \end{aligned} \quad (15)$$

with $M = F_x G_2 - F_t G_1 \neq 0$.

For the given equation one has explicit forms for the functions A_i 's. Now solving equation (15) with the known A_i 's, one can get the linearizing transformation functions F and G . Once F and G are known then using (13) we can transform (14) to the free particle equation (4) and solving the latter one can get the first integral. However, it is difficult to integrate it further unambiguously to obtain the general solution due to the non-local nature of the transformation (13). We are able to overcome this problem also here and devise a general procedure to construct the general solution. In the following, we briefly describe the procedure.

Integrating the free particle equation (4) once, we get

$$\frac{dX}{dT} = I_1 = C(t, x, \dot{x}), \quad (16)$$

where I_1 is the first integral. Now rewriting (16) for \dot{x} , we get

$$\dot{x} = f(t, x, I_1), \quad (17)$$

where f is a function of the indicated variables. Due to non-local nature of the independent variable we need to consider only a particular solution for the free particle equation (4), that is

$$X(t, x) = I_1 T \quad (18)$$

from which we get

$$x = g(t, T, I_1), \quad (19)$$

where g is a function of t, T and I_1 . Making use of relations (17) and (19), equation (13) can be rewritten in the form

$$dT = h(t, T, I_1) dt, \quad (20)$$

where again h is a function of t , T and I_1 . We find that in the case of linearizable equations one can separate the variables T and t in equation (20) and integrate the resultant equation which in turn leads to the general solution.

In the above, we have demonstrated how to deduce linearizing transformation and the general solution for the given equation. On the other hand, one can construct both linearizing transformation and specific linearizable equations. To illustrate this let us analyse a particular but important case of equation (14), namely, $A_3 = 0$ and $A_2 = 0$ in equation (15). However, the other choices, for example $A_3 = A_1 = 0$ and A_2 and $A_0 \neq 0$, also lead to many new linearizable equations. These will be dealt with separately. Solving the first and second equation in (15) with this restriction, we obtain

$$G_1 = a(t)F_x, \quad G_2 = a(t)F_t - (a_t x + b(t))F_x, \quad (21)$$

where a and b are arbitrary functions of t . By using equation (21) in the last two equations in (15), we get

$$A_1 = S_x + \frac{a_t}{(a_t x + b)} S + \frac{(a_{tt} x + b_t)}{(a_t x + b)}, \quad (22)$$

$$A_0 = S_t + \frac{a_t}{(a_t x + b)} S^2 + \frac{(a_{tt} x + b_t)}{(a_t x + b)} S, \quad (23)$$

where

$$S(t, x) = \frac{F_t}{F_x}. \quad (24)$$

Solving equation (22), we get

$$S = \frac{(c(t) - b_t x - \frac{1}{2} a_{tt} x^2 + \int A_1 (a_t x + b) dx)}{(a_t x + b)}, \quad (25)$$

where $c(t)$ is an arbitrary function of t . Substituting equation (25) into (23), we obtain

$$A_0 = \frac{a_t (c - b_t x - \frac{1}{2} a_{tt} x^2 + \int (a_t x + b) A_1 dx)^2}{(a_t x + b)^3} + \frac{c_t - b_{tt} x - \frac{1}{2} a_{ttt} x^2 + (\int ((a_{tt} x + b_t) A_1 + (a_t x + b) A_{1t}) dx)}{(a_t x + b)}. \quad (26)$$

The explicit form of F can be determined by substituting the expression for S into (24) and solving the resultant first-order partial differential equation for F . Once F is known G_1 and G_2 can be fixed using the relation (21) which in turn provides us the GLT through (13). The associated linearizable equation assumes the form $\ddot{x} + A_1(t, x)\dot{x} + A_0(t, x) = 0$, where A_0 is given in equation (26) and A_1 is the given function in this analysis.

To illustrate the procedure with a simple but non-trivial example, let us consider the case $A_1 = kx^q$, where k and q are arbitrary parameters, and fix the arbitrary functions a , b and c such as $a(t) = t$, $b(t) = c(t) = 0$, so that the equation (25) gives us

$$S = \frac{k}{(q+2)} x^{q+1}. \quad (27)$$

Once S is known F and A_0 can be fixed through the relations (24) and (26) of the form

$$A_0 = \frac{k^2}{(q+2)^2} x^{2q+1} \quad \text{and} \quad F = \frac{k}{q+2} t - \frac{1}{qx^q}. \quad (28)$$

The forms of A_0 and A_1 fix the linearizable equation (14) to the form

$$\ddot{x} + kx^q \dot{x} + \frac{k^2}{(q+2)^2} x^{2q+1} = 0. \quad (29)$$

Since $a(t) = t$ and $b(t) = 0$, from (21) we have

$$G_1 = \frac{t}{x^{q+1}}, \quad G_2 = \frac{kt}{q+2} - \frac{1}{x^q}. \quad (30)$$

As a consequence the linearizing transformation turns out to be

$$X = \frac{k}{q+2}t - \frac{1}{qx^q}, \quad dT = \left[-t \left(\frac{k}{q+2} + \frac{\dot{x}}{x^{q+1}} \right) + \frac{1}{x^q} \right] dt. \quad (31)$$

It is easy to check that equation (29) can be linearized to the free particle equation (4) through the transformation (31).

Equation (29) and its sub-cases have been widely discussed in the contemporary literature. In particular, Mahomed and Leach [4] have shown that equation (29) with $q = 1$ is one of the SNODEs that can be linearized to the free particle equation (4) through the IPT $X = \frac{k}{3}t - \frac{1}{x}$ and $T = \frac{t}{x} - \frac{kt^2}{6}$. Consequently, the group invariance and integrability properties of this sub-case, namely, $q = 1$, and the general equation (29) have been studied extensively by different authors, see for example [12–16]. However in the literature, equation (29) has been shown to be linearizable to free particle equation only for the value $q = 1$. For other values of q , the linearization of this equation through IPT or NPT was not known. *But in the present work we have proved above that one can linearize the entire class of equation (29) under the one general transformation (31), irrespective of the value of q .* One may note that choosing $q = 1$ the GLT (31) coincides exactly with the point transformation for equation (29) with the same parametric restriction. This example further confirms the arguments that IPT is a sub-case of GLT.

In the following, we derive the general solution of (29) using our procedure discussed through equations (16)–(20). Using (31) into equation (16), we obtain the first integral in the form

$$I_1 = \frac{\left(\frac{k}{q+2}x^{q+1} + \dot{x} \right)}{-t \left(\frac{k}{q+2}x^{q+1} + \dot{x} \right) + x}. \quad (32)$$

Rewriting (32) for \dot{x} , we get

$$\dot{x} = -\frac{k}{q+2}x^{q+1} + \frac{I_1}{(1+I_1t)}x. \quad (33)$$

Making use of the particular solution for the free particle equation given in equation (18) and rewriting this for x in equation (31), we get

$$x = \left(\frac{1}{q \left(\frac{k}{q+2}t - I_1T \right)} \right)^{\frac{1}{q}}. \quad (34)$$

Substituting (33) and (34) in the second equation in (31), we obtain

$$dT = q \left(\frac{1}{1+I_1t} \right) \left(\frac{k}{q+2}t - I_1T \right) dt. \quad (35)$$

Rewriting equation (35) in the form

$$\frac{dT}{dt} + \frac{qI_1}{1+I_1t}T = \frac{kq}{q+2} \left(\frac{t}{1+I_1t} \right) \quad (36)$$

and integrating the resultant equation, (36), we get

$$T = (1+I_1t)^{-q} \left(I_2 + \frac{(1+I_1t)^q (qI_1t - 1)}{I_1^2(2+3q+q^2)} \right), \quad (37)$$

where I_2 is the second integration constant. Substituting the resultant expression for T into (34), we obtain the general solution of (29), that is

$$x(t) = \left(\frac{I_1(q+1)(q+2)(1+I_1t)^q}{q(k(1+I_1t)^{q+1} - I_1^2 I_2(2+3q+q^2))} \right)^{\frac{1}{q}}, \quad (38)$$

which is the same as the one obtained by Feix *et al* [14].

Equation (29) is not an isolated example that can be linearized through the GLT. In fact, one can linearize a larger class of equations through this GLT and obtain the general solution. This is mainly due to the presence of arbitrary functions, namely, $a(t)$, $b(t)$ and $c(t)$ in the determining equations for given $A_1(x, t)$. To demonstrate this, we consider a slightly more general form

$$A_1 = k_1 x^q + k_2, \quad A_3 = A_2 = 0, \quad (39)$$

where k_1 , k_2 and q are arbitrary constants, in equation (14). In the present example, we have included an additive constant, k_2 , in the function A_1 and left the other two functions A_2 and A_3 the same as before. However, this additive constant itself enlarges the class of linearizable equations considerably, as we see below.

Let us again fix the arbitrary functions a , b and c of the same form as in the previous example, that is $a(t) = t$, $b(t) = 0$ and $c(t) = 0$, so that we get $S = \frac{k_2}{2}x + \frac{k_1}{q+2}x^{q+1}$. The respective linearizable equation turns out to be

$$\ddot{x} + (k_1 x^q + k_2)\dot{x} + \frac{k_1^2}{(q+2)^2}x^{2q+1} + \frac{k_1 k_2}{q+2}x^{q+1} + \frac{k_2^2}{4}x = 0 \quad (40)$$

and the GLT becomes

$$X = \frac{2k_1}{q(q+1)k_2} - \left(\frac{1}{qx^q} + \frac{2k_1}{q(q+1)k_2} \right) \exp\left(-\frac{q}{2}k_2 t\right), \quad (41)$$

$$dT = \left[\frac{1}{x^q} \left(1 - \frac{k_2}{2}t\right) - t \left(\frac{\dot{x}}{x^{q+1}} + \frac{k_1}{(q+1)} \right) \right] \exp\left(-\frac{q}{2}k_2 t\right) dt.$$

One may note that in the limit $k_2 \rightarrow 0$ both the linearizing transformations, (41), and the linearizable equation, (40), reduce to the earlier example (vide equations (31) and (29), respectively).

The associated first integral reads

$$I_1 = \frac{dX}{dT} = \frac{\left(\frac{k_2}{2}x + \frac{k_1}{q+2}x^{q+1} + \dot{x}\right)}{-t\left(\frac{k_2}{2}x + \frac{k_1}{q+2}x^{q+1} + \dot{x}\right) + x}. \quad (42)$$

Repeating the same steps given in the previous example one can get the general solution for equation (40) in the form

$$x(t) = (I_1 + t) \exp\left(-\frac{k_2}{2}t\right) \left(I_2 + \frac{qk_1}{(q+2)} \int_0^t \exp\left(-\frac{qk_2}{2}t'\right) (I_1 + t')^q dt' \right)^{-\frac{1}{q}}, \quad (43)$$

where I_2 is the second integration constant.

Next, we choose the arbitrary function $a(t)$ in an exponential form, namely, $a(t) = e^{\alpha t}$, where α is a constant, with $b(t) = c(t) = 0$. In this case, we get

$$S = \left(\frac{k_2 + \alpha}{2} \right) x + \frac{k_1}{q+2} x^{q+1} \quad \text{and} \quad \alpha = \sqrt{k_2^2 - 4\lambda}, \quad (44)$$

where λ is an arbitrary parameter. The functions b , c and A_1 give A_0 through the relation (26) which in turn gives us a new linearizable equation of the form

$$\ddot{x} + (k_1 x^q + k_2) \dot{x} + \frac{k_1^2}{(q+2)^2} x^{2q+1} + \frac{k_1 k_2}{(q+2)} x^{q+1} + \lambda x = 0. \quad (45)$$

Proceeding further, we obtain the GLT in the form

$$X = \left(\frac{(\alpha - k_2)}{2q\lambda} - \frac{(q+2)}{qk_1 x^q} \right) \exp \left(-\frac{q}{2} (k_2 + \alpha) t \right), \quad (46)$$

$$dT = \left[\frac{(q+2)}{k_1 x^{q+1}} \dot{x} + \frac{(q+2)}{2k_1 x^q} (k_2 - \alpha) + 1 \right] \exp \left(-\left(\frac{q}{2} k_2 + \frac{(q-2)}{2} \alpha \right) t \right) dt.$$

Now one can check that the transformations (41) and (46) transform (40) and (45) into the free particle equation (4). The first integrals for equation (46) can be constructed of the form

$$I_1 = \frac{dX}{dT} = e^{-\alpha t} \left(\frac{\dot{x} + \frac{(k_2 + \alpha)}{2} x + \frac{k_1}{q+2} x^{q+1}}{\dot{x} + \frac{(k_2 - \alpha)}{2} x + \frac{k_1}{q+2} x^{q+1}} \right), \quad (47)$$

and the general solution takes the form

$$x(t) = (e^{\alpha t} - I_1) \exp \left(-\frac{1}{2} (k_2 + \alpha) t \right) \left(I_2 + \frac{qk_1}{(q+2)} \int_0^t \left(\frac{e^{\alpha t'} - I_1}{\exp \left(\frac{1}{2} (k_2 + \alpha) t' \right)} \right)^q dt' \right)^{-\frac{1}{q}}, \quad (48)$$

where I_2 is the second integration constant. Equations (43) and (48) can be integrated further explicitly using the standard method [17]. To our knowledge, the solutions (43) and (48) are new to the literature.

We note that in the case $q = 1$, the terms on the right-hand side in the second equation of (46) can be written as a perfect derivative term and consequently leads us to the same IPT for the equation (45), respectively, with $q = 1$ obtained in [15, 16].

In this paper, we have introduced a new generalized linearizing transformation which can be used to linearize a class of equations that cannot be linearized by either IPT or NPT. In fact, both IPT and NPT can be derived as sub-cases from the proposed GLT. Since the independent variable is in a non-local form in the GLT, we have devised an algorithm to rewrite the new variables in terms of old variables. Needless to say, this algorithm can also be used in the case of NPT also. Importantly, we have illustrated our theory with certain concrete examples which are of contemporary interest. Naturally, one can also construct GLTs involving more general forms of \dot{x} in (12) and identify new linearizable equations. The procedure can also be extended to higher-order ODEs. The details will be discussed separately.

Acknowledgments

The work of VKC is supported by CSIR in the form of a CSIR Senior Research Fellowship. The work of MS and ML forms part of a research project sponsored by Department of Science and Technology, Government of India.

References

- [1] Olver P J 1995 *Equivalence, Invariants, and Symmetry* (Cambridge: Cambridge University Press)
- [2] Ibragimov N H 1999 *Elementary Lie Group Analysis and Ordinary Differential Equations* (New York: Wiley)
- [3] Steeb W H 1993 *Invertible Point Transformations and Nonlinear Differential Equations* (London: World Scientific)

-
- [4] Mahomed F M and Leach P G L 1985 *Quaestiones Math.* **8** 241
Mahomed F M and Leach P G L 1989 *Quaestiones Math.* **12** 121
- [5] Duarte L G S, Duarte S E S and Moreira I C 1987 *J. Phys. A: Math. Gen.* **20** L701
Duarte L G S, Duarte S E S and Moreira I C 1989 *J. Phys. A: Math. Gen.* **22** L201
- [6] Duarte L G S, Euler N, Moreira I C and Steeb W H 1990 *J. Phys. A: Math. Gen.* **23** 1457
- [7] Duarte L G S, Moreira I C, Euler N and Steeb W H 1991 *Phys. Scr.* **43** 449
- [8] Euler N, Steeb W H, Duarte L G S and Moreira I C 1991 *Int. J. Theor. Phys.* **30** 1267
- [9] Duarte L G S, Moreira C and Santos F C 1994 *J. Phys. A: Math. Gen.* **27** L739
- [10] Euler N and Euler M 2004 *J. Nonlinear Math. Phys.* **11** 399
- [11] Euler N, Wolf T, Leach P G L and Euler M 2003 *Acta Appl. Math.* **76** 89
- [12] Davis H T 1962 *Introduction to Nonlinear Differential and Integral Equations* (New York: Dover)
- [13] Leach P G L, Feix M R and Bouquet S 1988 *J. Phys. A: Math. Gen.* **29** 2563
Leach P G L 1985 *J. Math. Phys.* **26** 2510
Lemmer R L and Leach P G L 1993 *J. Phys. A: Math. Gen.* **26** 5017
- [14] Feix M R, Geronimi C, Cairo L, Leach P G L, Lemmer R L and Bouquet S 1997 *J. Phys. A: Math. Gen.* **30** 7437
- [15] Chandrasekar V K, Senthilvelan M and Lakshmanan M 2005 *Proc. R. Soc. London A* **461** 2451
- [16] Chandrasekar V K, Senthilvelan M and Lakshmanan M 2005 *Chaos Solitons Fractals* **26** 1399
- [17] Gradshteyn I S and Ryzhik I M 1980 *Table of Integrals, Series and Products* (London: Academic)