

Clean positive operator-valued measures for qubits and similar cases

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2007 J. Phys. A: Math. Theor. 40 4817

(<http://iopscience.iop.org/1751-8121/40/18/009>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.109

The article was downloaded on 03/06/2010 at 05:09

Please note that [terms and conditions apply](#).

Clean positive operator-valued measures for qubits and similar cases

Jonas Kahn

Université Paris-Sud 11, Département de Mathématiques, Bât 425, 91405 Orsay Cedex, France

E-mail: jonas.kahn@math.u-psud.fr

Received 4 October 2006, in final form 4 February 2007

Published 17 April 2007

Online at stacks.iop.org/JPhysA/40/4817

Abstract

In a recent paper (Buscemi *et al* 2005 *J. Math. Phys.* **46** 082109) a notion of clean positive operator-valued measures (POVMs) was defined. We characterize those POVMs which are clean in a class that we call quasi-qubit POVMs, namely POVMs whose elements are all rank-1 or full-rank elements. We give an algorithm to check whether a given quasi-qubit POVM satisfies this condition. We describe explicitly all the POVMs that are clean for the qubit. Meanwhile, we give a sufficient condition for a general POVM to be clean.

PACS number: 03.65.–w

Mathematics Subject Classification: 47L05, 81T05

1. Introduction

The laws of quantum mechanics impose restrictions on what measurements can be carried out on a quantum system. All the possible measurements can be described mathematically by ‘positive operator-valued measures’, POVMs for short. Apart from measuring a state, we can also transform it via a quantum channel. Now suppose we have at our disposal a POVM \mathbf{P} and a channel \mathcal{E} . We may first send our state through \mathcal{E} and then feed the transformed state in our measurement apparatus \mathbf{P} . This procedure is a new measurement procedure, and can therefore be encoded by a POVM \mathbf{Q} . Now transforming the state with \mathcal{E} can be seen as a kind of noise on the POVM \mathbf{P} . We may then view \mathbf{Q} as a disturbed version of \mathbf{P} , and we say that \mathbf{P} is *cleaner* than \mathbf{Q} . Now, what are the maximal elements for this order relation?

The order relation ‘ \mathbf{P} is cleaner than \mathbf{Q} ’ has been introduced in a recent article of Buscemi *et al* [1]. Herein the authors try to understand which POVMs can be obtained from another, either by pre-processing (the situation we just described, where we first send our state through a channel) or by classical post-processing of the data. Especially, they try to find which POVMs are biggest for these order relations (in the former case, the POVM is said to be *clean*; there is no ‘extrinsic’ noise). For pre-processing they get a number of partial answers. One of

those is that a POVM on a d -dimensional space with n outcomes, with $n \leq d$, is clean if and only if it is an observable. They do not get a complete classification, though.

The object of the present paper is to characterize those POVMs which are clean in a special class of measurements. Namely, we are interested in POVMs whose elements (see the definition below) have either full rank or rank 1. We call these POVMs *quasi-qubit POVMs*. Note that all the POVMs for qubits satisfy this condition.

Meanwhile we prove a sufficient condition for a POVM to be clean, which is usable also for POVMs that are not quasi-qubit.

It turns out that cleanness for quasi-qubit POVMs can be read on the span of the rank-1 elements. Moreover, if a (non-necessarily quasi-qubit) POVM is cleaner than a clean quasi-qubit POVM, the latter was in fact obtained by a channel that is a unitary transform. In other words, for clean quasi-qubit POVMs, cleanness equivalence is unitary equivalence.

We give an algorithm to check whether a quasi-qubit POVM is clean or not. This algorithm may be the main contribution of the paper, as almost all the following theorems can be summed up by saying the algorithm is valid.

In the end we apply these results to the qubit, for which all POVMs are quasi-qubit. We are then left with a very explicit characterization of clean POVMs for qubits.

Section 2 gives precise definitions of all the objects we cited in the introduction.

We define the algorithm, give heuristically the main ideas and define the important notion ‘totally determined’ (definition 3.2) in section 3.

Section 4 gives a sufficient condition for a POVM to be clean, namely that the supports of the elements of the POVM ‘totally determine’ the space (see definition 3.2). We use this condition to show that when the algorithm exits with a positive result, the quasi-qubit POVM is really clean.

Section 5 proves that the above sufficient condition is in fact necessary for quasi-qubit POVMs. It checks that when the algorithm exits with a negative result, the POVM is truly not clean.

Section 6 gathers the results relative to quasi-qubit POVMs in theorem 6.1 and deals with the qubit case in corollary 6.2.

Finally, section 7 gives a very rough idea for making more explicit the sufficient condition for a POVM to be clean we have given in section 4.

If one wishes to look for the results of this paper without bothering with the technical proofs, the best would be to read the algorithm of section 3 and then to read theorem 6.1 and corollary 6.2. You would also need lemma 5.3 that you could use as a definition of ‘totally determined’ if you are only interested in quasi-qubit POVMs.

If you also want the supplementary results that apply to other POVMs, further read definitions 3.1 and 3.2, and theorem 4.1.

2. Definitions and notations

We consider POVMs on a Hilbert space \mathcal{H} of dimension $d \geq 2$. Dimension 2 is the qubit case. The set $\{|e_i\rangle\}_{1 \leq i \leq d}$ will be an orthonormal basis of \mathcal{H} . If \mathcal{V} is a subspace of \mathcal{H} then \mathcal{V}^\perp is the subspace orthogonal to \mathcal{V} in \mathcal{H} . If we are given vectors $\{v_i\}_{i \in I}$, we denote by $\text{Span}(v_i, i \in I)$ the space they generate. The set of operators on \mathcal{H} is denoted by $\mathcal{B}(\mathcal{H})$.

A POVM \mathbf{P} (with finite outcomes, the case to which we restrict) is a set $\{P_i\}_{i \in I}$ of *non-negative* operators on \mathcal{H} , with I being finite, such that $\sum_{i \in I} P_i = \mathbf{1}$. The P_i are called *POVM elements*. We write $\text{Supp}(P_i)$ for the support of the element P_i . This support is defined by its orthogonal. The set of $|\phi\rangle \in \text{Supp}(P_i)^\perp$ is exactly the set of $|\phi\rangle$ such that $\langle \phi | P_i | \phi \rangle = 0$. The rank of a POVM element is its rank as an operator. In particular, rank-1 elements are of

the form $\lambda_i |\psi_i\rangle\langle\psi_i|$ and full-rank POVMs are invertible. Special cases of POVMs are *rank-1 POVMs*, that is POVMs whose all elements have rank 1, and *full-rank POVMs*, that is POVMs whose all elements have full rank. We are especially interested in a class of POVMs that includes both.

Definition 2.1 (Quasi-qubits POVMs). *A POVM \mathbf{P} is a quasi-qubit POVM if all its elements P_i are either full-rank or rank-1.*

Similarly, we shall speak of strict quasi-qubit POVMs for quasi-qubit POVMs which are neither rank-1 nor full-rank.

A channel \mathcal{E} is a completely positive identity-preserving map on $\mathcal{B}(\mathcal{H})$, the set of bounded operators on \mathcal{H} (in this paper, channels are always intended as going from $\mathcal{B}(\mathcal{H})$ to the same $\mathcal{B}(\mathcal{H})$). As a remark, this implies that the subspace of self-adjoint operators $\mathcal{B}_{sa}(\mathcal{H})$ is stable by \mathcal{E} . We know that we can write it using Kraus decomposition [2], that is we can find a finite number of operators $R_\alpha \in \mathcal{B}(\mathcal{H})$ such that

$$\mathcal{E}(A) = \sum_{\alpha} R_{\alpha}^{*} A R_{\alpha}, \quad \text{with} \quad \sum_{\alpha} R_{\alpha}^{*} R_{\alpha} = \mathbf{1}. \quad (1)$$

Here the star is the adjoint.

We shall write $\mathcal{E} = \{R_{\alpha}\}_{\alpha}$. This decomposition is not unique.

Using the channel \mathcal{E} before the measurement \mathbf{P} is the same as using the POVM $\mathbf{Q} = \mathcal{E}(\mathbf{P})$ defined by its POVM elements $Q_i = \mathcal{E}(P_i)$.

Definition 2.2. *A POVM \mathbf{P} is cleaner than a POVM \mathbf{Q} if and only if there exists a channel \mathcal{E} such that $\mathcal{E}(\mathbf{P}) = \mathbf{Q}$. We shall also write $\mathbf{P} \succ \mathbf{Q}$.*

Definition 2.3 (Clean POVM). *A POVM \mathbf{P} is clean if and only if for any \mathbf{Q} such that $\mathbf{Q} \succ \mathbf{P}$ then $\mathbf{P} \succ \mathbf{Q}$ also holds.*

We shall further say that two POVMs are cleanness equivalent if both $\mathbf{Q} \succ \mathbf{P}$ and $\mathbf{P} \succ \mathbf{Q}$ hold. A special case of this (but not the general case, as proved in [1]) is *unitary equivalence*, when there is a unitary operator U such that for any $i \in I$, we have $U P_i U^{*} = Q_i$.

3. Algorithm and ideas

3.1. Algorithm

We propose the following algorithm to check whether a quasi-qubit POVM \mathbf{P} is clean or not.

- (i) We check whether \mathbf{P} has rank 1. If it is, exit with result ‘ \mathbf{P} is clean’. Otherwise:
- (ii) Write the rank-1 elements $P_i = \lambda_i |\psi_i\rangle\langle\psi_i|$ for $1 \leq i \leq n$. Check whether these $|\psi_i\rangle$ generate \mathcal{H} . If not, exit with result ‘ \mathbf{P} is not clean’. Else:
- (iii) We can find a basis of \mathcal{H} as a subset of those $|\psi_i\rangle$. We assume that this basis consists of $|\psi_i\rangle$ for $1 \leq i \leq d$. We define a variable $C = \{V_j\}_{j \in J}$, consisting in a collection of subspaces whose direct sum is the Hilbert space $\mathcal{H} = \bigoplus_j V_j$. We initialize C with $V_i = \text{Span}(|\psi_i\rangle)$ for $1 \leq i \leq d$.
- (iv) For i from $d + 1$ to n , do:
 - (v) Write $|\psi_i\rangle = \sum_j v_j$ with $v_j \in V_j$. Call $J(i) = \{j | v_j \neq 0\}$.
 - (vi) Update $\{V_j\}$: suppress all V_j for $j \in J(i)$. Add $V_i = \bigoplus_{j \in J(i)} V_j$.
 - (vii) Check whether $C = \{\mathcal{H}\}$. If so, exit with result ‘ \mathbf{P} is clean’. Otherwise:
 - (viii) End of the ‘For’ loop.
 - (ix) Exit with result ‘ \mathbf{P} is not clean’.

Note that the algorithm terminates: every stage is finite and we enter the loop a finite number of times.

3.2. Heuristics: what the algorithm really tests

In the Kraus decomposition (1), each of the terms $R_\alpha^* A R_\alpha$ is non-negative if A is non-negative, so that $\mathcal{E}(A) \geq R_\alpha^* A R_\alpha$ for any α . Hence if $\mathcal{E}(\mathbf{Q}) = \mathbf{P}$, then $R_\alpha^* Q_e R_\alpha$ must have support included in $\text{Supp}(P_e)$ for all α and $e \in E$.

The central idea of the paper is the following: the condition $\text{Supp}(R_\alpha^* Q_e R_\alpha) \subset \text{Supp}(P_e)$ yields $d\text{-dim}(\text{Supp}(P_e))$ homogeneous linear equations on the matrix entries of R_α , where you should remember that $d = \dim(\mathcal{H})$. Now R_α is determined up to a constant by $d^2 - 1$ homogeneous-independent linear equations. In such a case, the additional condition $\sum R_\alpha^* R_\alpha = \mathbf{1}$ yields; all R_α are proportional to the same unitary U , so that the channel \mathcal{E} is unitary, and $\mathbf{P} > \mathbf{Q}$.

There is still one difficulty: the equations mentioned above depend not only on \mathbf{P} , but also on \mathbf{Q} . We would then like conditions on the supports of P_e such that the system of equations mentioned above is at least of rank $d^2 - 1$ for all \mathbf{Q} . We formalize this requirement with the following definitions.

Definition 3.1 (Corresponding). *Let \mathcal{V} be a Hilbert space and $\{F_i\}_{i \in I}$ a collection of subspaces of \mathcal{V} . Let $\{v_i\}_{i \in I}$ be a collection of vectors of \mathcal{V} . This set of vectors corresponds to $\{F_i\}_{i \in I}$ if for any $i \in I$, there is a linear transform R_i such that $R_i(v_i) \neq 0$ and, for all $j \in I$, the transform is taking v_j within F_j , that is $R_i(v_j) \in F_j$.*

In the text, we usually drop the reference to $\{F_i\}_{i \in I}$ and write that the $\{v_i\}_{i \in I}$ are a corresponding collection of vectors.

Definition 3.2 (Totally determined). *Let \mathcal{V} be a Hilbert space and $\{F_i\}_{i \in I}$ a collection of subspaces of \mathcal{V} .*

If for all corresponding collections of vectors $\{v_i\}_{i \in I}$ there is only one (up to a complex multiplicative constant) linear transform R such that $R(v_i) \in F_i$ for all $i \in I$, we say that \mathcal{V} is totally determined by $\{F_i\}_{i \in I}$, or alternatively that $\{F_i\}_{i \in I}$ totally determines \mathcal{V} .

If F_i is one dimensional with support vector w_i , this means there is only one R such that $R(v_i)$ is colinear to w_i for all $i \in I$.

What the algorithm does is checking that a quasi-qubit POVM \mathbf{P} is rank 1 (stage (i)), or that \mathbf{P} totally determines \mathcal{H} .

More precisely, proposition 4.9 states that each of the V_j belonging to C (appearing at stage (iii) and updated at stage (vi)) is totally determined by the $|\psi_i\rangle$ such that $|\psi_i\rangle \in V_j$. When the algorithm exits at stage (vii), then $C = \{\mathcal{H}\}$, so \mathcal{H} is totally determined. If the algorithm does not exit at stage (vii), on the other hand, then C has at least two elements at the last stage, and each $|\psi_i\rangle$ is included in one of those two elements, which entails, from lemma 5.3, that $\{\text{Supp}(P_e)\}$ does not totally determine \mathcal{H} .

The equivalence with cleanness for quasi-qubit POVMs is still needed to get validity of the algorithm. This equivalence stems from theorems 4.1 and 5.1. The former is the sufficient condition, for any POVM, not necessarily quasi-qubit. We have given the intuition for this theorem at the beginning of the section. Complementarily, theorem 5.1 states that a strict quasi-qubit POVM is not clean if its supports do not totally determine \mathcal{H} .

The proof of theorem 5.1 features the last important idea of the paper. A channel \mathcal{E} which is near enough to the identity may be inverted as a positive map on $\mathcal{B}(\mathcal{H})$, even though \mathcal{E}^{-1} is

not a channel. Now if we denote $\mathbf{Q} = \mathcal{E}^{-1}(\mathbf{P})$, we have $\mathcal{E}(\mathbf{Q}) = \mathbf{P}$. We are then left with two questions: is \mathbf{Q} a POVM, and can we find a channel \mathcal{F} such that $\mathcal{F}(\mathbf{P}) = \mathbf{Q}$?

The main possible obstacle to \mathbf{Q} being a POVM is the need for each of the Q_i to be non-negative. Now, if \mathcal{E} is near enough to the identity, if P_i is full rank, then Q_i is still full-rank non-negative. The remaining case is $Q_i = \mathcal{E}^{-1}(P_i) = \lambda_i \mathcal{E}^{-1}(|\psi_i\rangle\langle\psi_i|)$. Now, we shall see that we may use the set of subspaces $C = \{V_j\}$ given by the algorithm to build channels ensuring that these Q_i are still rank-1 non-negative matrices. Furthermore, these Q_i will have a bigger first eigenvalue than P_i , so that we are sure \mathbf{Q} is strictly cleaner than \mathbf{P} , as channels are spectrum-width decreasing (see lemma 5.2).

We now turn to the fully rigorous treatment.

4. Sufficient condition

We start by proving the following theorem, announced in the previous section.

Theorem 4.1. *If the supports $\{\text{Supp}(P_i)\}_{i \in I}$ of the elements P_i of a POVM \mathbf{P} totally determine \mathcal{H} , then \mathbf{P} is clean and any cleanness-equivalent POVM \mathbf{Q} is in fact unitarily equivalent to \mathbf{P} .*

Proof. It is enough to prove that if $\mathbf{Q} > \mathbf{P}$, then \mathbf{Q} is unitarily equivalent to \mathbf{P} . Let \mathbf{Q} be a POVM and $\mathcal{E} = \{R_\alpha\}_\alpha$ a channel such that $\mathcal{E}(\mathbf{Q}) = \mathbf{P}$. For all $i \in I$, we may write $Q_i = \sum_k \mu_{i,k} |\phi_i^k\rangle\langle\phi_i^k|$. Then we have $P_i = \sum_\alpha \sum_k \mu_{i,k} R_\alpha^* |\phi_i^k\rangle\langle\phi_i^k| R_\alpha$. Now $\mu_{i,k} R_\alpha^* |\phi_i^k\rangle\langle\phi_i^k| R_\alpha \geq 0$ for all k and α , and consequently $\mu_{i,k} R_\alpha^* |\phi_i^k\rangle\langle\phi_i^k| R_\alpha \leq P_i$. Hence $R_\alpha^* |\phi_i^k\rangle \in \text{Supp}(P_i)$.

Moreover, P_i is nonzero. So there is at least one $k(i)$ and one $\alpha(i)$ for each i such that $R_{\alpha(i)}^* |\phi_i^{k(i)}\rangle$ is nonzero. Thus $\{|\phi_i^{k(i)}\rangle\}_{i \in I}$ corresponds to $\{\text{Supp}(P_i)\}_{i \in I}$. As $\{\text{Supp}(P_i)\}_{i \in I}$ totally determines \mathcal{H} , there is only one R , up to a constant, such that $R|\phi_i^{k(i)}\rangle \in \text{Supp}(P_i)$ for all i . So $R_\alpha = c(\alpha)R$ for all α . Since $\sum_\alpha R_\alpha^* R_\alpha = \mathbf{1}$, there is a constant such that λR_1 is unitary, and $\mathcal{E} = \{\lambda R_1\}$. So \mathbf{P} and \mathbf{Q} are unitarily equivalent. \square

Before proving in theorem 4.9 that ‘when the algorithm exits at stage (vii), then the supports of the POVM \mathbf{P} totally determine \mathcal{H} ’, we need a few more tools.

We first need the notion of a *projective frame*. Indeed, in the algorithm, we are dealing with supports of rank-1 POVMs, that is essentially projective lines. And we want them to totally determine the space, that is essentially fix it. Projective frames are the most basic mathematical object meeting these requirements. We redefine them here, and reprove what basic properties we need; further information on projective frames may be found in most geometry or algebra textbooks, e.g. [3].

Definition 4.2. *A projective frame $\{v_i\}_{1 \leq i \leq d+1}$ of a vector space \mathcal{V} is a set of $(\dim(\mathcal{V}) + 1)$ vectors in general position, that is, such that any subset of $\dim(\mathcal{V})$ vectors is a basis of \mathcal{V} .*

Remark 4.3. Equivalently we may say that $\{v_i\}_{1 \leq i \leq n}$ is a basis of \mathcal{V} and $v_{d+1} = \sum_{i=1}^n c_i v_i$ with all $c_i \neq 0$.

Proposition 4.4. *A projective frame $\Psi = \{e_i\}_{1 \leq i \leq (n+1)}$ of \mathcal{V} totally determines \mathcal{V} .*

Proof. First we prove that if $\Phi = \{v_i\}_{1 \leq i \leq (n+1)}$ is not a projective frame, the set of vectors $\{v_i\}_{1 \leq i \leq (n+1)}$ does not correspond to Ψ . Indeed, as Φ is not a projective frame, we may find n vectors, say the n first, such that $\sum_{i=1}^n a_i v_i = 0$ with at least one a_i nonzero, say a_1 . Then for any R such that $R(v_i)$ is colinear to e_i for all i , we still have $\sum_{i=1}^n a_i R(v_i) = 0$. As $\{e_i\}_{1 \leq i \leq n}$ is a basis, $a_i R(v_i) = 0$ for all i , so that $R(v_1) = 0$. Hence $\{v_i\}_{1 \leq i \leq n+1}$ does not correspond to $\{e_i\}_{1 \leq i \leq n+1}$.

Let now $\Phi = \{v_i\}_{1 \leq i \leq (n+1)}$ be corresponding to Ψ . Notably, this implies that Φ is a projective frame. Furthermore, there is a nonzero linear transform R such that $R(v_i)$ is colinear to e_i for all i . We must show that R is unique up to a constant.

We know that $\{e_i\}_{1 \leq i \leq n}$ and $\{v_i\}_{1 \leq i \leq n}$ are both bases of \mathcal{V} . Hence there is a unique transfer matrix X from the latter basis to the former. Since $R(v_i) = D_i e_i$ for some D_i , we know that R is of the form DX where D is a diagonal matrix with diagonal values D_i .

We have not yet used our $(n+1)$ th condition. We are dealing with projective frames, so that $e_{n+1} = \sum_{i=1}^n b_i e_i$ and $v_{n+1} = \sum_{i=1}^n c_i v_i$ with all b_i and c_i being nonzero. Now $R(v_{n+1}) = \sum_{i=1}^n c_i R(v_i) = \sum_{i=1}^n c_i D_i e_i$, so that $c_i D_i / b_i$ must be independent of i and D , and hence R is fixed up to a complex multiplicative constant. \square

We now turn to a few observations about totally determined spaces.

Remark 4.5. If $\{F_i\}_{i \in I}$ totally determines \mathcal{H} , and if $\{v_i\}_{i \in I}$ corresponds to $\{F_i\}$, then up to a constant unique nonzero R such that $Rv_i \in F_i$ for all $i \in I$ is invertible.

Proof. Let us define $\Pi_{(\ker R)^\perp}$ the projector on the orthogonal of the kernel of R along its kernel, and $\Pi_{\ker R}$ the projector on the kernel of R along $(\ker R)^\perp$. We have $R = R\Pi_{(\ker R)^\perp}$, so that $R\Pi_{(\ker R)^\perp} v_i = Rv_i$. Thus $\{\Pi_{(\ker R)^\perp} v_i\}_{i \in I}$ is corresponding to $\{F_i\}_{i \in I}$. On the other hand, $\Pi_{\ker R} \Pi_{(\ker R)^\perp} = 0$, so that $(R + \Pi_{\ker R})(\Pi_{(\ker R)^\perp} v_i) = R(\Pi_{(\ker R)^\perp} v_i) \in F_i$. As $\{\Pi_{(\ker R)^\perp} v_i\}$ is corresponding to $\{F_i\}$, the latter equality implies that R is proportional to $(R + \Pi_{\ker R})$. This is only possible if $\Pi_{\ker R} = 0$. Hence R is invertible. \square

Remark 4.6. If $\{v_l\}_{l \in I \cup J}$ is corresponding to $\{F_l\}_{l \in I \cup J}$, then $\{v_i\}_{i \in I}$ (resp. $\{v_j\}_{j \in J}$) is corresponding to $\{F_i\}_{i \in I}$ (resp. $\{F_j\}_{j \in J}$).

Proof. The set I is a subset of $I \cup J$; thus, for all $i \in I$, there is an R_i such that $R_i v_i \neq 0$ and $R_i v_l \in F_l$ for all $l \in I \cup J$. *A fortiori* $R_i v_k \in F_k$ for all $k \in I$. Hence $\{v_i\}_{i \in I}$ is corresponding to $\{F_i\}_{i \in I}$. The same proof yields the result for J . \square

Remark 4.7. If $\{v_i\}_{i \in I}$ is corresponding to $\{F_i\}_{i \in I}$, then there exists R such that $Rv_i \in F_i$ and $Rv_i \neq 0$ for all i simultaneously.

Proof. By the definition of ‘corresponding to’, we have a set $\{R_i\}_{i \in I}$ of transforms such that $R_i v_i \neq 0$ and $R_i v_j \in F_j$ for all $j \in I$. Now, for any set of coefficients $\{a_i\}_{i \in I}$ the matrix $R = \sum_i a_i R_i$ fulfils $Rv_i \in F_i$ for all i . If we choose appropriately $\{a_i\}$ we also have $Rv_i \neq 0$. For example, we may write all the $R_i v_i$ in the same basis, take note of all coordinates, and choose the a_i as any real numbers algebraically independent of those coordinates. \square

Lemma 4.8. If \mathcal{V} and \mathcal{W} are both totally determined by sets of subspaces $\{F_i\}_{i \in I}$ and $\{F_j\}_{j \in J}$ and if \mathcal{V} and \mathcal{W} intersect (apart from the null vector), then their sum $\mathcal{U} = \mathcal{V} + \mathcal{W}$ is totally determined by $\{F_l\}_{l \in I \cup J}$.

Proof. Let $\{u_l\}_{l \in I \cup J}$ vectors of \mathcal{U} correspond to $\{F_l\}_{l \in I \cup J}$. In other words, there is an R^* such that $R^* u_l \in F_l$ for all $l \in I \cup J$. By remark 4.7, we may assume that $R^* u_l \neq 0$ for all l . We must show that R^* is unique up to a constant. Note that the restriction $R^* u_l \neq 0$ does not play a role: if we find another R non-proportional to R^* , such that $Ru_l \in F_l$ for all l , then $R^* + aR$ for appropriate a also fulfils $0 \neq (R^* + aR)u_l \in F_l$ for all l , and is not proportional to R^* .

We need a few notations. First, we consider the space $\mathcal{X} = \mathcal{V} \cap \mathcal{W}$. We also define \mathcal{Y} by $\mathcal{V} = \mathcal{Y} \oplus \mathcal{X}$ and \mathcal{Z} by $\mathcal{W} = \mathcal{Z} \oplus \mathcal{X}$. We write $I_{\mathcal{V}}$ and $I_{\mathcal{W}}$ for the natural inclusions of \mathcal{V} and \mathcal{W} in \mathcal{U} . We also denote by $\Pi_{\mathcal{V}}$ the projector on \mathcal{V} along \mathcal{Z} , by $\Pi_{\mathcal{W}}$ the projector on \mathcal{W} along \mathcal{Y} and by $\Pi_{\mathcal{X}}$ the projector on \mathcal{X} along $\mathcal{Y} + \mathcal{Z}$.

Be aware that we do not define $\Pi_{\mathcal{V}}$ and $\Pi_{\mathcal{W}}$ as endomorphisms of \mathcal{U} , but as applications from \mathcal{U} to \mathcal{V} and \mathcal{W} , respectively. The corresponding endomorphisms are $I_{\mathcal{V}}\Pi_{\mathcal{V}}$ and $I_{\mathcal{W}}\Pi_{\mathcal{W}}$.

As a first step, we show that $I_{\mathcal{V}}\Pi_{\mathcal{V}}R^*$ is unique up to a constant.

The rank of $I_{\mathcal{V}}\Pi_{\mathcal{V}}R^*$ is at most $\dim(\mathcal{V})$, so we can factorize it by \mathcal{V} : there exist two linear applications $L_{\mathcal{V}}^{\mathcal{U}}$ from \mathcal{U} to \mathcal{V} and $L_{\mathcal{U}}^{\mathcal{V}}$ from \mathcal{V} to \mathcal{U} , such that $I_{\mathcal{V}}\Pi_{\mathcal{V}}R^*L_{\mathcal{U}}^{\mathcal{V}}L_{\mathcal{V}}^{\mathcal{U}} = I_{\mathcal{V}}\Pi_{\mathcal{V}}R^*$.

Now for all $i \in I$, we have $R^*u_i \in F_i \subset \mathcal{V}$, so that $R^*u_i = I_{\mathcal{V}}\Pi_{\mathcal{V}}R^*u_i = I_{\mathcal{V}}\Pi_{\mathcal{V}}R^*L_{\mathcal{U}}^{\mathcal{V}}L_{\mathcal{V}}^{\mathcal{U}}u_i$, so that for all $i \in I$ we have the inclusion $0 \neq (\Pi_{\mathcal{V}}R^*L_{\mathcal{U}}^{\mathcal{V}})(L_{\mathcal{V}}^{\mathcal{U}}u_i) \in F_i$, where we have used $R^*u_i \neq 0$. Thus $\{L_{\mathcal{V}}^{\mathcal{U}}u_i\}_{i \in I}$ is corresponding to $\{F_i\}_{i \in I}$. On the other hand, we know that $\{F_i\}_{i \in I}$ totally determine \mathcal{V} . Hence there is a nonzero constant $\lambda_{\mathcal{V}}$, and a $R_{\mathcal{V}}$ depending only on $\{F_i\}_{i \in I}$, such that $\Pi_{\mathcal{V}}R^*L_{\mathcal{U}}^{\mathcal{V}} = \lambda_{\mathcal{V}}R_{\mathcal{V}}$. Moreover, by remark 4.5, $R_{\mathcal{V}}$ is invertible. So finally $I_{\mathcal{V}}\Pi_{\mathcal{V}}R^* = \lambda_{\mathcal{V}}I_{\mathcal{V}}R_{\mathcal{V}}L_{\mathcal{U}}^{\mathcal{V}}$, with image $\text{im}(\lambda_{\mathcal{V}}I_{\mathcal{V}}R_{\mathcal{V}}L_{\mathcal{U}}^{\mathcal{V}}) = \mathcal{V}$. Replacing \mathcal{V} with \mathcal{W} , we similarly get $I_{\mathcal{W}}\Pi_{\mathcal{W}}R^* = \lambda_{\mathcal{W}}I_{\mathcal{W}}R_{\mathcal{W}}L_{\mathcal{U}}^{\mathcal{W}}$.

The last step consists in proving that the two constants $\lambda_{\mathcal{V}}$ and $\lambda_{\mathcal{W}}$ are proportional, independently of R^* .

We note that $\Pi_{\mathcal{X}}I_{\mathcal{V}}\Pi_{\mathcal{V}} = \Pi_{\mathcal{X}} = \Pi_{\mathcal{X}}I_{\mathcal{W}}\Pi_{\mathcal{W}}$. Hence $\lambda_{\mathcal{V}}\Pi_{\mathcal{X}}I_{\mathcal{V}}R_{\mathcal{V}}L_{\mathcal{U}}^{\mathcal{V}} = \lambda_{\mathcal{W}}\Pi_{\mathcal{X}}I_{\mathcal{W}}R_{\mathcal{W}}L_{\mathcal{U}}^{\mathcal{W}}$. As $\mathcal{X} \subset \mathcal{V}$ and $\text{im}(\lambda_{\mathcal{V}}I_{\mathcal{V}}R_{\mathcal{V}}L_{\mathcal{U}}^{\mathcal{V}}) = \mathcal{V}$, we know that $\lambda_{\mathcal{V}}\Pi_{\mathcal{X}}I_{\mathcal{V}}R_{\mathcal{V}}L_{\mathcal{U}}^{\mathcal{V}} \neq 0$. The equality $\lambda_{\mathcal{V}}\Pi_{\mathcal{X}}I_{\mathcal{V}}R_{\mathcal{V}}L_{\mathcal{U}}^{\mathcal{V}} = \lambda_{\mathcal{W}}\Pi_{\mathcal{X}}I_{\mathcal{W}}R_{\mathcal{W}}L_{\mathcal{U}}^{\mathcal{W}}$ then yields the proportionality of $\lambda_{\mathcal{W}}$ and $\lambda_{\mathcal{V}}$.

We conclude by recalling that $\mathcal{V} + \mathcal{W} = \mathcal{U}$, so that knowing both $I_{\mathcal{V}}\Pi_{\mathcal{V}}R^*$ and $I_{\mathcal{W}}\Pi_{\mathcal{W}}R^*$ is equivalent to knowing R^* . As our only free parameter is the multiplicative constant $\lambda_{\mathcal{V}}$, we have proved uniqueness of R^* , up to a constant. \square

Lemma 4.8 and proposition 4.4 are the two ingredients for proving the following proposition, central for the validity of the algorithm.

Proposition 4.9. *In the algorithm, the spaces in the set $C = \{V_j\}_{j \in J}$ are always totally determined by the supports $K(j) = \{\text{Span}(|\psi_i\rangle) : |\psi_i\rangle \in V_j\}$ of the one-dimensional POVM elements they contain.*

Proof. We prove the proposition by induction on the stronger property Prop = ‘all V_j is totally determined by $K(j)$, and they are spanned by vectors of the initial basis, that is, they are of the form $\text{Span}(|\psi_i\rangle : i \in I(j))$, where $I(j)$ is a subset of $\{1, \dots, d\}$ ’.

Initialization. We initialize C at step (iii). At this stage V_j is defined for $j \in \{1, \dots, d\}$ by $V_j = \text{Span}(|\psi_j\rangle)$. So on the one hand V_j is of the form $\text{Span}(|\psi_i\rangle : i \in I(j))$, where $I(j)$ is a subset of $\{1, \dots, d\}$, and on the other hand V_j is totally determined by $K(j)$, as it is one dimensional and $|\psi_j\rangle$ is nonzero.

Update. We update C at stage (vi). We must prove that $V_i = \bigoplus_{j \in J(i)} V_j$ still fulfils the induction property.

For one thing, the space V_i is a sum of spaces of the form $\text{Span}(|\psi_i\rangle : i \in I(j))$, where $I(j)$ is a subset of $\{1, \dots, d\}$; hence V_i is also of this form with $I(i) = \bigcup_{j \in J(i)} I(j)$.

Now let us consider the set $I_{\text{int}} = \{j : j \in \{1, \dots, d\}, \langle \psi_i | \psi_j \rangle \neq 0\}$, and the space $V_{\text{int}} = \text{Span}(|\psi_j\rangle : j \in I_{\text{int}})$. Since the $|\psi_j\rangle$ for $j \in I_{\text{int}}$ are part of the initial basis $\{|\psi_j\rangle\}_{1 \leq j \leq d}$, they are independent. The definition of I_{int} also ensures $|\psi_i\rangle = \sum_{j \in I_{\text{int}}} c_j |\psi_j\rangle$ with j being nonzero; hence, by remark (9), the set $\{|\psi_k\rangle : k \in I_{\text{int}} \cup \{i\}\}$ is a projective frame of V_{int} . So, by proposition 4.4, the space V_{int} is totally determined by $\{|\psi_j\rangle\}_{j \in I_{\text{int}} \cup \{i\}}$. We initialize $K_{\text{int}} = I_{\text{int}} \cup \{i\}$.

Finally, by the definition of $J(i)$, we know that $V_{\text{int}} \cap V_j \neq 0$ for all $j \in J(i)$. Both are totally determined by $K(j)$ and K_{int} . Hence by lemma 4.8, $V_{\text{int}} \cup V_j$ is totally determined by

$K(j) \cup K_{\text{int}}$. We update $V_{\text{int}} = V_{\text{int}} \cup V_j$ and $K_{\text{int}} = K_{\text{int}} \cup K(j)$. We iterate the latter step for all $j \in J(i)$ and we end up with $V_{\text{int}} = V_i$, totally determined by $\bigcup_{j \in J(i)} K(j) \cup I_{\text{int}} \cup \{i\} \subset I(i)$. \square

Corollary 4.10. *When the algorithm ends at stage (vii), the POVM \mathbf{P} is clean.*

Proof. The algorithm ends at stage (vii) only if $C = \{\mathcal{H}\}$. By the above proposition, this condition implies that \mathcal{H} is totally determined by $\{\text{Span}(|\psi_j\rangle) : |\psi_j\rangle \in \mathcal{H}\}$. This amounts at saying that \mathcal{H} is totally determined by the supports of the POVM elements P_i , and we conclude by theorem 4.1. \square

This section aims at giving sufficient conditions for a POVM to be clean, and at proving that one of these conditions is fulfilled if the algorithm exits with result ‘ \mathbf{P} is clean’. We thus conclude the section with the case when the algorithm exits at stage (i). In other words, we must show that a rank-1 POVM is clean. Now, this has already been proved as theorem 11.2 of [1].

Theorem 4.11. *If \mathbf{P} is rank-1, then $\mathbf{Q} \succ \mathbf{P}$ if and only if \mathbf{P} and \mathbf{Q} are unitarily equivalent. Thus, rank-1 POVMs are clean [1].*

For a quasi-qubit POVM \mathbf{P} , we prove in the following section that \mathbf{P} is clean only if it fulfils the conditions either of theorem 4.11 or of theorem 4.1.

5. Necessary condition for quasi-qubit POVMs

This section proves that a clean quasi-qubit POVM either is rank-1, or the supports of its elements totally determine the space.

Theorem 5.1. *A non-rank-1 quasi-qubit POVM where $\{\text{Supp}(P_i)_{i \in I}\}$ does not determine \mathcal{H} is not clean.*

We need a few more tools to prove the theorem.

To begin with, we need a way to prove in specific situations that a POVM is not cleaner than another. Using the fact that channels are *spectrum-width decreasing* is the easiest method. This is lemma 3.1 of [1]:

Lemma 5.2. *If the minimal (resp. maximal) eigenvalue of X is denoted as $\lambda_m(X)$ (resp. $\lambda_M(X)$), then $\lambda_m(X) \leq \lambda_m(\mathcal{E}(X)) \leq \lambda_M(\mathcal{E}(X)) \leq \lambda_M(X)$ for any channel \mathcal{E} .*

This lemma implies that existence of $\mathbf{Q} \succ \mathbf{P}$ such that for some $i \in I$, either $\lambda_m(Q_i) < \lambda_m(P_i)$ or $\lambda_M(Q_i) > \lambda_M(P_i)$, entails that \mathbf{Q} is strictly cleaner than \mathbf{P} , so that \mathbf{P} is not clean.

We now give a characterization of the fact that \mathcal{H} is totally determined by $\{F_j\}_{j \in J}$ when all the F_j are one dimensional, that is when the F_j can be seen as vectors. This characterization applies to $\{\text{Supp}(P_i)\}_{i \in I}$ for quasi-qubit POVMs, and may be more intuitive than definition 3.2. Moreover, it is more adapted to our strategy of proof.

Lemma 5.3. *A set of vectors $\{|\psi_j\rangle\}_{j \in J}$ totally determines the space \mathcal{H} if and only if, for any two supplementary proper subspaces \mathcal{V} and \mathcal{W} , there is a $j \in J$ such that $|\psi_j\rangle \notin \mathcal{V}$ and $|\psi_j\rangle \notin \mathcal{W}$.*

Moreover, when the algorithm exits with result ‘ \mathbf{P} is not clean’, the supports of \mathbf{P} do not totally determine \mathcal{H} .

Proof. The proof consists of four steps.

- (a) For any finite set of vectors $\{|\psi_j\rangle\}_{j \in J}$, there is a POVM whose supports of the rank-1 elements are these vectors.
- (b) If we feed into the algorithm a non-rank-1 quasi-qubit POVM whose supports of rank-1 elements are the $|\psi_j\rangle$ and if $\{|\psi_j\rangle\}$ does not totally determine \mathcal{H} , then the algorithm exits with result ‘**P** is not clean’.
- (c) If the algorithm exits with result ‘**P** is not clean’, then we can find two supplementary proper subspaces such that $|\psi_j\rangle \in \mathcal{V}$ or $|\psi_j\rangle \in \mathcal{W}$ for all supports of rank-1 elements.
- (d) Finding two supplementary proper subspaces such that $|\psi_j\rangle \in \mathcal{V}$ or $|\psi_j\rangle \in \mathcal{W}$ for all $j \in J$ implies that $\{|\psi_j\rangle\}_{j \in J}$ does not totally determine \mathcal{H} .

The equivalence in the lemma is then proved by contraposition, and the last statement by combining (c) and (d).

Step (a). A valid example is given by $P_j = \frac{1}{2\#J} |\psi_j\rangle\langle\psi_j|$ for $j \in J$ and $P_{\#J+1} = \mathbf{1} - \sum_j P_j$. Indeed the latter element is positive since $\sum_j P_j \leq \frac{1}{2\#J} \#J \mathbf{1} = \frac{1}{2} \mathbf{1}$.

Step (b). Since the quasi-qubit POVM is assumed not to be of rank 1, we do not exit at stage (i). The only other possible exit with result ‘**P** is clean’ is at stage (vii). Now the proof of corollary 4.10 states that the algorithm exits at stage (vii) only if the supports of the rank-1 elements totally determine \mathcal{H} . Hence, the algorithm exits with result ‘**P** is not clean’.

Step (c). Exiting at stage (ii) means that the $|\psi_j\rangle$ do not generate \mathcal{H} . Then, if $J = \emptyset$, we may choose any two supplementary proper subspaces \mathcal{V} and \mathcal{W} . Anyhow $|\psi_j\rangle \in \mathcal{V}$ for all $j \in J$. If $J \neq \emptyset$, then $\mathcal{V} = \text{Span}(|\psi_i\rangle, i \in I)$ is a proper subspace of \mathcal{H} . Since $|\psi_j\rangle \in \mathcal{V}$ for all $j \in J$, any supplementary subspace \mathcal{W} of \mathcal{V} will turn the trick.

If the algorithm does not exit at stage (ii), then there is a basis included in $\{|\psi_j\rangle\}_{j \in J}$. We assume that it corresponds to $1 \leq j \leq d$.

Since the algorithm exits with result ‘**P** is not clean’, it exits at stage (ix). We end the algorithm with a collection $C = \{V_k\}$ of subspaces such that $\bigoplus_k V_k = \mathcal{H}$. Since we have not exited at stage (vii), we know that $C \neq \{\mathcal{H}\}$. Hence C counts at least two non-trivial elements. We take $\mathcal{V} = V_1$ and $\mathcal{W} = \bigoplus_{k \neq 1} V_k$.

The V_k are direct sums of the original $V_j = \text{Span}(|\psi_j\rangle)$ for $1 \leq j \leq d$. Hence, for $1 \leq j \leq d$, either $|\psi_j\rangle \in \mathcal{V}$ or $|\psi_j\rangle \in \mathcal{W}$. On the other hand, if $|\psi_j\rangle$ is not one of the original basis vectors, it was used in the ‘For’ loop. At the end of this loop, C was then containing a space $V = \bigoplus_{k \in J(j)} V_k$. And $|\psi_j\rangle$ was included in this space. This V is then included in one of the final V_j and *a fortiori* either in \mathcal{V} or in \mathcal{W} . We have thus proved that when the algorithm exits with a negative value we may find two supplementary proper subspaces \mathcal{V} and \mathcal{W} such that for all $i \in I$, either $|\psi_i\rangle \in \mathcal{V}$ or $|\psi_i\rangle \in \mathcal{W}$.

Step (d). Since $\mathbf{1}|\psi_j\rangle = |\psi_j\rangle$ for all j , by definition 3.1 the set of vectors $\{|\psi_j\rangle\}_{j \in J}$ is corresponding to the subspaces $\{|\psi_j\rangle\}_{j \in J}$. On the other hand, denoting by $\Pi_{\mathcal{V}}$ the projection on \mathcal{V} parallel to \mathcal{W} , we get that $\Pi_{\mathcal{V}}|\psi_j\rangle$ is colinear to $|\psi_j\rangle$ for all $j \in J$. Moreover, $\Pi_{\mathcal{V}}$ is not proportional to $\mathbf{1}$; so, by definition 3.2, the set of vectors $\{|\psi_j\rangle\}$ does not totally determine \mathcal{H} . \square

Finally, as explained in section 3, we want to build our cleaner POVMs as $\mathcal{E}^{-1}(\mathbf{P})$ where the channel is inverted as a positive map. We need to know some conditions under which a channel can be inverted. This is the purpose of lemma 5.4, for which we need the following norms.

The Hilbert–Schmidt norm on $\mathcal{B}(\mathcal{H})$ is defined as $\|M\|_{\text{HS}}^2 = \text{Tr}(MM^*)$. Notably, in any orthogonal basis,

$$\|M\|_{\text{HS}}^2 = \sum_{1 \leq i, j \leq d} |M_{i,j}|^2.$$

Moreover $\|M\|_{\text{HS}} = \|M^*\|_{\text{HS}}$.

We also define a norm on $\mathcal{B}(\mathcal{B}(\mathcal{H}))$, space to which the channels belong:

$$\|\mathcal{O}\|_1 = \sup_{\{M \mid \|M\|_{\text{HS}}=1\}} \|\mathcal{O}(M)\|_{\text{HS}}.$$

Lemma 5.4. *If in the Kraus representation of a channel $\mathcal{E} = \{R_\alpha\}$ one of the R_α fulfils*

$$\|\mathbf{1} - R_\alpha\|_{\text{HS}} \leq \epsilon,$$

then

$$\|\mathbf{1} - \mathcal{E}\|_1 \leq 2(1 + \sqrt{d})\epsilon + 2\epsilon^2 = f(\epsilon) \xrightarrow{\epsilon \rightarrow 0} 0. \tag{2}$$

As a consequence, if $f(\epsilon) < 1$, then \mathcal{E} is invertible (as a map on $\mathcal{B}(\mathcal{H})$) and $\|\mathcal{E}^{-1} - \mathbf{1}\|_1 \leq f(\epsilon)/(1 - f(\epsilon))$. This inverse lets $\mathcal{B}_{sa}(\mathcal{H})$ stable.

This in turn shows that for any $X \in \mathcal{B}_{sa}(\mathcal{H})$ such that $\lambda_m(X) \geq 0$, the spectrum of the image by the inverse is bounded through

$$\lambda_m(X) - \lambda_M(X) f(\epsilon) \sqrt{d} / (1 - f(\epsilon)) \leq \lambda_m(\mathcal{E}^{-1}(X)). \tag{3}$$

So for all $X > 0$, when ϵ is small enough, $\mathcal{E}^{-1}(X) \geq 0$.

Remark. The bound (2) is probably far from sharp, but sufficient for our needs.

Proof. Without loss of generality, we assume that

$$\|\mathbf{1} - R_1\|_{\text{HS}} \leq \epsilon.$$

We write $S = R_1 - \mathbf{1}_{\mathcal{H}}$ and $\mathcal{O} = \mathcal{E} - \mathbf{1}_{\mathcal{B}(\mathcal{H})}$.

Then

$$\mathcal{O} : M \mapsto S^*MS + S^*M + MS + \sum_{\alpha \neq 1} R_\alpha^*MR_\alpha.$$

And

$$\begin{aligned} \|\mathcal{O}\|_1 &= \sup_{\{M \mid \|M\|_{\text{HS}}=1\}} \left\| S^*MS + S^*M + MS + \sum_{\alpha \neq 1} R_\alpha^*MR_\alpha \right\|_{\text{HS}} \\ &\leq \sup_{\{M \mid \|M\|_{\text{HS}}=1\}} (\|S^*\| \|M\| \|S\| + \|S^*\| \|M\| + \|M\| \|S\| + \sum_{\alpha \neq 1} \|R_\alpha^*\| \|M\| \|R_\alpha\|) \\ &= \|S\|_{\text{HS}}^2 + 2\|S\|_{\text{HS}} + \sum_{\alpha \neq 1} \|R_\alpha\|_{\text{HS}}^2. \end{aligned}$$

Now, for one thing, by hypothesis, $\|S\|_{\text{HS}} \leq \epsilon$. Furthermore

$$\sum_{\alpha \neq 1} \|R_\alpha\|_{\text{HS}}^2 = \sum_{\alpha \neq 1} \text{Tr}(R_\alpha^*R_\alpha) = \text{Tr}(\mathbf{1} - R_1^*R_1) = -\text{Tr}(S^*S + S + S^*).$$

We finish our proof of (2) with the observation that $-\text{Tr}(S + S^*) \leq 2\sqrt{d}\|S\|_{\text{HS}} = 2\sqrt{d}\epsilon$.

If $\|\mathcal{O}\|_1 < 1$, we know that $\mathcal{E} = \mathbf{1} + \mathcal{O}$ is invertible and $\mathcal{E}^{-1} = \sum_{n \geq 0} (-\mathcal{O})^n$. By taking the norm, $\|\mathcal{E}^{-1} - \mathbf{1}\|_1 \leq \sum_{n \geq 1} \|\mathcal{O}\|_1^n = f(\epsilon)/(1 - f(\epsilon))$.

Channels stabilize $\mathcal{B}_{sa}(\mathcal{H})$; as \mathcal{E} is furthermore invertible, equality of dimension shows that $\mathcal{E}(\mathcal{B}_{sa}(\mathcal{H})) = \mathcal{B}_{sa}(\mathcal{H})$ and $\mathcal{E}^{-1}(\mathcal{B}_{sa}(\mathcal{H})) = \mathcal{B}_{sa}(\mathcal{H})$.

Now, X is positive, so that $\|X\|_{HS} \leq \sqrt{d}\lambda_M(X)$. This implies $\|(\mathcal{E}^{-1} - \mathbf{1})(X)\|_{HS} \leq \sqrt{d}\lambda_M(X)f(\epsilon)/(1 - f(\epsilon))$, and in turn $\mathcal{E}^{-1}(X) \geq X - \sqrt{d}\lambda_M(X)f(\epsilon)/(1 - f(\epsilon))\mathbf{1}$. Taking the bottom of the spectrum ends the proof. \square

We are now ready to prove theorem 5.1.

Proof of theorem 5.1. We aim at exhibiting a channel \mathcal{E} and a POVM \mathbf{Q} such that $\mathcal{E}(\mathbf{Q}) = \mathbf{P}$ and Q_i has a wider spectrum than P_i for some $i \in I$. Then lemma 5.2 proves that \mathbf{Q} is strictly cleaner than \mathbf{P} , and in turn that \mathbf{P} is not clean.

The building blocks are the subspaces supplied by lemma 5.3. Since \mathcal{H} is not determined by $\{\text{Supp}(P_i)\}_{i \in I}$, there are two supplementary proper subspaces \mathcal{V} and \mathcal{W} such that each rank-1 element has support included either in \mathcal{V} or in \mathcal{W} .

We shall write explicitly several matrices in the forthcoming proof. All of them shall be written on an orthonormal basis $\{e_j\}_{1 \leq j \leq d}$ of \mathcal{H} , chosen so that $\{e_j\}_{1 \leq j \leq \dim(\mathcal{V})}$ is a basis of \mathcal{V} . We shall express the matrices as two-by-two block matrices, the blocks corresponding to the subspaces \mathcal{V} and \mathcal{V}^\perp .

We study separately the following cases.

- (a) All POVM elements P_i are proportional to the identity, that is $P_i = \mu_i \mathbf{1}$.
- (b) The POVM is not full-rank, each rank-1 element has support either in \mathcal{V} or in \mathcal{V}^\perp , and all POVM elements are block-diagonal in \mathcal{V} and \mathcal{V}^\perp .
- (c) Each rank-1 element has support either in \mathcal{V} or \mathcal{V}^\perp , and at least one POVM element is not block-diagonal.
- (d) At least one rank-1 element has support neither in \mathcal{V} nor in \mathcal{V}^\perp .

As a sanity check, let us prove we did not forget any case. Either our POVM is full-rank, or it is not. In the latter situation, either there is a rank-1 element whose support is not included in \mathcal{V} nor in \mathcal{V}^\perp —and we are in case (d)—or all rank-1 elements are included in \mathcal{V} or \mathcal{V}^\perp . Then either there is a POVM element that is not block-diagonal—and we are in case (c)—or all POVM elements are block-diagonal—and we are in case (b). On the other hand, if \mathbf{P} is full-rank, we may choose the subspaces \mathcal{V} and \mathcal{W} any way we like. Notably, if one POVM element P_i is not proportional to the identity, so that it has non-trivial eigenspaces, we may choose \mathcal{V} such that P_i is not block-diagonal in \mathcal{V} and \mathcal{V}^\perp —and we are in case (c). Finally, if in contrast, all POVM elements are proportional to the identity, we are in case (a).

Case (a). If all POVM elements are of the form $P_i = \mu_i \mathbf{1}$, then, for any $\mathcal{E} = \{R_\alpha\}$, we have $\mathcal{E}(P_i) = \sum_\alpha R_\alpha^* (\mu_i \mathbf{1}) R_\alpha = \mu_i \sum_\alpha R_\alpha^* R_\alpha = \mu_i \mathbf{1} = P_i$. No channel can change the wholly uninformative measurement \mathbf{P} .

On the other hand, many POVMs can be degraded to \mathbf{P} . Consider for example the POVM given by $Q_1 = \mu_1 |e_1\rangle\langle e_1| + \sum_{j=2}^d |e_j\rangle\langle e_j|$ and $Q_i = \mu_i |e_1\rangle\langle e_1|$ for $i > 1$. Then $\mathbf{Q} \neq \mathbf{P}$, so that $\mathbf{P} \not\prec \mathbf{Q}$. Yet, with $R_\alpha = |e_1\rangle\langle e_\alpha|$ for $1 \leq \alpha \leq d$, we have $\mathcal{E}(\mathbf{Q}) = \mathbf{P}$, and $\mathbf{Q} \succ \mathbf{P}$. Hence \mathbf{P} is not clean.

Case (b). Since all rank-1 elements are included either in \mathcal{V} or in \mathcal{V}^\perp , we take $\mathcal{W} = \mathcal{V}^\perp$. We further choose \mathcal{V} to be the smaller of the two subspaces, that is $\dim(\mathcal{V}) \leq d/2 \leq \dim(\mathcal{W})$. Then there is a matrix $A : \mathcal{V} \rightarrow \mathcal{W}$ such that $AA^* = \mathbf{1}_\mathcal{V}$. If all rank-1 elements have support in \mathcal{W} , we further impose that at least one of these supports is not included in the kernel of A .

We then define $R_\mathcal{V}^*$ and $R_\mathcal{W}^*$ as

$$R_\mathcal{V}^*(\epsilon) = \left[\begin{array}{c|c} \mathbf{1}_\mathcal{V} & \epsilon A \\ \hline 0 & 0 \end{array} \right], \quad R_\mathcal{W}^*(\epsilon) = \left[\begin{array}{c|c} 0 & 0 \\ \hline 0 & \mathbf{1}_\mathcal{W} \end{array} \right].$$

Their images are respectively \mathcal{V} and \mathcal{W} .

From $R_{\mathcal{V}}(\epsilon)$ and $R_{\mathcal{W}}(\epsilon)$, we define the channel $\mathcal{E}_\epsilon = \{R_1(\epsilon), R_2(\epsilon), R_3(\epsilon)\}$:

$$\begin{aligned} R_1^*(\epsilon) &= \sqrt{\frac{\epsilon^2}{1+\epsilon^2}} R_{\mathcal{V}}^*(\epsilon) + \sqrt{\frac{1-\epsilon^2}{1+\epsilon^2}} R_{\mathcal{W}}^*(\epsilon) = \left[\begin{array}{c|c} \sqrt{\frac{\epsilon^2}{1+\epsilon^2}} \mathbf{1}_{\mathcal{V}} & \sqrt{\frac{\epsilon^4}{1+\epsilon^2}} A \\ \hline 0 & \sqrt{\frac{1-\epsilon^2}{1+\epsilon^2}} \mathbf{1}_{\mathcal{W}} \end{array} \right], \\ R_2^*(\epsilon) &= \sqrt{\frac{\epsilon^2}{1+\epsilon^2}} R_{\mathcal{W}}^*(\epsilon) = \left[\begin{array}{c|c} 0 & 0 \\ \hline 0 & \sqrt{\frac{\epsilon^2}{1+\epsilon^2}} \mathbf{1}_{\mathcal{W}} \end{array} \right], \\ R_3^*(\epsilon) &= \sqrt{\frac{1-\epsilon^2}{1+\epsilon^2}} R_{\mathcal{V}}^*(\epsilon) - \sqrt{\frac{\epsilon^2}{1+\epsilon^2}} R_{\mathcal{W}}^*(\epsilon) = \left[\begin{array}{c|c} \sqrt{\frac{1-\epsilon^2}{1+\epsilon^2}} \mathbf{1}_{\mathcal{V}} & \sqrt{\frac{\epsilon^2-\epsilon^4}{1+\epsilon^2}} A \\ \hline 0 & -\sqrt{\frac{\epsilon^2}{1+\epsilon^2}} \mathbf{1}_{\mathcal{W}} \end{array} \right]. \end{aligned}$$

Since $AA^* = \mathbf{1}_{\mathcal{V}}$, we have $\sum_{\alpha} R_{\alpha}^* R_{\alpha} = \mathbf{1}$; hence these matrices $\{R_{\alpha}\}$ define a genuine channel. A few calculations show that the effect of this channel is

$$\mathcal{E}_\epsilon : \left[\begin{array}{c|c} B & C \\ \hline C^* & D \end{array} \right] \rightarrow \left[\begin{array}{c|c} \frac{1}{1+\epsilon^2} (B + \epsilon(AC^* + CA^*) + \epsilon^2 ADA^*) & 0 \\ \hline 0 & D \end{array} \right]. \quad (4)$$

Now, for any $w \in \mathcal{W}$, we have

$$\left[\begin{array}{c|c} -\epsilon Aw & \\ \hline w & \end{array} \right] \left[\begin{array}{c|c} -\epsilon Aw & \\ \hline w & \end{array} \right]^* = \left[\begin{array}{c|c} \epsilon^2 Aw w^* A^* & -\epsilon Aw w^* \\ \hline -\epsilon w w^* A^* & w w^* \end{array} \right],$$

so that for any sequence of $w_j \in \mathcal{W}$, the matrix $\sum_{j,k} \left[\begin{array}{c|c} \epsilon^2 A w_j w_k^* A^* & -\epsilon A w_j w_k^* \\ \hline -\epsilon w_j w_k^* A^* & w_j w_k^* \end{array} \right]$ is non-negative. As any non-negative endomorphism D of \mathcal{W} can be written as $\sum_{j,k} w_j w_k^*$ for appropriate w_j , we get that for any non-negative D , the matrix $\left[\begin{array}{c|c} \epsilon^2 ADA^* & -\epsilon AD \\ \hline -\epsilon DA^* & D \end{array} \right]$ is non-negative. Moreover, applying equation (4) yields that its image by \mathcal{E}_ϵ is $\left[\begin{array}{c|c} 0 & 0 \\ \hline 0 & D \end{array} \right]$.

Similarly, if $B \in \mathcal{B}(\mathcal{V})$ is non-negative, then $\left[\begin{array}{c|c} (1+\epsilon^2)B & 0 \\ \hline 0 & 0 \end{array} \right]$ is non-negative and its image by \mathcal{E}_ϵ is $\left[\begin{array}{c|c} B & 0 \\ \hline 0 & 0 \end{array} \right]$.

We use these observations to define a map (not a channel) \mathcal{F}_ϵ on the block-diagonal matrices

$$\mathcal{F}_\epsilon : \left[\begin{array}{c|c} B & 0 \\ \hline 0 & D \end{array} \right] \rightarrow \left[\begin{array}{c|c} (1+\epsilon^2)B + \epsilon^2 ADA^* & -\epsilon AD \\ \hline -\epsilon DA^* & D \end{array} \right]. \quad (5)$$

We get that $\mathcal{E}_\epsilon(\mathcal{F}_\epsilon(M)) = M$ for all block-diagonal M and that if furthermore $M \geq 0$ then $\mathcal{F}_\epsilon(M) \geq 0$.

We now isolate one full-rank element of \mathbf{P} , say P_1 . For all $i \neq 1$, we define $Q_i(\epsilon) = \mathcal{F}_\epsilon(P_i)$. They are non-negative and fulfil $\mathcal{E}_\epsilon(Q_i(\epsilon)) = P_i$. Define now $Q_1(\epsilon) = \mathbf{1} - \sum_{i \neq 1} Q_i(\epsilon)$. The closure relation ensures that $\mathcal{E}_\epsilon(Q_1(\epsilon)) = P_1$. What's more, recalling that $\sum_i B_i = \mathbf{1}_{\mathcal{V}}$ and $\sum_i D_i = \mathbf{1}_{\mathcal{W}}$, we obtain

$$\begin{aligned} Q_1(\epsilon) &= \left[\begin{array}{c|c} \mathbf{1}_{\mathcal{V}} - (1+\epsilon^2)\sum_{i \neq 1} B_i - \epsilon^2 A(\sum_{i \neq 1} D_i)A^* & \epsilon A \sum_{i \neq 1} D_i \\ \hline -\epsilon \sum_{i \neq 1} D_i A^* & \mathbf{1}_{\mathcal{W}} - \sum_{i \neq 1} D_i \end{array} \right] \\ &= \left[\begin{array}{c|c} (1+\epsilon^2)B_1 + \epsilon^2 A D_1 A^* - 2\epsilon^2 \mathbf{1}_{\mathcal{V}} & \epsilon A(\mathbf{1}_{\mathcal{W}} - D_1) \\ \hline \epsilon(\mathbf{1}_{\mathcal{W}} - D_1)A^* & D_1 \end{array} \right] \xrightarrow{\epsilon \rightarrow 0} \left[\begin{array}{c|c} B_1 & 0 \\ \hline 0 & D_1 \end{array} \right] \\ &= P_1. \end{aligned}$$

Since P_1 is positive, this convergence entails the non-negativity of $Q_1(\epsilon)$ for ϵ small enough. As $Q_1(\epsilon)$ has been chosen so that $\sum_i Q_i(\epsilon) = \mathbf{1}$, we have defined a genuine POVM $\mathbf{Q}(\epsilon) = \{Q_i(\epsilon)\}_{i \in I}$ such that $\mathcal{E}_\epsilon(\mathbf{Q}(\epsilon)) = \mathbf{P}$, hence $\mathbf{Q} \succ \mathbf{P}$.

We end the study of this case by considering a rank-1 element $P_i = \mu_i |\psi_i\rangle\langle\psi_i|$ whose support is not in the kernel of A . Using formula (5), if $|\psi_i\rangle \in \mathcal{V}$, we get $\text{Tr}(Q_i(\epsilon)) = (1 + \epsilon^2) \text{Tr}(P_i) > \text{Tr}(P_i)$, else $|\psi_i\rangle \in \mathcal{W}$ and we get $\text{Tr}(Q_i(\epsilon)) = \text{Tr}(P_i) + \epsilon^2 \text{Tr}(A|\psi_i\rangle\langle\psi_i|A^*) > \text{Tr}(P_i)$. In both cases, bigger trace implies that the spectrum of $Q_i(\epsilon)$ is wider than that of P_i and lemma 5.2 yields $\mathbf{P} \not\prec \mathbf{Q}$. So \mathbf{P} is not clean.

Case (c). Since all rank-1 elements are included either in \mathcal{V} or in \mathcal{V}^\perp , we take $\mathcal{W} = \mathcal{V}^\perp$. We now define the channel \mathcal{E}_ϵ through

$$R_1(\epsilon) = \epsilon \Pi_{\mathcal{V}}, \quad R_2(\epsilon) = \epsilon \Pi_{\mathcal{W}} = \epsilon \Pi_{\mathcal{V}^\perp}, \quad R_3(\epsilon) = \sqrt{1 - \epsilon^2} \mathbf{1},$$

where Π denotes here orthogonal projection.

For ϵ small enough, by lemma 2, the channel is invertible as a positive map. We then define $Q_i = \mathcal{E}_\epsilon^{-1}(P_i)$.

Through the formula $\mathcal{E}_\epsilon(Q_i) = P_i$, we check that

$$P_i = \left[\begin{array}{c|c} B & C \\ \hline C^* & D \end{array} \right], \quad \text{then} \quad Q_i(\epsilon) = \left[\begin{array}{c|c} B & (1 - \epsilon^2)^{-1} C \\ \hline (1 - \epsilon^2)^{-1} C^* & D \end{array} \right]. \quad (6)$$

The first remark is that the closure relation ensures $\sum Q_i(\epsilon) = \mathbf{1}$.

We also note that, since rank-1 elements have support either in \mathcal{V} or in $\mathcal{W} = \mathcal{V}^\perp$, the rank-1 elements are block-diagonal and $Q_i(\epsilon) = P_i$.

We know that at least one POVM element is not block-diagonal. So there is an $i \in I$ such that P_i is full-rank and C is nonzero (say $[C]_{j,k} \neq 0$). Then, writing $n = \dim(\mathcal{V})$, there is an $\epsilon_+ \in (0, 1)$ such that

$$\begin{aligned} [Q_i(\epsilon_+)]_{j,j} [Q_i(\epsilon_+)]_{n+k,n+k} &= [B]_{j,j} [D]_{k,k} < \frac{1}{1 - \epsilon_+^2} |[C]_{j,k}|^2 \\ &= [Q_i(\epsilon_+)]_{j,n+k} [Q_i(\epsilon_+)]_{n+k,j} \end{aligned}$$

so that we cannot have positivity of $Q_i(\epsilon_+)$.

We define the bottom of the spectrum of the images Q_i of the full-rank elements of \mathbf{P} :

$$\lambda_m(\epsilon) = \inf_{i|P_i \text{ full-rank}} \lambda_m(Q_i(\epsilon)).$$

Equation (6) implies that the matrix $Q_i(\epsilon)$ is a continuous function of ϵ for $\epsilon \in [0, 1)$. Hence its spectrum is also a continuous function of ϵ . Accordingly, the function $\lambda_m(\epsilon)$ is the minimum of a finite number of continuous function of ϵ ; therefore $\lambda_m(\epsilon)$ is continuous. Its value in 0 is the bottom of the spectrum of the full-rank elements of \mathbf{P} , that is $\lambda_m(0) = \inf_{i|P_i \text{ full-rank}} \lambda_m(P_i) > 0$. Moreover, we have just proved that $\lambda_m(\epsilon_+) < 0$. Thus, by the intermediate value theorem, there is an $\epsilon_+ > \epsilon > 0$ such that $0 < \lambda_m(\epsilon) < \lambda_m(0)$.

As $\lambda_m(\epsilon) > 0$, the $Q_i(\epsilon) = \mathcal{E}_\epsilon(P_i)$ for full-rank P_i are non-negative, and valid POVM elements. Likewise, we already know that $Q_i(\epsilon) = P_i$ is a valid POVM element if P_i is rank 1. Since we have also shown that $\sum Q_i(\epsilon) = \mathbf{1}$, we have proved that $\mathbf{Q}(\epsilon)$ is a POVM. Furthermore, $\mathcal{E}_\epsilon(\mathbf{Q}(\epsilon)) = \mathbf{P}$; thus $\mathbf{Q}(\epsilon) \succ \mathbf{P}$.

As $\lambda_m(\epsilon) < \lambda_m(0)$, there is a full-rank element P_i such that $\lambda_m(Q_i(\epsilon)) < \lambda_m(P_i)$. Hence, using lemma 5.2, we get $\mathbf{P} \not\prec \mathbf{Q}(\epsilon)$ and \mathbf{P} is not clean.

Hence $\lambda_m(\epsilon_+) \leq 0 < \lambda_m$. By the intermediate value theorem, we can find an $\epsilon_0 \in (0, \epsilon_+)$ such that $\lambda_m(\epsilon_0) = 0$. As $0 \leq \lambda_m(\epsilon_0) < \lambda_m$ we have proved that $\mathbf{Q}(\epsilon_0) \succ \mathbf{P}$ and that \mathbf{P} is not clean.

Case (d). As \mathcal{V} and \mathcal{W} are supplementary we may choose a matrix $A \in M_{\dim(\mathcal{V}), d - \dim(\mathcal{V})}(\mathbb{C})$ such that the nonzero columns of the following block matrix form an orthogonal (though not

orthonormal) basis of \mathcal{W} :

$$R_{\mathcal{W}}^* = \begin{bmatrix} 0 & A \\ 0 & \mathbf{1} \end{bmatrix}.$$

We know that the image of a matrix is spanned by its columns, so the image of $R_{\mathcal{W}}^*$ is \mathcal{W} .

We then define

$$B(\epsilon) = \sqrt{\mathbf{1} - \left(\frac{\epsilon^4}{1 - \epsilon^2} + \frac{\epsilon^2}{(1 - \epsilon^2)^2} \right) AA^*}. \tag{7}$$

This definition is valid if the matrix under the square root is positive. Now $\left(\frac{\epsilon^4}{1 - \epsilon^2} + \frac{\epsilon^2}{(1 - \epsilon^2)^2} \right)$ is going to 0 with ϵ , so that

$$\lim_{\epsilon \rightarrow 0} \mathbf{1} - \left(\frac{\epsilon^4}{1 - \epsilon^2} + \frac{\epsilon^2}{(1 - \epsilon^2)^2} \right) AA^* = \mathbf{1}.$$

From this we conclude that $\mathbf{1} - \left(\frac{\epsilon^4}{1 - \epsilon^2} + \frac{\epsilon^2}{(1 - \epsilon^2)^2} \right) AA^*$ is positive for ϵ small enough.

Accordingly, we can define

$$R_{\mathcal{V}}^*(\epsilon) = \begin{bmatrix} B(\epsilon) & -\frac{A}{1 - \epsilon^2} \\ 0 & 0 \end{bmatrix}.$$

Note that the image of $R_{\mathcal{V}}^*$ is included in \mathcal{V} .

We may now define our channel \mathcal{E}_ϵ by

$$R_1^*(\epsilon) = \epsilon R_{\mathcal{V}}^*(\epsilon) = \begin{bmatrix} \epsilon B(\epsilon) & -\frac{\epsilon A}{1 - \epsilon^2} \\ 0 & 0 \end{bmatrix} \tag{8}$$

$$R_2^*(\epsilon) = \epsilon R_{\mathcal{W}}^* = \begin{bmatrix} 0 & \epsilon A \\ 0 & \epsilon \mathbf{1} \end{bmatrix} \tag{9}$$

$$R_3^*(\epsilon) = \sqrt{1 - \epsilon^2} (R_{\mathcal{V}}^*(\epsilon) + R_{\mathcal{W}}^*) = \begin{bmatrix} \sqrt{1 - \epsilon^2} B(\epsilon) & -\frac{\sqrt{1 - \epsilon^2} A}{\sqrt{1 - \epsilon^2}} \\ 0 & \sqrt{1 - \epsilon^2} \mathbf{1} \end{bmatrix}. \tag{10}$$

Note that $\sum_{\alpha=1}^3 R_{\alpha}^*(\epsilon) R_{\alpha}(\epsilon) = \mathbf{1}$ so that $\mathcal{E}(\epsilon)$ is indeed a channel.

Moreover $\lim_{\epsilon \rightarrow 0} R_3(\epsilon) = \mathbf{1}_{\mathcal{H}}$. Hence, for ϵ small enough, $\|R_3 - \mathbf{1}\|_{\text{HS}}$ is as small as we want. So lemma 5.4 allows us to invert the channel \mathcal{E}_ϵ as a map on $\mathcal{B}_{sa}(\mathcal{H})$. We define $\mathbf{Q}(\epsilon)$ by its elements $Q_i(\epsilon) = \mathcal{E}_\epsilon^{-1}(P_i)$. Let us check that for ϵ small enough, $\mathbf{Q}(\epsilon)$ is still a *bona fide* POVM.

First the closure relation still holds, as $\sum_{i \in I} Q_i = \sum_{i \in I} \mathcal{E}^{-1}(P_i) = \mathcal{E}^{-1}(\mathbf{1})$. Now $\mathcal{E}(\mathbf{1}) = \sum_{\alpha} R_{\alpha}^* R_{\alpha} = \mathbf{1}$ and taking the inverse $\mathcal{E}^{-1}(\mathbf{1}) = \mathbf{1}$.

It then remains to be shown that all $Q_i(\epsilon)$ are non-negative.

If P_i is full-rank, then its spectrum is included in $[\lambda_m, 1]$, with $\lambda_m > 0$. If R_3 is near enough to the identity, that is, if ϵ is small enough, inequality (3) then ensures that $Q_i(\epsilon)$ is still positive.

If P_i is rank-1 $P_i = \lambda_i |\psi_i\rangle\langle\psi_i|$, then by hypothesis $|\psi_i\rangle \in \mathcal{V}$ or $|\psi_i\rangle \in \mathcal{W}$. As R_3 is invertible for ϵ small enough, we may consider $|\phi_i\rangle$ nonzero colinear to $(R_3^*(\epsilon))^{-1} |\psi_i\rangle$. Then $R_3^*(\epsilon) |\phi_i\rangle$ is colinear to $|\psi_i\rangle$, and nonzero. Note that $|\phi_i\rangle$ depends on ϵ , even if we drop it in the notation. Now

$$R_3(\epsilon)^* |\varphi\rangle = \sqrt{1 - \epsilon^2} (R_{\mathcal{V}}^*(\epsilon) |\varphi\rangle + R_{\mathcal{W}}^* |\varphi\rangle) \quad \text{with} \quad R_{\mathcal{V}}^*(\epsilon) |\varphi\rangle \in \mathcal{V} \quad \text{and} \quad R_{\mathcal{W}}^* |\varphi\rangle \in \mathcal{W}.$$

Since \mathcal{V} and \mathcal{W} are supplementary, the latter equality implies that $R_{\mathcal{V}}^*(\epsilon) |\varphi\rangle = 0$ when $R_3^*(\epsilon) |\varphi\rangle \in \mathcal{W}$ and $R_{\mathcal{W}}^*(\epsilon) |\varphi\rangle = 0$ when $R_3^*(\epsilon) |\varphi\rangle \in \mathcal{V}$. Definitions (8), (9), (10) then yield

$\mathcal{E}_\epsilon(|\phi_i\rangle\langle\phi_i|) = R_{\mathcal{W}}^*(|\phi_i\rangle\langle\phi_i|)R_{\mathcal{W}}$ if $|\psi_i\rangle \in \mathcal{W}$ and $\mathcal{E}_\epsilon(|\phi_i\rangle\langle\phi_i|) = R_{\mathcal{V}}^*(\epsilon)(|\phi_i\rangle\langle\phi_i|)R_{\mathcal{V}}(\epsilon)$ if $|\psi_i\rangle \in \mathcal{V}$. In both cases, the output matrix is of the form $\mathcal{E}_\epsilon(|\phi_i\rangle\langle\phi_i|) = C_i|\psi_i\rangle\langle\psi_i|$. So $Q_i(\epsilon) = (\lambda_i/C_i)|\phi_i\rangle\langle\phi_i|$ and is non-negative.

Thus, for ϵ small enough, all $Q_i(\epsilon)$ are non-negative. We have proved that $\mathbf{Q}(\epsilon)$ is a POVM. Furthermore, since $\mathcal{E}_\epsilon(\mathbf{Q}(\epsilon)) = \mathbf{P}$, we know $\mathbf{Q}(\epsilon) \succ \mathbf{P}$.

We must still show that $\mathbf{Q}(\epsilon)$ is strictly cleaner \mathbf{P} .

By hypothesis, there is a rank-1 element $P_i = \lambda_i|\psi_i\rangle\langle\psi_i|$ such that $|\psi_i\rangle \in \mathcal{W}$ and $|\psi_i\rangle \notin \mathcal{V}^\perp$. As above, we write $|\phi_i\rangle$ such that $Q_i(\epsilon) = (\lambda_i/C_i)|\phi_i\rangle\langle\phi_i|$. We start by proving that C_i is less than 1.

We write $|\phi_i\rangle = v_i + v_i^\perp$ with $v_i \in \mathcal{V}$ and $v_i^\perp \in \mathcal{V}^\perp$. Since $|\psi_i\rangle \in \mathcal{W}$, we get

$$\mathcal{E}_\epsilon(|\phi_i\rangle\langle\phi_i|) = R_{\mathcal{W}}^*(|\phi_i\rangle\langle\phi_i|)R_{\mathcal{W}} = \begin{bmatrix} Av_i^\perp \\ v_i^\perp \end{bmatrix} \begin{bmatrix} Av_i^\perp \\ v_i^\perp \end{bmatrix}^*.$$

As the latter expression is also equal to $C_i|\psi_i\rangle\langle\psi_i|$, we obtain that C_i is the square of the norm of $\begin{bmatrix} Av_i^\perp \\ v_i^\perp \end{bmatrix}$. Therefore $C_i = \|Av_i^\perp\|^2 + \|v_i^\perp\|^2$. Note that the squared norm of $|\phi_i\rangle$ is $1 = \|v_i\|^2 + \|v_i^\perp\|^2$. On the other hand, the image of $|\phi_i\rangle$ by $R_{\mathcal{V}}^*(\epsilon)$ is 0, so that $B(\epsilon)v_i - 1/(1 - \epsilon^2)Av_i^\perp = 0$. From this we get

$$Av_i^\perp = (1 - \epsilon^2)B(\epsilon)v_i.$$

Since $|\psi_i\rangle \notin \mathcal{V}^\perp$, this equality shows that $v_i \neq 0$. Now, as AA^* is non-negative we see by (7) that $B(\epsilon) \leq \mathbf{1}$. *A fortiori*, for any $\epsilon > 0$, we have $(1 - \epsilon^2)B(\epsilon) < \mathbf{1}$. So that

$$\|v_i\| > \|(1 - \epsilon^2)B(\epsilon)v_i\| = \|Av_i^\perp\|.$$

Thus, we finally obtain

$$C_i = \|Av_i^\perp\|^2 + \|v_i^\perp\|^2 < \|v_i\|^2 + \|v_i^\perp\|^2 = 1.$$

Hence the biggest eigenvalue of $Q_i(\epsilon) = (\lambda_i/C_i)|\phi_i\rangle\langle\phi_i|$, that is λ_i/C_i , is strictly bigger than the biggest eigenvalue of P_i , that is λ_i . Lemma 5.2 then gives $\mathbf{P} \not\prec \mathbf{Q}(\epsilon)$, and consequently \mathbf{P} is not clean. \square

6. Summary for quasi-qubit POVMs and a special case

We now gather all our results specific to quasi-qubit POVMs.

Theorem 6.1. *A quasi-qubit POVM \mathbf{P} is clean if and only if it is rank 1 or the supports of its rank-1 elements totally determine \mathcal{H} . The algorithm of section 3 figures out if this is the case. Moreover, if \mathbf{Q} is cleanness equivalent to \mathbf{P} , the two POVMs are even unitarily equivalent.*

Proof. Rank-1 POVMs are known to be clean (theorem 4.11). If the support of the rank-1 elements of \mathbf{P} totally determines \mathcal{H} , we also know that \mathbf{P} is clean by theorem 4.1. In both cases the theorems state that for these clean POVMs, cleanness equivalence is the same as unitary equivalence.

Conversely, if \mathbf{P} is neither rank 1 nor have rank-1 elements that totally determine \mathcal{H} , then theorem 5.1 applies and \mathbf{P} is not clean.

Stage (i) of the algorithm checks whether \mathbf{P} is rank 1, in which case it does say that \mathbf{P} is clean. If \mathbf{P} is not rank 1, the fact that whether it is clean or not depends on the support of its rank-1 elements. The only remaining positive exit of the algorithm is at stage (vii) and lemma 4.9 proves that in this case the rank-1 elements of \mathbf{P} totally determine \mathcal{H} .

Conversely, if the algorithm exits with a negative value, lemma 5.3 ensures that \mathcal{H} is not totally determined. \square

To get further feeling of these conditions we finish by making more explicit the qubit case, where the nice thing is that all POVMs are quasi-qubit.

Corollary 6.2. *A POVM \mathbf{P} for a qubit is clean if and only if it is rank 1 or if one can find three rank-1 elements whose supports are two-by-two non-colinear (that is if they make a projective frame). For these POVMs cleanness equivalence is the same as unitary equivalence.*

Proof. A POVM \mathbf{P} for a qubit has nonzero elements which can be either of rank 1, or of rank 2, as $d = 2$. In the latter case, they are full-rank, so we may apply theorem 6.1 to \mathbf{P} .

The only question is: when do the supports of the rank-1 elements totally determine \mathcal{H} ? They do by proposition 4.4 if they include a projective frame, that is a basis and a vector with all coefficients nonzero in this basis. As the space is of dimension 2, this amounts to saying a basis and a vector non-colinear to any basis vector, that is three vectors two-by-two non-colinear.

Conversely, if we cannot find a projective frame, then we can find two vectors v and w such that the support of any rank-1 element is v or w , and we can apply lemma 5.3 to obtain that \mathcal{H} is not totally determined by the supports of the rank-1 elements of \mathbf{P} . Thus \mathbf{P} is not clean. \square

7. Outlook

We have solved the problem of cleanness for quasi-qubit POVMs. The obvious continuation would be to solve it in the general case. However we do not think that the condition of theorem 4.1 is then necessary. Moreover it must be made explicit.

The heuristics in section 3.2 suggest that if the support of P_i are in ‘general position’, then it is sufficient for \mathbf{P} to be clean that $\sum_{e \in E} d - \dim[\text{Supp}(P_i)] \geq d^2 - 1$. Yet, we still need to appropriately define the ‘general position’ for general subspaces.

Acknowledgments

We thank Professor d’Ariano for introducing us to the notion of clean POVMs. We are also indebted to Sylvain Arlot for many general suggestions on writing. My failure at applying them is the source of all the remaining lack of clarity.

References

- [1] Buscemi F, d’Ariano G M, Keyl M, Perinotti P and Werner R F 2005 Clean positive operator valued measures *J. Math. Phys.* **46** 082109
- [2] Kraus K 1983 *States, Effects and Operations* (Berlin: Springer)
- [3] Audin M 2002 *Geometry* (Berlin: Springer)