

Fisher information, delocalization and the semiclassical description of molecular rotation

This article has been downloaded from IOPscience. Please scroll down to see the full text article.

2007 J. Phys. A: Math. Theor. 40 5127

(<http://iopscience.iop.org/1751-8121/40/19/012>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 171.66.16.109

The article was downloaded on 03/06/2010 at 05:10

Please note that [terms and conditions apply](#).

Fisher information, delocalization and the semiclassical description of molecular rotation

S Curilef¹, F Pennini^{1,2}, A Plastino² and G L Ferri³

¹ Departamento de Física, Universidad Católica del Norte, Av. Angamos 0610, Antofagasta, Chile

² Instituto de Física La Plata (CONICET–UNLP), Fac. de Ciencias Exactas, Universidad Nacional de La Plata, CC 67, 1900 La Plata, Argentina

³ Facultad de Ciencias Exactas, Universidad Nacional de La Pampa, Peru y Uruguay, Santa Rosa, La Pampa, Argentina

Received 3 January 2007, in final form 22 March 2007

Published 24 April 2007

Online at stacks.iop.org/JPhysA/40/5127

Abstract

We discuss phase-space delocalization for the rigid rotor within a semiclassical context by recourse to the Husimi distributions of both the linear and the 3D-anisotropic instances. The Husimi function can be viewed in a variety of ways. In particular, we obtain it as the expectation value of the density operator in a suitable basis of coherent states. Our treatment is based upon the concomitant Fisher information measures. The pertinent Wehrl entropy is also investigated in the linear case.

PACS numbers: 03.67.–a, 05.30.–d, 31.15.Gy

1. Introduction

Using Fisher's information measure as our main tool, we will be concerned here with the semiclassical description of the rotational dynamics of molecular systems. A pioneer analysis of such rotational dynamics is that of Morales *et al* in [1], who studied the connection between it and coherent states [2, 3].

The coherent states (CS) formulation is not unique and several authors have developed alternative descriptions for molecular quantum systems. We need CS in this work because one uses them so as to construct Husimi semiclassical distributions (see sections 2.2 and 3.1). With the latter, we determine Fisher's measure (see section 3.2):

Coherent states \rightarrow Husimi distribution \rightarrow Fisher's information measure.

We have chosen two traditional formulations for coherent states in order to discuss the semiclassical dynamics of molecular rotational systems. One of them is devoted to two-dimensional case [4] and the other to the three-dimensional one [1].

In [5, 6] the authors discuss, among other things, the advantages of the Husimi distribution for the interpretation of the electronic structure in hydrogen and nitrogen atoms. They suggested that their work may be extended to the molecular instance. Following this suggestion we address here the simplest applicable model, i.e., the rigid rotor. Its usefulness for describing diatomic molecules is well known [7].

Delocalization is an ‘energetically favourable’ process, since it distributes the wavefunction over a volume greater than the size of the sample. Thus, the net energy of the molecule is lowered, which results in resonance stabilization. The celebrated Fisher information \mathcal{F} [9–12] can be related (for example by Garbaczewski [8]) to the inverse of the delocalization measure as follows: if we take a wave package with a standard deviation σ , the Fisher information is given by $\mathcal{F} \geq 1/\sigma^2$, thus, in this sense, we realize that the Fisher information is a quite sensitive indicator of the delocalization of the wave package [9–11].

The rigid rotor is a system of a single particle whose quantum spectrum of energy is exactly known. Therefore, the study of typical thermodynamic properties can be analytically derived [13]. Applications lead to the treatment of important aspects of molecular systems [14] and several applications to materials [15–18].

The paper is organized as follows: in section 2, we introduce basic aspects and properties which must be satisfied by coherent states. In section 3, we explore the linear rigid rotor. We write the probability of finding a quantum state in a coherent state that is used to obtain an explicit expression for the (i) Husimi distribution, (ii) Wehrl entropy and (iii) Fisher Information. These results are of help in section 4, where we discuss a model for the three-dimensional rigid rotor. Finally, in section 5, we draw some concluding remarks.

2. Coherent states: harmonic oscillator

Quantum-mechanical phase-space distributions expressed in terms of the coherent states $|z\rangle$ of the harmonic oscillator [2] have been proved to be useful in different contexts. Particular reference is made to the important work of Anderson and Halliwell [19], who discuss, among other things, the concepts of Husimi distribution and Wehrl entropy. Coherent states are eigenstates of the annihilation operator \hat{a} , appropriate for the problem at hand, i.e.,

$$\hat{a}|z\rangle = z|z\rangle, \quad (1)$$

whose Hamiltonian reads $\hat{H}_o = \hbar\omega[\hat{a}^\dagger\hat{a} + 1/2]$ and the complex eigenvalues of the annihilation operator \hat{a} are given by

$$z = (m\omega/2\hbar)^{1/2}x + i(2\hbar\omega m)^{-1/2}p. \quad (2)$$

2.1. Requirements for a formulation of coherent states

It is well known that coherent states can be constructed in several ways and by recourse to different techniques, its formulation being of a non-unique character. Gazeau and Klauder suggest that a suitable formalism for coherent states should satisfy the following criteria: (i) continuity of labelling, (ii) resolution of unity, (iii) temporal stability and (iv) action identity [20], that we detail below.

- (i) Continuity of labelling refers to the map from the label space \mathcal{L} into Hilbert space. This condition means that the expression $||z'\rangle - |z\rangle| \rightarrow 0$ whenever $z' \rightarrow z$.
- (ii) Resolution of unity: a positive measure $\tau(z)$ on \mathcal{L} exists such that the unity operator admits the representation

$$\int_{\mathcal{L}} |z\rangle\langle z| d\tau(z) = 1, \quad (3)$$

where $|z\rangle\langle z|$ denotes a projector, which takes a state vector into a multiple of the vector $|z\rangle$.

- (iii) Temporal stability: the evolution of any coherent state $|z\rangle$ always remains a coherent state, which leads to a relation of the form

$$|z(t)\rangle = e^{-i\hat{H}t/\hbar}|z\rangle, \quad (4)$$

where $z(0) = z$, for all $z \in \mathcal{L}$ and t .

- (iv) Action identity: this property requires that

$$\langle z|\hat{H}|z\rangle = |z|^2\hbar\omega. \quad (5)$$

At this point, we remark that requirements (iii) and (iv) are satisfied when the Hamiltonian \hat{H} of the system and its spectrum has the form $E_n \sim n\hbar\omega$, where n is the quantum number and ω is the frequency of the oscillator [20]. In addition, there are some shortcomings about these requirements; for instance, Klauder's states cannot be used for degenerate systems. Furthermore, it is questionable that the action identity can lead to the classical action-angle variable interpretation [21].

2.2. Applications to Husimi function and Wehrl entropy

Wehrl's entropy W is a very useful measure of localization in phase space [22]. It is built up using coherent states [2, 19, 22] and constitutes a powerful tool in statistical physics. The pertinent definition reads

$$W = - \int \frac{dx dp}{2\pi\hbar} \mu(x, p) \ln \mu(x, p), \quad (6)$$

where $\mu(x, p) = \langle z|\hat{\rho}|z\rangle$ is a 'semiclassical' phase-space distribution function associated with the density matrix $\hat{\rho}$ of the system [2, 3]. μ is often referred to as the Husimi distribution [23]. The distribution $\mu(x, p)$ is normalized in the fashion

$$\int (dx dp/2\pi\hbar)\mu(x, p) = 1. \quad (7)$$

These last two equations clearly indicate that the Wehrl entropy is simply the 'classical entropy' (6) of a Wigner distribution. Indeed, $\mu(x, p)$ is a Wigner-distribution D_W smeared over an \hbar -sized region of phase space [19]. The smearing renders $\mu(x, p)$ a positive function, even if D_W does not have such a character. The semiclassical Husimi probability distribution refers to a special type of probability: that for simultaneous but approximate location of position and momentum in phase space [19]. The uncertainty principle manifests itself through the inequality $W \geq 1$, which was first conjectured by Wehrl [22] and later proved by Lieb [24].

The usual treatment of equilibrium in statistical mechanics makes use of the Gibbs's canonical distribution, whose associated, 'thermal' density matrix is given by

$$\hat{\rho} = Z^{-1} e^{-\beta\hat{H}}, \quad (8)$$

with $Z = \text{Tr}(e^{-\beta\hat{H}})$ being the partition function, $\beta = 1/k_B T$ the inverse temperature (T) and k_B the Boltzmann constant. In order to conveniently write an expression for W consider an arbitrary Hamiltonian \hat{H} of eigenenergies E_n and eigenstates $|n\rangle$ (n stands for a collection of all the pertinent quantum numbers required to label the states). One can always write [19]

$$\mu(x, p) = \langle z|\hat{\rho}|z\rangle = \frac{1}{Z} \sum_n e^{-\beta E_n} |\langle z|n\rangle|^2. \quad (9)$$

A useful route to W starts then with equation (9) and continues with equation (6).

3. Linear rigid rotor

We start the present study by exploring a simple model, the linear rigid rotor (LRR), based on the excellent discussion concerning the coherent states for angular momenta given in [25]. The LRR Hamiltonian is [7]

$$\hat{H} = \frac{\hat{L}^2}{2I_{xy}}, \quad (10)$$

where $\hat{L}^2 = \hat{L}_x^2 + \hat{L}_y^2$ is the angular momentum operator and I_x and I_y are the associated moments of inertia. We have assumed that $I_{xy} \equiv I_x = I_y$. Calling $|IK\rangle$ the set of H -eigenstates, we recall that they verify the relations

$$\begin{aligned} \hat{L}^2|IK\rangle &= I(I+1)\hbar^2|IK\rangle \\ \hat{L}_z|IK\rangle &= K\hbar|IK\rangle, \end{aligned} \quad (11)$$

with $I = 0, 1, 2, \dots$, for $-I \leq K \leq I$, the eigenstates' energy spectrum being given by

$$\varepsilon_I = \frac{I(I+1)\hbar^2}{2I_{xy}}. \quad (12)$$

Coherent states are constructed in [4, 26] for the lineal rigid rotor, using Schwinger's oscillator model of angular momentum, in the fashion

$$|IK\rangle = \frac{(\hat{a}_+^\dagger)^{I+K} (\hat{a}_-^\dagger)^{I-K}}{\sqrt{(I+K)!(I-K)!}} |0\rangle, \quad (13)$$

with \hat{a}_+ , \hat{a}_- being the pertinent creation and annihilation operators, respectively, and $|0\rangle \equiv |0, 0\rangle$ the vacuum state. The states $|IK\rangle$ are orthogonal and satisfy the closure relation, i.e.,

$$\langle I'K'|IK\rangle = \delta_{I',I} \delta_{K',K}, \quad (14)$$

$$\sum_{I=0}^{\infty} \sum_{K=-I}^I |IK\rangle \langle IK| = \hat{1}. \quad (15)$$

Since we deal with two degrees of freedom the ensuing coherent states are of the tensor product (denoted by \otimes) form that involves $|z_1\rangle$ and $|z_2\rangle$ [25, 27]:

$$|z_1z_2\rangle = |z_1\rangle \otimes |z_2\rangle, \quad (16)$$

where

$$\hat{a}_+|z_1z_2\rangle = z_1|z_1z_2\rangle, \quad (17)$$

$$\hat{a}_-|z_1z_2\rangle = z_2|z_1z_2\rangle. \quad (18)$$

Therefore, the coherent state $|z_1z_2\rangle$ writes [25]

$$|z_1z_2\rangle = e^{-\frac{|z_1|^2}{2}} e^{z_1\hat{a}_+^\dagger} e^{z_2\hat{a}_-^\dagger} |0\rangle, \quad (19)$$

with

$$|z_1\rangle = e^{-\frac{|z_1|^2}{2}} e^{z_1\hat{a}_+^\dagger} |0\rangle, \quad (20)$$

$$|z_2\rangle = e^{-\frac{|z_2|^2}{2}} e^{z_2\hat{a}_-^\dagger} |0\rangle. \quad (21)$$

We have introduced the convenient notation

$$|z|^2 = |z_1|^2 + |z_2|^2. \quad (22)$$

Using equations (13) and (19) we easily calculate $|z_1 z_2\rangle$ and, after a bit of algebra, find

$$|z_1 z_2\rangle = e^{-\frac{|z|^2}{2}} \sum_{n_+, n_-} \frac{z_1^{n_+}}{\sqrt{n_+!}} \frac{z_2^{n_-}}{\sqrt{n_-!}} |IK\rangle \quad (23)$$

where $n_+ = I + K$ and $n_- = I - K$. Therefore, the probability of observing the state $|IK\rangle$ in the coherent state $|z_1 z_2\rangle$ is of the form

$$|\langle IK | z_1 z_2 \rangle|^2 = e^{-|z|^2} \frac{|z_1|^{2n_+}}{n_+!} \frac{|z_2|^{2n_-}}{n_-!}. \quad (24)$$

The present coherent states satisfy resolution of unity:

$$\int \frac{d^2 z_1}{\pi} \frac{d^2 z_2}{\pi} |z_1 z_2\rangle \langle z_1 z_2| = 1. \quad (25)$$

Furthermore, z_1 and z_2 are continuous variables. We are at this point ready to begin presenting our new results.

3.1. Husimi distribution

Following the procedure developed by Anderson *et al* [19], we can readily calculate the pertinent Husimi distribution [23]. For our system this is defined, from equation (9), as

$$\mu(z_1, z_2) = \langle z_1, z_2 | \hat{\rho} | z_1, z_2 \rangle, \quad (26)$$

where the density operator is

$$\hat{\rho} = Z_{2D}^{-1} \exp(-\beta \hat{H}). \quad (27)$$

Here β is the ordinary inverse temperature parameter (this temperature is the ordinary thermodynamic one). The concomitant rotational partition function Z_{2D} is given in [7]

$$Z_{2D} = \sum_{I=0}^{\infty} (2I+1) e^{-I(I+1)\frac{\Theta}{T}}, \quad (28)$$

with $\Theta = \hbar^2 / (2I_x I_y k_B)$. Remark that in the present context speaking of the ‘trace operation’ entails performing the sum $\text{tr} \equiv \sum_{I=0}^{\infty} \sum_{K=-I}^I$. Inserting now the closure relation into equation (26), and using equation (24), we finally get our Husimi distribution in the fashion

$$\mu(z_1, z_2) = e^{-|z|^2} \frac{\sum_{I=0}^{\infty} \frac{|z|^{4I}}{(2I)!} e^{-I(I+1)\frac{\Theta}{T}}}{\sum_{I=0}^{\infty} (2I+1) e^{-I(I+1)\frac{\Theta}{T}}}. \quad (29)$$

It is easy to show that this distribution is normalized to unity

$$\int \frac{d^2 z_1}{\pi} \frac{d^2 z_2}{\pi} \mu(z_1, z_2) = 1, \quad (30)$$

where z_1 and z_2 are given by equations (17), (18) and (22). Note that we must deal with the binomial expression $(|z_1|^2 + |z_2|^2)^{4I}$ firstly and then integrate over the whole complex plane (in two dimensions) in order to verify the normalization condition (30). The differential element of area in the $z_1(z_2)$ plane is $d^2 z_1 = dx dp_x / 2\hbar$ ($d^2 z_2 = dy dp_y / 2\hbar$) [2]. Moreover, we have




Figure 1. Linear rotor: the Husimi function $\mu(z_1, z_2)$ is plotted as a function of $|z|$ for several values of the temperature: from $T/\Theta = 0.1$ to $T/\Theta = 10$. The behaviour of the Husimi function resembles that of a Gaussian distribution. The peak of the distribution increases as the temperature decreases.

the phase-space relationships

$$|z_1|^2 = \frac{1}{4} \left(\frac{x^2}{\sigma_x^2} + \frac{p_x^2}{\sigma_{p_x}^2} \right), \quad (31)$$

$$|z_2|^2 = \frac{1}{4} \left(\frac{y^2}{\sigma_y^2} + \frac{p_y^2}{\sigma_{p_y}^2} \right), \quad (32)$$

where $\sigma_x \equiv \sigma_y = \sqrt{\hbar/2m\omega}$ and $\sigma_{p_x} \equiv \sigma_{p_y} = \sqrt{m\omega\hbar/2}$. In figure 1, we depict the behaviour of the Husimi distribution $\mu(z_1, z_2)$ as a function of $|z|$ at fixed temperature. The profile of the Husimi function is similar to that of a Gaussian distribution.

3.2. Wehrl entropy and Fisher information

The Wehrl entropy is a semiclassical measure of localization [22] (so is Fisher's one [12] as well). Indeed, Wehrl's measure is simply a logarithmic Shannon measure built up with Husimi distributions. For the present bi-dimensional model this entropy reads

$$\mathcal{W} = - \int \frac{d^2 z_1}{\pi} \frac{d^2 z_2}{\pi} \mu(z_1, z_2) \ln \mu(z_1, z_2), \quad (33)$$

where $\mu(z_1, z_2)$ is given by equation (29). The so-called phase-space, shift-invariant Fisher measure [12] is a particular instance of the general Fisher one [10, 11] that can be regarded as an (also semiclassical) counterpart of Wehrl entropy [12]. Extending now to the present

2D case the ideas developed in [12] for the case of the 1D-harmonic oscillator, we define the (phase-space) shift-invariant Fisher measure in the fashion

$$\mathcal{F} = \int \frac{d^2 z_1}{\pi} \frac{d^2 z_2}{\pi} \mu(z_1, z_2) \mathcal{A}, \quad (34)$$

with

$$\begin{aligned} \mathcal{A} = & \sigma_x^2 \left[\frac{\partial \ln \mu(z_1, z_2)}{\partial x} \right]^2 + \sigma_{p_x}^2 \left[\frac{\partial \ln \mu(z_1, z_2)}{\partial p_x} \right]^2 \\ & + \sigma_y^2 \left[\frac{\partial \ln \mu(z_1, z_2)}{\partial y} \right]^2 + \sigma_{p_y}^2 \left[\frac{\partial \ln \mu(z_1, z_2)}{\partial p_y} \right]^2. \end{aligned} \quad (35)$$

It is easy to prove that the quantity \mathcal{A} has the following form, see details in the appendix:

$$\mathcal{A} = \eta(z_1, z_2)^2, \quad (36)$$

where

$$\eta(z_1, z_2) = \frac{\sum_{I=0}^{\infty} \left[\frac{|z|^{4I-1}}{(2I-1)!} - \frac{|z|^{4I+1}}{(2I)!} \right] e^{-I(I+1)\Theta/T}}{\sum_{I=0}^{\infty} \frac{|z|^{4I}}{(2I)!} e^{-I(I+1)\Theta/T}}. \quad (37)$$

Therefore, the corresponding Fisher measure acquires the simpler appearance

$$\mathcal{F} = \int \frac{d^2 z_1}{\pi} \frac{d^2 z_2}{\pi} \mu(z_1, z_2) \eta(z_1, z_2)^2, \quad (38)$$

i.e.,

$$\mathcal{F} \equiv \langle \eta(z_1, z_2)^2 \rangle, \quad (39)$$

where with the notation

$$\langle \mathcal{G} \rangle = \int \frac{d^2 z_1}{\pi} \frac{d^2 z_2}{\pi} \mu(z) \mathcal{G}, \quad (40)$$

we allude to the *semiclassical expectation value* of \mathcal{G} .

In figure 2, we plot both Fisher's information and Wehrl's entropy as a function of T/Θ . They behave in different manner. If the temperature $T \rightarrow 0$, Fisher's information measure (inverse delocalization) takes its maximum value and Wehrl's its minimum. This behaviour is reversed for high temperatures, with the degree of delocalization becoming larger and larger. In addition, it compares the rotor results with those for the harmonic oscillator. It depicts, as a function of the temperature, Wehrl's and Fisher's measures in the case of the two models. The overall trend is the same in the two different physical cases, a rather remarkable result.

3.3. Uncertainty relation

We calculate now

$$\langle |z|^n \rangle = \int \frac{d^2 z_1}{\pi} \frac{d^2 z_2}{\pi} \mu(z) |z|^n, \quad (41)$$

where $n = 1, 2$ and we find explicitly

$$\langle |z|^n \rangle = \frac{\sum_{I=0}^{\infty} \frac{\Gamma(2I+2+n/2)}{(2I)!} e^{-I(I+1)\frac{\Theta}{T}}}{\sum_{I=0}^{\infty} (2I+1) e^{-I(I+1)\frac{\Theta}{T}}}. \quad (42)$$

We see that $\Delta_z = \langle |z|^2 \rangle - \langle |z| \rangle^2$. With this result one can plot the Cramer-Rao product $\mathcal{F} \Delta_z$ versus temperature, depicted in figure 3, which indicates how phase-space localization




Figure 2. Trends for Fisher's (\mathcal{F}) and Wehrl's (\mathcal{W}) measures (inverse delocalization) for the linear rotor in comparison with those for the harmonic oscillator. The horizontal axis is the temperature. If it increases, the Fisher information decreases while Wehrl entropy increases. It must be noted that, in the case of harmonic oscillator, $t = kT/\hbar\omega$, and in the case of rotor, $t = kT(2I_{xy})/\hbar^2$.




Figure 3. Trends for Fisher's (\mathcal{F}) measure (inverse delocalization), $\Delta_z = \langle |z|^2 \rangle - \langle |z| \rangle^2$ and the Cramer-Rao product $\mathcal{F}\Delta_z$ as a function of the temperature for the linear rotor. The localization information is approximately constant over a wide temperature range.

ability changes with temperature. The strange behaviour near $T = 0$ should not cause undue worries because the semiclassical approximation is not valid near $T = 0$. Curiously enough, our localization information remains relatively constant over a wide temperature range. The inescapable conclusion is that the Cramer–Rao localization estimator is not affected by temperature. This independence from T makes the Cramer–Rao estimator a much better one than Wehrl’s entropy that strongly depends on the temperature.

4. Rigid rotor in three dimensions

In the present section we consider a more general problem, the 3D-rigid rotor model, whose Hamiltonian writes [1]

$$\hat{H} = \frac{\hat{L}_x^2}{2I_x} + \frac{\hat{L}_y^2}{2I_y} + \frac{\hat{L}_z^2}{2I_z}, \quad (43)$$

where I_x , I_y and I_z are the associated moments of inertia. A complete set of rotor eigenstates is $\{|IMK\rangle\}$. The following relations apply:

$$\begin{aligned} \hat{L}^2|IMK\rangle &= I(I+1)\hbar^2|IMK\rangle \\ \hat{L}_z|IMK\rangle &= K\hbar|IMK\rangle \\ \hat{J}_z|IMK\rangle &= M\hbar|IMK\rangle, \end{aligned} \quad (44)$$

where $I = 0, \dots, \infty$, $-I \leq K \leq I$ and $-I \leq M \leq I$. The states $|IMK\rangle$ satisfy orthogonality and closure relation [1]:

$$\langle I'M'K'|IMK\rangle = \delta_{I',I}\delta_{M',M}\delta_{K',K}, \quad (45)$$

$$\sum_{I=0}^{\infty} \sum_{M=-I}^I \sum_{K=-I}^I |IMK\rangle\langle IMK| = \hat{1}. \quad (46)$$

If we take $\hat{L}^2 = \hat{L}_x^2 + \hat{L}_y^2 + \hat{L}_z^2$ and assume axial symmetry, i.e., $I_{xy} \equiv I_x = I_y$, we can recast the Hamiltonian as

$$\hat{H} = \frac{1}{2I_{xy}} \left[\hat{L}^2 + \left(\frac{I_{xy}}{I_z} - 1 \right) \hat{L}_z^2 \right], \quad (47)$$

where \hat{L}^2 is the angular momentum operator and \hat{L}_z is its projection on the rotation axis z . The concomitant spectrum of energy becomes

$$\varepsilon_{I,K} = \frac{\hbar^2}{2I_{xy}} \left[I(I+1) + \left(\frac{I_{xy}}{I_z} - 1 \right) K^2 \right], \quad (48)$$

where $I = 0, 1, 2, \dots$ and it represents the eigenvalue of the angular momentum operator \hat{L}^2 , the numbers $m = -I, \dots, -1, 0, 1, \dots, I$ stand for the projections on the intrinsic rotation axis of the rotor. All states exhibit a $(2I+1)$ degeneracy. The parameters $I_x = I_y \equiv I_{xy}$ and I_z are the inertia momenta. Different ‘geometrical’ instances are characterized through the I_{xy}/I_z ratio. For example, the value $I_{xy}/I_z = 1$ corresponds to the spherical rotor. Limiting cases can also be considered. This is, $I_{xy}/I_z = 1/2$ and $I_{xy}/I_z \rightarrow \infty$, that correspond to the extremely oblate and prolate cases, respectively.

4.1. Coherent states for the rigid rotor in three dimensions

In order to obtain the Husimi distribution for this problem we need first of all to have the associated coherent states. Morales *et al* have constructed them in [1] and discussed their mathematical foundations. First, they introduced the auxiliary quantity

$$X_{I,M,K} = \sqrt{I!(I+M)!(I-M)!(I+K)!(I-K)!}, \quad (49)$$

and then write [1]

$$|z_1 z_2 z_3\rangle = e^{-\frac{|u|^2}{2}} \sum_{IMK} \frac{[(2I)!]^2 z_1^{I+M} z_2^I z_3^{I+K}}{X_{I,M,K}} |IMK\rangle, \quad (50)$$

where the following supplementary variable were introduced by Morales *et al* in [1]:

$$|u|^2 = |z_2|^2(1+|z_1|^2)^2(1+|z_3|^2)^2. \quad (51)$$

All coherent states share at least two requirements. Continuity of labelling and resolution of unity. In relation to the last property we add

$$\int d\Gamma |z_1 z_2 z_3\rangle \langle z_1 z_2 z_3| = 1 \quad (52)$$

where $d\Gamma$ is the measure of integration given by [1]

$$d\Gamma = d\tau \{4[(1+|z_1|^2)(1+|z_3|^2)]^4 |z_2|^4 - 8[(1+|z_1|^2)(1+|z_3|^2)]^2 |z_2|^2 + 1\} \quad (53)$$

with

$$d\tau = \frac{d^2 z_1}{\pi} \frac{d^2 z_2}{\pi} \frac{d^2 z_3}{\pi}, \quad (54)$$

and, of course, in this case we have three degrees of freedom. The present formulation satisfies the weaker version of the second requirement, because the measure is defined nonpositive [1].

4.2. Husimi function, Wehrl entropy and Fisher measure

Using now equation (50) we find

$$|\langle IMK | z_1 z_2 z_3 \rangle|^2 = \frac{e^{-|u|^2}}{X_{I,M,K}^2} [(2I)!]^2 |z_1|^{2(I+M)} |z_2|^{2I} |z_3|^{2(I+K)} \quad (55)$$

and determine that, in this case, the rotational partition function reads

$$Z_{3D} = \sum_{I=0}^{\infty} \sum_{K=-I}^I \sum_{M=-I}^I e^{-\beta \varepsilon_{I,K}}, \quad (56)$$

i.e.,

$$Z_{3D} = \sum_{I=0}^{\infty} (2I+1) e^{-I(I+1)\frac{\Theta}{T}} \sum_{K=-I}^I \exp\left(-\left(\frac{I_{xy}}{I_z} - 1\right) K^2 \frac{\Theta}{T}\right). \quad (57)$$

Remark that if we take the ‘extremely prolate’ limiting case $I_{xy}/I_z \rightarrow \infty$ just one term survives in the right sum of the right-hand side in equation (57), that for $K = 0$, while all terms for $K \neq 0$ vanish. In this special instance case Z_{2D} is recovered from Z_{3D} . The pertinent Husimi distribution becomes

$$\mu(z_1, z_2, z_3) = \frac{e^{-|u|^2}}{Z_{3D}} \sum_{I=0}^{\infty} \frac{(2I)!}{I!} |v|^{2I} e^{-I(I+1)\frac{\Theta}{T}} \times g(I), \quad (58)$$

where

$$g(I) = \sum_{K=-I}^I \frac{|z_3|^{2(I+K)}}{(I+K)!(I-K)!} \exp\left(-\left(\frac{I_{xy}}{I_z} - 1\right) K^2 \frac{\Theta}{T}\right), \quad (59)$$

with

$$|v|^2 = (1 + |z_1|^2)^2 |z_2|^2, \quad (60)$$

$$|u|^2 = |v|^2 (1 + |z_3|^2)^2. \quad (61)$$

We can easily verify that $\mu(z_1, z_2, z_3)$ is normalized in the fashion

$$\int d\Gamma \mu(z_1, z_2, z_3) = 1. \quad (62)$$

We compute now (i) the Wehrl entropy in the form

$$\mathcal{W} = \int d\Gamma \mu(z_1, z_2, z_3) \ln \mu(z_1, z_2, z_3), \quad (63)$$

and (ii) the Fisher measure as follows:

$$\mathcal{F} = \int d\Gamma \mu(z_1, z_2, z_3) \mathcal{A}_{3D}, \quad (64)$$

where in this case we define the three-dimensional quantity \mathcal{A}_{3D} in the fashion

$$\begin{aligned} \mathcal{A}_{3D} = & \sigma_x^2 \left[\frac{\partial \ln \mu(z_1, z_2)}{\partial x} \right]^2 + \sigma_{p_x}^2 \left[\frac{\partial \ln \mu(z_1, z_2)}{\partial p_x} \right]^2 + \sigma_y^2 \left[\frac{\partial \ln \mu(z_1, z_2)}{\partial y} \right]^2 \\ & + \sigma_{p_y}^2 \left[\frac{\partial \ln \mu(z_1, z_2)}{\partial p_y} \right]^2 + \sigma_z^2 \left[\frac{\partial \ln \mu(z_1, z_2)}{\partial z} \right]^2 + \sigma_{p_z}^2 \left[\frac{\partial \ln \mu(z_1, z_2)}{\partial p_z} \right]^2, \end{aligned} \quad (65)$$

with the phase-space relationships (31), (32) and

$$|z_3|^2 = \frac{1}{4} \left(\frac{z^2}{\sigma_z^2} + \frac{p_z^2}{\sigma_{p_z}^2} \right), \quad (66)$$

where $\sigma_z = \sqrt{\hbar/2m\omega}$ and $\sigma_{p_z} = \sqrt{m\hbar\omega/2}$. In this instance $d^2z_3 = dz dp_z/2\hbar$. Finally, we are led to

$$\mathcal{F} = \int d\Gamma \mu(z_1, z_2, z_3) \left\{ \gamma^2 (|z_1|^2 |z_2|^2 + \frac{1}{4} (1 + |z_1|^2)^2) + 4|u|^2 |z_3|^2 \right\}, \quad (67)$$

i.e.,

$$\mathcal{F} = \langle \gamma^2 (|z_1|^2 |z_2|^2 + \frac{1}{4} (1 + |z_1|^2)^2) + 4|u|^2 |z_3|^2 \rangle, \quad (68)$$

where

$$\begin{aligned} \gamma = & \frac{-(1 + |z_3|^2)^2 \sum_{I=0}^{\infty} \frac{2(2I)!}{I!} |v|^{2I+1} e^{-I(I+1)\frac{\Theta}{T}} \times g(I)}{\sum_{I=0}^{\infty} \frac{(2I)!}{I!} |v|^{2I} e^{-I(I+1)\frac{\Theta}{T}} \times g(I)} \\ & + \frac{\sum_{I=0}^{\infty} \frac{2(2I)!}{(I-1)!} |v|^{2I-1} e^{-I(I+1)\frac{\Theta}{T}} \times g(I)}{\sum_{I=0}^{\infty} \frac{(2I)!}{I!} |v|^{2I} e^{-I(I+1)\frac{\Theta}{T}} \times g(I)}. \end{aligned} \quad (69)$$




Figure 4. 3D-anisotropic rotor. We plot Wehrl's entropy \mathcal{W} as a function of T/Θ for several values of the anisotropy ratio ($I_{xy}/I_z = 1/2, 1, 5, 10$). Trends for the Wehrl entropy are not quite similar to those for the linear case. A maximum is seen at $T/\Theta = 0$ and a minimum at $T/\Theta \geq 2$, whose exact values depend on the ratio I_{xy}/I_z .

In the special instance $I_{xy}/I_z = 1$, that corresponds to the spherical rotor, we explicitly obtain

$$\mu(z_1, z_2, z_3) = e^{-|u|^2} \frac{\sum_{l=0}^{\infty} \frac{|u|^{2l}}{l!} e^{-l(l+1)\frac{\Theta}{T}}}{\sum_{l=0}^{\infty} (2l+1)^2 e^{-l(l+1)\frac{\Theta}{T}}}. \quad (70)$$

Having the Husimi functions the Wehrl entropy is straightforwardly computed.

In order to emphasize some special cases associated with possible applications we consider several possibilities.

- (i) The spherical rotor $I_{xy} = I_x = I_y = I_z$, thus $I_{xy}/I_z = 1$ (e.g. CH_4).
- (ii) The oblate rotor $I_{xy} = I_x = I_y < I_z$, specifically $1/2 \leq I_{xy}/I_z < 1$ (e.g. C_6H_6).
- (iii) The prolate rotor $I_{xy} = I_x = I_y > I_z$, which corresponds to $I_{xy}/I_z > 1$ (e.g. PCl_5).
- (iv) The extremely prolate rotor is equivalent to the linear case (all diatomic molecules) $I_z = 0$, this is $I_{xy}/I_z \rightarrow \infty$ (e.g. $\text{CO}_2, \text{C}_2\text{H}_2$).

In figure 4, we depict the Wehrl entropy \mathcal{W} , as a function of T/Θ , for several values of I_{xy}/I_z . This is the extremely oblate rotor $I_{xy}/I_z = 1/2$, the prolate rotor $I_{xy}/I_z = 5, 10$ and the spherical case $I_{xy}/I_z = 1$.

5. Concluding remarks

A study has been performed on the basis of Schwinger's angular momenta construction [4] and following Morales *et al* [1] coherent states' rotor formulation. Possible applications to specific molecular problems have been suggested.

We have concentrated our efforts on the study of the semiclassical behaviour of the rigid rotor and have obtained in analytical fashion the form of the appropriate semiclassical Husimi distribution for two cases, namely, the linear and the axially symmetric rigid rotor. The linear case can be reobtained as a particular instance of the three-dimensional formulation. We have also obtained an analytical expression of the shift-invariant Fisher measure built up with Husimi distributions for the rigid rotor model, concluding that Fisher's measure is a better indicator of the delocalization than Wehrl's entropy.

Acknowledgments

S Curilef and F Pennini would like to thank partial financial support by FONDECYT 1051075 and DGIP-UCN 2006.

Appendix

First of all, we carry out the differentiation of the Husimi distribution (29) with respect to the variable x , obtaining the following result:

$$\frac{\partial \ln \mu(z_1, z_2)}{\partial x} = 2\eta(z_1, z_2) \frac{\partial |z|}{\partial x}, \quad (\text{A.1})$$

where the quantity $\eta(z_1, z_2)$ was defined in equation (37). Moreover, from equations (31) and (32) we have

$$\frac{\partial |z|}{\partial x} = \frac{x}{4|z|\sigma_x^2}, \quad (\text{A.2})$$

and we are led to

$$\frac{\partial \ln \mu(z_1, z_2)}{\partial x} = \frac{\eta(z_1, z_2)x}{2|z|\sigma_x^2}. \quad (\text{A.3})$$

We arrive at a similar expression differentiating with respect to p_x ,

$$\frac{\partial \ln \mu(z_1, z_2)}{\partial p_x} = \frac{\eta(z_1, z_2)p_x}{2|z|\sigma_{p_x}^2}. \quad (\text{A.4})$$

Analogous expressions are obtained by replacing x and p_x with y and p_y , respectively. Finally, substituting these results into equation (35) we thus arrive at

$$\mathcal{A} = \eta(z_1, z_2)^2. \quad (\text{A.5})$$

References

- [1] Morales J A, Deumens E and Öhrn Y 1999 *J. Math. Phys.* **40** 776
- [2] Glauber R J 1963 *Phys. Rev.* **131** 2766
- [3] Klauder J R and Skagerstam B S 1985 *Coherent States* (Singapore: World Scientific)
- [4] Sakurai J J 1994 *Modern Quantum Mechanics* ed S F Tuan (Reading, MA: Addison-Wesley) p 217
- [5] Casida M E, Harriman J E and Anshell J L 1987 *Int. J. Quantum Chem.* **32** 435
- [6] Harriman J E and Casida M E 1993 *Int. J. Quantum Chem.* **45** 263
- [7] Pathria R K 1993 *Statistical Mechanics* (Exeter: Pergamon)
- [8] Garbaczewski P 2005 *Entropy* **7** 253
- [9] Frieden B R and Soffer B H 1995 *Phys. Rev. E* **52** 2274
- [10] Frieden B R 1998 *Physics from Fisher Information* (Cambridge: Cambridge University Press)
- [11] Frieden B R 2003 *Science from Fisher Information* (Cambridge: Cambridge University Press)
- [12] Pennini F and Plastino A 2004 *Phys. Rev. E* **69** 057101

- [13] Ullah N 1994 *Phys. Rev. E* **49** 1743
- [14] Arranz F J, Borondo F and Benito R M 1996 *Phys. Rev. E* **54** 2458
- [15] Titantah J T and Hounkonnou M N 1999 *J. Phys. A: Math. Gen.* **32** 897
- [16] Titantah J T and Hounkonnou M N 1997 *J. Phys. A: Math. Gen.* **30** 6327
- [17] Titantah J T and Hounkonnou M N 1997 *J. Phys. A: Math. Gen.* **30** 6347
- [18] Titantah J T and Hounkonnou M N 1995 *J. Phys. A: Math. Gen.* **28** 6345
- [19] Anderson A and Halliwell J J 1993 *Phys. Rev. D* **48** 2753
- [20] Gazeau J P and Klauder J R 1999 *J. Phys. A: Math Gen.* **32** 123
- [21] Thaik M and Inomata A 2005 *J. Phys. A: Math Gen.* **38** 1767
- [22] Wehrl A 1979 *Rep. Math. Phys.* **16** 353
- [23] Husimi K 1940 *Proc. Phys. Math. Soc. Japan* **22** 264
- [24] Lieb E H 1978 *Commun. Math. Phys.* **62** 35
- [25] Nieto M M 1980 *Phys. Rev. D* **22** 391
- [26] Schwinger J 1965 *In Quantum Theory of Angular Momentum* ed L C Biedenharn and H van Dam (New York: Academic) p 229
- [27] Cohen-Tannoudji C, Diu B and Laloë F 1977 *Quantum Mechanics* (Paris: Hermann)